



Northern Hemisphere ice sheet and ocean interactions during the last glacial period in a coupled ice sheet–climate model

Louise Abot¹, Aurélien Quiquet², and Claire Waelbroeck¹

¹LOCEAN/IPSL, CNRS, IRD, MNHN, Sorbonne Université, UMR 7159, 75005 Paris, France

²LSCE/IPSL, CEA, CNRS, UVSQ, Université Paris-Saclay, UMR 8212, 91190 Saint-Aubin, France

Correspondence: Louise Abot (louise.abot@locean.ipsl.fr)

Received: 2 July 2024 – Discussion started: 7 August 2024

Revised: 27 March 2025 – Accepted: 8 April 2025 – Published: 2 July 2025

Abstract. This study examines the interactions between the Northern Hemisphere ice sheets and the ocean during the last glacial period. Using the iLOVECLIM climate model of intermediate complexity and the GRISLI ice sheet model, we explore the consequences of an amplification of the melt rates beneath ice shelves on ice sheet dynamics and the associated feedbacks. First, the amplification of oceanic basal melt rates leads to significant freshwater release from both increased calving and basal melt fluxes. Grounding line retreat and dynamic thinning occur over the Eurasian and Ice-land ice sheets, while the oceanic perturbation fails to trigger a grounding line migration over the coasts of Greenland and the eastern part of the Laurentide ice sheet. Second, similar to hosing experiments with no coupling between the climate and the ice sheets, the influx of fresh water temporarily increases sea-ice extent; reduces convection in the Labrador Sea; weakens the Atlantic meridional overturning circulation; lowers surface temperatures in the Northern Hemisphere, especially over the North Atlantic Ocean; and increases the subsurface temperatures in the Nordic Seas. Third, the freshwater release and latent heat effect on ocean temperatures lead to a decrease in ice sheet discharge (negative feedback) for the Greenland and Eurasian ice sheets. The Laurentide ice sheet does not feature significant volume variations in the experiments. On the one hand, the amplification of the shelf melt rates produces a weak perturbation due to low background temperatures and salinity at shelf drafts in Baffin Bay and the Labrador Sea according to the model. On the other hand, the Laurentide ice sheet in the fully coupled model may be overly stable. We show that we are able to force a grounding line retreat and a North American ice sheet volume decrease by imposing ad hoc con-

stant oceanic melt rates. However, in both sets of perturbation experiments, the Hudson Strait ice stream does not exhibit the past dynamic instability indicated by the presence of Laurentide-origin ice-rafted debris in the North Atlantic sediment records. This suggests the possibility that the model is too stable, specifically in the Hudson Bay region. Different ice sheet geometries or modeling choices regarding the basal dynamics beneath the ice sheet could help address this issue. In summary, this study found that an episode of sub-surface warming may trigger dynamical instabilities and ice discharges along the coasts of the Nordic Seas but subsequent ocean–ice sheet interactions may be characterized by negative feedback thus dampening ice discharges. This study also emphasizes the need for further research using fully coupled models to explore the triggering mechanisms of massive iceberg discharges and to clarify the role of the ocean in these events.

1 Introduction

Recent studies of the Antarctic ice sheet show that warm oceanic water plays an important role in continental ice thinning, impacting not only floating but also grounded ice (e.g., Pritchard et al., 2012; Reese et al., 2018; Gudmundsson et al., 2019). The increase in ocean sub-shelf melt rates induces a thinning and eventual break up of the ice shelves, reducing the buttressing effect. This results in an acceleration of upstream ice flow and a dynamic thinning of the ice sheet inland. The ongoing warming of the ocean (Johnson and Lyman, 2020) could lead to the collapse of the West Antarctic Ice Sheet, among other dramatic consequences, amplifying

the sea level rise (Joughin and Alley, 2011; Naughten et al., 2023). To the north, the Greenland ice sheet's last floating ice tongue also undergoes a thinning driven by warmer ocean temperatures (Wekerle et al., 2024). Similarly, a number of studies show that ocean subsurface warming during the last glacial period could have destabilized ice sheets and led to massive iceberg discharges (e.g., Shaffer et al., 2004; Marcott et al., 2011; Max et al., 2022). Indeed, it was suggested that massive iceberg discharges occur after an initial reduction in the Atlantic meridional overturning circulation (AMOC) and the buildup of a heat reservoir at the subsurface in the North Atlantic, based on both observations and modeling studies (e.g., Jonkers et al., 2010; Max et al., 2022; Mignot et al., 2007).

The last glacial period refers to the portion of the last glacial cycle between the end of the last interglacial period and the transition into the present interglacial period. It spans the period from about 75 kyr before present (BP) (e.g., Martinson et al., 1987) to about 15 or 12 kyr BP (e.g., Rasmussen et al., 2014). An intriguing feature of the last glacial period is its millennial-scale variability, first revealed in Greenland ice cores, known as the Dansgaard–Oeschger (DO) cycles (Dansgaard et al., 1984, 1993). These are transitions between relatively cold (stadial) and warm (interstadial) periods in the Northern Hemisphere (Johnsen et al., 1992). The rapid warming (or DO event) generally takes place over a period of several decades and has a magnitude of around 10 °C at the North Greenland Ice Core Project (NGRIP) site (Kindler et al., 2014). Such events occurred repeatedly during the last glacial period. They are followed by a slower cooling that takes approximately 100 to 1000 years, and the cycle ends with a sudden cooling (Kindler et al., 2014).

Some of the stadials are accompanied by massive iceberg discharges from the Northern Hemisphere ice sheets in the North Atlantic Ocean, at a frequency of several thousand years (Heinrich, 1988; Bond and Lotti, 1995). The iceberg discharges are identified in marine sediment cores by one or several layers of ice-rafted debris, the materials being entrapped and transported by drifting ice (Bond et al., 1992; Bond and Lotti, 1995; Elliot et al., 1998). The iceberg discharges are referred to as Heinrich events (HEs) when the continental ice is originating from the Laurentide ice sheet (i.e., the central and eastern parts of the North American ice sheet). The Heinrich events mostly occur after several DO cycles and are followed by a DO event (Bond et al., 1993). Barker et al. (2015), through the examination of a site southwest of Iceland, suggested that massive iceberg discharges were a consequence of prolonged stadial conditions.

Nonetheless, there is no scientific consensus on the ultimate cause behind the massive iceberg discharges to date. For instance, some studies have shown that self-sustained ice sheet oscillations could be at play (MacAyeal, 1993), while others propose that the discharges were triggered by an external forcing such as a change in ocean conditions (Alvarez-Solas et al., 2010). The focus of our study is to test the latter

hypothesis, using a coupled climate–ice sheet model, namely the iLOVECLIM–GRISLI model.

More specifically, several studies show that a rise in sub-surface ocean temperature during a stadial in the North Atlantic could have led to massive iceberg discharges (e.g., Shaffer et al., 2004; Marcott et al., 2011; Max et al., 2022). Alvarez-Solas et al. (2010) have shown that fluctuations in oceanic temperatures could have triggered Northern Hemisphere ice sheet surges (i.e., an unsteady state of the ice stream associated with rapid ice flow and dynamic thinning of the ice sheet inland) and iceberg discharges during the last glacial period. Forcing the Laurentide ice sheet and Hudson stream with varying basal melt rates results in the collapse of buttressed ice shelves and the subsequent acceleration of inland ice streams (Alvarez-Solas et al., 2013). Iso-static adjustment may then allow the regrowth of the ice sheet (Bassis et al., 2017). Modeling results show that ocean temperature increases can also trigger or amplify ice discharges from the Eurasian and Greenland ice sheets (Alvarez-Solas et al., 2019; Tabone et al., 2019). However, in these studies, the ocean is considered a forcing and this consideration does not allow for a full understanding of the interplay between ocean and ice sheets. The novelty of this study lies in the use of a climate–ice sheet model: the iLOVECLIM model (of intermediate complexity) coupled with the GRISLI ice sheet model.

Indeed, grasping the response of the entire ice sheet to an oceanic perturbation or the subsequent oceanic changes following an influx of continental water is complex due to the presence of numerous feedback mechanisms at the ocean–ice sheet interface (Goosse et al., 2018; Holland et al., 2020). For instance, under warming conditions, initial ice sheet melting releases cold and fresh water at the surface, which can lead to negative feedback on the climate (Swingedouw et al., 2008; Li et al., 2024). Additionally, ocean-driven ice shelf thinning may raise the depth-dependent freezing temperature at the ice–ocean interface, reducing oceanic basal melt rates – a process that represents another form of negative feedback (Van Achter et al., 2023). Following an oceanic warming, the shelves' melted water released into the ocean could also lead to decreased vertical mixing and favor sea-ice expansion, reducing the initial ocean warming and basal melt rates, thus inducing negative feedback as well (Goosse et al., 2018). There is also numerous possible positive feedback that could contribute to the destabilization of the grounded ice sheet. For instance, freshwater release can increase surface stratification, leading to transient subsurface warming that amplifies oceanic basal melting (Moorman et al., 2020; Li et al., 2024). Another form of positive feedback is the marine ice sheet instability, which is linked to the dynamics of the grounding line. The mathematical formulation of the ice flux at the grounding line shows that this flux increases with the ice thickness (Schoof, 2007; Tsai et al., 2015). When a marine-terminating ice sheet sits on an upward-sloping bed toward the ocean, an initial retreat of the grounding line causes ice

thickening and acceleration of the ice flow at the grounding line, which in turn drives further retreat. This process may have played a role in Heinrich events given the bedrock's shape at the Hudson Strait mouth (Schoof, 2007). Additional feedback mechanisms could emerge from ice-sheet-cavity-scale processes, such as tidal forcing or buoyancy-driven circulation beneath ice shelves (e.g., Makinson et al., 2011; Gwyther et al., 2016). For example, a rise in ocean temperature at the base of the ice shelves could enhance the meltwater production, thereby strengthening the circulation within a cavity, leading to an increased heat supply and further melting (Gwyther et al., 2016). However, uncertainties related to these processes remain large since very few cavity-enabled ocean models can operate at the global scale. For this reason and given our coarse-resolution ocean model, we do not discuss cavity-scale feedback any further in this paper.

In our setup, under constant or transient forcings corresponding to the last glacial period, the iLOVECLIM–GRISLI coupled model does not exhibit abrupt internal oscillations, neither in the ice sheets nor in the ocean circulation. Therefore, in this study, we chose to conduct perturbation experiments at the ice–ocean interface. With the fully coupled model, we can examine whether a perturbation at the interface is being amplified or dampened.

Here, we take advantage of the fully coupled model to address the following questions: how does ocean subsurface warming at characteristic ice shelf drafts affect ice sheet dynamics? How does the continental ice melt release affect in turn oceanic circulation, hydrography and ice sheet dynamics? In this work, we impose amplified oceanic melt rates on the 40 kyr BP Northern Hemisphere ice sheets to get insights into both past climate variability and mechanisms at play at the ocean–ice sheet interface.

2 Model and methods

2.1 Climate and ice sheet model

2.1.1 iLOVECLIM

For this study, we use the coupled climate model of intermediate complexity iLOVECLIM, which is a code fork of LOVECLIM version 1.2 (Goosse et al., 2010). The core of the model includes oceanic, atmospheric and vegetation components (namely CLIO, EcBilt and VECODE). The model has been used for a wide range of climate studies, from the last million years to the Holocene (e.g., Caley et al., 2014; Bouttes et al., 2018; Arthur et al., 2023). In this section, we will emphasize some of its main features. VECODE is a dynamic vegetation and carbon allocation model. CLIO is a free surface ocean general circulation model solved on a spherical grid with 3° latitude and longitude resolution and 20 irregular levels in z coordinates. It includes a thermodynamic–dynamic sea-ice component (Fichefet and Maqueda, 1997). Despite its coarse resolution, CLIO is able to represent large-

scale features. Taking the example of the North Atlantic Ocean, these features include the main gyres (North Atlantic and subpolar gyres) as well as the main currents. The Gulf Stream, the North Atlantic Current, the Irminger Current and the East Greenland Current are depicted, although the depictions of the currents are wider in CLIO than in higher-resolution models (Fig. S1 in the Supplement). EcBilt is a quasi-geostrophic atmospheric model solved on a T21 spectral grid with a resolution of 5.6° in both latitude and longitude. It includes three vertical levels at 800, 500 and 200 hPa. Given that precipitation biases can alter the performance of CLIO to reproduce large-scale oceanic circulation, our model includes a flux correction of precipitation rates since Goosse et al. (2010). Specifically, precipitation is enhanced over the North Pacific while reduced over the Arctic Ocean and Atlantic Ocean (Fig. S2 in the Supplement). Major model parameters are listed in Table S1a and b in the Supplement. We use the default values from the iLOVECLIM source archive, except for the Bering Strait scaling factor and the greenhouse gas (GHG) concentration radiative forcing factor. The first is set to zero to ensure a closed Bering Strait. The second is doubled (Timm and Timmermann, 2007) to artificially increase the climate sensitivity and to achieve a relatively cool climate in order to maintain the ice sheets at 40 kyr BP. This modeling choice is informed by the relatively low equilibrium climate sensitivity of iLOVECLIM, which is approximately 2°C (Bouttes et al., 2024), in comparison to the IPCC AR6 best estimate of 3°C and likely range of $2.5\text{--}4^\circ\text{C}$ (IPCC, 2023).

2.1.2 GRISLI

iLOVECLIM is coupled with the dynamic ice sheet model GRISLI, which is a 3D thermomechanically coupled ice sheet model that is used here on a Cartesian grid of the Northern Hemisphere at a 40 km resolution. GRISLI is an appropriate tool to address ice sheet dynamics and interactions with the oceanic component as it explicitly calculates ice stream velocities, the position of the grounding line and the behavior of ice shelves. For instance, GRISLI has shown a good ability to reproduce grounding line migrations and ice volume changes in Antarctica during the last 400 kyr BP (Quiquet et al., 2018a; Crotti et al., 2022). The equations are described in Ritz et al. (2001) and Quiquet et al. (2018a). They are obtained from shallow ice and shallow shelf approximations. Major model parameters are listed in Table S1c in the Supplement.

The deformation is calculated using the Glen flow law with an enhancement factor (E_f) to account for anisotropy.

When the ice sheet's base is at the pressure melting point, the basal drag is a linear function of the basal velocities over the grounded ice sheet (Weertman, 1957). When the ice sheet's base is below the pressure melting point, the sliding velocity is set to zero (infinite friction). The friction is set to zero for the floating ice shelves. Following the Weertman

law, the basal drag is equal to the opposite of the basal sliding velocity times a basal drag coefficient β . This coefficient β is defined as the effective pressure at the base of the ice sheet times a tuning parameter (C_f) and is bounded by a minimum ($\beta_{\min}$) and a maximum ($\beta_{\max}$) value. Over thick sediment layers (more than 200 m), the basal drag is reduced by a factor of 20. We use the soft-hard basal drag mask (i.e., the sediment map) from Laske (1997) and the geothermal heat flux from Shapiro and Ritzwoller (2004).

The calving of icebergs is not modeled explicitly. Instead, the floating ice calves when its thickness falls below a critical thickness threshold, which varies with oceanic depth (Fig. S4 in the Supplement). This way, the shelves cannot extend over the deep ocean. It is to be noted that, after calving, the ice shelves are allowed to grow back, as a result of upstream ice flow and surface accumulation.

The grounding line formulation follows Tsai et al. (2015), and its position is determined by interpolating between the last grounded and the first floating grid cells.

Glacial isostatic adjustment is accounted for using an elastic–lithosphere–relaxed–asthenosphere model with a relaxation time of 3000 years (Le Meur and Huybrechts, 1996).

2.1.3 Coupling

The coupling procedure is described in Roche et al. (2014) and Quiquet et al. (2021b). In iLOVECLIM, the atmospheric model ECBilt includes an online downscaling (Quiquet et al., 2018b). At each atmospheric time step, temperature and precipitation are computed on the finer-resolution GRISLI grid using the standard ECBilt energy and moisture equations. This subgrid information is fed into an Insolation Temperature Melt model (Pollard, 1980; Van Den Berg et al., 2008) to compute surface mass balance every 4 h. This time step allows for numerical model stability, as the ECBilt model does not resolve the diurnal cycle. The yearly average surface mass balance and surface temperature are then used by GRISLI. In turn, ECBilt receives the yearly topography and ice mask data from GRISLI.

The continental ice melt fluxes are computed in GRISLI each year. These are distributed to the oceanic model over the following year. For the grounded ice sheet, surface melt is routed to the nearest oceanic grid cell following surface topography, while the oceanic basal melt and the calving flux are transferred from where they occur to the ocean surface. The local latent heat flux, associated with the calving flux and corresponding to the melting of icebergs by the ocean, is taken into account and the ocean temperatures are adjusted. More details about the freshwater coupling are given in Fig. S2 in the Supplement.

The oceanic basal melt rate (OBM) is parameterized as a quadratic function of ocean temperatures (Holland et al., 2008; Favier et al., 2019). When the local ocean temperature is above the freezing temperature, the oceanic basal melt rate

is derived as follows:

$$\text{OBM} = \gamma_T \cdot A \cdot (T - T_f)^2, \quad (1)$$

where the local freezing temperature $T_f = T_f(S, z = 200 \text{ m})$ is a function of salinity, following the formulation of Millero (1978). The freezing temperature is computed at a characteristic shelf draft of 200 m. $A = (\frac{\rho_0 C_{p0}}{L \rho_i})^2 \text{ K}^{-2}$ is a constant depending on physical parameters, with $\rho_0 = 1030 \text{ kg m}^{-3}$ (the density of seawater), $C_{p0} = 4002 \text{ J kg}^{-1} \text{ K}^{-1}$ (the specific heat capacity of seawater) and $L \rho_i = 300.33 \times 10^6 \text{ J m}^{-3}$ (the latent heat of fusion of ice times the ice density). γ_T is a tuned parameter that accounts for the heat transfer velocity, which has been calibrated for this study to $1.09 \times 10^{-5} \text{ m s}^{-1}$. This value for γ_T ensures that the ice sheet volume remains intermediate between that of the pre-industrial (PI) period and the last glacial maximum (LGM), preventing excessive expansion or disintegration. The OBM is computed in the ocean model each day and integrated over each year, as the time step for the dynamic ice sheet model is annual. For each horizontal grid point, the rate is calculated for all vertical levels in the ocean model, and the ice sheet model retains only the value corresponding to its actual shelf draft. When the ice sheet model cell is not covered by the ocean model, the value of the nearest ocean cell is retained.

The fluctuations in ice sheet volume have no impact on the global volume of the ocean in the model. As a result, sea level and bathymetry in the ocean model remain unchanged.

2.2 Experimental setup

2.2.1 Initial condition

We conduct a long coupled ice sheet–climate simulation under constant external forcing to derive the initial state. This approach assumes climate equilibrium, which is not realistic. However, the relatively small variations in greenhouse gas concentrations and insolation at 65° N around 40 kyr BP support the use of an equilibrium simulation for this specific time slice. Furthermore, this period aligns with Heinrich Stadial 4 (HS4), which extends from 39.85 to 38.17 kyr BP (Waelbroeck et al., 2019).

The climate model is forced with insolation (Berger, 1978) and greenhouse gas concentration (Lüthi et al., 2008). Forcings are held constant at their 40 kyr BP values (namely 0.013, 23.61° and 0.004 for eccentricity, obliquity and precession index, respectively, and 220 ppm for the CO_2 concentration). The ice sheet model is forced with sea level reconstruction (-64 m at 40 kyr BP; Waelbroeck et al., 2002). The latter is used by GRISLI to determine which areas of land are located above the sea level (i.e., where the continental ice can grow). The long spin-up starts from a previous equilibrium simulation of the last glacial maximum (LGM) similar to the one described in Quiquet et al. (2021b). We use a LGM bathymetry for the ocean model (Lhardy et al., 2021). However, since sea level and bathymetry at 40 kyr BP have

intermediate values between the LGM and the pre-industrial values, we also test the sensitivity of the results to the pre-industrial bathymetry in the discussion section. Thus, two extreme bathymetry possibilities are investigated.

For this long spin-up simulation, we use an asynchronous coupling between the ice sheets and the rest of the climate system. Since the ice sheets feature slower dynamics than the ocean and the atmosphere, we run the ice sheet model for 10 years every year of the rest of the climate model. The spin-up simulation runs for 8000 years for the ocean and the atmosphere but 80 000 years for the ice sheets. During the spin-up phase, the freshwater fluxes from the ice sheet model to the ocean are ignored. This approach is necessary because we cannot conserve both total ice mass change and the rate of mass change when using an acceleration factor (Fig. S2 in the Supplement). However, at the end of the spin-up phase, the ice sheet is in equilibrium with the rest of the climate system. Therefore, the ice sheet volume is quasi stable and produces a net zero freshwater flux to the ocean. There is thus no significant discontinuity in our control experiment with ice sheet freshwater flux included when branched from the spin-up simulation.

At the end of the spin-up simulation, we obtain a total volume of $24 \times 10^6 \text{ km}^3$ for the Northern Hemisphere ice sheets, which corresponds to 48 mSLE (mean sea level equivalent; directly calculated from the volume of grounded ice that is above sea level and compared to the present-day ocean area). In comparison, the reconstruction from Gowan et al. (2021) presents volumes of around 12 and $16 \times 10^6 \text{ km}^3$ for minimal and maximal scenarios, corresponding to 28 and 37 mSLE, respectively. However, Gowan et al. (2021) acknowledge that their total ice sheet reconstructions for the period between 57 and 29 kyr BP (MIS 3) do not align with proxy-based sea level reconstructions. Specifically, they estimate a global sea level that is 20 m higher (maximum scenario) than the reconstructed value of approximately -60 to -90 m at 40 kyr BP (Waelbroeck et al., 2002; Arz et al., 2007; Siddall et al., 2008). Also, the evolution of the Northern Hemisphere ice sheets remains uncertain prior to the last glacial maximum. Reconstructions often rely on inverse methods that use observational data and estimates of global mean sea level, sometimes supported by numerical ice sheet modeling (e.g., Marshall et al., 2000; Kleman et al., 2010; Pico et al., 2018; Gowan et al., 2021; Dalton et al., 2022). Significant discrepancies exist between the different reconstructions. For example, the question of whether the Hudson Bay area was glaciated around 40 kyr BP and to what extent remains debated (e.g., Batchelor et al., 2019; Miller and Andrews, 2019). Given these uncertainties, we conclude that our simulated ice sheet volume falls within an acceptable range for the 40 kyr BP time slice.

The ice sheet conditions at the end of the 40 kyr BP equilibrium simulation are depicted in Fig. 1. Ice is located over North America, Greenland, Iceland, Svalbard and Fennoscandia. Our North American ice sheet is larger than

the one presented in Gowan et al. (2021) for the same period. In contrast to this study, we obtain a glaciated Iceland and a marine-terminating Fennoscandian ice sheet. Also, the DATED-1 reconstruction for the Eurasian ice sheet (Hughes et al., 2015) presents a reduced extent of the Fennoscandian ice sheet and no Svalbard ice sheet compared to our simulated ice sheet. However, this reconstruction corresponds to the 38–34 kyr BP period, which is slightly later than the time slice examined in this study. The ice shelf area of our simulated ice sheet is relatively small, with no extensive regions that are entirely covered by ice, in contrast to what is observed in Antarctica today.

2.2.2 Perturbation experiments

Each of the following experiments are branched on the 40 kyr BP equilibrium. External forcings are held constant at their 40 kyr BP values. Differing from the spin-up simulation, the climate and the ice sheets are annually coupled. The acceleration is only used for the spin-up simulation in order to equilibrate the ice sheets with the climate, as the ice sheets take a long time to adjust. Here, freshwater fluxes from the ice sheet to the ocean are taken into account. The control experiment (CTRL) is the continuation of the equilibrium simulation described in Sect. 2.2.1 but with the addition of freshwater flux to the ocean resulting from ice sheet melting.

Each perturbation experiment consists of multiplying the oceanic basal melt rates (defined in Eq. 1) at each grid point by an amplification factor X , with X ranging from 5 to 300, for 500 years. This scaling (X) of the basal melt is equivalent to imposing a temperature anomaly, which depends on the water masses and background state of the ocean, at the shelf drafts. Specifically, we apply a perturbation whose intensity is proportional to the local temperature anomaly with respect to the freezing temperature, T_f . Indeed, when $T > T_f$, adding a perturbation term δT to T defined by $\delta T = \epsilon(T - T_f)$ (with $\epsilon > 0$) leads to the following oceanic basal melt rate: $\text{OBM} = \gamma_T A(T + \epsilon(T - T_f) - T_f)^2 = \gamma_T A(1 + \epsilon)^2(T - T_f)^2$. And we rewrite $X = (1 + \epsilon)^2$, so to demonstrate a direct correspondence between X and the temperature anomaly intensity ϵ seen by the ice shelves. For example, a doubling of the temperature anomaly, with respect to the freezing point, corresponds to $\epsilon = 1$ and $X = 4$. A 10 time increase in the temperature anomaly corresponds to $\epsilon = 9$ and $X = 100$.

This way, the perturbation experiments aim to represent an increased heat flux in the ice shelf drafts, reflecting subsurface ocean warming in the North Atlantic during a Dansgaard–Oeschger (DO) stadial (e.g., Rasmussen and Thomsen, 2004). The perturbation experiments were designed to be spatially consistent with the physics of the water masses. For instance, our design allows us to account for the fact that an abrupt change in AMOC likely leads to temperature changes in the AMOC's main areas of influence rather than a uniform temperature change over the North Atlantic and Nordic Seas.




Figure 1. The 40 kyr equilibrium. (a) Ice sheet elevation (shades of blue) and elevation above sea level (shades of gray) [m] at the end of the equilibrium simulation. Light-blue contour is the mean position of the grounding line at the same time. Yellow contours are the minimal and maximal ice sheets extent in the reconstruction from Gowan et al. (2021), and these differ only over the central North American ice sheet. Areas used to derive regional ice volumes in Fig. 2 for North American, Greenland–Iceland and Eurasian ice sheets are contoured in pink. (b) Ice shelf drafts [m] and (c) basal melt rates beneath the shelves [m yr^{-1}] at the end of the equilibrium simulation.

Here, we amplify the basal melt rates instantaneously, as this seems to be the most direct way to maximize the ice sheet response to the perturbation and to evaluate the signs of feedback at the ice–ocean interface. Note that the impact of a gradual increase in basal melt rates on ice loss is similar to that of an instantaneous increase (Fig. S3 in the Supplement). Still, a limitation of the basal melt scaling approach is that the experiment results are dependent on the initial representation of the temperature in the ocean and model biases. To overcome this limitation, we perform additional experiments by applying a constant temperature offset within the formulation of the basal melt rate (Sect. 4.1).

From the ice sheet point of view, increasing the oceanic basal melt rates is equivalent to imposing a subsurface warming, yet this has no effect on initial ocean temperatures. Initially, the ocean structure is the same for the control and perturbation experiments, as the simulations are branched on the same spin-up. Since we use a coupled setup, the ice sheet retreat induced by the perturbation impacts the ocean through the resulting freshwater flux. The ocean model adjusts its temperatures and salinity in response to the freshwater input, with associated impacts on the density profile, vertical convection and ocean circulation. Thus, the ocean model response can modulate the perturbation in the vicinity of the ice shelves. Therefore, in our perturbation experiments, the freshwater feedback that could arise from the ocean–ice sheet interactions is taken into account, unlike traditional hosing experiments.

3 Results

3.1 Ice sheet response

The oceanic perturbation causes a decrease in the total volume of the Northern Hemisphere ice sheets. The relative contributions of the regional ice sheets show that the North American one contributes the least to the total volume change, while the Greenland, Iceland and Eurasian ice sheets contribute the most (Fig. 2). The volume of the Greenland–Iceland ice sheets decreases rapidly, over the first 100 years, before reaching a new state with minimal ice volume change, for experiments with perturbation factors above 20. The Eurasian ice sheet loses volume throughout the perturbation period for all experiments. Once the perturbation is halted, Eurasia and Greenland regain mass at a slower pace. The North American ice volume remains relatively unaffected by the oceanic disturbance. For the highest perturbation factors, it even increases slightly for ~ 500 years and decreases once the perturbation is halted (Fig. 2a).

Despite its low resolution (40 km), GRISLI is capable of representing large-scale features such as major ice streams at the LGM (e.g., Quiquet et al., 2021a). The major ice streams, such as the Hudson Strait, Lancaster Sound and Amundsen Gulf ice streams (nomenclature follows Margold et al., 2015), remain present at 40 kyr BP (Fig. 3a). On the contrary, near the southern margin of the North American ice sheet, there are no well-identified ice streams. There are also large velocities (above 500 m yr^{-1}) along the southeast coast of Greenland and around Iceland (Fig. 3a and b). Active regions




Figure 2. Continental ice volume response to oceanic perturbation (i.e., amplified OBM by a coefficient X). Time series of grounded ice sheet volume [10^6 km^3] for the (a) North American, (b) Greenland–Iceland and (c) Eurasian ice sheets in experiments with different perturbation factors. The thin black lines are the free evolution, once the perturbation is released, of the experiments, where $X = 20$ and 100. Note that the same y-axis spacing is applied on the plots ($0.6 \times 10^6 \text{ km}^3$).

with velocities that are weaker yet still above 100 m yr^{-1} also include both Greenland coasts and the Fennoscandian coast.

The ice sheet evolution in the experiment with a large perturbation factor is depicted in Fig. 3c–f. After 200 years, the thinning is significant on the coasts bordering the Nordic Seas and to the south of the Greenland ice cap (Fig. 3c), corroborating the volume time series (Fig. 2). A smaller thinning is located inland to the south of Greenland. Minor thickening ($< -100 \text{ m}$; note that the color bar scale for Fig. 3c, e, and f is logarithmic) occurs over the Greenland ice sheet, especially along the east coast, upstream of coastal thinning. The latter is associated with the largest velocity decreases after 200 years ($< -100 \text{ m yr}^{-1}$; yellow contour in Fig. 3d). In this area, the perturbation fails to trigger a grounding line retreat. In contrast, ice flow velocities increased with the grounding line inland retreats on the Fennoscandian, Svalbard and Iceland coasts, sometimes so much that there are no more observable connections between ocean and land, as is the case for Iceland after 200 years (blue contours; Fig. 3b vs. d). These contrasting responses reflect regional differences in the initial basal melt rates (Fig. 1c) that arise from regional differences in ocean temperature and salinity (through freezing temperature) at the shelf drafts, as we see in Sect. 3.2.

To get further insight into the nature of the volume variations, we decompose the ice thickness changes using mass conservation (Eq. 2) as follows:

$$H(t) - H(0) = \int_0^t (\text{SMB} - \text{BM} - C) dt - \int_0^t \nabla \cdot (\mathbf{u}H) dt, \quad (2)$$

with H representing the ice thickness, SMB representing the surface mass balance, BM representing the basal melt rate, C representing the calving rate and $\nabla \cdot (\mathbf{u}H)$ representing the ice flow divergence; dt is 1 year. We refer to the first term as the climate term (although it contains the calving, which is not, strictly speaking, a climate term). The residual is the lowering or thickening due to ice flow (dynamical effect).

The perturbation induces a loss of volume through both climatic and dynamic thinning (Fig. 3e and f.) The oceanic

melt rate perturbation is visible through negative thickness anomalies along the coasts in Fig. 3e. The inland thinning over Greenland is also caused by the climate term. The latter is due to a change in surface mass balance (less accumulation over the Greenland when the fresh water is released; Fig. S4 in the Supplement).

Thinning due to the dynamic term is most important in areas such as the west coast and southern part (proglacial lake grounding line instability) of Fennoscandia, Svalbard, Iceland, and the southern tip of the Greenland ice sheet. Also, the minor thickening in coastal areas is due to the dynamic term. In the model, the ice flux at the grounding line connects the ice thickness and velocities in the grounding zone through a power-law relationship (Tsai et al., 2015). Where the oceanic perturbation fails to trigger a grounding line retreat, the perturbation only causes a reduction in thickness that leads to a reduction in the ice flux and to upstream convergence (Fig. 3f). However, the resulting ice elevation increase in these areas is insufficient to produce a discharge through an increase in the ice sheet surface slope. This thickening effect also appears north of the North American ice sheet, at its connection with the Arctic Ocean, and east, at its connection with Baffin Bay. This effect, combined with positive anomalies of the surface mass balance to the south, contributes to the small thickening of the North American ice sheet, for experiments with at least a 10-times increase in the temperature anomaly with respect to the freezing temperature, at the shelf drafts ($X \geq 100$; Fig. 2).

Still, the overall volume loss experienced by the ice sheets is equivalent to the freshwater release toward the ocean (Fig. 4a). This freshwater flux is particularly large during the first few years in all experiments and results from both abrupt increased calving and increased basal melt fluxes in the first few years (Fig. 4b and c). The initial amplification of the melt rates leads to large losses in the form of a basal melt flux (Fig. 4b). The initial amplification of the melt rates also leads to an abrupt increase in the calving flux (Fig. 4c). Indeed, some of the shelves are rather thin before the perturbation is triggered (Fig. 1b), so the perturbation causes these




Figure 3. Ice sheet history in the perturbation experiment with factor $X = 100$. Ice sheet (a) velocity [m yr^{-1}] and (b) thickness [m] averaged over the control simulation. Sky-blue contour indicates the position of the grounding line. PERT100–CTRL anomalies of (c) thickness and (d) ice velocities as well as (e) thickness due to climate anomalies and (f) ice convergence, after 200 years. Only significant values to 1 standard deviation from those of the control are shown for (c) and (d). Please note that the color bar for the ice sheet thickness is linear, while all the other ones are logarithmic.

shelves' thickness to fall below the calving threshold (Fig. S4 in the Supplement). Then, the calving flux drops toward the control value after 10 years while the basal melt flux slowly decreases toward the control value. This results in a freshwater flux that decreases with time (but remains significant). This decrease is partly attributed to the reduced ice volume exposed to basal melting, as ice shelves thin or vanish in certain regions, such as Svalbard and Iceland. It also suggests the presence of ice sheet stabilizing mechanisms within the model, i.e., negative feedback that mitigates ice volume loss. This feedback involves a decrease in the temperature at shelf drafts, which in turn reduces the basal melt rates.

At this point, this study suggests the possibility that an episode of subsurface warming may have triggered dynamical instabilities for the Eurasian, Svalbard and Iceland ice sheets. However, after about a decade, the simulated freshwater release to the ocean decreases with time. The freshwater release has the ability to modify climate, ocean hydrography and circulation, which can in turn affect the ice sheets. In the following section, we investigate climate and ocean changes and examine how they are connected with regional ice sheet changes.

3.2 Climate and ocean changes

Subsequent to the continental meltwater release, the Atlantic meridional overturning circulation (AMOC) weakens for several years in all simulations. There are no off modes in our simulations, even with the largest perturbation factors. At the same time, the extent of the sea-ice cover increases and the surface temperatures decrease over the Northern Hemisphere (Fig. 5). The AMOC rapid decrease is followed by a slower recovery toward the control simulation values, corroborating the reduction in the freshwater inflow with time (Fig. 4a).

Although one could expect that the transient Northern Hemisphere cooling would allow ice sheet regrowth during the perturbation period, this is not the case. Indeed, the Northern Hemisphere surface cooling is essentially located over the North Atlantic Ocean (especially over the Nordic and Labrador seas), while only limited cooling occurs over Greenland and Europe, and no notable temperature changes take place over North America.

For most of these metrics, there are no past estimates available. Greenland temperatures at the NGRIP sites are the only exception (Kindler et al., 2014). At this location, the authors obtain around -40 to -45 °C around 40 kyr BP, while we




Figure 4. Time series of (a) freshwater output from GRISLI to iLOVECLIM [Sv] (as a total ice volume change) in experiments with different perturbation factors. Diamonds are the freshwater release for the first year of perturbation, light colors are the yearly outputs and saturated colors are the smoothed time series (running mean over 21 years, centered). Time series of (b) calving flux [Sv] and (c) basal melt flux [Sv], also smoothed (running mean over 21 years, centered).




Figure 5. Times series of (a) AMOC intensity in the North Atlantic [Sv], (b) Northern Hemisphere sea-ice extent [10^{12} km²], (c) Northern Hemisphere and (d) NGRIP surface temperatures [°C] in perturbation experiments with different factors. The thin black lines represent the unperturbed extension of the simulations with factors 20 and 100. Light colors are the yearly outputs and saturated colors are the smoothed time series (running mean over 21 years, centered).

simulate warmer temperatures around -35 to -40 . Paleoproxy records show that the AMOC was in a much weaker regime during stadials than interstadials but do not provide quantitative estimates in Sv (Gottschalk et al., 2015; Henry et al., 2016; Waelbroeck et al., 2018).

Differences in timings of the different variables are difficult to establish, as the time series are noisy. Considering smoothed time series (running mean over 21 years, centered), the maximum changes for the sea-ice extent ($+10^{12}$ km² in comparison to the control), the AMOC (-5 Sv), the NGRIP (-0.53 °C) and Northern Hemisphere (-0.49 °C) temperatures are reached after 14, 18, 21 and 33 years, respectively, for the experiment, with a large temperature anomaly intensity at the shelf drafts (i.e., $X = 100$; Fig. 5).

In the unperturbed simulation, convective areas are located south of the sea-ice edge in the Labrador Sea and south of the Nordic Seas (Fig. 6g). Following the freshwater release, major cooling and freshening occur at the surface, and the sea-ice extent moves southward (Fig. 6a, b, d and e). This produces a decrease in the surface density over most of the convective sites resulting in convection reduction in the beginning of the perturbation period (Fig. 6h), except east of Iceland where convection increases. We note here that the

use of a higher-resolution model for the ocean might lead to a reduced amount of freshwater advected over the convective areas and, therefore, to a dampened reduction in the AMOC or Northern Hemisphere surface temperatures, in comparison to the results presented in Fig. 5. However, the latter is not very clear because other types of interactions arising in higher-resolution models (such as freshwater transport by eddies) might also amplify the freshwater exchanges between the boundary currents and convective sites (e.g., Swingedouw et al., 2022).

At the end of the perturbation period, the hydrographic variables almost return to their control values and the ice-edge is close to the one in the control simulation. Nevertheless, the end of the perturbation period presents a slight warming at the surface in the northeast Atlantic and a negative salinity anomaly remains at the surface in the northern Nordic Seas, while a slightly positive salinity anomaly is visible south of the sea-ice edge (Fig. 6c and f).

South of the sea-ice edge, the positive temperature and salinity anomalies are present not only at the surface but also at the subsurface along the Atlantic Water pathway (Figs. 6c and f and 7c and f). The positive anomalies, therefore, suggest a resumption of the AMOC and renewed influx of heat




Figure 6. Hydrography at 5 m depth horizon, in the perturbation experiment with factor $X = 100$. Maps of (a–c) conservative temperature [°C] over the control simulation and associated anomaly over 10–40 years and over 470–500 years comparing PERT100 and the control simulation. (d–f) Same for practical salinity [psu]. (g–i) Same for convection depth [m]. Only significant values to 1 standard deviation from those of the control simulation are shown for anomalies. Sky-blue contour is the grounding line position in the control, after 40 years and after 500 years (resp. in a–c). Turquoise and dark-blue contours are annual 30 % sea-ice contours in the control and the perturbation experiment, respectively.

and salt of Atlantic origin following the period of maximum AMOC slowdown.

North of the sea-ice edge, the freshening pattern at the surface is caused by the accumulation of continental meltwater through the duration of the perturbation (Fig. 6f). Additionally, the weak meridional circulation at these latitudes does not promote the advection of the freshwater supply out of the area (Fig. S5 in the Supplement). The slight warming is driven from below, as suggested by temperature patterns at 300 m water depth (Fig. 7c).

At 300 m depth (characteristic shelf draft), the subsurface warms in several locations following the freshwater release at the surface. These locations include east and south of Newfoundland, Baffin Bay, north of the Greenland Sea and the Arctic Ocean (Fig. 7b). At the end of the perturbation, only the warming at the Nordic Seas remains significant, with values above 1 °C (Fig. 7c). This area, where the positive tem-

perature anomaly has developed and amplified, corresponds to the area where the ice flux is the most sustained over the perturbation period. The salinity also increases in the subsurface over the Nordic Seas with time, probably as the result of sea-ice production and accumulation of saline Atlantic waters. The competing effects of temperature and salinity produce an overall density decrease at this depth that is smaller than the overall density decrease at the surface (Fig. S6 in the Supplement). However, after 100 years, the vertical density gradient sporadically reverts, due to accumulation of heat in the subsurface, and the warm subsurface waters are mixed with surface waters during strong intermittent convective events (Figs. S6 and S7 in the Supplement), leading to a positive surface temperature anomaly north of the sea-ice edge (Fig. 6c).

To summarize, the amplification of the melt rates results in surface freshening and heat accumulation at the subsur-




Figure 7. Hydrography at 300 m depth horizon, in the perturbation experiment with factor $X = 100$. Maps of (a) conservative temperatures [$^{\circ}\text{C}$] over the control simulation, conservative temperature anomalies (b) over 10–40 years and conservative temperature anomalies (c) over 470–500 years between the simulation with perturbation factor 100 and the control. (d–f) Same for absolute salinity [g kg^{-1}]. Only significant values to 1 standard deviation from those of the control simulation are shown for anomalies. Yellow contours indicate 0, $+1^{\circ}\text{C}$, and 36.6 g kg^{-1} iso-contours in (a), (c) and (d), respectively. Turquoise and dark-blue contours are annual 30 % sea-ice contours in the control and the perturbation experiment, respectively.

face (around 300 m depth) in the Nordic Seas, corroborating the sustained ice discharges in this region. In Baffin Bay, the amplification of melt rates also leads to surface freshening and subsurface warming, yet only for a few decades, which is consistent with less ice discharge over this area. The reason for these regional differences can be found in background oceanic temperatures and salinity. Both these quantities are quite small in Baffin Bay when compared to the same latitudes in the Nordic Seas (yellow contours Fig. 7a and b). Therefore, the subsurface temperature anomaly that we apply to the ice sheets, even using a large perturbation factor, may be insufficient to perturb the ice sheet dynamics there. This is discussed in Sect. 4.1.

3.3 Interactions between ice sheets and oceans

We perform another set of simulations maintaining the oceanic perturbation and cutting off freshwater fluxes from the ice sheet model to the ocean model, while the hydrological cycle of the climate model remains activated and precipitation that falls over land is still routed toward the ocean. In other words, here the ocean does not respond to the continental ice meltwater and associated oceanic feedback on the ice sheets is suppressed. Oceanic and atmospheric circulations could still vary from the control simulation, in response to changes in the ice sheet elevation (that, for instance, could perturb the atmospheric physics and dynamics)

or extent of continental ice (that, for instance, could induce albedo changes).

When the freshwater fluxes are suppressed, the ice sheet response to the perturbation is amplified during the first few decades with respect to the simulation where the freshwater fluxes are taken into account (Fig. 8a and b). The Norwegian and Iceland ice sheets experience more losses (red areas; Fig. 8c). In other areas, such as the northern Labrador Sea (bounded by the east coast of the Laurentide and from the west coast to the south of the Greenland ice sheets), an exacerbated thinning is also visible, yet it remains secondary.

To quantify and see the temporal evolution of freshwater feedback on ice sheets, we compute a feedback factor γ_X (Eq. 3).

$$\gamma_X = \frac{\delta V_X - \delta V_{X,\text{noFWF}}}{\delta V_X}, \quad (3)$$

where δV_X is the continental ice volume variation with respect to time 0 in the simulation with perturbation factor X , and $\delta V_{X,\text{noFWF}}$ is the associated simulation with no freshwater flux. This definition is adapted from Goosse et al. (2018), the difference being that here we do not consider equilibrium states but do consider transient states.

For the Greenland–Iceland and Eurasian ice sheets, δV_X is negative (Fig. 2). Thus, a negative γ_X means that $0 > \delta V_X > \delta V_{X,\text{noFWF}}$ and indicates that there is more ice loss when the freshwater fluxes are not taken into account. Concerning the Greenland–Iceland ice sheet, freshwater feedback on the vol-




Figure 8. Thickness anomalies [m] between (a) perturbation experiment with factor $X = 100$ and the control simulation, (b) same with cut freshwater fluxes and the difference between the two (c) after 40 years and (f) after 500 years. Time series of feedback factor calculations, as defined in (Eq. 3), applied to (d) the Greenland–Iceland ice sheet and (e) the Eurasian ice sheet in the perturbation experiments. Lightly colored lines are the yearly time series. Boldly colored lines are the smoothed time series (running mean over 21 years, centered) with values plotted when δV_X is above a critical value of $0.25 \times 10^{14} \text{ m}^3$.

ume variation is maximal at the beginning of the simulations (Fig. 8a, b, and d). After ~ 100 years, the ice evolution becomes stable in both experiments (Figs. 2b and 8d). This is because of two different mechanisms: (1) the Iceland grounding line retreats inland until there is no more sensitivity to the perturbation, and (2) the applied perturbation is not sufficient enough to destabilize most of the Greenland coasts, where initiated thinning only leads to reduced velocities, upstream convergence and minor thickening. The resulting difference in thickness is slightly negative in the Greenland interior at the end of the simulations (blue patch; Fig. 8e). As the thickness variation is also negative when the freshwater fluxes are taken into account, it means that there are fewer losses at this location when freshwater fluxes are not taken into account. This is similar to what we have already described in Sect. 3.1: the release of freshwater leads to less accumulation over the Greenland area.

Regarding the Eurasian ice sheet, the losses are amplified when the freshwater fluxes are not taken into account in the beginning of the simulation (Fig. 8e). After ~ 200 years, the time-varying feedback factor becomes rather stable yet still negative for all perturbation experiments. This non-zero freshwater feedback signifies that the ice sheet volume is still decreasing in both cases, with and without freshwater fluxes, but at a larger rate for the second (Figs. 2c and 8e). In both

cases, the Fennoscandian ice sheet experiences a grounding line mechanical instability that implies mass loss.

Either way, ice volume decrease is faster when freshwater fluxes are not taken into account than when they are. Meltwater fluxes into the ocean bring water that is fresher than the surrounding ocean and favor sea-ice expansion, increasing stratification and reducing vertical mixing. In addition, the local latent heat flux, due to oceanic melting of the calved ice, also helps to cool the upper part of the water column and to maintain a cold water layer that extends from the surface to the ice shelf drafts. The latent heat flux associated with sub-shelf melt is not taken into account in the model, yet including it would lead to a larger cooling of the upper water column. All this acts to temporarily protect the ice shelves from the underlying warm waters (Fig. 7c). Nevertheless, the grounding line of the Iceland and Fennoscandian ice sheets eventually retreats completely in both simulations (blue contours; Figs. 6c and 8f). This suggests the conclusion that freshwater fluxes are slowing down the grounding line retreat and dampening the ice volume losses rather than halting them.

4 Discussion

4.1 Laurentide ice sheet

The Hudson ice stream is often considered a major source of massive iceberg discharges during the last glacial period (e.g., Andrews and Tedesco, 1992; Bond et al., 1992; Calov et al., 2002), while the Laurentide ice sheet is overall stable in our coupled experiments. This raises the question of whether this stability is an inherent characteristic of this ice sheet simulated by the coupled model or if the disturbance applied is insufficient to cause destabilization.

On the one hand, the Laurentide ice sheet in the fully coupled model may be too stable. This could stem from the basal drag formulation, parameter values or spatial resolution. But it could also result from biases in the climate model, in the atmosphere (a warm bias could produce widespread temperate basal conditions, by thermal diffusion through the ice) or in the ocean (low thermal forcing in Baffin Bay).

On the other hand, we have pointed out that background temperatures are rather cold at the shelf drafts and salinity is low in Baffin Bay and the adjacent Labrador Sea in comparison to the same latitudes in the eastern part of the North Atlantic. Background oceanic basal melt rates are thus weaker (around 0.003 m yr^{-1}) at the mouth of the Hudson ice stream in the control simulation than around the Fennoscandian ice sheet (around 0.3 m yr^{-1}). Therefore, in our simulations, the oceanic perturbation imposed by subsurface temperature amplification at the shelf drafts is not able to destabilize the North American ice sheet/streams even with the highest multiplicative factor.

Inaccurate representations of the ocean circulation or hydrography may contribute to underestimated ocean temperatures in Baffin Bay area at 40 kyr BP. Such inaccuracies might result from the model's low resolution (for example, leading to the misrepresentation of the extent of the subpolar gyre and/or recirculation patterns along the west coast of Greenland) or from underestimated processes, such as the sinking of brines in the Southern Ocean, which could alter the meridional overturning circulation in the Atlantic (Bouttes et al., 2010). Improving the ocean representation is a key objective for future work. Indeed, our model can simulate carbon isotopes (Bouttes et al., 2015), allowing for direct comparison with observational data from marine sediment cores. This could serve as a constraint to improve ocean representation in subsequent studies.

To overcome these two potential issues (oceanic model biases and/or weak perturbation) and to force a different ice sheet response especially in the Hudson Bay region, we conduct an additional set of experiments using constant temperature offsets within the basal melt rate formulation rather than amplifying the local temperature anomalies relative to the freezing point seen by the ice shelves. In other words, we apply a δT to T in Eq. (1), defined by $\delta T = T_{\text{offset}}$ with $T_{\text{offset}} = 2, 6, 8$ and 10°C for 500 years.

This time, the North American ice sheet volume decreases (Fig. 9a) concurrently with the other ice sheet volume decrease. This results in increased freshwater input from the ice sheets to the climate model for the largest offsets in temperature, in comparison to the previous set of experiments (Fig. 9e). Consequently, the reduction in the AMOC is amplified, along with a more pronounced decline in the NGRIP temperatures (Fig. 9d and f). In the idealized and extreme case of a temperature offset of 10°C , the AMOC shuts down, with no recovery even after the end of the perturbation. In this case, surface cooling and freshening are so large that convection ceases in the North Atlantic Ocean and instead shifts to the North Pacific Ocean (not shown).

Nonetheless, dynamical ice losses are also confined to coastal areas in this set of experiments, despite a retreat of the grounding line (e.g., Fig. 9b and c). The Laurentide ice sheet does not exhibit dynamical instability or surges of the Hudson ice stream, suggesting that this area may be too stable in our model. The simulated behavior of the Laurentide ice sheet is inconsistent with the paleo-data and the presence of ice-rafted debris of Laurentide origin within the North Atlantic marine sediments (Broecker et al., 1992; Grousset et al., 2000). There are several possible reasons for this discrepancy between the simulated behavior and the observations. It could result from the presence of biases in the ice sheet basal temperatures. In our simulations, widespread basal conditions are temperate and constant over time, while it has been shown that transitions between basal states could lead to surges for the Laurentide ice sheet (Schannwell et al., 2024). Basal temperatures depend on ice thickness, geothermal heat flux and atmospheric temperatures. Thus, varying factors like geothermal forcing, for example, could help address this issue (Hank and Tarasov, 2024). Finally, different modeling choices could lead to different responses for the Laurentide ice sheet. Modeling choices regarding the basal dynamics or geothermal heat flux affect the ability of the ice sheet to slide over bedrock and could lead to different results (e.g., Joughin et al., 2019; Sun et al., 2020; Schannwell et al., 2023). For instance, a relatively low geothermal heat flux and/or the use of a regularized Coulomb sliding law (as opposed to a Weertman-type power law) can cause a transition from a continuously flowing Hudson Strait ice stream to long periods without flow punctuated by brief surges, as shown by Hank and Tarasov (2024).

4.2 Influence of the bathymetry choice on ice sheet sensitivity

Until now, we used bathymetry corresponding to the last glacial maximum. Here we conduct the same experiment but with a pre-industrial bathymetry, as the 40 kyr BP bathymetry lies between these two bathymetries. The initial condition is built the same way as described in Sect. 2.2.1. All other parameters remain unchanged.




Figure 9. Time series of (a) North American grounded ice sheet volume anomaly [10^6 km^3], (d) AMOC [Sv], (e) freshwater input to the ocean model [Sv] and (f) NGRIP temperatures [$^{\circ}\text{C}$] in perturbation experiments with $T_{\text{offset}} = 2, 6, 8$ and 10°C as well as the experiment with $X = 100$. Snapshot of (b) thickness anomaly [m] and (c) integrated ice convergence anomaly between the $T_{\text{offset}} = 8^{\circ}\text{C}$ data and the control experiment after 200 years. Only significant values to 1 standard deviation from those of the control are shown for (b).

The LGM and PI bathymetries produce different ocean currents and advection patterns of heat and salt, resulting in different distributions of water masses (Fig. S8 in the Supplement). Therefore, changing the bathymetry also produces a change in the ice sheet's geometry, leading to a reduced total volume of about $22 \times 10^6 \text{ km}^3$ (44 m SLE). Differences in continental ice distribution include a North American ice sheet that reaches lower latitudes on its eastern side and a Fennoscandian ice sheet of smaller extent. Additionally, the PI bathymetry also leads to a reduction in the southeast extent of the Greenland ice sheet (Fig. S8 in the Supplement).

Due to the differing geometries and ice exposure to warmer waters in the Nordic Seas and the Labrador Sea, the regions experiencing volume losses are not identical when we amplify the basal melt rates. The Greenland ice sheet loses significant volume along its southeast coast, particularly at the beginning of the experiment, with a large perturbation factor ($X = 100$, Fig. S9 in the Supplement). The Eurasian ice sheet still experiences substantial losses in its northern regions. The Laurentide ice sheet undergoes dynamical thinning with this geometry, but these losses are mostly confined to its southeastern region (Fig. S9 in the Supplement), where it is exposed to warmer waters (Fig. S8 in the Supplement). This produces a relatively small difference in the amount of freshwater input to the ocean following the onset of the perturbation, with respect to the experiment performed with the LGM bathymetry. The key climate variables (sea-ice extent, AMOC, Northern Hemisphere and NGRIP temperatures) follow similar trajectories to those obtained with the LGM bathymetry (Figs. 5 and S10).

Although we could expect a change in the fully coupled model response to the oceanic perturbation following the

change in bathymetry and distribution of the water masses, this is not the case here. The evolution of the feedback factor is also similar to what we obtained with the LGM bathymetry. The ice volume losses without freshwater input to the ocean are larger for both Greenland–Iceland and Eurasian ice sheets, especially at the beginning of the experiment. After ~ 50 years, the volume of the North American ice sheet remains stable in the experiment with the freshwater fluxes, while it continues to decrease with suppressed freshwater fluxes (Fig. S11 in the Supplement). Thus, the feedback factor is negative and the freshwater release dampens the ice volume losses with the pre-industrial bathymetry just as it does with the LGM bathymetry (Fig. 10). Note that γ_{100} is positive for the North American ice sheet with LGM bathymetry; this accounts for the mass gain described in Sect. 3.1.

4.3 Dansgaard–Oeschger and Heinrich events framework

Our simulations suggest ocean subsurface warming could amplify stadial conditions. Indeed, the increase in oceanic basal melt rates leads to a temporary increase in both calving fluxes and basal meltwater release. The resulting freshwater release induces a temporary slowdown of the AMOC and decrease in the Northern Hemisphere temperature.

According to the air temperature record of Kindler et al. (2014), the onset of Heinrich stadial 4 corresponds to a 6.5°C temperature decrease and it is followed by a 10°C increase at the NGRIP site. In comparison, our simulations present low temperature variation magnitudes, ranging between 0.5 and 2°C , for the smoothed time series (Fig. 5d). In




Figure 10. Time series of the feedback factor, as defined in (Eq. 3), in the perturbation experiments with factor $X = 100$, showing the pre-industrial setup and the LGM setup for the different ice sheets. The lightly colored lines are the yearly time series. The boldly colored lines are the smoothed time series (running mean over 21 years, centered) with values plotted when δV_X is above a critical value of $0.25 \times 10^{14} \text{ m}^3$.

the model, Greenland temperature changes are mainly produced by AMOC changes. Following the onset of the perturbation and the inflow of freshwater, the AMOC intensity temporarily reduces but remains relatively high (Fig. 5a). Consequently, Greenland temperatures also remain relatively high, leading to small anomalies.

In addition, in the paleoclimatic records, Heinrich events are followed by an atmospheric warming over the Greenland ice sheet and the North Atlantic region (Bond et al., 1993; Kindler et al., 2014), namely Dansgaard–Oeschger warming toward interstadial conditions. Here, our model results do not display such warming when the perturbation is released, which is inconsistent with observations (Fig. 5c and d). Some authors have shown that increased sea-ice extent during cold periods can lead to heat accumulation at the subsurface in the Nordic Seas (e.g., Rasmussen and Thomsen, 2004; Sadatzki et al., 2019). At some point, this accumulated heat destabilizes the water column, reversing the vertical density gradient. This reversal causes rapid oceanic mixing, bringing warm waters to the surface. The heat release to the atmosphere marks the onset of a Dansgaard–Oeschger interstadial (Dokken et al., 2013). In our simulation, the heat accumulated in the Nordic Seas (Fig. 7) dissipates once the perturbation is released (Fig. S12 in the Supplement). Part of this heat is transferred to the atmosphere as indicated by the sea-ice contour moving toward higher latitudes; the rising ocean surface temperatures during the first 30 years post-perturbation (Fig. S12 in the Supplement); and the strong, intermittent convective events that occurred during the first 100 years after the perturbation ends (Fig. S7 in the Supplement). However, this has no imprint on Greenland temperatures. One possibility is that the heat reservoir is insufficient to counteract the rapid re-stratification that follows the sur-

face heat release to the atmosphere (Fig. S6 in the Supplement), preventing the climate system from shifting to a new regime in our simulations. This could be explored in future studies.

The atmospheric circulation was not a primary focus of this study, yet biases in the atmosphere model could also influence the ice sheet’s volume dynamics and could be invoked to explain the absence of a DO-like event following the release of the perturbation. Climate biases have been assessed for the pre-industrial climate by Quiquet et al. (2021b): there is an overall cold bias around the Nordic Seas sector as well as the Greenland and Fennoscandian ice sheets, with underestimated precipitation. In contrast, there is a relatively warm bias over North America, with underestimated precipitation except in mountainous areas where it tends to be overestimated (Quiquet et al., 2021b). While it is unclear how these biases are translated at 40 kyr BP, they could affect the ice sheet’s ability for regrowth and modulate the rate of its volume decay. It is noteworthy that previous studies using the iLOVECLIM model have shown its ability to simulate abrupt temperature increases at the NGRIP site ($\sim +5^\circ\text{C}$) associated with a significant rise in AMOC strength ($\sim +15 \text{ Sv}$) (Quiquet et al., 2021b, Figs. 3 and 6). This suggests the finding that, in our study, it is rather a lack of AMOC variability that prevents sudden warming after the perturbation is halted.

We thus see that our simulations lack some of the key features of the paleoclimatic records. Beyond the absence of Laurentide ice sheet instabilities discussed in Sect. 4.1, these shortcomings include an overly weak intensity of temperature anomalies at the NGRIP site and the absence of rapid warming toward interstadial conditions. These limitations are probably related to the weak changes (in both intensity and duration) of the AMOC in response to freshwater flux originating from the ice sheets, this flux being likely underestimated over time due to the stability of the Laurentide ice sheet in our simulations.

5 Conclusions

In this study, we test the hypothesis that ocean subsurface warming triggers instabilities in the Northern Hemisphere ice sheets. We use an Earth model of intermediate complexity (iLOVECLIM) coupled with an ice sheet model (GRISLI) and apply an oceanic perturbation at the shelves draft by multiplying the background oceanic basal melt rates with different factors. Our numerical experiments lead to several conclusions:

1. Imposed amplified oceanic basal melt rates induce ice volume changes. Fresh water is released to the ocean through increased calving and oceanic basal meltwater discharges. For most places, the increase in oceanic basal melt rates leads to grounding line retreat, dynamical thinning and volume loss, while in some places the initial ice thickness decrease is small and leads to up-

stream minor dynamical thickening. Nevertheless, the influx of fresh water to the ocean induces a temporary increase in the sea-ice extent; a reduction in convection in the Labrador and Nordic Seas; and, hence, a reduction in the Atlantic meridional overturning circulation and surface temperatures in the Northern Hemisphere as well as subsurface warming in the Nordic Seas.

2. The release of fresh water and the effect of latent heat on ocean temperature dampen ice sheet discharges (negative feedback).
3. The Laurentide ice sheet volume does not vary and the Hudson Strait ice stream is stable in our experiments with amplified basal melt rates. This can be partly explained by weaker background temperature and salinity at shelf drafts in Baffin Bay and the Labrador Sea in comparison to the Nordic Seas. Imposing constant temperature offsets, up to 10 °C, produces a grounding line retreat and a volume decrease in the North American ice sheet, although the Hudson Strait region shows no dynamical instability. This suggests the possibility that the model is too stable in this region.
4. An episode of subsurface warming could trigger dynamical instabilities and ice discharges along the coasts of the Nordic Seas, affecting the Fennoscandian, Svalbard, and Iceland ice sheets. These events could lead to significant, albeit temporary, perturbation of the climate on a large scale, including sea-ice expansion, a weakening of the Atlantic meridional overturning circulation and Northern Hemisphere cooling.

While our modeled Laurentide ice sheet does not exhibit dynamical instability, this finding does not rule out the ocean as a potential trigger for Heinrich events. Instead, the limited inland propagation of oceanic perturbations would suggest an overall stability and/or a lack of nonlinear dynamics at the ice sheet–bedrock interface in our simulations. Different modeling choices regarding the basal dynamics beneath the ice sheet (i.e., a change in the sliding law) could help to address this issue. As we know from the presence of ice-rafted debris of Laurentide origin within the North Atlantic marine sediments (e.g., Broecker et al., 1992; Grousset et al., 2000), this ice sheet has been dynamically unstable in the past. This could be investigated in future work. Moreover, to better contextualize this study within the framework of paleo-observations of massive iceberg discharges and Dansgaard–Oeschger events, it is also key to represent the rapid warming toward interstadial conditions following Heinrich events. The effects of the Antarctic ice sheet, which are prescribed in this study, could be examined. Indeed, the extent of the Antarctic ice sheet may also fluctuate, leading to variations in the production of deep water around Antarctica (Paillard and Labeyrie, 1994). Accounting for these variations might result in more variability in the oceanic component than the

present study demonstrates. This is a potential area for future research. Lastly, our model can simulate ocean carbon isotopes (Bouttes et al., 2015), allowing direct comparison with observational data from marine sediment cores and providing additional constraints for the ocean component. This is another promising research direction.

Data availability. The source data of the figures in the main text of the paper are accessible on the Zenodo repository at <https://doi.org/10.5281/zenodo.12793237> (Abot et al., 2024). All color schemes are color-vision-deficiency-friendly and perceptually uniform, accessible from a freely available package (Crameri, 2023).

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/cp-21-1123-2025-supplement>.

Author contributions. LA carried out the simulations, analyzed the results and wrote the paper. All other authors contributed to designing the project and the simulations, analyzing the results and adding comments to improve the paper.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. We gratefully thank Casimir de Lavergne and Nathaëlle Bouttes for their help and early discussions. We also gratefully thank Matteo Willeit and an anonymous reviewer whose comments contributed to improving the paper. This work is a contribution to the French INSU (LEFE) project LASSO.

Financial support. Louise Abot received funding from a Sorbonne Université PhD scholarship.

Review statement. This paper was edited by Lev Tarasov and reviewed by Matteo Willeit and one anonymous referee.

References

Abot, L., Quiquet, A., and Waelbroeck, C.: Northern Hemisphere ice sheets and ocean interactions during the last glacial pe-

- riod in a coupled ice sheet–climate model, Zenodo [code], <https://doi.org/10.5281/zenodo.12793237>, 2024.
- Alvarez-Solas, J., Charbit, S., Ritz, C., Paillard, D., Ramstein, G., and Dumas, C.: Links between ocean temperature and iceberg discharge during Heinrich events, *Nat. Geosci.*, 3, 122–126, <https://doi.org/10.1038/ngeo752>, 2010.
- Alvarez-Solas, J., Robinson, A., Montoya, M., and Ritz, C.: Ice-berg discharges of the last glacial period driven by oceanic circulation changes, *P. Natl. Acad. Sci. USA*, 110, 16350–16354, <https://doi.org/10.1073/pnas.1306622110>, 2013.
- Alvarez-Solas, J., Banderas, R., Robinson, A., and Montoya, M.: Ocean-driven millennial-scale variability of the Eurasian ice sheet during the last glacial period simulated with a hybrid ice-sheet–shelf model, *Clim. Past*, 15, 957–979, <https://doi.org/10.5194/cp-15-957-2019>, 2019.
- Andrews, J. and Tedesco, K.: Detrital carbonate-rich sediments, northwestern Labrador Sea: implications for ice-sheet dynamics and iceberg rafting (Heinrich) events in the North Atlantic, *Geology*, 20, 1087–1090, [https://doi.org/10.1130/0091-7613\(1992\)020<1087:DCRSNL>2.3.CO;2](https://doi.org/10.1130/0091-7613(1992)020<1087:DCRSNL>2.3.CO;2), 1992.
- Arthur, F., Roche, D. M., Fyfe, R., Quiquet, A., and Renssen, H.: Simulations of the Holocene climate in Europe using an interactive downscaling within the iLOVECLIM model (version 1.1), *Clim. Past*, 19, 87–106, <https://doi.org/10.5194/cp-19-87-2023>, 2023.
- Arz, H. W., Lamy, F., Ganopolski, A., Nowaczyk, N., and Pätzold, J.: Dominant Northern Hemisphere climate control over millennial-scale glacial sea-level variability, *Quaternary Sci. Rev.*, 26, 312–321, <https://doi.org/10.1016/j.quascirev.2006.07.016>, 2007.
- Barker, S., Chen, J., Gong, X., Jonkers, L., Knorr, G., and Thornalley, D.: Icebergs not the trigger for North Atlantic cold events, *Nature*, 520, 333–336, <https://doi.org/10.1038/nature14330>, 2015.
- Bassis, J. N., Petersen, S. V., and Mac Cathles, L.: Heinrich events triggered by ocean forcing and modulated by isostatic adjustment, *Nature*, 542, 332–334, <https://doi.org/10.1038/nature21069>, 2017.
- Batchelor, C. L., Margold, M., Krapp, M., Murton, D. K., Dalton, A. S., Gibbard, P. L., Stokes, C. R., Murton, J. B., and Manica, A.: The configuration of Northern Hemisphere ice sheets through the Quaternary, *Nat. Commun.*, 10, 3713, <https://doi.org/10.1038/s41467-019-11601-2>, 2019.
- Berger, A.: Long-term variations of caloric insolation resulting from the Earth's orbital elements, *Quaternary Res.*, 9, 139–167, [https://doi.org/10.1016/0033-5894\(78\)90064-9](https://doi.org/10.1016/0033-5894(78)90064-9), 1978.
- Bond, G., Heinrich, H., Broecker, W., Labeyrie, L., McManus, J., Andrews, J., Huon, S., Jantschik, R., Clasen, S., Simet, C., Tedesco, K., Klas, M., Bonani, G., and Ivy, S.: Evidence for massive discharges of icebergs into the North Atlantic ocean during the last glacial period, *Nature*, 360, 245–249, <https://doi.org/10.1038/360245a0>, 1992.
- Bond, G., Broecker, W., Johnsen, S., McManus, J., Labeyrie, L., Jouzel, J., and Bonani, G.: Correlations between climate records from North Atlantic sediments and Greenland ice, *Nature*, 365, 143–147, <https://doi.org/10.1038/365143a0>, 1993.
- Bond, G. C. and Lotti, R.: Iceberg discharges into the North Atlantic on millennial time scales during the last glaciation, *Science*, 267, 1005–1010, <https://doi.org/10.1126/science.267.5200.1005>, 1995.
- Bouttes, N., Paillard, D., and Roche, D. M.: Impact of brine-induced stratification on the glacial carbon cycle, *Clim. Past*, 6, 575–589, <https://doi.org/10.5194/cp-6-575-2010>, 2010.
- Bouttes, N., Roche, D. M., Mariotti, V., and Bopp, L.: Including an ocean carbon cycle model into iLOVECLIM (v1.0), *Geosci. Model Dev.*, 8, 1563–1576, <https://doi.org/10.5194/gmd-8-1563-2015>, 2015.
- Bouttes, N., Swingedouw, D., Roche, D. M., Sanchez-Goni, M. F., and Crosta, X.: Response of the carbon cycle in an intermediate complexity model to the different climate configurations of the last nine interglacials, *Clim. Past*, 14, 239–253, <https://doi.org/10.5194/cp-14-239-2018>, 2018.
- Bouttes, N., Kwiatkowski, L., Bougeot, E., Berger, M., Brovkin, V., and Munhoven, G.: Projections of coral reef carbonate production from a global climate–coral reef coupled model, *EGU sphere* [preprint], <https://doi.org/10.5194/egusphere-2024-3738>, 2024.
- Broecker, W., Bond, G., Klas, M., Clark, E., and McManus, J.: Origin of the northern Atlantic's Heinrich events, *Clim. Dynam.*, 6, 265–273, <https://doi.org/10.1007/BF00193540>, 1992.
- Caley, T., Roche, D. M., and Renssen, H.: Orbital Asian summer monsoon dynamics revealed using an isotope-enabled global climate model, *Nat. Commun.*, 5, 5371, <https://doi.org/10.1038/ncomms6371>, 2014.
- Calov, R., Ganopolski, A., Petoukhov, V., Claussen, M., and Greve, R.: Large-scale instabilities of the Laurentide ice sheet simulated in a fully coupled climate-system model, *Geophys. Res. Lett.*, 29, 69–1–69–4, <https://doi.org/10.1029/2002GL016078>, 2002.
- Cramer, F.: Scientific colour maps (8.0.1), Zenodo [code], <https://doi.org/10.5281/zenodo.8409685>, 2023.
- Crotti, I., Quiquet, A., Landais, A., Stenni, B., Wilson, D. J., Severi, M., Mulvaney, R., Wilhelms, F., Barbante, C., and Frezzotti, M.: Wilkes subglacial basin ice sheet response to Southern Ocean warming during late Pleistocene interglacials, *Nat. Commun.*, 13, 5328, <https://doi.org/10.1038/s41467-022-32847-3>, 2022.
- Dalton, A. S., Stokes, C. R., and Batchelor, C. L.: Evolution of the Laurentide and Innuitian ice sheets prior to the Last Glacial Maximum (115 ka to 25 ka), *Earth-Sci. Rev.*, 224, 103875, <https://doi.org/10.1016/j.earscirev.2021.103875>, 2022.
- Dansgaard, W., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N., Hammer, C. U., and Oeschger, H.: North Atlantic climatic oscillations revealed by deep Greenland ice cores, *Climate Processes and Climate Sensitivity*, 29, 288–298, <https://doi.org/10.1029/GM029p0288>, 1984.
- Dansgaard, W., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N. S., Hammer, C. U., Hvidberg, C. S., Steffensen, J. P., Sveinbjörnsdóttir, A. E., Jouzel, J., and Bond, G.: Evidence for general instability of past climate from a 250-kyr ice-core record, *Nature*, 364, 218–220, <https://doi.org/10.1038/364218a0>, 1993.
- Dokken, T. M., Nisancioglu, K. H., Li, C., Battisti, D. S., and Kissel, C.: Dansgaard-Oeschger cycles: interactions between ocean and sea ice intrinsic to the Nordic seas, *Paleoceanography*, 28, 491–502, <https://doi.org/10.1002/palo.20042>, 2013.
- Elliot, M., Labeyrie, L., Bond, G., Cortijo, E., Turon, J.-L., Tisnerat, N., and Duplessy, J.-C.: Millennial-scale ice-

- berg discharges in the Irminger Basin during the last glacial period: relationship with the Heinrich events and environmental settings, *Paleoceanography*, 13, 433–446, <https://doi.org/10.1029/98PA01792>, 1998.
- Favier, L., Jourdain, N. C., Jenkins, A., Merino, N., Durand, G., Gagliardini, O., Gillet-Chaulet, F., and Mathiot, P.: Assessment of sub-shelf melting parameterisations using the ocean–ice-sheet coupled model NEMO(v3.6)–Elmer/Ice(v8.3), *Geosci. Model Dev.*, 12, 2255–2283, <https://doi.org/10.5194/gmd-12-2255-2019>, 2019.
- Fichefet, T. and Maqueda, M. M.: Sensitivity of a global sea ice model to the treatment of ice thermodynamics and dynamics, *J. Geophys. Res.-Oceans*, 102, 12609–12646, <https://doi.org/10.1029/97JC00480>, 1997.
- Goosse, H., Brovkin, V., Fichefet, T., Haarsma, R., Huybrechts, P., Jongma, J., Mouchet, A., Selten, F., Barriat, P.-Y., Campin, J.-M., Deleersnijder, E., Driesschaert, E., Goelzer, H., Janssens, I., Loutre, M.-F., Morales Maqueda, M. A., Opsteegh, T., Mathieu, P.-P., Munhoven, G., Pettersson, E. J., Renssen, H., Roche, D. M., Schaeffer, M., Tartinvill, B., Timmermann, A., and Weber, S. L.: Description of the Earth system model of intermediate complexity LOVECLIM version 1.2, *Geosci. Model Dev.*, 3, 603–633, <https://doi.org/10.5194/gmd-3-603-2010>, 2010.
- Goosse, H., Kay, J. E., Armour, K. C., Bodas-Salcedo, A., Chepfer, H., Docquier, D., Jonko, A., Kushner, P. J., Lecomte, O., Massonnet, F., Park, H.-S., Pithan, F., G., S., and Vancoppenolle, M.: Quantifying climate feedbacks in polar regions, *Nat. Commun.*, 9, 1919, <https://doi.org/10.1038/s41467-018-04173-0>, 2018.
- Gottschalk, J., Skinner, L. C., Misra, S., Waelbroeck, C., Menviel, L., and Timmermann, A.: Abrupt changes in the southern extent of North Atlantic Deep Water during Dansgaard–Oeschger events, *Nat. Geosci.*, 8, 950–954, <https://doi.org/10.1038/ngeo2558>, 2015.
- Gowan, E. J., Zhang, X., Khosravi, S., Rovere, A., Stocchi, P., Hughes, A. L., Gyllencreutz, R., Mangerud, J., Svendsen, J.-I., and Lohmann, G.: A new global ice sheet reconstruction for the past 80 000 years, *Nat. Commun.*, 12, 1199, <https://doi.org/10.1038/s41467-021-21469-w>, 2021.
- Grousset, F. E., Pujol, C., Labeyrie, L., Auffret, G., and Boelaert, A.: Were the North Atlantic Heinrich events triggered by the behavior of the European ice sheets?, *Geology*, 28, 123–126, 2000.
- Gudmundsson, G. H., Paolo, F. S., Adusumilli, S., and Fricker, H. A.: Instantaneous Antarctic ice sheet mass loss driven by thinning ice shelves, *Geophys. Res. Lett.*, 46, 13903–13909, <https://doi.org/10.1029/2019GL085027>, 2019.
- Gwyther, D. E., Cougnon, E. A., Galton-Fenzi, B. K., Roberts, J. L., Hunter, J. R., and Dinniman, M. S.: Modelling the response of ice shelf basal melting to different ocean cavity environmental regimes, *Ann. Glaciol.*, 57, 131–141, <https://doi.org/10.1017/aog.2016.31>, 2016.
- Hank, K. and Tarasov, L.: The comparative role of physical system processes in Hudson Strait ice stream cycling: a comprehensive model-based test of Heinrich event hypotheses, *Clim. Past*, 20, 2499–2524, <https://doi.org/10.5194/cp-20-2499-2024>, 2024.
- Heinrich, H.: Origin and consequences of cyclic ice rafting in the northeast Atlantic Ocean during the past 130,000 years, *Quaternary Res.*, 29, 142–152, [https://doi.org/10.1016/0033-5894\(88\)90057-9](https://doi.org/10.1016/0033-5894(88)90057-9), 1988.
- Henry, L., McManus, J., Curry, W., Roberts, N., Piotrowski, A., and Keigwin, L.: North Atlantic ocean circulation and abrupt climate change during the last glaciation, *Science*, 353, 470–474, <https://doi.org/10.1126/science.aaf5529>, 2016.
- Holland, D. M., Nicholls, K. W., and Basinski, A.: The southern ocean and its interaction with the Antarctic ice sheet, *Science*, 367, 1326–1330, <https://doi.org/10.1126/science.aaz5491>, 2020.
- Holland, P. R., Jenkins, A., and Holland, D. M.: The response of ice shelf basal melting to variations in ocean temperature, *J. Climate*, 21, 2558–2572, <https://doi.org/10.1175/2007JCLI1909.1>, 2008.
- Hughes, A. L., Gyllencreutz, R., Lohne, Ø. S., Mangerud, J., and Svendsen, J. I.: The last Eurasian ice sheets—a chronological database and time-slice reconstruction, *DATED-1*, *Boreas*, 45, 1–45, <https://doi.org/10.1111/bor.12142>, 2015.
- IPCC: Climate change 2023: synthesis report. Contribution of working groups I, II and III to the sixth assessment report of the intergovernmental panel on climate change, edited by: Core Writing Team, Lee, H. and Romero, J., Geneva, Switzerland, <https://doi.org/10.59327/IPCC/AR6-9789291691647>, 2023.
- Johnsen, S. J., Clausen, H. B., Dansgaard, W., Fuhrer, K., Gundestrup, N., Hammer, C. U., Iversen, P. I., Jouzel, J., Stauffer, B., and Steffensen, J. P.: Irregular glacial interstadials recorded in a new Greenland ice core, *Nature*, 359, 311–313, <https://doi.org/10.1038/359311a0>, 1992.
- Johnson, G. C. and Lyman, J. M.: Warming trends increasingly dominate global ocean, *Nat. Clim. Change*, 10, 757–761, <https://doi.org/10.1038/s41558-020-0822-0>, 2020.
- Jonkers, L., Moros, M., Prins, M. A., Dokken, T., Dahl, C. A., Dirks, N., Perner, K., and Brummer, G.-J. A.: A reconstruction of sea surface warming in the northern North Atlantic during MIS 3 ice-rafting events, *Quaternary Sci. Rev.*, 29, 1791–1800, <https://doi.org/10.1016/j.quascirev.2010.03.014>, 2010.
- Joughin, I. and Alley, R. B.: Stability of the West Antarctic ice sheet in a warming world, *Nat. Geosci.*, 4, 506–513, <https://doi.org/10.1038/ngeo1194>, 2011.
- Joughin, I., Smith, B. E., and Schoof, C. G.: Regularized Coulomb friction laws for ice sheet sliding: application to Pine Island Glacier, Antarctica, *Geophys. Res. Lett.*, 46, 4764–4771, <https://doi.org/10.1029/2019GL082526>, 2019.
- Kindler, P., Guillevic, M., Baumgartner, M., Schwander, J., Landais, A., and Leuenberger, M.: Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core, *Clim. Past*, 10, 887–902, <https://doi.org/10.5194/cp-10-887-2014>, 2014.
- Kleman, J., Jansson, K., De Angelis, H., Stroeven, A. P., Hättestrand, C., Alm, G., and Glasser, N.: North American Ice Sheet build-up during the last glacial cycle, 115–21 kyr, *Quaternary Sci. Rev.*, 29, 2036–2051, <https://doi.org/10.1016/j.quascirev.2010.04.021>, 2010.
- Laske, G.: A global digital map of sediment thickness, *Eos Trans. AGU*, 78, F483, 1997.
- Le Meur, E. and Huybrechts, P.: A comparison of different ways of dealing with isostasy: examples from modelling the Antarctic ice sheet during the last glacial cycle, *Ann. Glaciol.*, 23, 309–317, <https://doi.org/10.3189/S0260305500013586>, 1996.
- Lhardy, F., Bouttes, N., Roche, D. M., Crosta, X., Waelbroeck, C., and Paillard, D.: Impact of Southern Ocean surface conditions on deep ocean circulation during the LGM: a model analysis,

- Clim. Past, 17, 1139–1159, <https://doi.org/10.5194/cp-17-1139-2021>, 2021.
- Li, D., DeConto, R. M., Pollard, D., and Hu, Y.: Competing climate feedbacks of ice sheet freshwater discharge in a warming world, *Nat. Commun.*, 15, 5178, <https://doi.org/10.1038/s41467-024-49604-3>, 2024.
- Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura, K., and Stocker, T. F.: High-resolution carbon dioxide concentration record 650,000–800,000 years before present, *Nature*, 453, 379–382, <https://doi.org/10.1038/nature06949>, 2008.
- MacAyeal, D.: Binge/purge oscillations of the Laurentide ice sheet as a cause of the North Atlantic’s Heinrich events, *Paleoceanography*, 8, 775–784, <https://doi.org/10.1029/93PA02200>, 1993.
- Makinson, K., Holland, P. R., Jenkins, A., Nicholls, K. W., and Holland, D. M.: Influence of tides on melting and freezing beneath Filchner-Ronne Ice Shelf, Antarctica, *Geophys. Res. Lett.*, 38, 6, <https://doi.org/10.1029/2010GL046462>, 2011.
- Marcott, S. A., Clark, P. U., Padman, L., Klinkhammer, G. P., Springer, S. R., Liu, Z., Otto-Bliesner, B. L., Carlson, A. E., Ungerer, A., Padman, J., He, F., Cheng, J., and Schmittner, A.: Ice-shelf collapse from subsurface warming as a trigger for Heinrich events, *P. Natl. Acad. Sci. USA*, 108, 13415–13419, <https://doi.org/10.1073/pnas.1104772108>, 2011.
- Margold, M., Stokes, C. R., and Clark, C. D.: Ice streams in the Laurentide Ice Sheet: identification, characteristics and comparison to modern ice sheets, *Earth-Sci. Rev.*, 143, 117–146, <https://doi.org/10.1016/j.earscirev.2015.01.011>, 2015.
- Marshall, S. J., Tarasov, L., Clarke, G. K., and Peltier, W. R.: Glaciological reconstruction of the Laurentide Ice Sheet: physical processes and modelling challenges, *Can. J. Earth Sci.*, 37, 769–793, <https://doi.org/10.1139/e99-113>, 2000.
- Martinson, D. G., Pisias, N. G., Hays, J. D., Imbrie, J., Moore Jr., T. C., and Shackleton, N. J.: Age dating and the orbital theory of the ice ages: development of a high-resolution 0 to 300,000-year Chronostratigraphy 1, *Quaternary Res.*, 27, 1–29, [https://doi.org/10.1016/0033-5894\(87\)90046-9](https://doi.org/10.1016/0033-5894(87)90046-9), 1987.
- Max, L., Nürnberg, D., Chiessi, C. M., Lenz, M. M., and Mulitza, S.: Subsurface ocean warming preceded Heinrich Events, *Nat. Commun.*, 13, 4217, <https://doi.org/10.1038/s41467-022-31754-x>, 2022.
- Mignot, J., Ganopolski, A., and Levermann, A.: Atlantic subsurface temperatures: response to a shutdown of the overturning circulation and consequences for its recovery, *J. Climate*, 20, 4884–4898, <https://doi.org/10.1175/JCLI4280.1>, 2007.
- Miller, G. H. and Andrews, J. T.: Hudson Bay was not deglaciated during MIS-3, *Quaternary Sci. Rev.*, 225, 105944, <https://doi.org/10.1016/j.quascirev.2019.105944>, 2019.
- Millero, F.: Freezing point of sea water, eighth report of the Joint Panel of Oceanographic Tables and Standards, Appendix 6, UNESCO Tech. Pap. Mar. Sci., 6, 29–31, 1978.
- Moorman, R., Morrison, A. K., and Mc C. Hogg, A.: Thermal responses to Antarctic ice shelf melt in an eddy-rich global ocean–sea ice model, *J. Climate*, 33, 6599–6620, <https://doi.org/10.1175/JCLI-D-19-0846.1>, 2020.
- Naughten, K. A., Holland, P. R., and De Rydt, J.: Unavoidable future increase in West Antarctic ice-shelf melting over the twenty-first century, *Nat. Clim. Change*, 13, 1222–1228, <https://doi.org/10.1038/s41558-023-01818-x>, 2023.
- Paillard, D. and Labeyrie, L.: Role of the thermohaline circulation in the abrupt warming after Heinrich events, *Nature*, 372, 162–164, <https://doi.org/10.1038/372162a0>, 1994.
- Pico, T., Birch, L., Weisenberg, J., and Mitrovica, J.: Refining the Laurentide Ice Sheet at Marine Isotope Stage 3: a data-based approach combining glacial isostatic simulations with a dynamic ice model, *Quaternary Sci. Rev.*, 195, 171–179, <https://doi.org/10.1016/j.quascirev.2018.07.023>, 2018.
- Pollard, D.: A simple parameterization for ice sheet ablation rate, *Tellus*, 32, 384–388, <https://doi.org/10.3402/tellusa.v32i4.10593>, 1980.
- Pritchard, H., Ligtenberg, S. R., Fricker, H. A., Vaughan, D. G., van den Broeke, M. R., and Padman, L.: Antarctic ice-sheet loss driven by basal melting of ice shelves, *Nature*, 484, 502–505, <https://doi.org/10.1038/nature10968>, 2012.
- Quiquet, A., Dumas, C., Ritz, C., Peyaud, V., and Roche, D. M.: The GRISLI ice sheet model (version 2.0): calibration and validation for multi-millennial changes of the Antarctic ice sheet, *Geosci. Model Dev.*, 11, 5003–5025, <https://doi.org/10.5194/gmd-11-5003-2018>, 2018a.
- Quiquet, A., Roche, D. M., Dumas, C., and Paillard, D.: Online dynamical downscaling of temperature and precipitation within the iLOVECLIM model (version 1.1), *Geosci. Model Dev.*, 11, 453–466, <https://doi.org/10.5194/gmd-11-453-2018>, 2018b.
- Quiquet, A., Dumas, C., Paillard, D., Ramstein, G., Ritz, C., and Roche, D. M.: Deglacial ice sheet instabilities induced by proglacial lakes, *Geophys. Res. Lett.*, 48, e2020GL092141, <https://doi.org/10.1029/2020GL092141>, 2021a.
- Quiquet, A., Roche, D. M., Dumas, C., Bouttes, N., and Lhardy, F.: Climate and ice sheet evolutions from the last glacial maximum to the pre-industrial period with an ice-sheet–climate coupled model, *Clim. Past*, 17, 2179–2199, <https://doi.org/10.5194/cp-17-2179-2021>, 2021b.
- Rasmussen, S. O., Bigler, M., Blockley, S. P., Blunier, T., Buchardt, S. L., Clausen, H. B., Cvijanovic, I., Dahl-Jensen, D., Johnsen, S. J., Fischer, H., Gkinis, V., Guillevic, M., Hoek, W. Z., Lowe, J. J., Pedro, J. B., Popp, T., Seierstad, I. K., Stefensen, J. P., Svensson, A. M., Vallengaard, P., Vinther, B. M., Walker, M. J. C., Wheatley, J. J., and Winstrup, M.: A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy, *Quaternary Sci. Rev.*, 106, 14–28, <https://doi.org/10.1016/j.quascirev.2014.09.007>, 2014.
- Rasmussen, T. L. and Thomsen, E.: The role of the North Atlantic Drift in the millennial timescale glacial climate fluctuations, *Palaeogeogr. Palaeoclimatol.*, 210, 101–116, <https://doi.org/10.1016/j.palaeo.2004.04.005>, 2004.
- Reese, R., Gudmundsson, G. H., Levermann, A., and Winkelmann, R.: The far reach of ice-shelf thinning in Antarctica, *Nat. Clim. Change*, 8, 53–57, <https://doi.org/10.1038/s41558-017-0020-x>, 2018.
- Ritz, C., Rommelaere, V., and Dumas, C.: Modeling the evolution of Antarctic ice sheet over the last 420,000 years: implications for altitude changes in the Vostok region, *J. Geophys. Res.-Atmos.*, 106, 31943–31964, <https://doi.org/10.1029/2001JD900232>, 2001.
- Roche, D. M., Dumas, C., Bügelmayr, M., Charbit, S., and Ritz, C.: Adding a dynamical cryosphere to iLOVECLIM (version 1.0):

- coupling with the GRISLI ice-sheet model, *Geosci. Model Dev.*, 7, 1377–1394, <https://doi.org/10.5194/gmd-7-1377-2014>, 2014.
- Sadatzki, H., Dokken, T. M., Berben, S. M., Muschitiello, F., Stein, R., Fahl, K., Menviel, L., Timmermann, A., and Jansen, E.: Sea ice variability in the southern Norwegian Sea during glacial Dansgaard-Oeschger climate cycles, *Science Advances*, 5, eaau6174, <https://doi.org/10.1126/sciadv.aau6174>, 2019.
- Schannwell, C., Mikolajewicz, U., Ziemann, F., and Kapsch, M.-L.: Sensitivity of Heinrich-type ice-sheet surge characteristics to boundary forcing perturbations, *Clim. Past*, 19, 179–198, <https://doi.org/10.5194/cp-19-179-2023>, 2023.
- Schannwell, C., Mikolajewicz, U., Kapsch, M.-L., and Ziemann, F.: A mechanism for reconciling the synchronisation of Heinrich events and Dansgaard-Oeschger cycles, *Nat. Commun.*, 15, 2961, <https://doi.org/10.1038/s41467-024-47141-7>, 2024.
- Schoof, C.: Ice sheet grounding line dynamics: steady states, stability, and hysteresis, *J. Geophys. Res.-Earth*, 112, F3, <https://doi.org/10.1029/2006JF000664>, 2007.
- Shaffer, G., Olsen, S. M., and Bjerrum, C. J.: Ocean subsurface warming as a mechanism for coupling Dansgaard-Oeschger climate cycles and ice-rafting events, *Geophys. Res. Lett.*, 31, 24, <https://doi.org/10.1029/2004GL020968>, 2004.
- Shapiro, N. M. and Ritzwoller, M. H.: Inferring surface heat flux distributions guided by a global seismic model: particular application to Antarctica, *Earth Planet. Sc. Lett.*, 223, 213–224, <https://doi.org/10.1016/j.epsl.2004.04.011>, 2004.
- Siddall, M., Rohling, E. J., Thompson, W. G., and Waelbroeck, C.: Marine isotope stage 3 sea level fluctuations: data synthesis and new outlook, *Rev. Geophys.*, 46, 4, <https://doi.org/10.1029/2007RG000226>, 2008.
- Sun, S., Pattyn, F., Simon, E. G., Albrecht, T., Cornford, S., Calov, R., Dumas, C., Gillet-Chaulet, F., Goelzer, H., Golledge, N. R., Greve, R., Hoffman, M. J., Humbert, A., Kazmierczak, E., Kleiner, T., Leguy, G. R., Lipscomb, W. H., Martin, D., Molighem, M., Nowicki, S., Pollard, D., Price, S., Quiquet, A., Seroussi, H., Schlemm, T., Sutter, J., van de Wal, R. S. W., Winkelmann, R., and Zhang, T.: Antarctic ice sheet response to sudden and sustained ice-shelf collapse (ABUMIP), *J. Glaciol.*, 66, 891–904, <https://doi.org/10.1017/jog.2020.67>, 2020.
- Swingedouw, D., Fichet, T., Huybrechts, P., Goosse, H., Driesschaert, E., and Loutre, M.-F.: Antarctic ice-sheet melting provides negative feedbacks on future climate warming, *Geophys. Res. Lett.*, 35, 17, <https://doi.org/10.1029/2008GL034410>, 2008.
- Swingedouw, D., Houssais, M.-N., Herbaut, C., Blaizot, A.-C., Devilliers, M., and Deshayes, J.: AMOC recent and future trends: a crucial role for oceanic resolution and Greenland melting?, *Frontiers in Climate*, 4, 838310, <https://doi.org/10.3389/fclim.2022.838310>, 2022.
- Tabone, I., Robinson, A., Alvarez-Solas, J., and Montoya, M.: Impact of millennial-scale oceanic variability on the Greenland ice-sheet evolution throughout the last glacial period, *Clim. Past*, 15, 593–609, <https://doi.org/10.5194/cp-15-593-2019>, 2019.
- Timm, O. and Timmermann, A.: Simulation of the last 21 000 years using accelerated transient boundary conditions, *J. Climate*, 20, 4377–4401, <https://doi.org/10.1175/JCLI4237.1>, 2007.
- Tsai, V. C., Stewart, A. L., and Thompson, A. F.: Marine ice-sheet profiles and stability under Coulomb basal conditions, *J. Glaciol.*, 61, 205–215, <https://doi.org/10.3189/2015JoG14J221>, 2015.
- Van Achter, G., Fichet, T., Goosse, H., Pelletier, C., Haubner, K., and Pattyn, F.: Ocean–ice sheet coupling in the Totten Glacier area, East Antarctica: analysis of the feedbacks and their response to a sudden ocean warming, *Geosciences*, 13, 106, <https://doi.org/10.3390/geosciences13040106>, 2023.
- Van Den Berg, J., van de Wal, R., and Oerlemans, H.: A mass balance model for the Eurasian Ice Sheet for the last 120,000 years, *Global Planet. Change*, 61, 194–208, 2008.
- Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J.-C., McManus, J. F., Lambeck, K., Balbon, E., and Labracherie, M.: Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records, *Quaternary Sci. Rev.*, 21, 295–305, [https://doi.org/10.1016/S0277-3791\(01\)00101-9](https://doi.org/10.1016/S0277-3791(01)00101-9), 2002.
- Waelbroeck, C., Pichat, S., Böhm, E., Loughheed, B. C., Faranda, D., Vrac, M., Missiaen, L., Vazquez Riveiros, N., Burckel, P., Lippold, J., Arz, H. W., Dokken, T., Thil, F., and Dapigny, A.: Relative timing of precipitation and ocean circulation changes in the western equatorial Atlantic over the last 45 kyr, *Clim. Past*, 14, 1315–1330, <https://doi.org/10.5194/cp-14-1315-2018>, 2018.
- Waelbroeck, C., Loughheed, B. C., Vazquez Riveiros, N., Missiaen, L., Pedro, J., Dokken, T., Hajdas, I., Wacker, L., Abbott, P., Dumoulin, J.-P., Thil, F., Eynaud, F., Rossignol, L., Fersi, P., Albuquerque, A. L., Arz, H., Austin, W. E. N., Came, R., Carlson, A. E., Collins, J. A., Dennielou, B., Desprat, S., Dickson, A., Elliot, M., Farmer, C., Giraudeau, J., Gottschalk, J., Henderiks, J., Huguen, K., Jung, S., Knutz, P., Lebreiro, S., Lund, D. C., Lynch-Stieglitz, J., Malaizé, B., Marchitto, T., Martínez-Méndez, G., Mollenhauer, G., Naughton, F., Nave, S., Nürnberg, D., Oppo, D., Peck, V., Peeters, F. J. C., Penaud, A., da Costa Portilho-Ramos, R., Repschläger, J., Roberts, J., Rühlemann, C., Salgueiro, E., Sanchez Goni, M. F., Schönfeld, J., Scussolini, P., Skinner, L. C., Skonieczny, C., Thornalley, D., Toucanne, S., Van Rooij, D., Vidal, L., Voelker, A. H. L., Wary, M., Syee, W., and Ziegler, M.: Consistently dated Atlantic sediment cores over the last 40 thousand years, *Scientific Data*, 6, 165, <https://doi.org/10.1038/s41597-019-0173-8>, 2019.
- Weertman, J.: On the sliding of glaciers, *J. Glaciol.*, 3, 33–38, <https://doi.org/10.3189/S0022143000024709>, 1957.
- Wekerle, C., McPherson, R., von Appen, W.-J., Wang, Q., Timmermann, R., Scholz, P., Danilov, S., Shu, Q., and Kanzow, T.: Atlantic Water warming increases melt below Northeast Greenland's last floating ice tongue, *Nat. Commun.*, 15, 1336, <https://doi.org/10.1038/s41467-024-45650-z>, 2024.