



Supplement of

Continental shelf glaciations off Northeast Greenland since the Late Miocene

Frank W. Jakobsen et al.

Correspondence to: Frank W. Jakobsen (frank.jakobsen@uit.no)

The copyright of individual parts of the supplement might differ from the article licence.




Figure S1: a) seismic profile KAN94-3 tied with Kanumas site 13. b) Core recovery (black). c) Lithological description by T. Lærdal modified from StatoilHydro (2008). M=mud, z=silt, S=sand. Black triangles show location of core photographs in (e). d) Borehole correlated to respective seismic mega units (this study). e-i) A selection of core photographs modified from StatoilHydro (2008). Seismic courtesy of GEUS.




Figure S3: Composite seismic profile. See Fig. 1 for location. a) composite seismic profile from Norske Trough to Greenland Basin. Tied to Kanumas site 13 in Norske Trough and ODP Site 913 in Greenland Basin. b) Composite seismic profile with interpreted regional reflections NEG-1-NEG-3 (this study). Seismic courtesy of AWI, TGS and GEUS.




Figure S4: Isochron thickness maps in two-way travel time (TWT; units in thickness maps are roughly equal to metres assuming an average subsurface velocity of 2 km/s), shown with linear scales with contours every 250 ms (TWT). Note the different scale for the individual maps. a) Total thickness measured from NEG-1 to seafloor. b) Thickness map of MU-1, measured from NEG-1 to NEG-2. c) Thickness map of MU-2, measured from NEG-1+NEG-2 to NEG-3. d) Thickness map of MU-3, measured from NEG-3 to the seafloor. MU=Mega Unit (-1, 2, 3).