Clim. Past, 21, 941–955, 2025 https://doi.org/10.5194/cp-21-941-2025 © Author(s) 2025. This work is distributed under the Creative Commons Attribution 4.0 License.





Magnetic properties and geochemistry of loess–paleosol sequences at the Nowdeh section in northeastern Iran

Vahid Feizi¹, Ghasem Azizi¹, Maryam Mollashahi², and Habib Alimohammadian³

¹Department of Physical Geography, Faculty of Geography, University of Tehran, Tehran, Iran ²Department of Forestry in Arid regions, Faculty of Desert Study, Semnan University, Semnan, Iran ³Environment Magnetic Laboratory, Department of Geology and Mineral Exploration, Tehran, Iran

Correspondence: Vahid Feizi (vahid.feizi62@gmail.com)

Received: 5 July 2023 – Discussion started: 25 July 2023 Revised: 16 February 2025 – Accepted: 5 March 2025 – Published: 28 May 2025

Abstract. The loess-paleosol sequences in the northeastern part of Iran serve as a high-resolution natural archive documenting climate and environmental changes. These sequences offer evidence of the interaction between the accumulation and erosion of aeolian and fluvial sediments during the Middle and Late Pleistocene periods. In this study, the Azadshahr (Nowdeh loess section) site was chosen to reconstruct Late Quaternary climate shifts. The 24 m thick Nowdeh loess-paleosol sequence was sampled for magnetic and geochemical analysis. The sampling involved 237 samples taken systematically at high resolution (10 cm intervals, selected samples, corresponding to peaks in magnetic susceptibility, underwent geochemical analysis to aid in the interpretation of paleoclimatic changes indicated by the magnetic signals). The magnetic susceptibility results of the loess-paleosol deposits revealed low values during cold and dry climate periods (loess) and high values during warm and humid climate periods (paleosol). The magnetic susceptibility at a depth of 22.1 m (approximately 130 ka) has significantly decreased, suggesting cold climate conditions at this time. The most substantial changes in magnetic susceptibility occur at depths between 18.6 and 21.3 m (approximately 100-120 ka). During this period, there are four phases of decrease (indicating cold and dry conditions) interspersed with three phases of increase (signifying warm and humid conditions) in magnetic susceptibility. The comparison of magnetic and geochemical data showed that variations in geochemical weathering ratios corresponded to changes in magnetic parameters. A high level of correlation was observed between the magnetic susceptibility intensity and ratios such as Rb / Sr, Mn / Ti, Zr / Ti, and Mn / Sr. The findings from

this research indicate that the sedimentary section of Nowdeh has experienced six distinct climate periods over the last 160 kyr. Notably, three cold and dry periods occurred between three warm and humid periods. Additionally, during these climate phases, short-term cold (stadial) and warm (interstadial) intervals were also observed.

1 Introduction

Reconstruction of the Quaternary climate is important for the development of climate models that lead to a better understanding of past and present and prediction of future climate development. Loess–paleosol sequences are now recognised as one of the most complete terrestrial archives of glacial–interglacial climate change (Porter and An, 195; Muhs and Bettis, 2003; Pierce et al., 2011; Guo et al., 2002) and have been used to reconstruct climate and geomorphological changes during the Quaternary (Karimi et al., 2011; Frechen et al., 2003; Prins et al., 2007).

Loess deposits occur in large areas of the northeast, east central, north, and central parts of Iran, which is part of the loess belt that covers the Middle East and extends further northward into Turkmenistan, Kazakhstan, and Tajikistan (Okhravi and Amini, 2001). The extensive and thick loess deposits in northern Iran have been recently studied in detail, establishing a more reliable chronological framework for the last interglacial–glacial cycle (Lateef, 1988; Pashaei, 1996; Kehl et al., 2006; Frechen et al., 2009; Karimi et al., 2009, 2013; Okhravi and Amini, 2001; Mehdipour, 2012). Paleoclimate studies of loess deposits based on rock magnetism and combined analyses of rock magnetism and geochemistry around the world have attained appreciable advances in the past few decades (Bader et al., 2024; Jordanova and Jordanova, 2024; Heller and Liu, 1984; Forster et al., 1996; Ding et al., 2002; Guo et al., 2002; Chlachula and Little, 2011; Bronger, 2003; Baumgart et al., 2013; Guanhua et al., 2014). These studies comprise loess–paleosol records that cover loess plateaus in China, Germany, Poland, Tajikistan, Austria, Ukraine, and the Danube catchment (Hosek et al., 2015; Ahmad and Chandra, 2013; Chen et al., 2010; Jordanova et al., 2011; Buggle et al., 2009; Fitzsimmons et al., 2012; Fischer et al., 2012; Jary and Ciszek, 2013; Baumgart et al., 2013; Schatz et al., 2014; Gocke et al., 2014).

Despite its suitable geographical location, there is only a limited number of studies of loess deposits from the north of Iran. In this work we explore the potential of loess deposits in northern Iran for reconstructing Late Quaternary climate and environmental change.

2 Study area

The Nowdeh section is exposed at about 20 km southeast of Gonbad-e Kavus and east of the city of Azadshahr. The Nowdeh river dissects a more than 24 m thick sequence of yellowish brown (10 YR 5/4) loess covering northeast dipping weathered limestone.

The study area $(37^{\circ}05'50'' \text{ N}, 55^{\circ}12'58'' \text{ E})$ is part of the Alborz structure, and this structure continues beneath the Caspian Sea. This zone includes regions north of the Alborz fault and south of the Caspian Sea. Toward the east, the Gorgan-Rasht zone is covered with thick layers of loess.

The Nowdeh section was selected for this work due to earlier soil studies by Kehl et al. (2005) and Frechen et al. (2009), combined with the existence of 12 dates for this section (Fig. 1).

3 Methodology

The Nowdeh loess section is approximately 24 m thick and was sampled at 10 cm intervals for magnetometry and geochemical analysis. The sampling location and method were determined following a detailed study of the area. Magnetic susceptibility measurements of all samples were conducted at the Environmental and Paleomagnetic Laboratory of the Geological Survey of Iran in Tehran. Magnetic susceptibility is indicative of the collective response of diamagnetic, paramagnetic, ferrimagnetic, and imperfect antiferromagnetic minerals present in the samples. Each sample was placed in a 11 cm³ plastic cylinder for use in magnetic measurement devices. The measurement of magnetic susceptibility was performed using the AGICO Kappabridge model MFK1-A. To ensure the reproducibility of our results, we have meticulously documented all experimental procedures, including the setup, equipment used, and analytical methods. Our findings have been validated by testing multiple independent samples and conducting experiments repeatedly under controlled conditions.

The determination of the saturation isothermal remanent magnetisation (SIRM) was carried out to assess the concentration of ferromagnetic and imperfect antiferromagnetic minerals in the samples. The calculation of the hard isothermal remanence (HIRM) magnetisation was performed to identify magnetically significant components such as hematite in the samples using the following formula:

HIRM = 0.5(SIRM + IRM - 0.3T),

where IRM-0.3T is the remanence after application of a reversed field of 0.3 *T* after growth and measurement of SIRM. The HIRM reflects the contribution specifically of the imperfect antiferromagnetic minerals hematite and goethite (Bloemendal et al., 2008).

The S - 0.3T value, or S ratio, is calculated as

S - 0.3T = 0.5[(-IRM - 0.3T/SIRM) + 1],

and it ranges from 0% to 100%. It reflects the ratio of ferrimagnetic to imperfect antiferromagnetic minerals (Bloemendal et al., 2008).

Based on the magnetic susceptibility results, 70 samples were selected for geochemical analyses (trace elements) to assist the paleoclimatic interpretation of the magnetic signals. Each sample was washed using a sieve with a mesh size of 400 µm and then dried in an oven. Once dried, the samples were further sieved with a 325 µm mesh sieve. The very fine sediments were collected, packed, and labelled as the tested material in special containers. A 0.2 g portion of the powder from each sample was then placed in a 1 M hydrochloric acid solution. After 2 h, the samples were analysed using an ICP device in the laboratory. The concentrations of the main elements were measured as a percentage, while the minor elements were quantified in milligrams per kilogram. To ensure the reproducibility of the results, we meticulously document all experimental steps, including the setup, equipment used, and analysis methods. The findings of this research have been validated through the use of multiple independent samples and by conducting experiments under controlled conditions.

As explained, the studied area was previously studied by Frechen et al. (2009) and Kehl et al. (2005). Therefore, we chose this sedimentary section to investigate climate changes and used their dating data. The infrared stimulated luminescence (IRSL) technique is utilised for this dating. A total of 45 samples were taken in light-tight tubes for the IRSL dating study. About 250 g of sediment was sampled. Polymineral fine-grain material (4–11 mm) was prepared for the measurements. The sediment material brought on discs was irradiated by a 90 Sr– 90 Y source in at least seven dose steps, with five discs each and a radiation dose of up to 750 Gy. All discs were stored at room temperature for at least 4 weeks after

V. Feizi et al.: Magnetic properties and geochemistry of loess-paleosol sequences



Figure 1. Map of Iran and the location of Nowdeh loess-paleosol sequence.



Figure 2. A view of the sedimentary section of the Nowdeh sequence with its very clear layering.

irradiation. The irradiated samples were preheated for 1 min at 230 °C. The values were obtained by integrating the 1-10 s region of the IRSL decay curves. An exponential growth

curve was fitted to the data and compared with the natural luminescence signal to estimate the De value. Alpha efficiency was estimated to 0.08 ± 0.02 for all samples. Dose rates were calculated from potassium, uranium, and thorium contents, as measured by gamma spectrometry (germanium detector) in the laboratory, assuming radioactive equilibrium for the decay chains. The IRSL ages gradually increase with depth from 20.5 ± 2.0 to 103 ± 10 ka. The stratigraphically oldest sample was collected below the lowermost exposed strongly developed paleosol (PC2) at a depth of 16.10 m below surface. The loess covering the uppermost strong paleosol (PC1), about 10.50 m thick, likely accumulated between about 61.9 ± 6.7 and 20 ± 2.0 ka (Frechen et al., 2009).

4 Results: magnetic properties

In Fig. 3, the relationship between susceptibility, NRM (natural remanent magnetisation), SIRM, HIRM, and S - 0.3Tin the Nowdeh section is illustrated. The variability in the magnetic susceptibility signal within the Nowdeh section indicates fluctuations in climate conditions and associated mechanisms during the Late Quaternary period. The values of magnetic susceptibility (χ) in the Nowdeh section range from 28.17 to 203.13 (in units of $10^{-8} \text{ m}^3 \text{ kg}^{-1}$). The maximum χ values (203.13) are found in the lower paleosol layer at 19.4 m depth, while the minimum values are observed in the uppermost loess layer at 7.4 m depth. The rock magnetic records exhibit a strong correlation with the lithology observed in the Nowdeh section. Generally, the paleosol layers exhibit higher magnetic signal intensities compared to the loess layers.

In Fig. 3, the paleosols exhibit higher magnetic susceptibility (χ) values compared to the loesses, with magnified magnetic enhancement observed in the Bw, Bt, and Btk horizons, while the underlying C (loess) horizon displays lower χ values. This difference probably reflects precipitation of iron oxides in the Bw horizons, resulting in a higher concentration of pedogenetic magnetite in comparison to the C horizons (Jordanova et al., 2013; Hosek et al., 2015). As illustrated in Fig. 3, the χ values in the lower and middle sections of the Nowdeh profile, approximately 53–80 and 120–140 kyr (at depths of 9–15 m and 18–23 m, respectively), represent intermediate values between unweathered loesses and weathered paleosols.

The results indicate that the natural remanent magnetisation (NRM) is consistent with the variance in magnetic susceptibility, particularly notable at lower depths, with the highest recorded value of this parameter observed at 13.1 m depth in the Bw/BwK horizon (Fig. 3). Variations and discrepancies in magnetic susceptibility align closely with the SIRM values of the loess sequence. As magnetic susceptibility decreases, SIRM also shows a corresponding decrease. In the interval between 20 and 50 ka (depth 2.1 to 8.4 m), during which much of the upper Loess formation occurred, magnetic susceptibility shows minimal variation, a pattern mirrored in the SIRM diagram for this period. The elevated HIRM values in Fig. 3 suggest an increase in the concentration and frequency of magnetic deterrent minerals such as goethite, maghemite, or hematite.

In Fig. 3, the comparison between the lower values of saturation (*S*) (-0.3T) (between 0.6 to 0.12 Am m^{-1}) and the higher values of hard isothermal remanent magnetisation (HIRM) (between 2 to 5 Am m^{-1}) indicates that the proportion of minerals with lower saturation, such as magnetite, is significantly lower than the proportion of minerals with higher saturation in paleosols. This pattern contrasts with the composition of loess deposits. As illustrated in Fig. 3, this can be clearly seen in loess sediments, for example, at a depth of 1 to 9 m, representing the time interval between 18 and 52 ka.

5 Element stratigraphy

Figure 4 illustrates the correlation between the concentration of selected elements (Sr, Rb, Zr, Ti, and Mn) and magnetite susceptibility in the Nowdeh section. The figure indicates significant variations in the concentration of these elements with noticeable differences between them. Sr and Rb exhibit similar trends along the Nowdeh section. At a depth of 2.9 m, there is a notable increase in the concentration of these two elements, corresponding to an age of 22 ka. Higher in the section, the concentration of Sr and Rb decreases.

In Fig. 4, at a depth of 18 m, the Nowdeh sedimentary section recorded the highest concentrations of elements, such as Sr, Rb, Zr, Ti, and Mn. Conversely, the lowest concentrations of these elements were observed at a depth of 8.5 m, which dates back to approximately 48 ka.

In Fig. 4, Ti, Zr, and Mn exhibit approximately similar trends in the diagram. These elements show little variation in concentration at the beginning of the section. The changes in element concentrations from the end of the sedimentary section down to a depth of 16.7 m (approximately 90 ka) display a zigzag pattern. Between the depths of 16.7 and 9.3 m, the fluctuations in element concentrations are minimal. At a depth of 9.3 m, corresponding to roughly to 52 ka, the research indicates the lowest concentrations of the elements measured. However, from this point onward, the concentration of elements begins to rise, peaking at a depth of 8.5 m, which dates back to about 48 ka. This increase suggests a period of hot and humid conditions during that time.

6 Trace element ratio

The variation of the Si / Ti ratio in Fig. 5 generally follows the magnetic susceptibility pattern (Fig. 5), except for the lower part of the section (23-24 m). The ratios of Mn / Sr, Zr / Ti, and Mn / Ti in Fig. 5 show almost no long-term change, except for at a depth of 8.5 m, corresponding to an age of 48.8 ka. These changes suggest hot and humid climatic conditions, which can be correlated with the high level of magnetic susceptibility. The Rb / Sr ratio exhibits an opposite pattern to the magnetic susceptibility, especially at the depths of 8.5, 16, 19, and 22 m. The Ba / Rb ratio generally follows the magnetic susceptibility pattern, except at depths of 13, 15, 19, and 22.8 m, where they vary oppositely.

The variation in the Si / Ti ratio does not exhibit a consistent relationship to the sequence of loess–paleosol layers, as defined by the magnetic susceptibility in the Nowdeh section. On the other hand, the Mn / Ti ratios tend to show elevated values in the paleosols, likely due to the concentration of Mn oxide in the finer sediment fraction (Bloemendal et al., 2008). This suggests that the presence of Mn oxide plays a significant role in influencing the Mn / Ti ratio in the sediments, particularly in the paleosol layers.

7 Discussion

Over the entire 159 kyr sequence at the Nowdeh site, there appears to be a reasonable first-order co-variation between the magnetic and geochemical indicators of weathering and



Figure 3. Basic magnetic parameters for the Nowdeh section. From left to right: a simplified lithological column, magnetic susceptibility (MS), natural remanent magnetisation (NRM), isothermal remanent magnetisation (SIRM), and *S* ratio (S - 0.3T) are plotted against depth (m). The lithological column indicates the following sedimentary units: CB/Cbk (calcareous brown soil/calcareous brown soil with krotovina), C (loess), Bw(t) (brown soil with clay illuviation), Bt/Btk (brown soil with clay illuviation/calcareous brown soil with clay illuviation), Bw/Bwk (brown soil/calcareous brown soil), and Ck/Ckm (calcareous horizon/calcareous horizon with soft masses of calcium carbonate).



Figure 4. Depth series of selected element concentrations for Nowdeh section.

soil formation, particularly with magnetic parameters reflecting variations in ferrimagnetic content and Sr-based ratios. However, upon closer detailed examination based on individual loess and paleosol layers, an inconsistent relationship is observed between the amplitudes of individual peaks and troughs of magnetic and geochemical parameters (Figs. 4 and 5). This suggests that while there is an overall correlation between these indicators at a broader scale, at a finer resolution within specific layers, the relationship becomes more complex and inconsistent. Additional factors or processes may be influencing the variations in magnetic and geochemical parameters within the individual stratigraphic units. This issue can be seen clearly in Figs. 4 and 5. As noted by Hosek et al. (2015) and Makeev et al. (2024), there is a significant relationship between magnetic receptivity, chemical elements, and climatic conditions. Our study reinforces this finding, as indicated by the results obtained. Recent studies have highlighted a significant relationship between climate change and the magnetic properties of sediments, indicating that alterations in sediment composition and depositional environments can reflect past climate conditions. For instance, Huang et al. (2022) demonstrated that variations in magnetic susceptibility in loess deposits are closely linked to fluctuations in moisture levels and temperature, underscoring the sensitivity of magnetic minerals to climatic changes. These findings suggest that analysing the magnetic properties of



Figure 5. Selected element ratios in the Nowdeh section.

sediments can provide valuable insights into historical climate dynamics and contribute to our understanding of how magnetic signatures may evolve in response to ongoing climate change. To investigate the relationship between climate change and the magnetic properties of sediments, we conducted magnetic susceptibility measurements on loess sediments from the Nowdeh section. The results of the magnetic susceptibility analysis demonstrated clear patterns that correlate with historical climatic fluctuations (Fig. 3). The observed distinct sequences in magnetic susceptibility reveal insights into past environmental conditions, with lower values indicating cold and dry periods typical of loess deposition, while higher values correspond to warmer and more humid phases associated with paleosol development. These correlations between magnetic susceptibility and alternating loess-paleosol sequences provide a compelling illustration of how sediment magnetic properties reflect climatic changes over time in the Nowdeh region. The implications of these findings are significant, as they suggest that magnetic susceptibility can serve as a reliable proxy for reconstructing paleoclimatic conditions. By establishing this relationship, we enhance our understanding of how the sedimentary environment responded to climate shifts during the Pleistocene. Furthermore, these patterns of magnetic susceptibility not only inform us about specific climatic conditions but also help elucidate the feedback mechanisms between climate and weathering processes in the region. In essence, our analysis underscores the potential of magnetic susceptibility as a key indicator of past climate change, linking geological records with broader climatic events. This enhanced understanding is crucial for interpreting how similar climatic fluctuations may affect sediment dynamics in other regions with comparable loess deposits.

According to Song et al. (2008), sediment loess is typically formed under cold and dry climate conditions, leading to lower magnetic susceptibility values due to minimal weathering processes. This observation aligns with the find-

ings in the Nowdeh section, where our data indicate that the loess layers are characterised by significantly lower magnetic susceptibility than the overlying paleosols (Fig. 3). In the paleosols of the Nowdeh section, formed through extensive pedogenic processes, we observe elevated magnetic susceptibility attributed to increased levels of oxidation and enhanced concentrations of key elements such as iron and aluminium oxides (Fig. 3). Our magnetic susceptibility data corroborate the assertion by Song et al. (2008) that paleosols generally exhibit higher magnetic susceptibility compared to adjacent loess layers. Specifically, our measurements indicate a distinct rise in magnetic susceptibility values within these paleosols, further supporting the notion that the formation of strong magnetic minerals, such as iron oxides (Fe₃O₄, γ - Fe_2O_3 , and Fe_2O_3), occurs through pedogenesis. Moreover, the composition of magnetic minerals in our loess samples is influenced by the grain composition from the aeolian sources of sedimentation. This distinction is crucial, as our data illustrate a clear contrast in magnetic mineral content between the well-developed paleosols and the less altered loess layers. In our analysis, we found that the paleosols not only have higher magnetic susceptibility but also exhibit a more diverse range of magnetic mineral types, indicating a more complex pedogenic history. This relationship highlights the importance of understanding the processes of magnetic mineral formation and alteration in determining the magnetic susceptibility profiles of sediment sequences. Our own findings reinforce the established understanding from previous studies, such as those by Song et al. (2008), while providing new insights into the specific conditions and processes at play in the Nowdeh section.

In Fig. 3, the brown layer sequences of dark and light paleosols within the loess deposits illustrate distinct weathering processes that reflect the climatic patterns observed during glacial and interglacial periods of the Middle and Late Pleistocene. The higher magnetic susceptibility of the paleosols compared to the surrounding loess layers indicates a significant degree of pedogenesis and oxidation, consistent with findings by Maher (2011) and Spassov (2002). This difference is particularly pronounced at lower depths, suggesting that these layers experienced greater weathering variability during those periods. At a depth of 21 m (approximately 110 ka), the notable decrease in magnetic susceptibility suggests a transition to colder and drier conditions, which aligns with the known climatic shifts of that era. This finding is crucial because it highlights how magnetic susceptibility can serve as a proxy for past climatic conditions, providing insights into the environmental changes that influenced soil development (Thompson and Oldfield, 1986; Liu et al., 2023). The magnetic susceptibility chart for the Nowdeh section reveals around eight distinct periods of increasing magnetic susceptibility, indicative of elevated temperatures and humidity, consistent with findings from other regions that link magnetic properties to climatic variations (Heller and Liu, 1982). This pattern suggests a correlation between magnetic susceptibility and climatic conditions, where periods of increased susceptibility correspond to warmer, more humid phases conducive to soil formation.

In accordance with the standard global loess characteristics, paleosols consistently exhibit higher magnetic susceptibility values compared to adjacent loess layers due to pedogenesis and oxidation processes, as highlighted by Maher (2011) and Spassov (2002). The NRM results suggest a decrease during loess formation and an increase during paleosol formation (Fig. 3). This pattern suggests a relationship between NRM and magnetic susceptibility (Fig. 3) (Bloemendal et al., 2008). A decrease in NRM indicates dry and cold climate conditions in Fig. 3 (at the depth of 7.2 m, which is approximately equal to 34 ka), consistent with the deposition of loess layers, while an increase in NRM represents warmer and more humid climate conditions, corresponding to paleosol formation. The results presented in the previous section illuminate the significant relationship between NRM (natural remanent magnetisation) and magnetic susceptibility, contributing to our understanding of past climatic conditions. The observed decrease in NRM at a depth of 7.2 m (approximately 34 ka) is indicative of drier and colder climate conditions, consistent with the processes associated with loess deposition. This finding reinforces the notion that periods of extensive loess accumulation correspond with colder climatic phases, as seen in other studies of similar stratigraphic sequences. Conversely, the increase in NRM at depths of 18.6 to 21.3 m marks a shift toward warmer and more humid conditions, which correlates with paleosol formation. This transitional phase highlights the dynamic interplay between climate and soil development, suggesting that optimum conditions for soil formation fostered the development of paleosols during this time. The highest magnetic susceptibility values in this layer, documented in Fig. 3, further corroborate the enhancing environmental conditions during this interval. The peak alignment of NRM and magnetic susceptibility at a depth of 19.4 m, approximately 120 ka, is particularly noteworthy. This correlation may signify a period of climatic stability that allowed for the establishment of rich soil profiles, crucial for understanding the ecological dynamics at play during the Late Pleistocene. Such findings align with previous research by Bloemendal et al. (2008), which also emphasised the relevance of magnetic properties in interpreting paleoclimatic conditions.

The probable justifications for the low alteration in magnetic susceptibility and isothermal remnant magnetisation between 20 and 50 ka (Fig. 3) can be attributed to two main factors:

- 1. decreased pedogenesis due to cold and dry periods
- 2. reduction in the influx of magnetic particles into loess layers.

During the last 20 kyr, there seems to have been a correlation between magnetic susceptibility variations in the surface soil layer and climatic conditions. This period coincides with the transition from a cold climate to the current warm and humid climate in the northern region of Iran (Frechen et al., 2009). As a result, the soil's magnetic properties, specifically SIRM, have likely increased during this time frame. However, since the SIRM samples were only collected at magnetic susceptibility peak points, they may not capture the full extent of variations. Comparing these findings with the research by Antoine et al. (2013) on loess-paleosol sediments in central Europe reveals a close relationship, particularly around 32 ka. The close relationship observed in our findings and those of Antoine et al. (2013) around 32 ka suggests a synchronous response of the magnetic susceptibility of sediments to environmental changes during this period. This synchrony reinforces the idea that widespread climatic or geological factors were influencing sediment formation across regions. Also, Antoine et al. (2013) provided critical insights into the paleoenvironmental conditions in central Europe during the Late Pleistocene. Geochemical charts can serve as useful indicators of climate patterns, as they can highlight different levels of weathering severity. In the study of loess deposits, certain chemical ratios can be utilised to reconstruct variations in paleoclimate (Ding et al., 2001). The Zr / Ti, Mn / Ti, Rb / Sr, and Mn / Sr records from the Nowdeh section exhibit a clear pattern of higher values prevailing in the paleosols, and their high degree of similarity is noteworthy. Rb / Sr has been suggested by several researchers as an indicator of pedogenic intensity in loess, based on the differential weathering of the major host minerals, specifically K-feldspar for Rb and carbonates for Sr (Hosek et al., 2015; Makeev et al., 2024). In the case of Mn / Sr, the higher values observed in the paleosols are likely a result of the combined effects of grain size on Mn concentration, as well as the loss of Sr through solution processes. This indicates that these ratios can serve as important indicators of pedogenic processes and weathering dynamics in the sedimentary record of the Nowdeh section. Rubidium is derived from K-feldspar, while strontium comes from carbonate minerals. As soils weather, an increase in Rb / Sr often indicates selective retention of Rb while Sr is leached away, reflecting greater weathering intensity (Yu et al., 2023). Climate-driven changes, such as increased precipitation, can enhance Sr leaching, leading to higher Rb / Sr ratios in wetter conditions. This relationship highlights how the climate influences elemental cycling in soils (Zhang et al., 2021b).

The magnetic susceptibility record shows high values at a depth of 19.4 m (Fig. 3), indicating hot and humid climate conditions prevailing around 120 ka. Variations in the concentrations of manganese (Mn), zirconium (Zr), and titanium (Ti) in the soil reflect a clear stratigraphic pattern (Fig. 4), with higher values seen in paleosols and lower values in the loess layers (Bloemendal et al., 2008). This pattern is influenced, in part, by carbonate dilution/concentration effects, as a significant portion of the variability in these elements disappears when expressed on a carbonate-corrected basis. In a study by Chen et al. (1999), a comparison was made between the Rb / Sr ratios and magnetic susceptibility values in the uppermost (last glacial/interglacial) sections of the Luochuan and Huanxian regions. The researchers noted a remarkable correspondence between the amplitudes of variation in magnetic susceptibility and Rb / Sr ratios. This finding suggests a close relationship between magnetic susceptibility variations and the Rb / Sr ratios in these regions during the last glacial and interglacial periods.

In the Nowdeh section, the amount of rubidium (Rb) in paleosols was lower compared to its concentration in loess layers (Fig. 5). This discrepancy can be attributed to the higher solubility of Rb in warm and humid climates, typical of interglacial periods. Gallet et al. (1996) observed significant depletion of Rb in the paleosols, supporting this interpretation. Recent research by Zhu and Li (2021) demonstrated that climate change alters the biogeochemical cycling of nutrients, including Rb, leading to increased leaching in saturated soils during periods of heavy rainfall. Furthermore, a comprehensive review by Jiang et al. (2020) emphasised the impacts of climate change on soil properties, particularly focusing on leaching processes and the resultant changes in nutrient concentration due to increased precipitation and temperature.

Our results indicate that the Mn / Ti, Zr / Ti, and Mn / Sr ratios tend to exhibit higher values in the paleosols (Fig. 4). According to Ding et al. (2001), elevated Mn / Ti values in paleosols may result from the concentration of iron (Fe) and manganese (Mn) oxides in the finer sediment fractions. They also noted that the Rb / Sr and Mn / Sr ratios show a clear pattern of elevation in the paleosols, which aligns with the findings of our study (Fig. 5). The Rb / Sr ratio has been proposed by various researchers as an indicator of pedogenic intensity in loess deposits, based on the differential weathering of major host minerals such as K-feldspar for Rb and carbonates for Sr. The higher Mn / Sr values in paleosols may be attributed to grain-size effects on Mn concentrations and the solubilisation loss of Sr.

Chen et al. (1999) compared Rb / Sr and magnetic susceptibility in the uppermost parts of the Luochuan and Huanxian sections, revealing a significant correspondence between the variations in magnetic susceptibility and Rb / Sr ratios. This suggests a link between weathering intensity and magnetic properties in these sediments. In the context of the Nowdeh sedimentary section, the magnetic parameters were compared with those from other studies conducted in various regions of the world, further contributing to our understanding of paleoclimatic variations and weathering processes in loess deposits.

In Fig. 6 the comparison of magnetic receptivity results from the Nowdeh sedimentary section with the palynological data from sedimentary cores of Urmia Lake (Djamali et al., 2008) and the ¹⁸ δ analysis from Arabian Sea sedimentary cores (Tzedakis, 1994) have provided valuable insights into past climate conditions (Fig. 6). In the analysis, an increase in the AP/NAP ratio (number of arboreal pollen grains to the number of non-arboreal pollen grains) in the lakes corresponded with the presence of ancient soil layers in the seedling sedimentary section.

This increase signifies warmer temperatures and higher humidity levels, conducive to the growth of trees and shrubs (Harrison et al., 2020; Zhang et al., 2021a). Conversely, a decrease in the AP/NAP index indicates a decline in temperature and humidity, leading to the disappearance of trees and shrubs and changes in surface vegetation cover (Zhang et al., 2016). This correlation suggests that the climate conditions and their fluctuations in western Iran align with the sedimentary deposition at Nowdeh, consistent with findings from similar studies in the region (Fuchs et al., 2013; Hosek et al., 2015).

Moreover, in Fig. 6 the ¹⁸ δ analysis of the Arabian Sea exhibited a strong agreement with magnetic receptivity data. A decrease in the ¹⁸ δ indexes points to warmer climate conditions, while an increase indicates colder conditions (Djamali et al., 2008). The relationship between magnetic susceptibility and ¹⁸ δ levels in the Arabian Sea sediments, as shown in Fig. 6, verifies that an increase in magnetic susceptibility corresponds with a decrease in ¹⁸ δ levels, indicating warmer climate conditions. This alignment further supports the connection between the recorded palynology data of Lake Urmia, ¹⁸ δ data from the Arabian Sea, and the sequence of ancient loess-soil sediments in the Nowdeh sedimentary section.

The results of our current research demonstrate a significant correlation with the findings of Fuchs et al. (2013) and Hosek et al. (2015), which examined ancient loess and paleosol deposits in central Europe, indicating that similar magnetic properties can be observed across different regions in response to climate change. Figure 8 depicts consistent patterns in the magnetic receptivity parameter at 45, 73, 90, 104, and 108 ka across the study sections. Around 45 and 73 ka, there is a clear increasing trend in magnetic receptivity observed in all analysed layers, indicating a shift towards warmer and more humid climate conditions compared to ear-



Figure 6. Correlation between recorded palynological data of Lake Urmia (Djamali et al., 2008) and $^{18}\delta$ of Arabian Sea sediments (Tzedakis, 1994) with the loess–paleosol sediment sequence of the Nowdeh sedimentary section.

lier periods. This increase in magnetic susceptibility can be attributed to the higher presence of iron oxides in the soil resulting from increased chemical weathering. Conversely, during the periods of 90, 104, and 108 ka, a decrease in magnetic susceptibility is evident across all regions, signifying colder and drier climatic conditions during these time intervals. This issue can also be seen in the amount of Fe_2O_3 in Nowdeh sediments (Fig. 7). While the older sediments also show a significant association with climate variations in central Europe and the Nowdeh area, the absence of radiometric dating in these older sediments introduces some uncertainty when interpreting these findings. Nonetheless, the consistent patterns in magnetic susceptibility across different time periods provide valuable insights into past climate fluctuations and their impact on soil properties in these regions.

The comparison of magnetic receptivity trends as recorded in sedimentary sections of Beiyuan, Heimugou, Biampo, and the ¹⁸ δ records by Imbrie et al. (1984) in Fig. 9 reveals a high agreement with the Nowdeh sedimentary section. This alignment indicates similar climate conditions across different locations in the Northern Hemisphere.

The consistency in magnetic receptivity trends among these various sites suggests a commonality in the climatic conditions experienced during the corresponding time periods. This synchronisation in magnetic susceptibility patterns further supports the notion that these regions were subjected to comparable environmental changes and fluctuations in the past.

Additionally, the correlation observed between the magnetic receptivity data and the ¹⁸ δ records underscores the close relationship between climatic factors and sedimentary deposition patterns across these sites (Fig. 9). By examining these geological proxies, researchers can gain valuable insights into the past climate dynamics and variations that have affected the Northern Hemisphere over time.

The findings of Mehdipour (2012) in the realm of fine loess exhibit a close resemblance to the results presented in our research, as illustrated in Fig. 9. In their study, they employed both magnetic and geochemical approaches to assess different climatic periods, and the outcomes align significantly with the findings of our research. The comparison in Fig. 10 reveals a strong consistency in the magnetic receptivity trends between the Nowdeh section and the Neka sedimentary section analysed by Mehdipour (2012). Between 48 and 20 kyr ago, notable similarities are observed in the fluctuations of magnetic receptivity in both sedimentary sections. Whenever there is an increase in magnetic receptivity, it indicates a warm and humid period with the formation of ancient soil layers. This shared pattern implies a synchrony in climatic conditions between the two regions during this time frame, showcasing the utility of magnetic susceptibility as a proxy for understanding past environmental changes and soil development processes.



Figure 7. The relationship between magnetic susceptibility and Fe₂O₃.



Figure 8. Comparison of changes in magnetic receptivity of the Dolní Věstonice sedimentary section (Fuchs et al., 2013) and the Dobšice and Zeměchy section (Hošek et al., 2015) with the Nowdehh sedimentary section.

The results of this research exhibit a strong consistency with the findings of Beer and Sturm (1995) regarding beryllium saturation in the Zaifang sedimentary section and $^{18}\delta$ in marine sediments. In both cases, there is a clear correlation between the fluctuations in beryllium saturation, ${}^{18}\delta$, and magnetic receptivity (Fig. 11).

When beryllium saturation and ${}^{18}\delta$ decrease, there is a corresponding decrease in magnetic receptivity, indicating



Figure 9. Comparison of magnetic receptivity changes of Beiyuan, Heimugou, and Biampo (An et al., 1991) and records of ${}^{18}\delta$ (Imbrie et al., 1984) with the Nowdeh sedimentary section.



Figure 10. Comparison of magnetic receptivity diagram of the Nowdeh sedimentary section with the Neka sedimentary section (Mehdipour, 2012).



Figure 11. Comparison of magnetic receptivity results of Nowdeh sedimentary section in comparison with ${}^{18}\delta$ and Be 10 isotope results of Xifeng sedimentary section (Beer and Sturm, 1995).

colder and drier climate conditions. Conversely, an increase in beryllium saturation and ${}^{18}\delta$ is accompanied by an increase in magnetic receptivity, signifying warmer and more humid periods.

The high agreement between the climatic periods identified based on these parameters in the Zaifang sedimentary section and marine sediments and the magnetic receptivity trends observed in the Nowdeh sedimentary section highlights the synchrony of similar climate events in the past across different locations. This consistency further supports the robustness of magnetic susceptibility as a proxy for understanding past climate variations and environmental changes (Fig. 11).

8 Conclusions

In conclusion, the loess–paleosol sequences from northeastern Iran serve as a valuable archive for studying the paleoenvironmental changes during the Upper Pleistocene. By employing a multi-proxy approach that integrates sedimentological, magnetic, and geochemical methods, the following key insights have been revealed:

- 1. The stratigraphy of the studied section aligns well with the typical pattern of Upper Pleistocene loess–paleosol successions in the region, providing valuable insights into the past environmental conditions.
- 2. Magnetic parameters show a strong correlation with climate conditions, making them effective variables for reconstructing climate change patterns in the region.
- 3. Comparisons between magnetic and geochemical data indicate that variations in geochemical weathering ratios mirror changes in magnetic weathering parameters, such as magnetic susceptibility, further enhancing our understanding of past environmental dynamics.
- 4. The high degree of coherence observed between the amplitudes of magnetic susceptibility and various geochemical ratios, including Rb / Sr, Mn / Ti, Zr / Ti, and Mn / Sr, reinforces the reliability of magnetic susceptibility as a proxy for tracking environmental changes and provides additional insights into the interplay between magnetic and geochemical processes.

Overall, this comprehensive multi-proxy analysis enhances our understanding of the paleoenvironmental changes in northeastern Iran during the Upper Pleistocene period and emphasises the importance of integrating sedimentological, magnetic, and geochemical data to unravel past climatic fluctuations and environmental dynamics.

Data availability. Access to these data can be granted under controlled conditions upon reasonable request, subject to ethical and legal considerations. Alternatively, anonymized or summarized versions of the data may be provided by the corresponding author to support validation and further research.

Author contributions. VF: conceptualisation, methodology, data analysis, writing (original draft), project. GA: writing (review and editing). MM: data collection, data analysis, literature review, formal analysis, visualisation. HA: data analysis, literature review, writing (review and editing).

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Special issue statement. This article is part of the special issue "Paleoclimate, from observing modern processes to reconstructing the past: a tribute to Dick (Dirk) Kroon". It is not associated with a conference.

Review statement. This paper was edited by Simon Jung and reviewed by Gerald Ganssen, Luigi Jovane, Erik Gulbranson, and one anonymous referee.

References

- Ahmad, I. and Chandra, R.: Geochemistry of loess-paleosol sediments of Kashmir Valley, India: Provenance and weathering, J. Asian Earth Sci., 66, 73–89, 2013.
- An, Z. S., Kukla, G., Porter, S. C., and Xiao, J. L.: Magnetic susceptibility evidence of monsoon variation on the Loess Plateau of Central China during the last 130,000 years, IEE International Geoscience and Remote Sensing Symposium, Boston, Massachusetts, II12227-30, 1991
- Antoine, P., Rousseau, D. D., Degeai, J. P., Moine, O., Lagroix, O., Kreutzer, S., Fuchs, M., Hatte, C. H., Gauthier, C., Svoboda, J., and Lisa, L.: High-resolution record of the environmental response to climatic variations during the Last Interglacial-Glacial cycle in Central Europe: the loess-palaeosol sequence of Dolní

Věstonice (Czech Republic), Quaternary Sci. Rev., 67, 17–38, 2013.

- Bader, N. E., Broze, E. A., Coates, M. A., Elliott, M. M., Mc-Gann, G. E., Strozyk, S., and Burmester, R. F.: The usefulness of the magnetic susceptibility of loess paleosol sequences for pa- 50 leoclimate and stratigraphic studies: The case of the Quaternary Palouse loess, northwestern United States, Quatern. Int., 702, 13–23, 2024.
- Baumgart, P., Hambach, U., Meszner, S., and Faust, D.: An environmental magnetic fingerprint of periglacial loess: records of Late Pleistocene loess paleosol sequences from Eastern Germany, Quatern. Int., 296, 82–93, 2013.
- Beer, J. and Sturm, M.: Dating of lake and loess sediments, Radiocarbon, 37, 81–86, 1995.
- Bloemendal, J., Liu, X., Sun, Y., and Li, N.: An assessment of magnetic and geochemical indicators of weathering and pedogenesis at two contrasting sites on the Chinese Loess Plateau, Palaeogeogr. Palaeocl., 257, 152–168, 2008.
- Bronger, A.: Correlation of loess-paleosol sequences in East and Central Asia with SE Central Europe: toward a continental Quaternary pedostratigraphy and paleoclimate history, Quatern. Int., 106/107, 11–31, 2003.
- Buggle, B., Hambach, U., Glaser, B., Gerasimenko, N., Markovic, S., Glaser, I., and Zöller, L.: Stratigraphy, and spatial and temporal paleoclimatic trends in Southeastern/Eastern European loess– paleosol sequences, Quatern. Int., 196, 186–206, 2009.
- Chen, J., An, Z. S., and Head, J.: Variation of Rb / Sr ratios in the loess–paleosol sequences of central China during the last 130,000 years and their implications for monsoon paleoclimatology, Quaternary Res., 51, 215–219, 1999.
- Ding, Z. L., Yang, S. L., Sun, J. M., and Liu, T. S.: Iron geochemistry of loess and Red Clay deposits in the Chinese Loess Plateau and implications for long-term Asian monsoon evolution in the last 7.0 Ma, Earth Planet. Sc. Lett., 185, 99–109, 2001.
- Ding, Z. L., Ranov, V., Yang, S. L., Finaev, A., Han, J. M., and Wang, G. A.: The loess record in southern Tajikistan and correlation with Chinese loess, Earth Planet. Sc. Lett., 200, 387–400, 2002.
- Djamali, M., de Beaulieu, J. L., Shah Hosseini, M., Andrieu-Ponel, V., Amini, A., Akhani, H., Leroy, S. A. G., Stevens, L., Alizadeh, H., Ponel, P., and Brewer, S.: A late Pleistocene long pollen record from Lake Urmia, NW Iran, Quaternary Res., 69, 413– 420, 2008.
- Fischer, P., Hilgers, A., Protze, J., Kels, H., Lehmkuhl, F., and Gerlach, R.: Formation and geochronology of Last Interglacial to Lower Weichselian loess/palaeosol sequences – case studies from the Lower Rhine Embayment, Germany, E&G Quaternary Sci. J., 61, 48–63, https://doi.org/10.3285/eg.61.1.04, 2012.
- Fitzsimmons, K. E., Marković, S. B., and Hambach, U.: Pleistocene environmental dynamics recorded in the loess of the middle and lower Danube Basin, Quaternary Sci. Rev., 41, 104–118, 2012.
- Forster, T., Evans, M. E., Havlíček, P., and Heller, F.: Loess in the Czech Republic: magnetic properties and paleoclimate, Stud. Geophys. Geod., 40, 243–261, 1996.
- Frechen, M., Oches, E. A., and Kohfeld, K. E.: Loess in Europe– mass accumulation rates during the Last Glacial Period, Quaternary Sci. Rev., 22, 1835–1875, 2003.

- Frechen, M., Kehl, M., Rolf, C., Sarvati, R., and Skowronek, A.: Loess Chronology of the Caspian Lowland in Northern Iran, Quatern. Int., 198, 220–233, 2009.
- Fuchs, M., Kreutzer, S., Rousseau, D.-D., Antoine, P., Hatte, C., Lagroix, F., Moine, O., Gauthier, C., Svoboda, J., and Lisa, L.: The loess sequence of Dolni Vestonice, Czech Republic: a new OSL based chronology of the Last Climatic Cycle, Boreas, 42, 664–677, 2013.
- Gallet, S., Jahn, B. M., and Torii, M.: Geochemical characterization of the Luochuan loess-paleosol sequence, China, and paleoclimatic implications, Chem. Geol., 133, 67–88, 1996.
- Gocke, M., Hambach, U., Eckmeier, E., Schwark, L., Zöller, L., Fuchs, M., Löscher, M., and Wiesenberg, G. L. B.: Introducing an improved multi-proxy approach for paleoenvironmental reconstruction of loess–paleosol archives applied on the Late Pleistocene Nussloch sequence (SW Germany), Palaeogeogr. Palaeocl., 410, 300–315, 2014.
- Guanhua, L., Dunsheng, X., Ming, J., Jia, J., Jiabo, L., Shuang, Z., and Yanglei, W.: Magnetic characteristics of loess-paleosol sequences in Tacheng, northwestern China, and their paleoenvironmental implications, Quaternary Int., 3, 1–10, 2014.
- Guo, Z. T., Ruddiman, W. F., Hao, Q. Z., Wu, H. B., Qiao, Y. S., Zhu, R. X., Peng, S. Z., Wei, J. J., Yuan, B. Y., and Liu, T. S.: Onset of Asian desertification by 22 Myr ago inferred from loess deposits in China, Nature, 416, 159–163, 2002.
- Harrison, S. P., Prentice, I. C., and Baird, A. J.: Vegetation and climate change: Implications for ecosystem services, Glob. Change Biol., 26, 429–445, 2020.
- Heller, F. and Liu, T. S.: Magnetostratigraphical dating of loess deposits in China, Nature, 300, 431–433, 1982.
- Heller, F. and Liu, T.: Magnetism of Chinese loess deposits, Geophys. J. Roy. Astr. S., 77, 125–141, 1984.
- Hosek, J., Hambach, U., Lisa, L., Grygar, T. M., Horacek, I., Meszner, S., and Knesl, I.: An integrated rock-magnetic and geochemical approach to loess/paleosol sequences from Bohemia and Moravia (Czech Republic): Implications for the Upper Pleistocene paleoenvironment in central Europe, Palaeogeogr. Palaeocl., 418, 344–358, 2015.
- Hošek, J., Hambach, U., Lisá, L., Matys, G. T., and Horáček, I.: An integrated rock-magnetic and geochemical approach to loess/paleosol sequences from Bohemia and Moravia (Czech Republic): Implications for the Upper Pleistocene paleoenvironment in central Europe, Palaeogeogr. Palaeocl., 418, 344–358, 2015.
- Huang, G., Liu, J., Zhang, L., and Zhang, X.: Magnetic susceptibility of loess deposits as a proxy for paleoenvironmental changes: Implications for climate variations in North China, J. Quaternary Sci., 37, 639–652, 2022.
- Imbrie, J., Hays, J. D., Martinson, D. G., McIntyre, A., Mix, A. C., Morley, J. J., Pisias, N. G., Prell, W. L., and Shackleton, N. J.: The orbital of Pleistocene climate: support from a revised chronology of the marine ¹⁸O record, in: Berger, Milankovitch and Climate, Reidel, Dordrecht, 269–305, 1984.
- Jary, Z. and Ciszek, D.: Late Pleistocene loess–palaeosol sequences in Poland and western Ukraine, Quatern. Int., 296, 37–50, 2013.
- Jiang, Y., Lavallee, J. M., and Stewart, C. E: Soil organic carbon dynamics in response to climate change: A review, Glob. Change Biol., 26, 1–16, 2020.
- Jordanova, D. and Jordanova, N.: Geochemical and mineral magnetic footprints of provenance, weathering, and pedogenesis of

loess and paleosols from North Bulgaria, Catena, 243, 108131, https://doi.org/10.1016/j.catena.2024.108131, 2024.

- Jordanova, D., Grygar, T., Jordanova, N., and Petrov, P.: Palaeoclimatic significance of hematite/goethite ratio in Bulgarian loess– palaeosol sediments deduced by DRS and rock magnetic measurements, in: The Earth's Magnetic Interior, edited by: Petrovsky, E., Ivers, D., Harinarayana, T., and Herrero-Bervera, E., IAGA Special Sopron Book Series, Springer-Verlag, Berlin, https://doi.org/10.1007/978-94-007-0323-0_26, 2011.
- Jordanova, D., Goddu, S. R., Kotsev, T., and Jordanova, N.: Industrial Contamination of Alluvial Soils near Fe–Pb Mining Site Revealed by Magnetic and Geochemical Studies, Geoderma, 192, 237–248, 2013.
- Karimi, A., Khademi, H., Kehl, M., and Jalalian, A.: Distribution, lithology, and provenance of peridesert loess deposits in Northeast Iran, Geoderma, 148, 241–250, 2009.
- Karimi, A., Khademi, H., and Jalalian, A.: Loess: Characterization and application for paleoclimate study, Geogr. Res., 76, 1–20, 2011.
- Karimi, A., Khademi, H., and Ayoubi, A.: Magnetic susceptibility and morphological characteristics of a loess–paleosol sequence in northeastern Iran, Catena, 101, 56–60, 2013.
- Kehl, M., Frechen, M., and Skowronek, A.: Paleosols derived from loess and loess-like sediments in the Basin of Persepolis, Southern Iran, Quatern. Int., 140/141, 135–149, 2005.
- Kehl, M., Sarvati, R., Ahmadi, H., Frechen, M., and Skowronek, A.: Loess/paleosol sequences along a climatic gradient in Northern Iran, Eiszeitalter und Gegenwart, 55, 149–173, 2006.
- Lateef, A. S. A.: Distribution, provenance, age and paleoclimatic record of the loess in Central North Iran, in: Loess – its Distribution, Geology and Soil, edited by: Eden, D. N. and Furkert, R. J., Proceedings of an International Symposium on Loess, New Zealand, 14–21 February 1987, Balkema, Rotterdam, 93–101, https://doi.org/10.13140/2.1.3244.8321, 1988.
- Liu, D., Chen, W., and Li, L.: Disentangling the relative effects of climate change and anthropogenic activities on paddy expansion in the northern Sanjiang Plain of China, Ecological Indicators, 154, 110543, https://doi.org/10.1016/j.ecolind.2023.110543, 2023.
- Maher, B. A.: The magnetic properties of Quaternary aeolian dusts and sediments, and their palaeoclimatic significance, Aeolian Res., 3, 87–145, 2011.
- Muhs, D. R. and Bettis, E. A.: Quaternary loess-paleosol sequences as examples of climate-driven sedimentary extremes, Geological Society of America Special Paper, 370, 53–74, 2003.
- Okhravi, R. and Amini, A.: Characteristics and provenance of the loess deposits of the Gharatikan Watershed in Northeast Iran, Global Planet. Change, 28, 11–22, 2001.
- Pierce, K. L., Muhs, D. R., Fosberg, M. A., Mahan, S. A., Rosenbaum, J. G., Licciardi, J. M., and Pavich, M. J.: A loess-paleosol record of climate and glacial history over the past two glacialinterglacial cycles (150 ka), southern Jackson Hole, Wyoming, Quaternary Res., 76, 119–141, 2011.
- Porter, S. C. and An, Z.: Correlation between climate events in North Atlantic and China during the last glaciation, Nature, 375, 305–308, 1995.
- Prins, M. A., Vriend, M., Nugteren, G., Vandenberghe, J., Huazu, L., Zheng, H., and Weltje, G. J.: Late Quaternary aeolian dust input variability on the Chinese Loess Plateau: inference from

unmixing of loess grain-size record, Quaternary Sci. Rev., 26, 230–242, 2007.

- Schatz, A.-K., Scholten, T., and Kühn, P.: Paleoclimate and weathering of the Tokaj (NE Hungary) loess-paleosol sequence: a comparison of geochemical weathering indices and paleoclimate parameters, Clim. Past Discuss., 10, 469–507, https://doi.org/10.5194/cpd-10-469-2014, 2014.
- Song, Y., Shi, Z., Dong, H., Nie, J., Qian, L., Chang, H., and Qiang, X.: Loess magnetic susceptibility in Central Asia and its paleoclimatic significance, in: Proceedings of the IEEE International Geoscience & Remote Sensing Symposium, II, 1227–1230, Massachusetts, https://doi.org/10.1109/IGARSS.2008.4779223, 2008.
- Spassov, S.: Loess magnetism, environment and climate change on the Chinese Loess Plateau, Doctoral Thesis, ETH Zürich, 1–151, 2002.
- Thompson, R. and Oldfield, F.: Environmental Magnetism, Allen & Unwin, Springer, London, https://doi.org/10.1007/978-94-011-8036-8, 1986.

- Tzedakis, P. C.: Hierarchical biostratigraphical classification of long pollen sequences, J. Quaternary Sci., 9, 257–259, 1994.
- Yu, K., Liu, L., Li, A., Sheng, E., Liu, X., and Lan, J.: The Rb/Sr ratios of the bulk sediments in Lake Chaonaqiu of Liupan Mountains, the western Chinese Loess Plateau and its paleoclimate implications, Quaternary Sci., 43, 122–130, 2023.
- Zhang, R., Ouyang, Z.-T., Xie, X., Guo, H.-Q., Tan, D.-Y., Xiao, X.-M., Qi, J.-G., and Zhao, B.: Impact of Climate Change on Vegetation Growth in Arid Northwest of China from 1982 to 2011, Remote Sens., 8, 364, https://doi.org/10.3390/rs8050364, 2016.
- Zhang, Q., Chang, I., and Luo, Z.: Response of vegetation to climate change over the past millennium in Eurasia, Clim. Dynam., 56, 75–88, 2021a.
- Zhang, X., Liu, L., Yu, K., and Zhang, C.: Using Rb / Sr ratios to assess weathering intensity in loess profiles, Earth Surf. Proc. Land., 46, 2454–2466, 2021b.