



Climate change drove Late Miocene to Pliocene rise and fall of C₄ vegetation at the crossroads of Africa and Eurasia (Anatolia, Türkiye)

Maud J. M. Meijers^{1,2}, Tamás Mikes^{2,3,✉}, Bora Rojay⁴, H. Evren Çubukçu⁵, Erkan Aydar^{5,6}, Tina Lüdecke^{2,7}, and Andreas Mulch^{2,8}

¹Department of Earth Sciences, NAWI Graz Geocenter, University of Graz, Heinrichstraße 26, 8010 Graz, Austria

²Senckenberg Biodiversity and Climate Research Centre, Senckenberganlage 25, 60325 Frankfurt am Main, Germany

³Independent geological consultant, Arnljot Gellines vei 35, 0657 Oslo, Norway

⁴Department of Geological Engineering, Middle East Technical University, 06800 Çankaya, Ankara, Türkiye

⁵Department of Geological Engineering, Hacettepe University, Beytepe Campus, 06800 Ankara, Türkiye

⁶Laboratoire Magmas et Volcans, CNRS, IRD, OPGC, Université Clermont Auvergne, 63000 Clermont-Ferrand, France

⁷Emmy Noether Group for Hominin Meat Consumption, Max Planck Institute for Chemistry, Hahn-Meitner-Weg 1, 55128 Mainz, Germany

⁸Institute of Geosciences, Goethe University Frankfurt, Altenhöferallee 1, 60438 Frankfurt am Main, Germany

✉deceased

Correspondence: Maud J. M. Meijers (maud.meijers@uni-graz.at)

Received: 4 December 2024 – Discussion started: 14 January 2025

Revised: 5 December 2025 – Accepted: 7 April 2026 – Published: 12 May 2026

Abstract. Life on Earth has been capitalizing on the C₃ photosynthetic pathway for 2.8 billion years. However, in the world's grasslands that emerged since the Paleogene, C₄ vegetation expanded dramatically between 8 and 3 Ma in response to climatic changes. Here we present the first comprehensive Late Miocene to Holocene $\delta^{13}\text{C}$ soil carbonate record from the Eastern Mediterranean region (Anatolia) to reconstruct long-term geographic distributions of C₃ and C₄ plants, a region with patchy records compared to parts of Africa and Asia. Our results show a colonization of Anatolian floodplains by C₄ biomass by 9.9 Ma, similar to regions in NW and E Africa, followed by a transition from this mixed C₃–C₄ vegetation to C₄ dominance between ca. 7.1 and 4.9 Ma. The transition to C₄ in Anatolia coincides with a similar shift from C₃ to C₄ vegetation in southern Asia and is generally attributed to the Late Miocene Cooling in response to decreasing atmospheric $p\text{CO}_2$. However, the Anatolian paleoecosystem patterns are unique due to a rapid and permanent return to C₃ dominance in the Early Pliocene, which is not observed elsewhere and occurs simultaneously with the disappearance of the open environment-adapted large mam-

mal Pliocene chronofauna. We propose that this return to C₃ vegetation was caused by paleoclimatic processes that regionally shifted precipitation from the warm to the cool season, resembling the modern Mediterranean climate. In conclusion, changes in rainfall seasonality under subhumid climate, rather than increased aridity, drove the demise of C₄-dominated floodplains and the open-environment adapted Pliocene chronofauna at the Eurasian-African crossroads.

1 Introduction

The earliest grassland ecosystems that emerged in the Paleogene were occupied by vegetation using the C₃ oxygenic photosynthetic pathway (Strömberg, 2011). However, many of the grassland environments that occupy ca. 40 % of Earth's landmass today are dominated by C₄ vegetation, including the Great Plains of North America, eastern South America, sub-Saharan Africa, southeast Asia, and northern Australia (Still et al., 2003). C₄ biomass consists of grasses (ca. 60 %), sedges, and dicots, and is adapted to conditions of drought,

low atmospheric $p\text{CO}_2$, and high temperatures (Sage, 2004). Although C₄ vegetation has been present since the Oligocene (Peppe et al., 2023; Urban et al., 2010), the rise to dominance of C₄ at the expense of C₃ vegetation in much of the world's grassland environments was delayed until the Late Miocene or even the Pliocene (Edwards et al., 2010; Strömberg, 2011).

Grassland expansion led to the evolution of open environment-adapted and hypsodont large herbivorous mammal communities of the Old World savannah paleobiome (OWSP) in Asia, East Africa, and southern Europe during the Late Miocene (Kaya et al., 2018) until its Early Pliocene fragmentation (Böhme et al., 2021). The western Eurasian branch of the OWSP, the Pikermian chronofauna, reached its greatest geographic extent around ca. 8.0–6.6 Ma-ago and occupied large parts of Europe and Central Asia, including Anatolia (Eronen et al., 2009) when C₄ vegetation became dominant in certain grasslands of the Old World. Reconstructing the spread of C₄ vegetation through geologic time is crucial for identifying the global and regional climatic drivers behind faunal turnover, including the spread and decline of the Pikermian chronofauna. Direct evidence for the presence of C₄ vegetation in southern Europe and the Turkish-Iranian plateau region, however, is spatially and temporally patchy and incomplete (Böhme et al., 2017; Butiseacă et al., 2022; Urban et al., 2010).

Here, we reconstruct proportions of C₄ biomass in south-central Anatolia (Türkiye; Fig. 1), which is crucially located at the crossroads of well-studied regions in terms of vegetation dynamics in Africa and Asia (Behrensmeyer et al., 2007; Uno et al., 2011) and is presently characterized by a low amount of C₄ vegetation (< 10 %). We therefore analyze carbon isotopic compositions ($\delta^{13}\text{C}$) of ca. 10 Ma to Holocene pedogenic carbonates from Anatolia. Based on the fundamentally different fractionation of carbon isotopes during photosynthesis of C₃ ($\delta^{13}\text{C} = -26.7 \pm 12.3\text{‰}$) and C₄ vegetation ($\delta^{13}\text{C} = -12.5 \pm 1.1\text{‰}$; Cerling et al., 1997) the transfer of this signal into materials such as pedogenic carbonates, mammalian tooth enamel, or leaf waxes can be used to reconstruct contributions of C₄ vegetation in deep time (Tippie and Pagani, 2007).

Our results of 447 pedogenic carbonate samples from sixteen sites show that C₄ plants spread to dominance in Anatolian floodplains during the Late Miocene and diminished to similar-to-modern proportions during the Pliocene. We compare the timing (Quade and Cerling, 1995) of floral overturn with the few available Anatolian (Türkiye) and Aegean (Greece) soil carbonate $\delta^{13}\text{C}$ records, as well as with records of faunal overturn. We then assess the drivers of vegetation dynamics in terms of components of C₄ and C₃ vegetation in Anatolia and the Aegean by comparing their timing with other soil carbonate $\delta^{13}\text{C}$ records from the Old World, as well as global marine climate records.

2 Material and methods

2.1 Pedogenic carbonate sampling

In Anatolia, eight pedogenic carbonate sites were sampled within the Central Anatolian Volcanic Province (CAVP) and two sites each adjacent to the Tuz Gölü and Ecemiş fault zones, as well as the Ermenek basin. Two additional sites were sampled within the Adana basin near the Mediterranean Sea (Fig. 1b, Table 1; see Table S1 and Text S1 for detailed site information). Site elevations from this study range between ca. 110 and 1650 m a.s.l. and the sampled soil carbonates range in age from ca. 10 Ma to modern (Fig. 1b; Table S1). Modern pedogenic carbonate samples consist of friable to weakly consolidated calcareous nodules typically 1 to 3 cm in diameter, from the B horizon of well-drained soils with a pink to red hackly appearance. Miocene to Pleistocene carbonate subsoil horizons developed within floodplain deposits. They comprise moderately consolidated to hard calcareous nodules ranging from 3 to 8 cm in diameter, or representative domains of dm-scale nodular carbonate layers. Whenever possible, pedogenic carbonate nodules and in some cases casts of taproots and fibrous roots, were collected from Bk horizons of cream to red-colored subsurface soils. Sampling choices were based on the availability of pedogenic carbonates in outcrops, which were taken over several months of fieldwork. Where identifiable, pedogenic carbonate was sampled at a minimum soil depth of 30 cm to reduce the influence of atmospheric CO₂ (Cerling and Quade, 1993). Representative pictures of sampled outcrops and levels are shown in Fig. S1.

2.2 $\delta^{13}\text{C}$ of pedogenic carbonates

Powdered material from pedogenic carbonate samples was extracted with a diamond-tip dental drill. Only micritic parts devoid of clasts were drilled for analysis. A total of 447 $\delta^{13}\text{C}$ values were obtained for pedogenic carbonate samples from sixteen sites using a Thermo V mass spectrometer at the Institute of Geology (University of Hanover, Hanover, Germany) and a Thermo MAT 253 mass spectrometer at the joint Goethe University-Senckenberg BiK-F Stable Isotope Facility (Frankfurt, Germany). Data were collected during multiple sessions in both labs. $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values were evaluated simultaneously and are reported in Table S1. The $\delta^{18}\text{O}$ values that were obtained during the measurements were published in Meijers et al. (2025) and interpreted with respect to the surface uplift history of Anatolia. To assess intra- and inter-nodule variability, multiple measurements were performed on select single pedogenic carbonate nodules which were cross-sectioned from three sites: 10TG55 ($n = 16$, five nodules, three to four analyses per nodule), 10FD ($n = 48$, 25 nodules, of which three nodules with six to 13 analyses per nodule), and 12C047–052 ($n = 24$, 16 nodules, of which two nodules with five measure-

Table 1. Summary of pedogenic carbonate δ¹³C records from Anatolia (Türkiye, this study).

Basin/area	Locality	Site/samples	Number of samples	Age (Ma)	Δ Age (Ma)	Latitude (°N)	Longitude (°E)	Average δ ¹³ C (‰ V-PDB)	1σ SD (‰)	2σ SD (‰)	Median δ ¹³ C (‰ V-PDB)	25th perc. (‰)	75th perc. (‰)
Holocene (5 ± 5 ka)													
Ermenek basin-Türkiye	Yalıncıl	08Erm003A-004A ^b	2	0.005	0.005	36.50844	32.95255	-9.0	0.4	0.9	-9.0	-9.2	-8.9
Ermenek basin-Türkiye	Güneyyurt	08Erm006A-008A ^b	3	0.005	0.005	36.68815	32.76500	-8.8	0.7	1.4	-8.9	-9.2	-8.5
Tuz Gölü-Türkiye	Altınkaya	10TG55 ^b	16	0.005	0.005	38.63093	33.80005	-8.5	0.1	0.2	-8.5	-8.5	-8.4
EFZ-Türkiye	Bulanlık	10BL01-07 + 09BL02-06 ^b	12	0.005	0.005	37.89825	35.10614	-6.3	0.6	1.2	-6.0	-6.8	-5.8
EFZ-Türkiye	Fındıklı	10FD ^b	48	0.005	0.005	37.54615	34.94563	-8.4	0.4	0.8	-8.4	-8.7	-8.2
Adana basin-Türkiye	Gıldirli	V01-128-136 ^a	6	0.005	0.005	37.34890	35.05038	-7.4	2.1	4.3	-8.6	-8.7	-5.6
Average: -8.1 ± 1.0‰ (n = 6; 1σ SD)													
Ca. 3.9 to 1.4 Ma													
Adana basin-Türkiye	Baklılıand Misis	11AD01-95 ^b	89	1.35	1.25		^c	-8.2	0.6	1.1	-8.3	-8.6	-7.9
CAVP-Türkiye	Taşhan	14 MM 05-07 ^b	3	3.81	1.08	38.90621	35.45562	-7.4	0.5	1.0	-7.3	-7.6	-7.1
Tuz Gölü-Türkiye	Cerit	10TG80, 87-91, 93-97 ^b	96	3.86	0.20	38.61480	33.88557	-7.2	0.5	1.1	-7.0	-7.7	-6.8
Average: -7.6 ± 0.5‰ (n = 3; 1σ SD)													
Ca. 7.1 to 4.9 Ma													
CAVP-Türkiye	Güzölöz	12C-070-084 ^b	15	4.86	0.16	38.39440	34.97231	1.0	2.1	4.3	1.1	0.3	2.6
CAVP-Türkiye	Taşkımpaşa SW	12C-099-101 ^b	3	5.68	0.66	38.48550	34.93776	0.1	1.1	2.3	-0.3	-0.6	0.5
CAVP-Türkiye	Şahinefendi	10CKK ^b	86	5.68	0.66	38.46783	34.93887	-1.3	1.6	3.3	-1.3	-2.1	-0.4
CAVP-Türkiye	Orta Tepe W	12C-002-023 ^b	22	6.21	0.14	38.47888	34.95045	-2.5	2.0	4.1	-2.6	-3.8	-1.1
CAVP-Türkiye	Taşkımpaşa S	12C-063-069 ^b	6	6.21	0.14	38.48662	34.94468	-1.6	2.6	5.1	-0.6	-3.1	0.2
CAVP-Türkiye	Orta Tepe S	12C-025-046 ^b	22	7.05	0.15	38.47325	34.95856	-1.1	2.1	4.1	-0.7	-2.6	0.8
Average: -0.9 ± 1.3‰ (n = 6; 1σ SD)													
Ca. 9.9 Ma													
CAVP-Türkiye	Mustafapaşa	12C-047-062 ^b	24	9.88	0.75	38.57471	34.91145	-5.2	2.6	5.2	-6.0	-6.9	-3.9

SD = standard deviation; perc. = percentile; CAVP = Central Anatolian Volcanic Province; EFZ = Eceemis fault zone. For detailed geochronologies we refer to Text S1. ^a δ¹⁸O values published in: Meijers et al. (2025); ^b Locality 11AD consists of three sublocalities from time-equivalent deposits in a small area and were therefore merged. Baklılı 1: 37.00761° N, 35.63193° E; Baklılı 2: 37.01079° N, 35.63286° E; Baklılı 3: 37.00203° N, 35.62834° E; Misis (Yakapınar): 36.97569° N, 35.62544° E. GPS coordinates in WGS84.




Figure 1. (a) Topographic map of the studied Eastern Mediterranean region including sampling sites from this study and published studies in the Aegean (Greece) and Anatolia (Türkiye). Circles: this study; triangle Meijers et al. (2018); star: Lepetit (2010); diamonds: Böhme et al. (2017); inverted triangles: Quade et al. (1994). Geographic sampling areas correspond to those in Table S1 and S3 in the Supplement. CAVP = Central Anatolian Volcanic Province. (b) Location of the study area (red box) in panel (a) within the Old World.

ments each; Table S1). Powdered carbonate samples were digested in 100 % H₃PO₄ and analyzed as CO₂ in continuous flow mode using above mentioned mass spectrometers interfaced with a Thermo GasBench II. Analytical procedures followed those of Spötl and Vennemann (2003), which includes corrections for scale, linearity, and drift. Raw isotopic ratios were calibrated against an in-house standard (Carrara marble; $\delta^{13}\text{C} = 2.01\text{‰}$ (V-PDB), $\delta^{18}\text{O} = -1.74\text{‰}$ (V-SMOW)), a synthetic Merck standard ($\delta^{13}\text{C} = -35.69\text{‰}$ (V-PDB), $\delta^{18}\text{O} = 12.38\text{‰}$ (V-SMOW)) and international carbonate reference material (NBS18). The in-house Carrara marble standard was weighed in for four different sample sizes for the linearity correction. Final isotopic ratios are reported against Vienna Pee Dee Belemnite (V-PDB), with analytical uncertainties that are typically better than 0.07‰. In the text, we refer to average $\delta^{13}\text{C}$ values with 1 σ standard deviations; averages, 1 σ , and 2 σ standard deviations, as well as medians, and 25th and 75th percentiles are displayed in Tables 1 and S1. Pedogenic carbonate that formed in equilibrium with soil-respired CO₂ is enriched by ca. 14‰–17‰ in $\delta^{13}\text{C}$ compared to CO₂ derived from root respiration and microbial decomposition of organic matter (e.g., Tipple and Pagani, 2007).

2.3 Pedogenic carbonate chronology

Age constraints for the paleosols containing the analyzed pedogenic carbonates from the CAP and the Tuz Gölü area are based on radiometrically dated ignimbrite intercalations of the CAVP (Aydar et al., 2012; Özsayın et al., 2013). The stratigraphic framework for the sampled horizons is based on the ignimbrite stratigraphy and the published geochronological data (see Text S1 and Table S1). A composite section

of the area of the southern CAVP where six sites were sampled shows the stratigraphic relationship between the sampled intervals (Fig. S1). The Quaternary age for site 11AD is based on the 1 : 500 000 geological map of Adana (Ulu, 2002). Modern soil carbonate horizons were identified based on field relationships and assigned a Holocene age (5 ± 5 ka).

2.4 Reconstruction of climatic parameters

A reconstruction of paleobotany-derived climatic parameters of Anatolia (Fig. 2b, c) is based on the coexistence approach (Mosbrugger and Utescher, 1997). All data are listed in Table S2.

2.5 Compilation of $\delta^{13}\text{C}$ values from Old World soil carbonates

The compilation of 12 Ma to Holocene soil carbonate $\delta^{13}\text{C}$ values from Türkiye and Greece (Fig. 2a; Table S3) includes data from Böhme et al. (2017), Lepetit (2010), and Quade et al. (1994). The compilation of 12 Ma to Holocene soil carbonate $\delta^{13}\text{C}$ values from Asia and Africa (Fig. 3d, e) was retrieved from Fox et al. (2018). Datasets lacking age constraints were excluded. We also excluded elevated $\delta^{13}\text{C}$ values from one multiproxy study on the Qaidam basin, which are attributed to low soil respiration rates in response to increased aridification (Zhuang et al., 2011). As such, the $\delta^{13}\text{C}$ values may not be only interpreted in terms of variations in the dominating photosynthetic pathways (see also Sect. 4.1). Studies included in the Africa compilation (Fig. 3e): Aronson et al. (2008), Bestland and Krull (1999), Cerling and Hay (1986), Cerling et al. (1988, 1991, 2003, 2011), Kingston (1992), Levin et al. (2004, 2011), Plummer et al. (1999,




Figure 2. 12 to 0 Ma Anatolian and Aegean pedogenic carbonate $\delta^{13}\text{C}$ records and Anatolian climatic, faunal, and surface uplift records. **(a)** Pedogenic carbonate $\delta^{13}\text{C}$ values and their averages per site for this (orange) and published (purple, yellow) studies (Böhme et al., 2017; Lepetit, 2010; Meijers et al., 2018; Quade et al., 1994) from Anatolia and the Aegean region. Small symbols with black stroke for sites with a single measurement. Global “Late Miocene Cooling” (blue shading) according to Herbert et al. (2016). MSC: Messinian Salinity Crisis. **(b, c)** Published paleobotanical data-based (Table S2) cold month mean temperature (CMT) and mean annual temperature (MAT) reconstructions for Anatolia. Sites were subdivided in plateau interior and (near-)coastal because not only climate change but also surface uplift affected CMT and MAT on the plateau. Light green circles indicate sites that are currently within in the CAP interior (ca. 1–1.5 km elevation), dark green circles indicate (near-)coastal sampling sites. **(d)** Periods during which the Pikermian chronofauna roamed Anatolia and surface uplift of the CAP occurred (onset: ca. 11 Ma; Meijers et al., 2018). The greatest geographic extent refers to the western Eurasian distribution of the Pikermian fauna (Eronen et al., 2009).

2009), Quade et al. (2004), Quinn et al. (2007), Sahnouni et al. (2011), Sikes (1994), Sikes et al. (1999), Sikes and Ashley (2007), WoldeGabriel et al. (2009), Wynn (2000, 2004), Wynn et al. (2006). Studies included in the Asia compilation (Fig. 3d): An et al. (2005), Behrensmeyer et al. (2007), Ding and Yang (2000), Ghosh et al. (2004), Kaakinen et al. (2006), Passey et al. (2009), Quade et al. (1994), Quade and Cerling (1995), Sanyal et al. (2004), Yao et al. (2010).

3 Results

$\delta^{13}\text{C}$ values from the sixteen 10 Ma to Holocene sampled soil carbonate locations range from -9.5‰ to 3.4‰ , with an average of $-5.4 \pm 3.4\text{‰}$ (Fig. 2a, Tables 1 and S1). Based on their $\delta^{13}\text{C}$ values (uncertainties reported at 1 sigma standard deviation (1σ SD)), we categorize the pedogenic carbonate record into four time intervals:

- 9.9 Ma: $\delta^{13}\text{C}$ values of a single 9.9 Ma site ($n = 24$) in the Central Anatolian Volcanic Province (CAVP) average $-5.2 \pm 2.6\text{‰}$.

- 7.1 to 4.9 Ma: The average $\delta^{13}\text{C}$ value for the six sections ($n = 154$) from the CAVP is $-1.2 \pm 2.0\text{‰}$, which is ca. 4‰ more positive than the 9.9 Ma dataset.
- 3.9 to 1.4 Ma: Three pedogenic carbonates sites were sampled in the southern part of the CAP (CAVP and Tuz Gölü fault (TGF)), the Tauride Mountains (Ecemiş fault zone (EFZ) and Ermenek basin), and the Adana basin. The average $\delta^{13}\text{C}$ value ($n = 188$) is $-7.7 \pm 0.8\text{‰}$, which is nearly 7‰ more negative than the 7.1 to 4.9 Ma dataset.
- Holocene (5 ± 5 ka) $\delta^{13}\text{C}$ values from the six sections containing Holocene soil carbonates ($n = 87$) average $-8.1 \pm 1.0\text{‰}$, which is within uncertainty identical to the 3.9 to 1.4 Ma datasets.

In summary, all soil carbonates younger than ca. 3.9 Ma yield nearly 7‰ lower $\delta^{13}\text{C}$ values than the 7.1 to 4.9 Ma soil carbonates, as well as ca. 2.5‰ lower $\delta^{13}\text{C}$ values than the ca. 9.9 Ma-old soil carbonate dataset (Fig. 2a).

4 Discussion

4.1 Soil carbonate $\delta^{13}\text{C}$ reveals variations of C₃ and C₄ vegetation in Anatolia and the Aegean

The 12 Ma to Holocene $\delta^{13}\text{C}$ record from Anatolia shows large changes between the four time intervals (see Sect. 3 Results) and covers a wide range from -9.5‰ to 3.4‰ (Fig. 2). This variation in $\delta^{13}\text{C}$ values (up to 12.9‰) may reflect temporal changes in soil respiration rates, plant water stress, and/or the dominating photosynthetic pathway (C₃ vs. C₄) of biomass in Anatolia. We interpret obtained $\delta^{13}\text{C}$ values (and associated $\delta^{18}\text{O}$ values; Meijers et al., 2025) as primary (i.e. reflecting the isotopic ratios upon the formation of the carbonates), because the values are within a range typical of soil carbonates (e.g., Cerling and Quade, 1993). Additionally, the paleoaltimetry study by Meijers et al., (2025) that is based on the $\delta^{18}\text{O}$ values associated with the $\delta^{13}\text{C}$ values of this study includes two dual clumped isotope (Δ_{47} , Δ_{48}) temperatures. First, results for the ca. 5.4 Ma pedogenic carbonate sample (from CAVP site 10CKK, see also this study) and lake carbonate sample (from the Adana basin in southern Turkey) plot within error of the dual clumped isotope equilibrium line, which implies that their isotopic compositions are devoid of significant kinetic biases. Second, the obtained Δ_{47} temperatures of $15.6 \pm 3.3\text{ °C}$ (CAVP) and $24.1 \pm 3.5\text{ °C}$ (Adana basin) show that the carbonates precipitated under near-surface conditions rather than at elevated diagenetic temperatures. More evidence for the primary origin of the sampled soil carbonates comes from the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values and thin sections of lake carbonate that was sampled in the same or similar fluvio-lacustrine host lithologies, which show no evidence of diagenetic alteration (Meijers et al., 2018, 2020). Moreover, all sites within the Central

Anatolian Plateau interior were buried only 170 m or less (Table S1), except for Mustafapaşa (ca. 9.9 ± 0.8 Ma; burial depth ca. 400 m). Low burial depths result from relatively low sedimentation rates (ca. 25–80 m Myr⁻¹; Table S1) and rapid drainage integration and incision of the basins within maximal 2.5 Myr after the latest deposition of the soil carbonate host lithologies (Brocard et al., 2021; Meijers et al., 2020).

Under conditions of low soil CO₂ production, soil carbonates tend to form at shallower depths and may incorporate varying proportions of atmospheric CO₂ (Cerling, 1984; Cerling and Quade, 1993), resulting in higher $\delta^{13}\text{C}$ values (Caves et al., 2016; Licht et al., 2020). Particularly in dry ecosystems, C₃ plants may yield increased $\delta^{13}\text{C}$ values (Kohn, 2010). Although Anatolia is presently characterized by a climate with dry summers, all available paleobotanic datasets from the region suggest subhumid climatic conditions during the Late Miocene to Pliocene (Table S2). We therefore interpret the variations in $\delta^{13}\text{C}$ values of our 10 Ma to recent soil carbonate record to reflect significant changes in the relative contribution of C₃ and C₄ components of vegetation.

The large range of soil carbonate $\delta^{13}\text{C}$ values of over 10‰ at ca. 9.9 Ma in Anatolia is consistent with a heterogeneous vegetation cover that includes both C₃ and C₄ plants, albeit with a dominance of C₃ vegetation given an average $\delta^{13}\text{C}$ of -5.2‰ . Pedogenic carbonate $\delta^{13}\text{C}$ values between 7.1 and 4.9 Ma, which average 4.0‰ more positive compared to $\delta^{13}\text{C}$ values at 9.9 Ma indicate central Anatolian floodplain environments dominated by C₄ vegetation (Fig. 2a). The dominance of C₄ biomass between 7.1 and 4.9 Ma is consistent with published CAP $\delta^{13}\text{C}$ values of pedogenic carbonate from this time interval (6.7 to 4.8 Ma; Fig. 2a; Lepetit, 2010), some of which were obtained from the same stratigraphic intervals. For the time intervals from 3.9 to 1.4 Ma and the Holocene our $\delta^{13}\text{C}$ averages are significantly lower – by nearly 7‰ – than those from the 7.1 to 4.9 Ma interval, and by ca. 2.5‰ compared to the 9.9 Ma interval. This indicates the presence of floodplains dominated by C₃ vegetation after 3.9 Ma.

$\delta^{13}\text{C}$ values of ca. 9.3 to 5.3 Ma fossil equid tooth enamel (Rey et al., 2013), and ca. 10 to 6.5 Ma pedogenic carbonate $\delta^{13}\text{C}$ records (Böhme et al., 2017; Quade et al., 1994; Table S3) from the Aegean region (Greece) indicate C₃ vegetation. However, phytoliths, as well as $\delta^{13}\text{C}$ values of published pedogenic carbonate and fossil herbivore tooth enamel, imply the presence of C₄ vegetation between ca. 9 and 7 Ma in the CAVP (Kayseri-Özer et al., 2017; Lepetit, 2010), Pikerimi and Samos (Aegean, Greece; Fig. 1b; Böhme et al., 2017; Quade et al., 1994), and Maragheh (Bernor et al., 2016; Biasatti et al., 2015; Strömberg et al., 2007). On Crete and Cyprus (Greece) ca. 5‰ variations in $\delta^{13}\text{C}$ values of leaf waxes (long-chain n-alkanes produced by terrestrial plants), superimposed on an overall 4‰ increase in $\delta^{13}\text{C}$ values between ca. 7 and 6 Ma, indicate the expansion of C₄ vegeta-




Figure 3. Climate and carbon isotope records spanning the last 12 Ma. **(a)** Cenozoic Global Reference (CENOGRID) benthic foraminifera $\delta^{13}\text{C}$ curves, resampled and smoothed over 20 kyr (blue) and 1 Myr (red) periods (Westerhold et al., 2020). **(b)** SST anomalies for different hemispheric and latitude bins (Herbert et al., 2016). **(c–e)** Compilation of soil carbonate $\delta^{13}\text{C}$ values from Anatolia and the Aegean **(c)**, Asia **(d)**, and Africa **(e)**. Asian and African datasets were retrieved from Fox et al. (2018). See Sect. 2.5. **(f)** Onset of the initial C₄ expansion in the Old World from a compilation by Polissar et al. (2019) and including Anatolia (this study). Blue shading indicates time interval of Late Miocene Cooling (Herbert et al., 2016). NH = Northern Hemisphere, SH = Southern Hemisphere. MSC: Messinian Salinity Crisis (5.96–5.33 Ma; e.g. Roveri et al., 2014), during which the restriction of the Mediterranean Basin led to drastic changes in local hydroclimate; AMOC+: strengthened Atlantic Meridional Overturning Circulation (AMOC) in response to shoaling of the Central American Seaway (ca. 4.8–4.0 Ma; Haug and Tiedemann, 1998) resulted in higher North Atlantic SSTs (Karas et al., 2017). We suggest that associated hydroclimatic changes over Europe and the Mediterranean region resulted in the decline of C₄ vegetation in Anatolia (see Sect. 4.4).

tion (Butiseacă et al., 2022; Mayser et al., 2017) in response to Messinian climate cycles. Collectively, our data in combination with published datasets indicate that the geographic extent of C₄ expansion in the Eastern Mediterranean region during the Late Miocene and earliest Pliocene may not have been restricted to Anatolia, but extended into the Aegean and the Iranian plateau. After 3.9 Ma (late Early Pliocene), $\delta^{13}\text{C}$ values from Anatolian soil carbonates are similar to those derived from the Aegean (Fig. 1b; Quade et al., 1994) and indicate vegetation dominated by C₃ biomass, which is the observed dominant vegetation type in Anatolia and the Aegean region today (< 10 % C₄; Still et al., 2003).

4.2 Late Miocene to Pliocene circum-Anatolian ecosystem reconstructions

Pedogenic carbonates from ca. 10 Ma to Holocene floodplain deposits in Anatolia indicate that the region has been characterized by rainfall seasonality since the Late Miocene, as their formation requires periodic soil drying (e.g., Zamanian et al., 2016). Furthermore, the dominance of C₄ vegetation between 7.1 and 4.9 Ma in Anatolia, as reconstructed from our pedogenic carbonate $\delta^{13}\text{C}$ values, suggests that open-habitat grasslands characterized portions of the landscape. This is supported by (a) paleobotanical (macrofossils, pollen, and spores) data that suggest the introduction of steppe elements in Anatolia between 9 and 6 Ma (Denk et al., 2018), (b) the (Middle-)Late Miocene rise of open-habitat grasslands in Anatolia and nearby Greece and Iran reconstructed from phytoliths (Strömberg et al., 2007), and (c) the presence of the open-environment adapted Pliocene chronofauna in aforementioned regions (Eronen et al., 2009). However, macrofossil, pollen, and spore records from Anatolia, Greece, and Bulgaria (Denk et al., 2018) also sketch Late Miocene landscapes covered by evergreen needleleaf forests and mixed forests. As such, Denk et al. (2018) reject a cohesive savannah biome. We propose that the results from the paleobotanical and paleontological studies, as well as our stable isotope-based vegetation reconstructions, can be reconciled by heterogeneous Late Miocene Anatolian landscapes with largely interconnected forested as well as savannah-like environments (see also Fortelius et al., 2019).

The paleobotanical, paleontological, and soil carbonate records were all retrieved from ca. 11 to 4 Ma low-relief floodplains where fluvial and lacustrine deposits accumulated. Similar to today, the low-relief areas were interrupted by relict mountain ranges and local fault-controlled relief, but were not connected to marine basins. The fluvial and lacustrine records are currently accessible as a result of rapid latest Miocene to Pliocene drainage integration of the Anatolian plateau and subsequent river incision (Brocard et al., 2021; Meijers et al., 2020). As such, our soil carbonate $\delta^{13}\text{C}$ records are biased towards partially preserved and incised intermontane floodplains and underrepresent vegetation dynamics of (potentially) forested and eroding to-

pographic highs. Simultaneously, wind-blown pollen from forested montane areas are partially preserved in the fluvial and lacustrine records. Our $\delta^{13}\text{C}$ soil carbonate record therefore highlights the importance of multi-proxy ecosystem reconstructions and solidifies evidence for a highly variable vegetation cover in Anatolia during the Late Miocene.

4.3 Timing and drivers of Late Miocene ecological change

Located at the crossroads of Africa, Asia, and Europe, the soil carbonate $\delta^{13}\text{C}$ record of Anatolia bears the potential to identify paleoenvironmental dynamics specific to the spread of C₄ vegetation in the Old World as well as to changing paleoclimatic conditions in the Mediterranean region. We compare the Anatolian soil carbonate $\delta^{13}\text{C}$ record with available leaf wax, fossil tooth enamel, and soil carbonate $\delta^{13}\text{C}$ records that constrain the initial expansion of C₄ vegetation (Polissar et al., 2019) and its rise to dominance in the Old World grasslands during the Late Miocene (Fig. 3d–f). We conclude that the onset of C₄ expansion in Anatolia during the Late Miocene (at the latest at ca. 9.9 Ma) is roughly coeval with the expansion of C₄ ecosystems in NW and E Africa and predates its rise in other Asian and African regions (Fig. 3f). However, C₄ vegetation dominated Anatolian floodplains and potentially parts of the Aegean by ca. 7.2 Ma, which roughly coincides with the start of the rapid rise to dominance of C₄ vegetation in southern Asian grasslands between 7.8 and 7.2 Ma (Fig. 3c, d; e.g., Behrensmeier et al., 2007; Quade and Cerling, 1995; Tauxe and Feakins, 2020) and predates the rise to dominance of C₄ vegetation in the grasslands of the southern East African Rift during the Pliocene (Fig. 3e; e.g., Cerling et al., 2011; Lüdecke et al., 2016).

The rise to dominance of C₄ vegetation in East Asian grassland ecosystems is hypothesized to have occurred in response to declining atmospheric $p\text{CO}_2$ levels that led to Late Miocene Cooling (LMC, ca. 7.5 to 5.5 Ma; Fig. 3b; Herbert et al., 2016) when low $p\text{CO}_2$ provided an evolutionary advantage for C₄ over C₃ vegetation despite decreasing temperatures (e.g., Polissar et al., 2019; Wen et al., 2023). Worldwide, LMC is manifested in a sharp drop in sea surface temperatures (SSTs; Fig. 3b; Herbert et al., 2016) and its onset is accompanied by a sharp decrease in $\delta^{13}\text{C}$ values of benthic foraminifera (Fig. 3a; Westerhold et al., 2020), which attests to a profound change in the global carbon cycle and consequently ocean circulation patterns (Holbourn et al., 2018). Climatic reconstructions from Anatolia based on the coexistence approach (Table S2) indicate a ca. 2–3 °C decrease in mean annual and an up to 10 °C decrease in cold month mean temperatures during the Late Miocene in the CAP interior, although uncertainties in both reconstructed temperatures and ages are large (see Fig. 2b, c). Because the rise to dominance of C₄ vegetation in Anatolian floodplains and possibly the Aegean occurred simultaneously with its rise in

southern Asian ecosystems and LMC (Fig. 3c, d) we suggest that it was caused by drivers that go beyond regional environmental changes such as surface uplift of the CAP since ca. 11 Ma (Fig. 2d; Meijers et al., 2018).

Whereas C₄ grassland expansion in southern Asia, the Chinese Loess Plateau, and Arabia coincided with aridification (Huang et al., 2007; Wen et al., 2023), the (gradual) expansion of C₄ vegetation in northern and eastern Africa was not driven by aridification (Crocker et al., 2022; Polissar et al., 2019). The latter also appears to hold for Anatolia and the Aegean, as indicated by Anatolian paleobotanical datasets suggesting subhumid conditions (e.g., Kayseri-Özer, 2017; Table S3) and mesic environments in Anatolia and the Aegean instead (Denk et al., 2018).

Around ca. 8.0–6.6 Ma, the Pliocene chronofauna peaked in terms of geographic extent (Fig. 2d), including large parts of Europe and Central Asia (Eronen et al., 2009). We suggest that the spread of C₄ grasslands, which started before 9.9 Ma, and their dominance in floodplain environments by 7.2 Ma led to the expansion of the hypsodont Pliocene chronofauna in Anatolia. A similar process is observed in southern Asian grasslands, where combined soil and fossil mammal tooth enamel $\delta^{13}\text{C}$ values show that long-term climate forcing changed the vegetation structure (C₃ to C₄) between 8.5 and 6.0 Ma and led to the disappearance of most mammalian lineages that fed on C₃ vegetation (Badgley et al., 2008).

4.4 Drivers and consequences of unique and persistent Early Pliocene C₄ decline

A marked decrease in our pedogenic carbonate $\delta^{13}\text{C}$ values by approximately 7‰ from 4.9 to 3.9 Ma (Fig. 2a) implies a major overturn of Anatolian and potentially Aegean ecosystems. This transition led to the reemergence of landscapes dominated by C₃ vegetation, a feature that persists to the present day. Plant leaf wax $\delta^{13}\text{C}$ values and pollen data from a sediment core in the Gulf of Aden reveal a brief reversal in the trend of increasing C₄ vegetation contributions between 4.9 and 4.6 Ma (Feakins et al., 2013). On and around the Chinese Loess Plateau, variations in C₄ and C₃ biomass have been linked to the strengthening and weakening of the East Asian summer monsoon (e.g., An et al., 2005; Passey et al., 2009; Yao et al., 2010). However, these fluctuations are not persistent over time. Therefore, the rapid and persistent Pliocene switch to C₃ vegetation is a feature that is so far unique to Anatolia. By contrast, southeast China (Fig. 3f) and regions in the New World (see e.g., Polissar et al., 2019) have been colonized by C₄ vegetation only after the return to C₃-dominated vegetation in Anatolia. Given that the return to C₃ vegetation is geographically limited we assess potential causes for C₄ decline in Anatolia within a regional framework.

The Messinian Salinity Crisis (MSC) in the Mediterranean basin (5.96–5.33 Ma; e.g., Roveri et al., 2014) had the poten-

tial to alter the regional hydrologic cycle and hence vegetation. However, considering that the MSC is relatively short-lived and predates the decline of C₄ grasslands in Anatolia between 4.9 and 3.9 Ma, it is unlikely to be the main driver of the prolonged changes in the vegetation structure of the Eastern Mediterranean region.

During the Early Pliocene, regions adjacent to Anatolia also experienced significant environmental changes. In Central Europe, increased aridity led to the disappearance of open woodland, broad-leaf evergreen, and humid sclerophyllous taxa (Mosbrugger et al., 2005), some of which found refuge in Anatolia and the Eastern Mediterranean region (Eronen et al., 2009). Concurrently, Eastern Europe experienced a transition to cooler, drier conditions, leading to the expansion of grasslands and an enhanced fire regime since ca. 4.4 Ma (Feurdean and Vasiliev, 2019). The divergence in climate between a drier Central Europe and a more humid Eastern Mediterranean region (Kovar-Eder, 2003) has been attributed to the deflection of the westerlies from Central Europe to the Mediterranean region (Eronen et al., 2009). Its timing coincides with strengthening of the Atlantic Meridional Overturning Circulation (AMOC) in response to shoaling of the Central American Seaway (CAS) around 4.8–4.0 Ma (e.g., Haug and Tiedemann, 1998). We therefore surmise that the increase in SSTs over the North Atlantic associated with AMOC strengthening (Karas et al., 2017) caused hydroclimatic changes over Europe and the Mediterranean region.

Regions with prevalent C₄ vegetation today are not only characterized by high growing season temperatures but also by warm season precipitation. In contrast, Anatolia presently experiences hot, dry Mediterranean summers and receives most of its precipitation from October through May (Schemmel et al., 2013; Türkeş and Erlat, 2005). Notably, leaf morphologies of Miocene species of evergreen oak (*Quercus* sect. *Ilex*) from southwest Türkiye and Aegean islands resemble modern Himalayan members, which suggests (summer-)humid climatic conditions (see Denk et al., 2018, but also Denk et al., 2014, and Velitzelos et al., 2014). Additionally, Tortonian (ca. 11.6–7.2 Ma) paleoclimate simulations of Europe indicate humid to subhumid summer conditions in the northern Mediterranean region (Quan et al., 2014). We therefore propose that the shift from warm to cold season precipitation and the emergence of a Mediterranean-style climate drove the demise of C₄ grasslands and a return to a C₃-dominated environment in Anatolia and the Aegean during the Early Pliocene (4.9 to 3.9 Ma). Such a higher proportion of precipitation falling during the cool season has been shown to result in a strong and rapid (3–4 years) effect on the relative abundance of C₃ and C₄ vegetation during the 1930s Dust Bowl and was successfully reproduced in experimental studies in modern grasslands in the Great Plains (United States; Knapp et al., 2020).

The Early Pliocene demise of C₄ biomass occurred simultaneously with significant large mammal faunal turnover

(Huang et al., 2019) and the vanishing of open-landscape adapted large mammals from Anatolia (MN14; Eronen et al., 2009), which formed the last stronghold of the Pliocene chronofauna (Eronen et al., 2009). Our study therefore identifies long-term vegetation dynamics and indicates that Early Pliocene climate change profoundly and irreversibly transformed vegetation structures and drove a turnover in mammalian populations.

5 Conclusions

C₄ vegetation became ecologically dominant in Anatolian grasslands by 7.2 Ma, contemporaneous with similar developments in southern Asia and the Aegean. In contrast to the common association of C₄ biomass expansion with arid conditions as the main driver, however, C₄ expansion in Anatolia occurred under relatively humid conditions as indicated by consistent paleobotanical and stable isotope paleoclimate records. This suggests that a reduction in atmospheric *p*CO₂ between ca. 7.5 and 5.5 Ma primarily drove C₄ expansion in this region. A distinctive, rapid decline of C₄ vegetation is observed in Anatolia between 4.9 and 3.9 Ma, leading to a C₃ vegetation-dominated environment that has persisted until today. We hypothesize that this shift, along with the disappearance of the Pliocene chronofauna, was influenced by the transition to a Mediterranean climate characterized by a change from warm to cold season precipitation.

Data availability. All supporting datasets are available in the Supplement.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/cp-22-975-2026-supplement>.

Author contributions. MJM, TM, and AM: Conceptualization; MJM and TM: Investigation; MJM, TM, TL, AM: Methodology; MJM, TM, BR, HEÇ, EA, TL, and AM: Resources; MJM: Visualization; MJM: Writing – original draft; MJM, TM, BR, HEÇ, EA, TL, and AM: Writing – review & editing.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. We dedicate this publication to Tamás Mikes, who sadly passed away last year at a much too young age. We will remember Tamás as a joyful, curious, and exceptionally kind person and a knowledgeable and valued colleague. Without his dedication to field work and passion for science we would not have been able to conduct the presented research.

We express our sincere appreciation to the ESF TopoEurope VAMP and US-NSF CD-CAT research consortia for their significant contributions. AM acknowledges funding through ESF/DFG MU2845/1-1; EA and HEÇ acknowledge funding through Tübitak 107Y333. The authors acknowledge financial support by the University of Graz. We particularly thank A. Çiner (Hacettepe University, Ankara) for generous support throughout much of the study. Special thanks go to P. Ballato and F. Schemmel for field assistance. We thank E. Krsnik, U. Treffert (Senckenberg BiK-F), C. Wenske (Univ. Hannover), and J. Fiebig (Goethe University Frankfurt) for invaluable laboratory support. We thank Mine Sezgül Kayseri Özer, William Lukens, and an anonymous reviewer for their constructive comments and suggestions.

Financial support. This research has been supported by the Deutsche Forschungsgemeinschaft (grant no. MU2845/1-1) and the Türkiye Bilimsel ve Teknolojik Araştırma Kurumu (grant no. 107Y333).

Review statement. This paper was edited by Denis-Didier Rousseau and reviewed by Mine Sezgül Kayseri Özer, William Lukens, and one anonymous referee.

References

- An, Z., Huang, Y., Liu, W., Guo, Z., Clemens, S., Li, L., Prell, W., Ning, Y., Cai, Y., Zhou, W., Lin, B., Zhang, Q., Cao, Y., Qiang, X., Chang, H., and Wu, Z.: Multiple expansions of C₄ plant biomass in East Asia since 7 Ma coupled with strengthened monsoon circulation, *Geology*, 33, 705–708, <https://doi.org/10.1130/G21423.1>, 2005.
- Aronson, J., Hailemichael, M., and Savin, S.: Hominid environments at Hadar from paleosol studies in a framework of Ethiopian climate change, *J. Hum. Evol.*, 55, 532–550, <https://doi.org/10.1016/j.jhevol.2008.04.004>, 2008.
- Aydar, E., Schmitt, A. K., Çubukçu, H. E., Akin, L., Ersoy, O., Sen, E., Duncan, R. A., and Atici, G.: Correlation of ignimbrites in the central Anatolian volcanic province using zircon and plagioclase ages and zircon compositions, *J. Volcanol. Geoth. Res.*, 213–214, 83–97, <https://doi.org/10.1016/j.jvolgeores.2011.11.005>, 2012.
- Badgley, C., Barry, J. C., Le, M., Morgan, E., Nelson, S. V., Behrensmeyer, A. K., Cerling, T. E., and Pilbeam, D.: Ecological changes in Miocene mammalian record show impact of prolonged climatic forcing, *P. Natl. Acad. Sci. USA* 105, 12145–12149, <https://doi.org/10.1073/pnas.0805592105>, 2008.
- Behrensmeyer, A. K., Quade, J., Cerling, T. E., Kappelman, J., Khan, I. A., Copeland, P., Roe, L., Hicks, J., Stubblefield, P., Willis, B. J., and Latorre, C.: The structure and rate of late Miocene expansion of C₄ plants: Evidence from lateral variation in stable isotopes in paleosols of the Siwalik Group,

- northern Pakistan, *Bull. Geol. Soc. Am.*, 119, 1486–1505, <https://doi.org/10.1130/B26064.1>, 2007.
- Bernor, R. L., Mirzaie Ataabadi, M., Meshida, K., and Wolf, D.: The Maragheh hipparions, late Miocene of Azarbaijan, Iran, *Paleobiodivers. Paleoenviron.*, 96, 453–488, <https://doi.org/10.1007/s12549-016-0235-2>, 2016.
- Bestland, E. A. and Krull, E. S.: Palaeoenvironments of Early Miocene Kisingiri volcano Proconsul sites: evidence from carbon isotopes, palaeosols and hydromagmatic deposits, *J. Geol. Soc. London*, 156, 965–976, <https://doi.org/10.1144/gsjgs.156.5.0965>, 1999.
- Biasatti, D., Bernor, R. L., and Cooper, L. W.: Insights on Late Miocene climate change and regional uplift in Maragheh Basin, eastern Azerbaijan Province, northwest Iran revealed by stable carbon and oxygen isotope analyses of fossil horse tooth enamel, in: 75th Annual Meeting, Society of Vertebrate Paleontology, 89–89, 2015.
- Böhme, M., Spassov, N., Ebner, M., Geraads, D., Hristova, L., Kirscher, U., Kötter, S., Linnemann, U., Prieto, J., Roussiakis, S., Theodorou, G., Uhlir, G., and Winklhofer, M.: Messinian age & savannah environment of the possible hominin *Graecopithecus* from Europe, *PLoS One*, 12, <https://doi.org/10.1371/journal.pone.0177347>, 2017.
- Böhme, M., Spassov, N., Majidifard, M. R., Gärtner, A., Kirscher, U., Marks, M., Dietzel, C., Uhlir, G., El Atfy, H., Begun, D. R., and Winklhofer, M.: Neogene hyperaridity in Arabia drove the directions of mammalian dispersal between Africa and Eurasia, *Commun. Earth Environ.*, 2, 85, <https://doi.org/10.1038/s43247-021-00158-y>, 2021.
- Brocard, G. Y., Meijers, M. J. M., Cosca, M. A., Salles, T., Wilenbring, J., Teyssier, C., and Whitney, D. L.: Fast Pliocene integration of the Central Anatolian Plateau drainage: Evidence, processes, and driving forces, *Geosphere*, 17, 739–765, <https://doi.org/10.1130/GES02247.1>, 2021.
- Butiseacă, G. A., van der Meer, M. T. J., Kontakiotis, G., Agiadi, K., Thivaïou, D., Besiou, E., Antonarakou, A., Mulch, A., and Vasiliev, I.: Multiple crises preceded the Mediterranean Salinity Crisis: Aridification and vegetation changes revealed by biomarkers and stable isotopes, *Glob. Planet Change*, 217, <https://doi.org/10.1016/j.gloplacha.2022.103951>, 2022.
- Caves, J. K., Moragne, D. Y., Ibarra, D. E., Bayshashov, B. U., Gao, Y., Jones, M. M., Zhamangara, A., Arzhannikova, A. V., Arzhannikov, S. G., and Chamberlain, C. P.: The Neogene de-greening of Central Asia, *Geology*, 44, 887–890, <https://doi.org/10.1130/G38267.1>, 2016.
- Cerling, T. E.: The stable isotopic composition of modern soil carbonate and its relationship to climate, *Earth Planet Sc. Lett.*, 71, 229–240, [https://doi.org/10.1016/0012-821X\(84\)90089-X](https://doi.org/10.1016/0012-821X(84)90089-X), 1984.
- Cerling, T. E. and Hay, R. L.: An Isotopic Study of Paleosol Carbonates from Olduvai Gorge, *Quat. Res.*, 25, 63–78, [https://doi.org/10.1016/0033-5894\(86\)90044-X](https://doi.org/10.1016/0033-5894(86)90044-X), 1986.
- Cerling, T. E. and Quade, J.: Stable Carbon and Oxygen Isotopes in Soil Carbonates, *Geophysical. Monograph.*, 78, 217–231, <https://doi.org/10.1029/GM078p0217>, 1993.
- Cerling, T. E., Bowman, J. R., and O’Neil, J. R.: An isotopic study of a fluvial-lacustrine sequence: The Plio-Pleistocene koobi fora sequence, East Africa, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 63, 335–356, [https://doi.org/10.1016/0031-0182\(88\)90104-6](https://doi.org/10.1016/0031-0182(88)90104-6), 1988.
- Cerling, T. E., Quade, J., Ambrose, S. H., and Sikes, N. E.: Fossil soils, grasses, and carbon isotopes from Fort Ternan, Kenya: grassland or woodland?, *J. Hum. Evol.*, 21, 295–306, [https://doi.org/10.1016/0047-2484\(91\)90110-H](https://doi.org/10.1016/0047-2484(91)90110-H), 1991.
- Cerling, T. E., Harris, J. M., MacFadden, B. J., Leakey, M. G., Quade, J., Eisenmann, V., and Ehleringer, J. R.: Global vegetation change through the Miocene/Pliocene boundary, *Nature*, 389, 153–158, <https://doi.org/10.1038/38229>, 1997.
- Cerling, T. E., Harris, J. M., and Leakey, M. G.: 12.2. Isotope Paleoecology of the Nawata and Nachukui Formations at Lothagam, Turkana Basin, Kenya, in: Lothagam, Columbia University Press, 605–624, <https://doi.org/10.7312/leak11870-024>, 2003.
- Cerling, T. E., Wynn, J. G., Andanje, S. A., Bird, M. I., Korir, D. K., Levin, N. E., Mace, W., Macharia, A. N., Quade, J., and Remien, C. H.: Woody cover and hominin environments in the past 6 million years, *Nature*, 476, 51–56, <https://doi.org/10.1038/nature10306>, 2011.
- Crocker, A. J., Naafs, B. D. A., Westerhold, T., James, R. H., Cooper, M. J., Röhl, U., Pancost, R. D., Xuan, C., Osborne, C. P., Beerling, D. J., and Wilson, P. A.: Astronomically controlled aridity in the Sahara since at least 11 million years ago, *Nat. Geosci.*, 15, 671–676, <https://doi.org/10.1038/s41561-022-00990-7>, 2022.
- Denk, T., Güner, T. H., and Grimm, G. W.: From mesic to arid: Leaf epidermal features suggest preadaptation in Miocene dragon trees (*Dracaena*), *Rev. Palaeobot. Palynol.*, 200, 211–228, <https://doi.org/10.1016/j.revpalbo.2013.09.009>, 2014.
- Denk, T., Zohner, C. M., Grimm, G. W., and Renner, S. S.: Plant fossils reveal major biomes occupied by the late Miocene Old-World Pikermian fauna, *Nat. Ecol. Evol.*, 2, 1864–1870, <https://doi.org/10.1038/s41559-018-0695-z>, 2018.
- Ding, Z. L. and Yang, S. L.: C₃/C₄ vegetation evolution over the last 7.0 Myr in the Chinese Loess Plateau: evidence from pedogenic carbonate $\delta^{13}\text{C}$, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 160, 291–299, [https://doi.org/10.1016/S0031-0182\(00\)00076-6](https://doi.org/10.1016/S0031-0182(00)00076-6), 2000.
- Edwards, E. J., Osborne, C. P., Strömberg, C. A. E., Smith, S. A., Bond, W. J., Christin, P. A., Cousins, A. B., Duvall, M. R., Fox, D. L., Freckleton, R. P., Ghannoum, O., Hartwell, J., Huang, Y., Janis, C. M., Keeley, J. E., Kellogg, E. A., Knapp, A. K., Leakey, A. D. B., Nelson, D. M., Saarela, J. M., Sage, R. F., Sala, O. E., Salamin, N., Still, C. J., and Tiplle, B.: The origins of C₄ Grasslands: Integrating evolutionary and ecosystem science, *Science*, 328, 587–591, <https://doi.org/10.1126/science.1177216>, 2010.
- Eronen, J. T., Ataabadi, M. M., Micheels, A., Karme, A., Bernor, R. L., and Fortelius, M.: Distribution history and climatic controls of the Late Miocene Pikermian chronofauna, *P. Natl. Acad. Sci. USA*, 106, 11867–11871, <https://doi.org/10.1073/pnas.0902598106>, 2009.
- Feakins, S. J., Levin, N. E., Liddy, H. M., Sieracki, A., Eglinton, T. I., and Bonnefille, R.: Northeast african vegetation change over 12 m.y., *Geology*, 41, 295–298, <https://doi.org/10.1130/G33845.1>, 2013.
- Feurdean, A. and Vasiliev, I.: The contribution of fire to the late Miocene spread of grasslands in eastern Eurasia (Black Sea re-

- gion), *Sci. Rep.*, 9, <https://doi.org/10.1038/s41598-019-43094-w>, 2019.
- Fortelius, M., Bibi, F., Tang, H., Žliobaitė, I., Eronen, J. T., and Kaya, F.: The nature of the Old World savannah palaeobiome, *Nat. Ecol. Evol.*, 3, 504, <https://doi.org/10.1038/s41559-019-0857-7>, 2019.
- Fox, D. L., Pau, S., Taylor, L., Strömberg, C. A. E., Osborne, C. P., Bradshaw, C., Conn, S., Beerling, D. J., and Still, C. J.: Climatic Controls on C₄ Grassland Distributions During the Neogene: A Model-Data Comparison, *Front. Ecol. Evol.*, 6, <https://doi.org/10.3389/fevo.2018.00147>, 2018.
- Ghosh, P., Padia, J. T., and Mohindra, R.: Stable isotopic studies of palaeosol sediment from Upper Siwalik of Himachal Himalaya: Evidence for high monsoonal intensity during late Miocene?, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 206, 103–114, <https://doi.org/10.1016/j.palaeo.2004.01.014>, 2004.
- Haug, G. H. and Tiedemann, R.: Effect of the formation of the Isthmus of Panama on Atlantic Ocean thermohaline circulation, *Nature*, 393, 673–676, <https://doi.org/10.1038/31447>, 1998.
- Herbert, T. D., Lawrence, K. T., Tzanova, A., Peterson, L. C., Caballero-Gill, R., and Kelly, C. S.: Late Miocene global cooling and the rise of modern ecosystems, *Nat. Geosci.*, 9, 843–847, <https://doi.org/10.1038/ngeo2813>, 2016.
- Holbourn, A. E., Kuhnt, W., Clemens, S. C., Kochhann, K. G. D., Jöhnck, J., Lübbers, J., and Andersen, N.: Late Miocene climate cooling and intensification of southeast Asian winter monsoon, *Nat. Commun.*, 9, 1584, <https://doi.org/10.1038/s41467-018-03950-1>, 2018.
- Huang, S., Meijers, M. J. M., Eyres, A., Mulch, A., and Fritz, S. A.: Unravelling the history of biodiversity in mountain ranges through integrating geology and biogeography, *J. Biogeogr.*, 46, 1777–1791, <https://doi.org/10.1111/jbi.13622>, 2019.
- Huang, Y., Clemens, S. C., Liu, W., Wang, Y., and Prell, W. L.: Large-scale hydrological change drove the late Miocene C₄ plant expansion in the Himalayan foreland and Arabian Peninsula, *Geology*, 35, 531, <https://doi.org/10.1130/G23666A.1>, 2007.
- Kaakinen, A., Sonninen, E., and Lunkka, J. P.: Stable isotope record in paleosol carbonates from the Chinese Loess Plateau: Implications for late Neogene paleoclimate and paleovegetation, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 237, 359–369, <https://doi.org/10.1016/j.palaeo.2005.12.011>, 2006.
- Karas, C., Nürnberg, D., Bahr, A., Groeneveld, J., Herrle, J. O., Tiedemann, R., and Demenocal, P. B.: Pliocene oceanic seaways and global climate, *Sci. Rep.*, 7, <https://doi.org/10.1038/srep39842>, 2017.
- Kaya, F., Bibi, F., Žliobaite, I., Eronen, J. T., Hui, T., and Fortelius, M.: The rise and fall of the Old World savannah fauna and the origins of the African savannah biome, *Nat. Ecol. Evol.*, 2, 241–246, <https://doi.org/10.1038/s41559-017-0414-1>, 2018.
- Kayseri-Özer, M. S.: Cenozoic vegetation and climate change in Anatolia – A study based on the IPR-vegetation analysis, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 467, 37–68, <https://doi.org/10.1016/j.palaeo.2016.10.001>, 2017.
- Kayseri-Özer, M. S., Karadenizli, L., Akgün, F., Oyal, N., Saraç, G., Şen, Ş., Tunçoğlu, C., and Tuncer, A.: Palaeoclimatic and palaeoenvironmental interpretations of the Late Oligocene, Late Miocene–Early Pliocene in the Çankırı–Çorum Basin, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 467, 16–36, <https://doi.org/10.1016/j.palaeo.2016.05.022>, 2017.
- Kingston, J.: Stable isotopic evidence for hominid paleoenvironments in East Africa, PhD thesis, Harvard University, 1–162, ISBN 9798641957470, 1992.
- Knapp, A. K., Chen, A., Griffin-Nolan, R. J., Baur, L. E., Carroll, C. J. W., Gray, J. E., Hoffman, A. M., Li, X., Post, A. K., Slette, I. J., Collins, S. L., Luo, Y., and Smith, M. D.: Resolving the Dust Bowl paradox of grassland responses to extreme drought, *P. Natl. Acad. Sci. USA*, 117, 22249–22255, <https://doi.org/10.1073/pnas.1922030117>, 2020.
- Kohn, M. J.: Carbon isotope compositions of terrestrial C₃ plants as indicators of (paleo)ecology and (paleo)climate, *Proceedings of the National Academy of Sciences*, 107, 19691–19695, <https://doi.org/10.1073/pnas.1004933107>, 2010.
- Kovar-Eder, J.: Vegetation dynamics in Europe during the Neogene, in: *Distribution and Migration of Tertiary Mammals in Eurasia. A volume in Honour of Hans de Bruijn*, edited by: Reumer, J. W. F. and Wessels, W., DEINSEA, Rotterdam, 373–392, ISSN: 0923-9308, 2003.
- Lepetit, P.: Kohlenstoff-Isotopie miozäner Calcretes in Kappadokien (Türkei), PhD thesis, Friedrich-Schiller-Universität Jena, Jena, 1–218, https://www.db-thueringen.de/receive/dbt_mods_00017261 (last access: 29 April 2026), 2010.
- Levin, N. E., Quade, J., Simpson, S. W., Semaw, S., and Rogers, M.: Isotopic evidence for Plio–Pleistocene environmental change at Gona, Ethiopia, *Earth Planet Sci. Lett.*, 219, 93–110, [https://doi.org/10.1016/S0012-821X\(03\)00707-6](https://doi.org/10.1016/S0012-821X(03)00707-6), 2004.
- Levin, N. E., Brown, F. H., Behrensmeier, A. K., Bobe, R., and Cerling, T. E.: Paleosol carbonates from the Omo Group: Isotopic records of local and regional environmental change in East Africa, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 307, 75–89, <https://doi.org/10.1016/j.palaeo.2011.04.026>, 2011.
- Licht, A., Dupont-Nivet, G., Meijer, N., Caves Rugenstein, J., Schauer, A., Fiebig, J., Mulch, A., Hoorn, C., Barbolini, N., and Guo, Z.: Decline of soil respiration in north-eastern Tibet through the transition into the Oligocene icehouse, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 560, <https://doi.org/10.1016/j.palaeo.2020.110016>, 2020.
- Lüdecke, T., Schrenk, F., Thiemeyer, H., Kullmer, O., Bromage, T. G., Sandrock, O., Fiebig, J., and Mulch, A.: Persistent C₃ vegetation accompanied Plio–Pleistocene hominin evolution in the Malawi Rift (Chiwondo Beds, Malawi), *J. Hum. Evol.*, 90, 163–175, <https://doi.org/10.1016/j.jhevol.2015.10.014>, 2016.
- Mayser, J. P., Flecker, R., Marzocchi, A., Kouwenhoven, T. J., Lunt, D. J., and Pancost, R. D.: Precession driven changes in terrestrial organic matter input to the Eastern Mediterranean leading up to the Messinian Salinity Crisis, *Earth Planet Sci. Lett.*, 462, 199–211, <https://doi.org/10.1016/j.epsl.2017.01.029>, 2017.
- Meijers, M. J. M., Brocard, G. Y., Cosca, M. A., Lüdecke, T., Teyssier, C., Whitney, D. L., and Mulch, A.: Rapid late Miocene surface uplift of the Central Anatolian Plateau margin, *Earth Planet Sci. Lett.*, 497, 29–41, <https://doi.org/10.1016/j.epsl.2018.05.040>, 2018.
- Meijers, M. J. M., Brocard, G. Y., Whitney, D. L., and Mulch, A.: Palaeoenvironmental conditions and drainage evolution of the central Anatolian lake system (Turkey) during Late Miocene to Pliocene surface uplift, *Geosphere*, 16, 490–509, <https://doi.org/10.1130/GES02135.1>, 2020.
- Meijers, M. J. M., Mikes, T., Rojay, B., Aydar, E., Çubukçu, H. E., Wagner, T., Lüdecke, T., Krsnik, E., Fiebig, J., and

- Mulch, A.: Miocene decoupling of surface uplift of the Central Anatolian Plateau interior (Türkiye) and its modern mountainous margins, *Earth Planet. Sci. Lett.*, 663, 119414, <https://doi.org/10.1016/j.epsl.2025.119414>, 2025.
- Mosbrugger, V. and Utescher, T.: The coexistence approach – a method for quantitative reconstructions of Tertiary terrestrial palaeoclimate data using plant fossils, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 134, 61–86, [https://doi.org/10.1016/S0031-0182\(96\)00154-X](https://doi.org/10.1016/S0031-0182(96)00154-X), 1997.
- Mosbrugger, V., Utescher, T., and Dilcher, D. L.: Cenozoic continental climatic evolution of Central Europe, *P. Natl. Acad. Sci. USA*, 102, 14964–14969, <https://doi.org/10.1073/pnas.0505267102>, 2005.
- Özsayin, E., Çiner, T. A., Rojay, F. B., Dirik, R. K., Melnick, D., Fernandez-Blanco, D., Bertotti, G., Schildgen, T. F., Garcin, Y., Strecker, M. R., and Sudo, M.: Plio-Quaternary extensional tectonics of the Central Anatolian Plateau: a case study from the Tuz Gölü Basin, Turkey, *Turk. J. Earth Sci.*, <https://doi.org/10.3906/yer-1210-5>, 2013.
- Passey, B. H., Ayliffe, L. K., Kaakinen, A., Zhang, Z., Eronen, J. T., Zhu, Y., Zhou, L., Cerling, T. E., and Fortelius, M.: Strengthened East Asian summer monsoons during a period of high-latitude warmth? Isotopic evidence from Mio-Pliocene fossil mammals and soil carbonates from northern China, *Earth Planet. Sci. Lett.*, 277, 443–452, <https://doi.org/10.1016/j.epsl.2008.11.008>, 2009.
- Peppe, D. J., Cote, S. M., Deino, A. L., Fox, D. L., Kingston, J. D., Kinyanjui, R. N., Lukens, W. E., MacLatchy, L. M., Novello, A., Strömberg, C. A. E., Driese, S. G., Garrett, N. D., Hillis, K. R., Jacobs, B. F., Jenkins, K. E. H., Kityo, R. M., Lehmann, T., Manthi, F. K., Mbua, E. N., Michel, L. A., Miller, E. R., Mugume, A. A. T., Muteti, S. N., Nengo, I. O., Oginga, K. O., Phelps, S. R., Polissar, P., Rossie, J. B., Stevens, N. J., Uno, K. T., and McNulty, K. P.: Oldest evidence of abundant C₄ grasses and habitat heterogeneity in eastern Africa, *Science*, 380, 173–177, <https://doi.org/10.1126/science.abq2834>, 2023.
- Plummer, T., Bishop, L. C., Ditchfield, P., and Hicks, J.: Research on Late Pliocene Oldowan Sites at Kanjera South, Kenya, *J. Hum. Evol.*, 36, 151–170, <https://doi.org/10.1006/jhev.1998.0256>, 1999.
- Plummer, T. W., Ditchfield, P. W., Bishop, L. C., Kingston, J. D., Ferraro, J. V., Braun, D. R., Hertel, F., and Potts, R.: Oldest Evidence of Toolmaking Hominins in a Grassland-Dominated Ecosystem, *PLoS One*, 4, e7199, <https://doi.org/10.1371/journal.pone.0007199>, 2009.
- Polissar, P. J., Rose, C., Uno, K. T., Phelps, S. R., and deMenocal, P.: Synchronous rise of African C₄ ecosystems 10 million years ago in the absence of aridification, *Nat. Geosci.*, 12, 657–660, <https://doi.org/10.1038/s41561-019-0399-2>, 2019.
- Quade, J. and Cerling, T. E.: Expansion of C₄ grasses in the Late Miocene of Northern Pakistan: evidence from stable isotopes in paleosols, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 115, 91–116, [https://doi.org/10.1016/0031-0182\(94\)00108-K](https://doi.org/10.1016/0031-0182(94)00108-K), 1995.
- Quade, J., Solounias, N., and Cerling, T. E.: Stable isotopic evidence from paleosol carbonates and fossil teeth in Greece for forest or woodlands over the past 11 Ma, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 108, 41–53, [https://doi.org/10.1016/0031-0182\(94\)90021-3](https://doi.org/10.1016/0031-0182(94)90021-3), 1994.
- Quade, J., Levin, N., Semaw, S., Stout, D., Renne, P., Rogers, M., and Simpson, S.: Paleoenvironments of the earliest stone tool-makers, Gona, Ethiopia, *Geol. Soc. Am. Bull.*, 116, 1529–1544, <https://doi.org/10.1130/B25358.1>, 2004.
- Quan, C., Liu, Y. S. C., Tang, H., and Utescher, T.: Miocene shift of European atmospheric circulation from trade wind to westerlies, *Sci. Rep.*, 4, <https://doi.org/10.1038/srep05660>, 2014.
- Quinn, R. L., Lepre, C. J., Wright, J. D., and Feibel, C. S.: Paleogeographic variations of pedogenic carbonate $\delta^{13}\text{C}$ values from Koobi Fora, Kenya: implications for floral compositions of Plio-Pleistocene hominin environments, *J. Hum. Evol.*, 53, 560–573, <https://doi.org/10.1016/j.jhevol.2007.01.013>, 2007.
- Rey, K., Amiot, R., Lécuyer, C., Koufos, G. D., Martineau, F., Fourel, F., Kostopoulos, D. S., and Merceron, G.: Late Miocene climatic and environmental variations in northern Greece inferred from stable isotope compositions ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$) of equid teeth apatite, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 388, 48–57, <https://doi.org/10.1016/j.palaeo.2013.07.021>, 2013.
- Roveri, M., Flecker, R., Krijgsman, W., Lofi, J., Lugli, S., Manzi, V., Sierro, F. J., Bertini, A., Camerlenghi, A., De Lange, G., Govers, R., Hilgen, F. J., Hübscher, C., Meijer, P. T., and Stonica, M.: The Messinian Salinity Crisis: Past and future of a great challenge for marine sciences, *Mar. Geol.*, 352, 25–58, <https://doi.org/10.1016/j.margeo.2014.02.002>, 2014.
- Sage, R. F.: The evolution of C₄ photosynthesis, *New Phytol.*, 161, 341–370, <https://doi.org/10.1111/j.1469-8137.2004.00974.x>, 2004.
- Sahnouni, M., Van der Made, J., and Everett, M.: Ecological background to Plio-Pleistocene hominin occupation in North Africa: the vertebrate faunas from Ain Boucherit, Ain Hanech and El-Kherba, and paleosol stable-carbon-isotope studies from El-Kherba, Algeria, *Quat. Sci. Rev.*, 30, 1303–1317, <https://doi.org/10.1016/j.quascirev.2010.01.002>, 2011.
- Sanyal, P., Bhattacharya, S. K., Kumar, R., Ghosh, S. K., and Sangode, S. J.: Mio-Pliocene monsoonal record from Himalayan foreland basin (Indian Siwalik) and its relation to vegetational change, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 205, 23–41, <https://doi.org/10.1016/j.palaeo.2003.11.013>, 2004.
- Schemmel, F., Mikes, T., Rojay, B., and Mulch, A.: The impact of topography on isotopes in precipitation across the central Anatolian plateau (Turkey), *Am. J. Sci.*, 313, 61–80, <https://doi.org/10.2475/02.2013.01.2013>.
- Sikes, N. E.: Early hominid habitat preferences in East Africa: Paleosol carbon isotopic evidence, *J. Hum. Evol.*, 27, 25–45, <https://doi.org/10.1006/jhev.1994.1034>, 1994.
- Sikes, N. E. and Ashley, G. M.: Stable isotopes of pedogenic carbonates as indicators of paleoecology in the Plio-Pleistocene (upper Bed I), western margin of the Olduvai Basin, Tanzania, *J. Hum. Evol.*, 53, 574–594, <https://doi.org/10.1016/j.jhevol.2006.12.008>, 2007.
- Sikes, N. E., Potts, R., and Behrensmeyer, A. K.: Early Pleistocene habitat in Member 1 Olororgesailie based on paleosol stable isotopes, *J. Hum. Evol.*, 37, 721–746, <https://doi.org/10.1006/jhev.1999.0343>, 1999.
- Spötl, C. and Vennemann, T. W.: Continuous-flow isotope ratio mass spectrometric analysis of carbonate minerals, *Rapid Commun. Mass Sp.*, 17, 1004–1006, <https://doi.org/10.1002/rcm.1010>, 2003.
- Still, C. J., Berry, J. A., Collatz, G. J., and DeFries, R. S.: Global distribution of C₃ and C₄ vegetation: Car-

- bon cycle implications, *Global Biogeochem. Cycles*, 17, <https://doi.org/10.1029/2001gb001807>, 2003.
- Strömberg, C. A. E.: Evolution of grasses and grassland ecosystems, *Annu. Rev. Earth Planet Sci.*, 39, 517–544, <https://doi.org/10.1146/annurev-earth-040809-152402>, 2011.
- Strömberg, C. A. E., Werdelin, L., Friis, E. M., and Saraç, G.: The spread of grass-dominated habitats in Turkey and surrounding areas during the Cenozoic: Phytolith evidence, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 250, 18–49, <https://doi.org/10.1016/j.palaeo.2007.02.012>, 2007.
- Tauxe, L. and Feakins, S. J.: A Reassessment of the Chronostratigraphy of Late Miocene C₃–C₄ Transitions, *Palaeogeogr. Palaeoclimatol.*, 35, <https://doi.org/10.1029/2020PA003857>, 2020.
- Tipple, B. J. and Pagani, M.: The early origins of terrestrial C₄ photosynthesis, *Annu. Rev. Earth Planet Sci.*, 35, 435–461, <https://doi.org/10.1146/annurev.earth.35.031306.140150>, 2007.
- Türkeş, M. and Erlat, E.: Climatological responses of winter precipitation in Turkey to variability of the North Atlantic Oscillation during the period 1930–2001, *Theor. Appl. Climatol.*, 81, 45–69, <https://doi.org/10.1007/s00704-004-0084-1>, 2005.
- Ulu, Ü.: 1 : 500 000 Geological Map of Turkey (Adana), General Directorate of Mining Research and Exploration (MTA), 2002.
- Uno, K. T., Cerling, T. E., Harris, J. M., Kunimatsu, Y., Leakey, M. G., Nakatsukasa, M., and Nakaya, H.: Late Miocene to Pliocene carbon isotope record of differential diet change among East African herbivores, *P. Natl. Acad. Sci. USA*, 108, 6509–6514, <https://doi.org/10.1073/pnas.1018435108>, 2011.
- Urban, M. A., Nelson, D. M., Jimenez-Moreno, G., Chateaufneuf, J.-J., Pearson, A., and Hu, F. S.: Isotopic evidence of C₄ grasses in southwestern Europe during the Early Oligocene-Middle Miocene, *Geology*, 38, 1091–1094, <https://doi.org/10.1130/G31117.1>, 2010.
- Velitzelos, D., Bouchal, J. M., and Denk, T.: Review of the Cenozoic floras and vegetation of Greece, *Rev. Palaeobot. Palyno.*, 204, 56–117, <https://doi.org/10.1016/j.revpalbo.2014.02.006>, 2014.
- Wen, Y., Zhang, L., Holbourn, A. E., Zhu, C., Huntington, K. W., Jin, T., Li, Y., and Wang, C.: CO₂-forced Late Miocene cooling and ecosystem reorganizations in East Asia, *P. Natl. Acad. Sci. USA*, 120, <https://doi.org/10.1073/pnas.2214655120>, 2023.
- Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnet, J. S. K., Bohaty, S. M., De Vleeschouwer, D., Florindo, F., Frederichs, T., Hodell, D. A., Holbourn, A. E., Kroon, D., Lauretano, V., Littler, K., Lourens, L. J., Lyle, M., Pälike, H., Röhl, U., Tian, J., Wilkens, R. H., Wilson, P. A., and Zachos, J. C.: An astronomically dated record of Earth's climate and its predictability over the last 66 million years, *Science*, 369, 1383–1387, <https://doi.org/10.1126/science.aba6853>, 2020.
- WoldeGabriel, G., Ambrose, S. H., Barboni, D., Bonnefille, R., Bremond, L., Currie, B., DeGusta, D., Hart, W. K., Murray, A. M., Renne, P. R., Jolly-Saad, M. C., Stewart, K. M., and White, T. D.: The Geological, Isotopic, Botanical, Invertebrate, and Lower Vertebrate Surroundings of *Ardipithecus ramidus*, *Science*, 326, 65, <https://doi.org/10.1126/science.1175817>, 2009.
- Wynn, J. G.: Paleosols, stable carbon isotopes, and paleoenvironmental interpretation of Kanapoi, Northern Kenya, *J. Hum. Evol.*, 39, 411–432, <https://doi.org/10.1006/jhev.2000.0431>, 2000.
- Wynn, J. G.: Influence of Plio-Pleistocene aridification on human evolution: Evidence from paleosols of the Turkana Basin, Kenya, *Am. J. Phys. Anthropol.*, 123, 106–118, <https://doi.org/10.1002/ajpa.10317>, 2004.
- Wynn, J. G., Alemseged, Z., Bobe, R., Geraads, D., Reed, D., and Roman, D. C.: Geological and palaeontological context of a Pliocene juvenile hominin at Dikika, Ethiopia, *Nature*, 443, 332–336, <https://doi.org/10.1038/nature05048>, 2006.
- Yao, Z., Xiao, G., Wu, H., Liu, W., and Chen, Y.: Plio-Pleistocene vegetation changes in the North China Plain: Magnetostratigraphy, oxygen and carbon isotopic composition of pedogenic carbonates, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 297, 502–510, <https://doi.org/10.1016/j.palaeo.2010.09.003>, 2010.
- Zamanian, K., Pustovoytov, K., and Kuzyakov, Y.: Pedogenic carbonates: Forms and formation processes, *Earth-Sci. Rev.*, 157, 1–17, <https://doi.org/10.1016/j.earscirev.2016.03.003>, 2016.
- Zhuang, G., Hourigan, J. K., Koch, P. L., Ritts, B. D., and Kent-Corson, M. L.: Isotopic constraints on intensified aridity in Central Asia around 12Ma, *Earth Planet Sci. Lett.*, 312, 152–163, <https://doi.org/10.1016/j.epsl.2011.10.005>, 2011.