

This discussion paper is/has been under review for the journal *Climate of the Past* (CP).
Please refer to the corresponding final paper in CP if available.

Changing climatic response: a conceptual model for glacial cycles and the Mid-Pleistocene Transition

I. Daruka¹ and P. D. Ditlevsen²

¹Johannes Kepler University, Institute of Semiconductor and Solid State Physics,
Altenbergerstrasse 69, 4040 Linz, Austria

²Centre for Ice and Climate, Niels Bohr Institute, University of Copenhagen,
Juliane Maries Vej 30, 2100 Copenhagen, Denmark

Received: 4 February 2014 – Accepted: 18 February 2014 – Published: 13 March 2014

Correspondence to: I. Daruka (istvan.daruka@jku.at) and P. D. Ditlevsen (pditlev@nbi.ku.dk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Milankovitch's astronomical theory of glacial cycles, attributing ice age climate oscillations to orbital changes in Northern Northern-Hemisphere insolation, is challenged by the paleoclimatic record. The climatic response to the variations in insolation is far from trivial. In general the glacial cycles are highly asymmetric in time, with slow cooling from the interglacials to the glacials (inceptions) and very rapid warming from the glacials to the interglacials (terminations). We shall refer to this fast-slow dynamics as the "saw-tooth" shape of the paleoclimatic record. This is non-linearly related to the time-symmetric variations in the orbital forcing. However, the most pronounced challenge to the Milankovitch theory is the Mid-Pleistocene Transition (MPT) occurring about one million years ago. During that event, the prevailing 41 kyr glacial cycles, corresponding to the almost harmonic obliquity cycle were replaced by longer saw-tooth shaped cycles with a time scale around 100 kyr. The MPT must have been driven by internal changes in climate response, since it does not correspond to any apparent changes in the orbital forcing. In order to identify possible mechanisms causing the observed changes in glacial dynamics, it is relevant to study simplified models with the capability of generating temporal behavior similar to the observed records. We present a simple oscillator type model approach, with two variables, a temperature anomaly and an ice volume analogous, climatic memory term. The generalization of the ice albedo feedback is included in terms of an effective multiplicative coupling between this latter climatic memory term (representing the internal degrees of freedom) and the external drive. The simple model reproduces the temporal asymmetry of the late Pleistocene glacial cycles and suggests that the MPT can be explained as a regime shift, aided by climatic noise, from a period 1 frequency locking to the obliquity cycle to a period 2–3 frequency locking to the same obliquity cycle. The change in dynamics has been suggested to be a result of a slow gradual decrease in atmospheric greenhouse gas concentration. The presence of chaos in

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



changes due to precession and obliquity. This is referred to as the 100 kyr problem of the Milankovich theory (Imbrie et al., 1993). The solution to this problem is probably that the 100 kyr world is not paced by eccentricity, which is also why the 400 kyr modulation to the eccentricity is not seen in the climatic response. This is referred to as the 400 kyr problem or the Stage 11 (MIS 11) problem: marine isotope stage 11 should not have been an interglacial if changes in eccentricity were the driver. It has been suggested that the 100 kyr world could rather be seen as multiplets of the 41 kyr obliquity cycle, such that they are approximately 80 and 120 kyr long, occurring in a more or less alternating way (Huybers, 2007; Ditlevsen, 2009). Note in Fig. 1 that the duration between the warm states around 200 kyr BP (MIS 7 and MIS 6) is approximately 40 kyr.

Since there are no apparent changes in the astronomical forcing at the MPT, the transition must be governed by the internal dynamical response to the forcing. The hypothesis is that a gradual change in some environmental parameter of the system led to a dynamical change in the response to the orbital forcing. Two main hypothesis have been put forward for the change in environment; either a slow decrease in atmospheric $p\text{CO}_2$ (Saltzman and Maasch, 1991) led to reduced greenhouse warming and the possibility of deeper glaciations, or slow glacial erosion of the regolith under the glaciers, such that the glaciers after the MPT would grow on the bedrock, which permits higher glaciers to be stable (Clark and Pollard, 1998). For a review see Clark et al. (2006). Here we shall argue by introducing a semi-conceptual driven climate oscillator model that the MPT could be a result of frequency locking to the orbital forcing, such that a slight change in parameters can alter the period of the frequency locking.

The phenomenon of phase-locking for explaining climatic response to orbital forcing has been suggested before. De Saedeleer et al. (2013) discuss in terms of the Van der Pol oscillator the possibility of synchronizing to spectral components of the 65°N summer solstice insolation. These are expressed in terms of a d'Alembert series resulting from the perturbative calculation of the orbital parameters (Berger, 1978). Here we are not so much concerned with the problem of which component of the orbital forcing the climate system is most sensitive. Since the different orbital parameters

influence the insolation very differently in both seasonality and latitudinal variation it is unlikely that a single time series (with annual or less resolution) is sufficient to give the full account of the Pleistocene glacial cycles.

Still it is an open problem to which extent the global stack marine isotope record itself is sufficient to discriminate between the suggested low dimensional or conceptual models of the Pleistocene glacial cycles. A decisive model of the MPT, probably have to wait for future comprehensive climate model simulations and strong improvements in the regional climate reconstructions. Exploring alternative mechanisms in simple and conceptual models are thus important both for identifying robust features and for discriminating through hypothesis testing.

2 Existing models

In order to put our model in perspective, we review in the following some of the proposed models and their limitations. This is by no means a comprehensive review. The models of glacial cycles can roughly be categorized in two types: firstly, in mono- or multi-stable state models where the periodicity is solely a result of the periodicity of the orbital forcing. Secondly, in climate oscillator models where the glacial cycles in one way or another result from the internal oscillator resonating with the orbital forcing. For a thorough review of oscillator type models see Crucifix (2012).

The classical energy balance models (Budyko, 1969; Sellers, 1969) belong to the first category. In these models the ice-albedo feedback results in a two-state system, a present climate state and a glaciated state, which today is considered more realistically to describe the Snowball Earth climate. The relative weakness in the magnitude of the 100 kyr eccentricity cycle in insolation led to the introduction of the concept of stochastic resonance (Benzi et al., 1982).

For contrasting the glacial dynamics before and after the MPT a semi empirical model was proposed by Paillard (1998). The model is a rule based threshold model with three possible stable climate states, an interglacial, a mild glacial and a full glacial. With

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changing climatic response

I. Daruka and
P. D. Ditlevsen

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

specific rules of transitions and a slow change of threshold, the model reproduces the Pleistocene record including the MPT as a response to the Milankovitch forcing. An empirical dynamical model, with a three stable state bifurcation diagram, explaining the specific rules of transitions in the Paillard model has been proposed (Ditlevsen, 2009). In this model a change of the bifurcation structure causes the MPT. A different approach was taken in Huybers and Wunsch (2005) for explaining the 100 kyr world as a result of the 41 kyr orbital forcing. This is a stochastic threshold model with a linear drift towards glaciation. When a threshold proportional to the obliquity is reached the glaciation terminates and the climate is reset to the interglacial condition. This model naturally reproduces the sawtooth shape of the climate curve. The linear drift assumes a very long internal time scale of the order 100 kyr for glaciation. By a simple rescaling of this time scale the model can also reproduce the 41 kyr world, however, with more time asymmetric (sawtooth shaped) glaciations, than observations indicate.

Many different types of oscillator models have been suggested: The dynamics of ice sheets is roughly: Higher temperature \rightarrow higher accumulation \rightarrow growing ice sheet \rightarrow high albedo \rightarrow lower temperature (Kallen et al., 1979; Tziperman and Gildor, 2003). This results in free oscillations of the order 5–15 kyr, which are too fast to account for the glacial cycles. The non-linear response in this model to beat periods (combination tones) between the 19 and the 23 kyr precessional frequencies has been suggested to explain the 100 kyr glacial cycles (Le Treut and Ghil, 1983). To obtain free climate oscillations of as long a duration as 100 kyr, the combined effect of isostatic rebound and reduced accumulation with a high ice sheet (the elevation desert effect) is another suggestion (Hyde and Peltier, 1985). None of these models attempt to explain the MPT shift in glacial periods. It was suggested by Saltzman and Maasch (1991) that the ice sheet growth is controlled by the deep-ocean temperature: low ocean temperature \rightarrow higher uptake of atmospheric CO_2 \rightarrow less greenhouse warming \rightarrow growth of ice sheets and sea ice \rightarrow reduced meridional oceanic heat transport \rightarrow higher ocean temperature. In the model this is a limit cycle, which is initiated at the MPT through a Hopf bifurcation as a result of a slow decrease in CO_2 from increased weathering.

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

This model is not concerned with the 41 kyr climate oscillations prior to the MPT. A switch mechanism involving the sea-ice was proposed by Gildor and Tziperman (GT) (Tziperman and Gildor, 2003). In this GT-model the feedback loop is: high sea-ice cover (limited by the warm mid-latitude ocean) → reduced atmospheric temperature and precipitation (accumulation) over the ice sheets → negative mass balance → rapid retreat of the ice sheets → reduced albedo → increased temperature → rapid retreat of the sea-ice → increased precipitation and a positive mass balance. This is a relaxation oscillator, where the 100 kyr cycle is internally driven and independent of the orbital forcing. The time scale for growth of the ice sheets estimated as V/Acc , where V is the volume of the ice sheet and Acc is the accumulation (in suitable units) is of the order 10–30 kyr. The time scale of 100 kyr glacial cycles in the GT-model is rather estimated from $V/(\text{Acc} - \text{Abl})$, where $(\text{Acc} - \text{Abl})$ is the difference between accumulation and ablation in the growth phase. In principle this estimate is not well constrained, since in the case of almost mass balance this growing time is infinite. If the deep ocean in the model is warm enough, the sea-ice switch is not active, and the model oscillates linearly with the orbital forcing. Gradual cooling of the deep ocean, crossing a threshold at MPT activates the sea-ice switch mechanism and the 100 kyr oscillations of the late Pleistocene period. Alternatively, the 41 kyr world could be self-sustained oscillations, perhaps locked to the obliquity cycles (Ashkenazy and Tziperman, 2004).

All the above suggested physical mechanisms are potentially at play in the climate system, thus it is difficult to assess the relative importance without realistic quantitative modelling, which at present is computationally prohibited. However, identifying possible dynamical mechanisms for explaining the observed record merits the more conceptual model approaches. Rial (2004) suggested a logistic-delayed differential equation for the ice volume coupled, through a carrying capacity, to a temperature. The temperature is determined by energy balance between incoming and outgoing radiation. By changing the coupling via the carrying capacity, the MPT is reproduced. The underlying physical mechanism is difficult to identify, partly because delay-equations are notoriously difficult to analyse. An even simpler approach was taken by Huybers (2009), in which the

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Pleistocene climate is described by a discrete map, where the ice volume depends on the ice volume with a lag of 9 kyr, which is approximately the lag between climate oscillations and their derivative in a 40 kyr harmonic cycle. The hypothesis is that the late Pleistocene cycles are purely chaotic, while the 41 kyr cycle prior to the MPT are results of coincidental oscillations near an unstable period 2 cycle in the map. However, within the model framework, the observed long sequence of 41 kyr cycles prior to the MPT seems highly unlikely.

The conceptual modelling is also our approach here, where we shall argue that the MPT could be a change in the internal dynamics leading to a change in frequency locking to the obliquity cycle. This would imply a fundamental limitation in climate predictability, such as inferences about the next inception. We do, however, find this type of un-falsifiable very long time predictions into the future rather academic, but the suggested critical dependence on model parameters is potentially an important guideline for more realistic future model simulations and theories of the Pleistocene glacial dynamics.

3 Model

As a minimal modelling approach we explore a two variable non-linear oscillator model, containing a scaled temperature anomaly $\tau = (T - T_0)/T_0$, where T is a large scale (Northern Hemisphere or global) mean surface temperature, and T_0 is the long term mean. The other variable x was inspired by the the global ice mass anomaly. However, the approaches of Saltzman and Maasch (1991) and that of Tziperman and Gildor (2003) clearly indicate that besides the ice volume, also the deep sea temperature and possible other factors play a decisive role in determining the internal climate dynamics, leading to coupled and nonlinear climatic memory effects. It is possible that if two (or more) interacting processes are involved in shaping the climatic memory, it might not possess a unique characteristic time, and also the climatic state dependent effective internal timescale might be much longer than any of the involved characteristic times

due to the potential multiplicative coupling. With the integral variable x we mimic such integrated climatic memory effects.

In the light of the above, we consider a simple integrative relation between the effective climatic memory term x and the temperature anomaly

$$\dot{x}(t) = \lambda\tau(t), \quad (1)$$

where λ^{-1} represents the timescale unit of the climatic memory. The paleoclimatic record strongly indicate that the climate can be in one of more possible stable states. Within a two variable model, not resolving this multi-state dynamics, the evolution of the temperature anomaly is represented by an effective climate potential $V(x)$ which could possess different local minima and a damping of the anomaly ($-\kappa\tau$). The interaction between the internal dynamics and the external drive is modeled as a multiplicative coupling between the solar insolation $A(t)$ and the climatic memory term $x(t)$:

$$\dot{x}(t) = -V'(x(t)) - \kappa\tau(t) - x(t)A(t) + \sigma\eta(t), \quad (2)$$

where $'$ denotes derivative with respect to the argument. The last term is a climatic noise term to account for unresolved processes, where $\eta(t)$ represents a white noise contribution with $\langle\eta(t)\eta(t')\rangle = \delta_{t,t'}$.

Furthermore, we define

$$V(x) = \alpha x - x^2/2 + x^4/4 \quad (3)$$

as the effective climate potential. This is the simplest non-trivial polynomial multiple state potential, which can contain two minima representing two distinct stable climate states. We note that the non-symmetric, non-harmonic nature of the implemented climate potential renders no characteristic timescale to the climatic response. Furthermore, note that this potential is not the potential derived from the classical Budyko–Sellers energy balance model (Budyko, 1969), now believed to describe the Snowball Earth dynamics. The energy balance model includes the ice albedo effect in

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the potential, while we include it in the forcing term $-x(t)A(t)$ involving a multiplicative coupling.

The model described by Eqs. (1)–(3) is a forced non-linear oscillator, including a multiplicative coupling between the internal dynamics (climatic memory effects) and the external forcing and can possess a two state effective climatic potential (Eq. 2).

We will however not be concerned with the strongly non-linear regime, where multiple steady states are possible. On the contrary, we will throughout the rest of the paper use $\alpha = 0.8$ in Eq. (2), thus $V(x)$ is a non-harmonic skewed potential with a single minimum. The major effect of the non-harmonic nature of the potential is that the frequency of free oscillations (in case of no damping and no forcing) depends on the amplitude of oscillation. Thus the system does not have a natural internal frequency of oscillation.

The external drive has a period (time-scale) $T_{\text{ext}} = 2\pi/\omega$, where ω is the dominant frequency of the orbital forcing. In this work we shall not so much be concerned with the multi-frequency nature of the orbital forcing (De Saedeleer et al., 2013). The orbital changes in insolation strongly depends on latitude and season, which is not directly incorporated in an effective low dimensional model as presented here.

In the following we demonstrate that the model is capable of reproducing the climate record including the MPT and the time asymmetry, the sawtooth shapes, of the glacial cycles in the 100 kyr world solely as a response to the obliquity cycles. In order to explore the dynamical features of the model, we shall first simplify by applying a pure sinusoidal forcing $A(t) = A \sin(\omega t)$ with a period of 41 kyr, which quite accurately represents the obliquity cycle.

The simple model presented here exhibits surprisingly complex dynamics. Our postulate is that the MPT is a result of a slow environmental change represented in a slow change in a model parameter. Here we restrict ourselves to changes in the climatic damping coefficient κ . This change results in a shift from a period-1 frequency locking to a period 2 and 3 frequency locking. Such a change in the dynamical response to the forcing results in a very long non-periodic transient response, which might explain why the timing of the MPT varies so much (from 1.2 Myr to 800 kyr).

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Basic properties of the model

In spite of the apparent simplicity of this conceptual model, it contains a rich dynamics in its basic mathematical and physical aspects. To illustrate this, first we apply a simple deterministic ($\sigma = 0$) sinusoidal external driving; $A(t) = A \sin(\omega t)$, taking $\omega = 2\pi(42 \text{ kyr})^{-1}$ corresponding to the obliquity cycle. With this choice, the free parameters of the model are: α , A , κ and λ . Again, we use the value $\alpha = 0.8$, corresponding to a climatic potential (V) with a single minimum. Also, if not stated otherwise, the parameter λ was set to 10 (in appropriate units; $(100 \text{ kyr})^{-1}$) throughout our investigations, corresponding to a characteristic timescale of 10 kyr. The Eqs. (1)–(3) were numerically integrated using a fourth-order Runge–Kutta scheme. As initial conditions, $\tau_0 = 0$ and $x_0 = -1.2756$ were chosen. The latter value corresponds to the minimum of the climatic potential for $\alpha = 0.8$.

We found the following types of temperature anomaly oscillations: linear response, the response frequency coincides with that of the driving; period doubling, period multiplication, with related “exotic” shaped oscillations; and chaotic behavior.

The period multiplication scenario is shown in Fig. 2 for a section of the (A – κ) parameter space. A standard procedure for identifying periodicity as multiple of the driving period, would be generating a Poincare map, and counting the number of intersections, regarding points within circles of some small radius ϵ as identical. Here we apply a slightly more robust method where the periodicity n is calculated as

$$n = \min_m: \int_{T_{\text{end}} - mT}^{T_{\text{end}}} |x(t) - x(t - mT)| dt < \epsilon.$$

Figure 2 demonstrates the occurrence of all sorts of periods and a period doubling route to chaos. We also note, that due to the existence of sharp phase boundary lines separating different period oscillations, a small change in the model parameters can lead to a remarkable transition in the climatic response. As an example, Fig. 3

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



demonstrates a sharp period doubling transition in the climatic response τ due to a very small, gradual change in the sinusoidal drive, which we otherwise consider constant in this paper. This way, the response to a slight change in the amplitude of the forcing results not only in a change in periodicity, but also in a change in the amplitude of the response.

We found that in general, the appearance of chaos was promoted at small values of ω , large values of A , large values of λ , and small values of κ . Figure 4 shows the Lyapunov-exponent for the temperature anomaly (defined as $\partial \ln(\delta\tau)/\partial t$) for a section of the $(\lambda-\kappa)$ phase space where chaotic behavior occurs. Interestingly, while in certain directions the Lyapunov exponent changes continuously from zero to a positive value, in other directions one observes an abrupt change.

The systematic mathematical investigation of the different period multiplied phases and the features of chaos in the current model points beyond the scope of the present paper and will be reported elsewhere. Instead, we focus on the climatic relevance of the possible nonlinear responses.

4.1 Possible mechanism for the Mid-Pleistocene Transition

The paleoclimatic record contains enough enigmas to realize that the Milankovitch theory of orbitally forced ice age cycles still has missing links. Especially explaining the MPT is a challenge. The 100 kyr time scale is very long in comparison to reasonable estimates for internal climate oscillators to be at play. In the very inspiring paper (De Saedeleer et al., 2013) it was recently proposed, based on a study of the Van de Pol oscillator, that the climatic response to orbital forcing is a phase locking of internal periods to periods of the orbital forcing. This was proposed as a way of assessing the determining periods of the multi period orbital forcing. Here we demonstrate a different scenario for the dynamical response: first and foremost, our model does not have any internal periods of oscillation. This is a simple consequence of the fact that the proposed “climate potential” is not harmonic, thus the frequency of oscillation depends continuously on the amplitude of oscillation. The system can thus resonate

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with the external forcing by adjusting the amplitude of the oscillation. Furthermore, as it was seen in the case of a simple harmonic forcing, by a slow change of parameters the system can show both period doubling transitions to a chaotic state and other transitions such as transition to period three response. We thus propose that throughout the Pleistocene epoch, a gradual change in the climate system, which is represented by a slow variation in the model parameters took place. Such a change is feasible to assume as there is a long term decreasing trend in the global temperature of the Earth in the past 50 Myr (with the exception of some isolated abrupt changes induced by extreme events related to volcanos, or sharp tectonic changes). This long term global cooling can possibly be attributed to the gradual decrease of the global CO₂ levels (Clark et al., 2006) and/or slow tectonic processes (Clark and Pollard, 1998). We may assume that the relevant driving is the 41 kyr obliquity cycle throughout the Pleistocene epoch (Huybers and Wunsch, 2005; Huybers, 2009), in the latter part slowly varying parameters changed the dynamical response to periods two or three, thus the last part of the Pleistocene climate is interpreted as a mixture of 80 and 120 kyr responses to the obliquity cycle.

To illustrate this scenario we incorporate the assumed long-term change in the Earth's climatic response into the model by introducing a gradual decrease of the climatic damping coefficient κ . In particular, we used the following implementation:

$$\kappa(t) = \kappa_1 + 0.5(\kappa_0 - \kappa_1)\{1.0 - \tanh((t - t_0)/t_s)\}, \quad (4)$$

where $\kappa_0 > \kappa_1$, t_0 , and t_s are positive constants.

Furthermore, we ran the model with the real obliquity data (Berger and Loutre, 1991), implementing $A(t) = 4(O(t) - O_{av})$, where $O(t)$ is the actual obliquity dataset and O_{av} is its time average. Figure 1 shows a MPT that was simulated using model parameters $\kappa_0 = 1.1$, $\kappa_1 = 0.3$, $t_0 = 3.5$ Myr, $t_s = 2$ Myr, $\lambda = 10$, and $x_0 = -1.2756$, $\tau_0 = 0$ were used as initial conditions. One can see that the period of the oscillations changed from 41 kyr to approximately 120 kyr after the MPT.

We also found that the maximum temperatures lagged 9 kyr to the maximum amplitude of the external drive, throughout the whole Pleistocene epoch (Fig. 6). This result is in a good agreement with the findings of Huybers (2009).

Beside the change in response at the MPT the model also shows the characteristic sawtooth shape of the oscillations after the MPT. The sawtooth shape is a typical signature of fast-slow dynamics. To further explore this feature of the model, Fig. 5 shows the details in the period three response to a sinusoidal forcing, repeating the third panel in Fig. 2. In the top panel of Fig. 5, the variables $\tau(t)$ (blue), $x(t)$ (green) and the forcing $A\sin(\omega t)$ (red) are shown. The grey bands indicate the rapid warmings (fast phase), while the broad white sections indicate the gradual cooling periods (slow phase). What is important to notice is that the forcing (red curve) differs strongly for the two fast phase periods, while $x(t)$ is almost identical for the two periods. The product $xA\sin(\omega t)$ is shown in second panel, red curve. The model thus indicates that glacial terminations do not have to be in phase with the change in orbital forcing. The cause of the fast transitions is solely encoded in the “potential drift” term $-dV/dx$, which is shown in the second panel, blue curve. To illustrate this further, a phase space portrait of the period three cycle is shown in the bottom left panel while the trajectory of the forcing vs. x is shown in the middle panel. The coloring of the curves is such that the red part corresponds to the fast phases and the blue part to the slow phases. The fast-slow dynamics is thus a consequence of the asymmetry in the internal dynamics, represented in the simple model by the skewed climate potential, shown in the bottom right panel. Again the red part corresponds to the fast periods while the blue part corresponds to the slow periods. The obvious physical interpretation is that the melting and collapse of ice sheets and perhaps the melting of sea ice is a much faster process than the buildup of ice sheets from precipitation.

The glacial oscillations after the MPT are apparently much less regular than the 41 kyr period one oscillations prior to the MPT. This could indicate either that the latter oscillations are such that they vary between a regime of period two and a regime of period three response. Or the response is simply a long transient response before

Changing climatic response

I. Daruka and
P. D. Ditlevsen

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the system settles into a period three regime. This points to the fundamental issue of climate predictability, since the model shows both critical dependence on initial conditions and chaotic regimes. These are not synonymous, since the first can arise due to non-trivial boundaries for basins of attraction of different solutions even in the non-chaotic regime.

To illustrate this, we implemented a small perturbation on the slow temporal change protocol of the climatic damping coefficient. We found that even such a small disturbance gets amplified and shifts a glacial by a whole obliquity cycle (41 kyr) after 0.5 Myr later, constituting a *climatic butterfly* effect (Fig. 7). This exponential amplification of a small perturbation is a sign of chaos in the system.

However, for this range of parameters, there is no chaos in the system when it is driven by a simple sinusoidal oscillation. To further investigate the underlying mechanism, instead of the real obliquity data, we forced the model with a simpler, two sinusoidal amplitude modulated sinusoidal oscillation:

$$A(t) = A_0 \{ 1 + A_1 \sin(2\pi t/T_1) + A_2 \sin(2\pi t/T_2) \} \sin(2\pi t/T_0), \quad (5)$$

where we used $T_0 = 41$ kyr, $T_1 = 212.24$ kyr, $T_2 = 478.22$ kyr, $A_0 = 2$, $A_1 = 0.2$, and $A_2 = 0.3$. In Eq. (4) we implemented $\kappa_0 = 2.1$, $\kappa_1 = 0.1$, $t_0 = 35$ Myr, and $t_s = 2$ Myr as the adiabatic temporal change protocol parameters of the climatic damping coefficient. Furthermore, we set the parameter $\lambda = 10$ and $x_0 = -1.2756$, $\tau_0 = 0$ were used as initial conditions. We found that even a small amount of climatic noise $\sigma = 0.01$ in Eq. (2) can drastically change the behavior and can induce an MPT (Fig. 8a). Thus, this scenario corresponds to a *noise aided MPT*. As one can see, the late-Pleistocene climatic oscillations have two types, a ~ 80 kyr ($\sim 2 \times 41$ kyr) and a ~ 120 kyr ($\sim 3 \times 41$ kyr) period, following each other in an alternating way. This sequence gives a 100 kyr period on average.

The increase of the modulation amplitudes in the above mentioned model external drive (Eq. 5) ($A_1 = 0.3$, and $A_2 = 0.8$) and the implementation of a somewhat different damping coefficient temporal protocol (using $\kappa_0 = 1.1$, $\kappa_1 = 0.001$, $t_0 = 35$ Myr, and $t_s =$

2 Myr in Eq. 4) lead to the appearance of an MPT even in the absence of added noise. This behavior corresponds to an *obliquity irregularities aided MPT scenario* (Fig. 8b).

5 Conclusions

We have formulated a semi-conceptual model to describe the Mid-Pleistocene Transition, which we propose to be a period two and period three response to the 41 kyr obliquity forcing. This resolves the “100 and 400 kyr problems” of the Milankovitch theory, since the eccentricity cycle is proposed to be insignificant, and is omitted all together in the forcing. The change at the MPT is caused by a long term, gradual decrease of some parameter of the system. With the conceptual model approach, we can only point to dynamical mechanisms and not to the real environmental change. However, the idea of a gradual change aligns with the decrease in global CO₂ or tectonic re-arrangements as proposed in the literature.

The model does not in itself possess an internal frequency of oscillation, thus suggesting that the time scale of the glacial cycles is determined solely by the non-linear response to the frequency of the obliquity pacing. We observe that the model also reproduces the observed saw-tooth shaped time reversal asymmetry observed in the record, which is not present in the forcing. In the model this is related to the larger amplitude of the glacial cycles after the MPT in comparison to the more symmetric cycles prior to the MPT. With the larger cycles, the system experience the non-linearity in the internal dynamical response (the skewed climate potential in the model) much stronger than in the case of small, almost harmonic oscillations. As we have shown, the modulations in the forcing amplitude can also lead to a change in the periodicity of the response, rendering an intrinsically non-linear climatic behaviour.

Despite its simplicity, the model shows a surprisingly wide range of behaviours depending on the forcing, the initial conditions and the values of the parameters. The observed variety of possible climatic responses in the model raises the perplexing question: How robust is the climatic history of the Earth? We are obviously not in the

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



position, where we can rerun the past. Thus we must ask in which sense, we should be able to model the past, by reproducing the evolution, which has been realised, or by reproducing the past in some statistical sense.

Furthermore, the presence of chaos in the present model suggests that this could be manifested in the climate system itself in a so-called climatic butterfly effect: a small perturbation, or even just the internal climatic noise, might cause a significant shift in the glacial cycles. Obviously, this could impose fundamental limitations on long-term climate predictability.

Acknowledgements. I. Daruka acknowledges the support of the Hungarian Science Foundation under the contract OTKA NK72037 and that of the Hungarian Meteorological Service.

References

- Augustin, L., Barbante, C., Barnes, P. R., and other EPICA community members: Eight glacial cycles from an Antarctic ice core, *Nature*, 429, 623–628, 2004. 1103
- Ashkenazy, Y. and Tziperman, E.: Are the 41 kyr oscillations a linear response to Milankovitch forcing?, *Quaternary Sci. Rev.*, 23, 1879–1890, 2004. 1107
- Benzi, R., Parisi, G., Sutera, A., and Vulpiani, A.: Stochastic resonance in climatic change, *Tellus*, 34, 10–16, 1982. 1105
- Berger, A. L.: Long-term variations of daily insolation and quaternary climatic changes, *J. Atmos. Sci.*, 35, 2362–2367, 1978. 1104
- Berger, A.: A brief history of the astronomical theories of paleoclimates, in: *Climate Change*, edited by: Berger, A., Mesinger, F., and Sijacki, D., Springer, Wien, 107–129, 2012. 1103
- Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million of years, *Quaternary Sci. Rev.*, 10, 297–317, 1991. 1113, 1120, 1126
- Budyko, M. I.: The effect of solar radiation changes on the climate of the Earth, *Tellus*, 21, 611–619, 1969. 1105, 1109
- Clark, P. U. and Pollard, D.: Origin of the middle Pleistocene transition by ice sheet erosion of regolith, *Paleoceanography*, 13, 1–9, 1998. 1104, 1113
- Clark, P. U., Archer, D., Pollard, D., Blum, J. D., Rial, J. A., Brovkin, V., Mix, A. C., Pisias, N. G., and Roy, M.: The middle Pleistocene transition: characteristics, mechanisms,

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changing climatic responseI. Daruka and
P. D. Ditlevsen[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and implications for long-term changes in atmospheric $p\text{CO}_2$, *Quaternary Sci. Rev.*, 25, 3150–3184, 2006. 1104, 1113

Crucifix, M.: Oscillators and relaxation phenomena in Pleistocene climate theory, *Philos. T. R. Soc. A*, 370 1140–1165, 2012. 1105

5 De Saedeleer, B., Crucifix, M., and Wiczorek, S.: Is the astronomical forcing a reliable and unique pacemaker for climate? A conceptual model study, *Clim. Dynam.*, 40, 273–294 2013, 2013. 1104, 1110, 1112

Ditlevsen, P. D.: Bifurcation structure and noise-assisted transitions in the Pleistocene glacial cycles, *Paleoceanography*, 24, PA3204, doi:10.1029/2008PA001673, 2009. 1104, 1106

10 Huybers, P.: Early Pleistocene glacial cycles and the integrated summer insolation forcing, *Science*, 313, 508–511, 2006. 1103

Huybers, P.: Glacial variability over the last two million years: an extended depth-derived age model, continuous obliquity pacing, and the Pleistocene progression, *Quaternary Sci. Rev.*, 26, 37–55, 2007. 1104

15 Huybers, P.: Pleistocene glacial variability as a chaotic response to obliquity forcing, *Clim. Past*, 5, 481–488, doi:10.5194/cp-5-481-2009, 2009. 1107, 1113, 1114

Huybers, P. and Wunsch, C.: Obliquity pacing of the late Pleistocene glacial terminations, *Nature*, 434, 491–494, 2005. 1106, 1113

20 Hyde, W. T. and Peltier, W. R.: Sensitivity experiments with a model of the ice age cycle. The response to harmonic forcing, *J. Atmos. Sci.*, 42, 2170–2188, 1985. 1106

Imbrie, J., Berger, A., Boyle, E. A., Clemens, S. C., Duffy, A., Howard, W. R., Kukla, G., Kutzbach, J., Martinson, D. G., McIntyre, A., Mix, A. C., Molfino, B., Morley, J. J., Peterson, L. C., Pisias, N. G., Prell, W. L., Raymo, M. E., Shackleton, N. J., and Toggweiler, J. R.: On the structure and origin of major glaciation cycles 2, the 100 000-year cycle, *Paleoceanography*, 8, 699–735, 1993. 1104

25 Kallen, E., Crafoord, C., and Ghil, M.: Free oscillations in a climate model with ice-sheet dynamics, *J. Atmos. Sci.*, 36, 2292–2303, 1979. 1106

LeTreut, H. and Ghil, M.: Orbital forcing, climate interactions, and glacial cycles, *J. Geophys. Res.*, 88, 5167–5190, 1983. 1106

30 Lisiecki, L. E. and Raymo, M. E.: A Pliocene–Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records, *Paleoceanography*, 20, PA1003, doi:10.1029/2004PA001071, 2005. 1103, 1120

**Changing climatic
response**I. Daruka and
P. D. Ditlevsen[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Paillard, D.: The timing of Pleistocene glaciations from a simple multiple-state climate model, *Nature*, 391, 378–381, 1998. 1105
- Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J.-M., Basile, I., Benders, M., Chappellaz, J., Davis, M., Delayque, G., Delmotte, M., Kotlyakov, V. M., Legrand, M., Lipenkov, V. Y., Lorius, C., Pépin, L., Ritz, C., Saltzman, E., and Stievenard, M.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429–436, 1999. 1103
- Rial, J. A.: Abrupt climate change: chaos and order at orbital and millennial scales, *Global Planet. Change*, 41, 95–109, 2004. 1107
- 10 Saltzman, B. and Maasch, K.: A first-order global model of late Cenozoic climate change II: a simplification of CO₂ dynamics, *Clim. Dynam.*, 5, 201–210, 1991. 1104, 1106, 1108
- Sellers, W. D.: A climate model based on the energy balance of the earth–atmosphere system, *J. Appl. Meteorol.*, 8, 392–400, 1969. 1105
- 15 Shackleton, N. J.: Deep-sea sediment record and the Pliocene–Pleistocene boundary, *Quatern. Int.*, 40, 33–35, 1997. 1103
- Tziperman, E. and Gildor, H.: On the mid-Pleistocene transition to 100 kyr glacial cycles and the asymmetry between glaciation and deglaciation times, *Paleoceanography*, 18, 1–8, 2003. 1106, 1107, 1108

Changing climatic response

I. Daruka and
P. D. Ditlevsen




Fig. 1. The Mid-Pleistocene Transition (MPT). The red curve corresponds to the composite deep sea foraminiferal isotope record (Lisiecki et al., 2005). The MPT is seen as a change around 1 MyrBP in period from 41 kyr in the early Pleistocene to approximately 100 kyr in the late Pleistocene. The former cycles are symmetric, while the latter cycles are asymmetric (“sawtooth shaped”). The blue curve shows the model results for the related temperature anomaly when driven by real obliquity data (Berger and Loutre, 1991), represented by the black curve. Both the foraminiferal record and the obliquity data were scaled and shifted vertically for better visual representation. The parameter values used for this plot are discussed in the text.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 2. The behavior of the model driven by a sinusoidal forcing $A\sin(\omega t)$, shown in red in the second panel. The top panel shows the periods in multiplets of the driving period ($2\pi/\omega$): White corresponds to periodic solutions with period 1, yellow: period 2, green: period 3, red: period 4, magenta: period 6, black: ≥ 8 , including non-periodic (chaotic) solutions. The bottom three panels show the solutions corresponding to the blue circles in the top panel left-to-right (periods 3, 2, 1 respectively). Changing parameters along a horizontal line shows a period doubling route to chaos. We used the parameters $\omega = 2\pi/10$, $\lambda = 0.1$, with $x_0 = 0$, and $\tau_0 = 1$ as initial conditions for this plot.

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changing climatic response

I. Daruka and
P. D. Ditlevsen




Fig. 3. Even a very small, visually unnoticeable gradual change in the sinusoidal driving (plotted in green) can lead to a sharp, remarkable period doubling in the climatic response τ (plotted in red). We used the parameters $\omega = 7$, $\lambda = 10$, $\kappa = 1.3$, and $A(t) = 0.008t \sin(\omega t)$ with $x_0 = -1.2756$, and $\tau_0 = 0$ as initial conditions for this plot.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 4. Plot of the Lyapunov-exponent (in units of kyr^{-1}) for the temperature anomaly (τ) oscillations for a section of the (λ - κ) parameter phase space, demonstrating the presence of chaos in the model. The parameter values are $\omega = 10$, $A = 1$, and $x_0 = 0$, $\tau_0 = 1$ were used as initial conditions.

Changing climatic response

I. Daruka and
P. D. Ditlevsen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changing climatic response

I. Daruka and
P. D. Ditlevsen




Fig. 5. The period three response to a sinusoidal forcing, as in Fig. 2. The parameter values are $\alpha = 0.8$, $\lambda = 0.087$, $\kappa = 0.078$, $A = 1.16$, $\omega = 2\pi/10$. Top panel shows the variables $\tau(t)$ (blue), $x(t)$ (green) and $A \sin(\omega t)$ (red). The grey bands with rapid increase in $\tau(t)$ are defined as the fast periods, while the rest are the slow periods. The middle panel shows the two dominant terms in the right hand side of the dynamical Eq. (2): $-dV/dx$ (blue) and $x A \sin(\omega t)$ (red). It is seen that the fast dynamics is governed by the former term, representing internal dynamics. The bottom left panel shows the phase space portrait of the period three solution, while the bottom middle panel shows the forcing $A \sin(\omega t)$ vs. τ . The bottom right panel shows the “climate potential”. The asymmetry of the potential is responsible for the fast-slow dynamics and the saw-tooth shape of the record. The red parts of the curves correspond to the fast periods.

Changing climatic response

I. Daruka and
P. D. Ditlevsen




Fig. 6. The temperature oscillations lag the external drive. The plot shows the temperature oscillations (red curves) shifted by 9 kyrs to match the phase of the obliquity oscillations (green curves). The 9 kyr lag persists through the whole model-MPT process. The parameter values used for this plot are identical with that of Fig. 1.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Changing climatic response

I. Daruka and
P. D. Ditlevsen




Fig. 7. A small perturbation in the climatic damping coefficient (green curve) leads to a remarkable shift of the glacial cycles, constituting a climatic butterfly effect. The blue and red curves correspond to the resulting temperature anomalies without and with the perturbation in the time protocol of the climatic damping coefficient, both driven by real obliquity data (Berger and Loutre, 1991) in the same way as described in the text for Fig. 1. The parameter values are $\lambda = 10$, $\kappa_0 = 1.1$, $\kappa_1 = 0.3$, $t_0 = 3.5$ Myr, $t_s = 2.008$ Myr, and $x_0 = -1.2756$, $\tau_0 = 0$ were used as initial conditions.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Changing climatic response

I. Daruka and
P. D. Ditlevsen




Fig. 8. (a) Climatic noise induced MPT. The figure demonstrates the 41 kyr pre-MPT and the (80+120) kyr post-MPT temperature oscillations cycles (red curve). The green curve represents the model external drive Eq. (5), scaled and vertically shifted for better visibility. (b) Obliquity irregularities induced MPT. The figure demonstrates the 41 kyr pre-MPT and the post-MPT “saw-tooth shaped” temperature oscillations (red curve). The green curve represents the model external drive Eq. (5), scaled and vertically shifted for better visibility. The parameter values used for these plots are discussed in the text.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)