



Seasonal variations
recorded in cave
monitoring results

M. Van Rampelbergh
et al.

Seasonal variations recorded in cave monitoring results and a 10 year monthly resolved speleothem $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ record from the Han-sur-Lesse cave, Belgium

M. Van Rampelbergh¹, S. Verheyden^{1,2}, M Allan³, Y. Quinif⁴, E. Keppens¹, and P. Claeys¹

¹Earth System Sciences, Vrije Universiteit Brussel (VUB), Pleinlaan, 1050, Brussels, Belgium

²Royal Belgian Institute of Natural Sciences, Geological Survey, Direction Earth and History of Life, Jennerstraat 13, 1000, Brussels, Belgium

³AGEs, Département de Géologie, Université de Liège, Allée du 6 Août, B18 Sart-Tilman, 4000, Liège, Belgium

⁴Faculté Polytechnique, Université de Mons, Rue de Houdain 9, 7000, Mons, Belgium

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 21 March 2014 – Accepted: 8 April 2014 – Published: 22 April 2014

Correspondence to: M. Van Rampelbergh (mvrampel@vub.ac.be)

Published by Copernicus Publications on behalf of the European Geosciences Union.

CPD

10, 1821–1856, 2014

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Speleothems provide paleoclimate information on multi-millennial to decadal scales in the Holocene. However seasonal or even monthly resolved records remain scarce. They require fast growing stalagmites and a good understanding of the proxy transfer function on very short time scales. The Proserpine stalagmite from the Han-sur-Less cave (Belgium) displays seasonal layers of 0.5 to 2 mm thickness that reconstruct paleoclimates at a monthly scale. Through a regular cave monitoring, we acquired a good understanding of how $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ signals in modern calcite reflect climate variations on sub-seasonal scale. Cave parameters vary seasonally in response to the activity of the vegetation cover and outside air temperature. From December to June, the cave remains in “winter-mode”. Outside temperatures are cold inducing low cave air and water temperatures. Bio-productivity in the soil is limited leading to low $p\text{CO}_2$, higher $\delta^{13}\text{C}$ composition of the CO_2 in the cave air and high discharge due to the inactivity of the plant coverage. From June to December, the cave switches to “summer-mode” and the measured factors display an opposite behavior. The $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ signals of fresh calcite precipitated on glass slabs vary seasonally. Lowest $\delta^{18}\text{O}$ values occur during the summer-mode when the $\delta^{13}\text{C}$ values are high. The $\delta^{18}\text{O}$ composition of the calcite is in equilibrium with the drip water $\delta^{18}\text{O}$ and display seasonal variations due to changes in the cave air and water temperature. In contrast to the $\delta^{18}\text{O}$ signal, $\delta^{13}\text{C}$ values of the calcite precipitated on the glass slabs do not reflect equilibrium conditions. Highest $\delta^{13}\text{C}$ values occur during summer, when discharge rates are low increasing the evaporation effect on the thin water film covering the stalagmite. This same antithetical behavior of the $\delta^{18}\text{O}$ vs. the $\delta^{13}\text{C}$ signals is seen in the monthly resolved speleothem record that covers the period between 1976 and 1985 AD. Dark layers are formed during summer, while light layers are formed during winter when calcite deposition occurs fast. The darker the color of a layer, the more compact its calcite structure, the more negative its $\delta^{18}\text{O}$ signal and the more positive its $\delta^{13}\text{C}$ signal.

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

In the past 25 years, speleothem records have provided important information on past climate systems on multi-millennial to decadal scales (Genty et al., 2003; McDermott, 2005; Verheyden et al., 2008b; Wang et al., 2008; Van Rampelbergh et al., 2013).

5 With the increasing number of studies on cave calcite deposition dynamics (Dreybrodt, 1999, 2008; Verheyden et al., 2008a; Lachniet, 2009; Scholz et al., 2009; Ruan and Hu, 2010; Oster et al., 2012) and with the help of modern analytical tools (Fairchild et al., 2006; Spotl and Matthey, 2006; Jochum et al., 2012), speleothems document climate at sub-seasonal and even monthly scales. However, this high resolution is reached in only
10 a few studies due to several limitations (Treble et al., 2003; Matthey et al., 2008).

To study paleoclimate at sub-seasonal scale from stalagmites, their growth rate needs to be significantly high (around 1 mm yr^{-1}) to deposit thick layers allowing monthly resolved time-series. Speleothem-growth rates vary according to different factors such as drip water rate and calcium ion concentration (Baker et al., 1998; Dreybrodt, 1999) rendering the estimation of an average rate difficult. Generally, stalagmites
15 increase at $10\text{--}100 \mu\text{m yr}^{-1}$ in cool temperate climates and at $300\text{--}500 \mu\text{m yr}^{-1}$ in subtropical climates (Fairchild et al., 2006), clearly showing that fast growing (more than 1 mm yr^{-1}) speleothems are truly exceptional.

A second limitation is that the transfer function of the measured proxies needs to be well understood for the studied cave and time frame. On classical multimillennial and centennial time scales, the transfer functions for the stable isotopes of oxygen and carbon are well established (Fairchild et al., 2006; Baker et al., 2007). However, local cave specific effects affect seasonally or even monthly resolved $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ signals. For studies at seasonal scales, a detailed study of the cave dynamics is required
20 (Dreybrodt, 1999) in order to understand which factors drive the isotopic signals, and at which intensity. Different cave monitoring systems have been set up all over the world to better understand these seasonal and sub-seasonal processes (Genty and Deflandre, 1998; Matthey et al., 2008; Riechelmann et al., 2011). Only few of them,
25

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



provided answers on the isotope fractionation processes occurring between the drip water and recent precipitated calcite (Verheyden et al., 2008a; Tremaine et al., 2011; Riechelmann et al., 2013).

Previous studies of the Han-sur-Lesse karst system show that the cave responds seasonally to external climate factors and that it is well suited for high-resolution speleothem based climate reconstructions (Genty and Quinif, 1996; Verheyden et al., 2006, 2008a). In the Han-sur-Lesse cave, the high growth rate (up to 2.1 mmyr^{-1}) and clear seasonal banding of the “Proserpine” stalagmite make it possible to reconstruct climate variations at monthly scale (Verheyden et al., 2006). In this study, we report results of a every two-week cave environment monitoring carried out for one year (2013) that shows how oxygen and carbon isotopes signals obtained from the Proserpine banding reflect climate variations at sub-seasonal scale. The results are then compared to high-resolution $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ signals measured on the 10 thickest layers from the upper 10 cm of the Proserpine, which cover the period from 1976 to 1985 AD. This approach improves the knowledge and accuracy for the use of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ signals in speleothems at seasonal scale.

2 Study area and hydrological setting

The Han-sur-Lesse cave is located within Givetian limestones of the Dinant synclinorium and is the largest and best-developed karst system in Belgium (Delvaux De Fenffe, 1985). The Lesse River carves the cave within a hill called the “Massif de Boine” entering the karst system at the “Gouffre de Belvaux” and exiting approximately 24 h later through the “Trou de Han” (Fig. 1). The cave is exploited since the mid 19th century as a touristic attraction and is characterized by large chambers and well developed speleothem formations. The cave monitoring and speleothem sampling for this study is carried out in the “Salle-Du-Dôme” chamber. This 150 m wide and 60 m high chamber formed by collapse and is the largest of the whole cave system. The Proserpine stalagmite is easily reachable following the tourist path into the cave for approximately

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

700 m from the cave's exit at the "Trou de Han" (Fig. 1). It has a large tabular shape with a relatively horizontal slightly undulating surface of about 1.5 m^2 and is fed by a continuous high drip water flow. The epikarst thickness above the stalagmite is estimated to be 40 m (Quinif, 1988). Based on a study comparing the isotopic signals of rainwater and drip water in the Pere Noel cave, formed in the same karst system as the Han-sur-Lesse cave, the residence time of the water in the epikarst is approximately 5 to 6 months (Verheyden et al., 2008a). After a heavy rainfall, the drip rate above the stalagmite reacts within 24 h suggesting a piston flow system above the studied stalagmite. Two passages connect the "Salle-Du-Dôme" to the neighboring chambers and allow air circulation in the chamber. The Lesse River flows at the bottom of the chamber enhancing air circulation and causing the chamber to remain well ventilated throughout the year.

The mean annual precipitation at the nearest meteorological station of Han-sur-Lesse is 844 mm yr^{-1} and the mean annual air temperature 10.3°C (Royal Meteorological Institute Belgium). While the temperature displays a well-marked seasonality with cool summers and mild winters, the rainfall is spread all over the entire year. For the studied period lasting between November 2012 and January 2014, air temperature was at its lowest between December and March and highest between July and September. The coldest temperature of -4.2°C was reached on 15 March 2013. The plant coverage above the cave consists of C3-type vegetation with oaks, beech and hazel trees. The soil is approximately 40 cm thick and consists of silty stony soil with more than 50 % limestone fragments (Belgian Geological Survey map). The area above the cave is part of a protected natural reserve, preserved from direct human influence for more than 50 years.

3 Methods

Between October 2012 and January 2014, the cave was visited every two weeks to record environmental parameters. Cave air and water temperature were measured with

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 a HANNA HI955501 thermometer with a precision of 0.2 °C. Air temperature was taken directly above the stalagmite. The drip water temperature and pH were determined in a small pool on the stalagmite's surface. The pH of the drip water was measured with a HANNA HI991300 sensor (precision of 0.01 pH). The concentration of CO₂ in the
10 cave air was obtained using an ACCURO 640000 manual Dräger pump with a standard deviation of 10 to 15 %. Three times per visit, *p*CO₂ values were measured and reported as an average of the three values. The drip water discharge (volumetric flow rate, here given in volume (mL) of water reaching the speleothem surface per minute) above the Proserpine stalagmite was measured in a graded cylinder after collecting
15 the drip water during 10 min in an inflatable soft plastic swimming pool that was placed on the stalagmite's surface and that covered a slightly larger area (1.77 m²). Drip water samples for δ¹⁸O and δD measurements were collected in fully filled glass bottles and stored in a cool and dark environment. Rainwater samples for δ¹⁸O and δD measurements were collected in a garden close to the cave using a thermos bottle and sampled
20 every 15 days between November 2012 and January 2014. To avoid evaporation processes, the rainwater was collected using a funnel with a raised edge connected to a tube reaching the bottom of the thermos bottle. The funnel was attached to the bottle trough a hermetic cap. Glass bottles were fully filled with the collected rainwater and stored in a dark fridge until being analyzed.

25 The δ¹⁸O and δD composition of the waters were measured using a PICARRO L2130-c Cavity Ring down Spectrometer at the Vrije Universiteit Brussel (VUB). For every sample, 1.4 mL of water was used for the measurements. Every sample was injected and analyzed 10 consecutive times. The measured values were then corrected using a three point correction method with three home made standards with isotopic composition ranging between 6.8 ‰ and –22.3 ‰ for δ¹⁸O and 32.5 ‰ and –127.3 ‰ for δD. All water samples were analyzed two times in different order. The reported values are the average of the two measurement rounds. Analytical uncertainties (2σ) equal 0.07 ‰ for the δ¹⁸O values and 0.5 ‰ for the δD values.

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The isotopic composition of the cave air CO₂ was measured from samples collected by filling vacuum 2 L glass containers. To avoid “human” contamination, these samples were taken at the beginning of every cave visit. The CO₂ was extracted from the container using a manual extraction line at the VUB. The extracted CO₂ was then analyzed for its isotopic composition on a Thermo Delta plus XL mass spectrometer in dual inlet mode. To test the reproducibility of the method, samples for the analyses of the δ¹³C composition of the dissolved inorganic carbon (DIC) in the water were collected following two different methods. The first method consisted of sampling the drip water by filling 12 mL gastight glass tubes all the way to the top to avoid air CO₂ contamination. A drop of HgCl₂ was immediately added and the bottle hermetically closed and stored in a dark and cool environment until being analyzed. The day before the analysis, a headspace was created in the bottle by taking out 3 mL of water, while bubbling He through the septum. Once the headspace was formed, H₃PO₄ was added and the sample shaken overnight to convert all DIC species into CO₂. The CO₂ gas was then extracted from the bottle and measured for its δ¹³C composition. Samples were duplicated and measured immediately after sampling and 1 month later to test if evaporation processes affect the DIC composition.

The second method is based on that described by Matthey et al. (2008). Gastight 12 mL glass vials were first flushed with He in the lab. In the cave, 1 mL of drip water was injected in these vials through the septum. Back in the lab, samples were acidified with H₃PO₄ and equilibrated for at least 2 h before being analyzed. Using this method, samples must be analyzed within 2 days to avoid possible evaporation of the CO₂ through the pierced septum. For both sampling methods, the δ¹³C composition of the DIC was measured on a Flash EA 1112 device connected to a Delta V plus mass spectrometer, with an analytical uncertainty of 0.4 ‰ (2σ).

Every visit, 3 glass slabs were placed on the surface of the stalagmite to test current calcite deposition conditions within the cave. The slabs were collected during the next visit. The freshly precipitated calcite was then scraped off the glass and 5 samples per

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

fore this “early summer” decrease, the discharge record shows a short maximum to 280 mL min^{-1} in early June, most probably related to the heavy rainfall period at the end of May 2013 (Fig. 2a). An increase in drip water flow above the Proserpine is observed within 24 h following a heavy rainfall event. The drip water pH (Fig. 2f) varies between 8.4 and 7.9, decreasing in spring (sharply in May) and gradually increasing back at the end of summer and throughout autumn (from September through January). The heavy rainfall period at the end of May 2013 (Fig. 2a) seems to correspond with a pH decrease below 8.0. The $p\text{CO}_2$ values (Fig. 2g) remain relatively stable around 500 ppm throughout much of the year except for a marked increase in the summer, reaching 1000 ppm in July and August. The $\delta^{13}\text{C}$ signature (Fig. 2h) of the cave air varies around an average of -19.5‰ displaying an opposite trend to that of the $p\text{CO}_2$ concentrations. During the high discharge event at the end of May, the $\delta^{13}\text{C}$ values of cave air fall to -20.7‰ , and remain around -20.0‰ or below during summer (JJA). The $\delta^{13}\text{C}$ composition of the DIC (Fig. 2i) stay constant throughout the year at an average of $-11.8 \pm 0.3\text{‰}$. The $\delta^{13}\text{C}$ compositions of the DIC measured using the new sampling method and preparation developed in this study (Table 1a) are the same as those obtained using the approach established by Matthey et al. (2008) (Table 1b). No meaningful differences were found between both methods as can be seen in Table 1b.

Rainwater $\delta^{18}\text{O}$ values (Fig. 3c) average $-8.18 \pm 0.07\text{‰}$ and the δD values (Fig. 3d) $-55.52 \pm 0.5\text{‰}$. Both the $\delta^{18}\text{O}$ and δD signals increase by 3‰ and 30‰ respectively during the summer months, presumably due to temperature effect. Both δ values also display 6 small drops (blue clouds and numbered 1 trough 6 in Fig. 3c and d) of approximately 2‰ for the $\delta^{18}\text{O}$ and 20‰ for the δD . They occur in November 2012, December 2012, May 2013, July 2013, December 2013 and January 2014. They appear to correspond to periods of higher rainfall recorded in the site (Fig. 3b). One larger drop (red arrow in Fig. 3c and d) of 9‰ for the $\delta^{18}\text{O}$ and of 90‰ for the δD signal occurs at the beginning of March and seems to correlate with unusually cold air temperatures of -6.3°C (Fig. 3a), measured at the RMI station of Han-sur-Lesse.

5 Discussion

5.1 Forcing of the rain and drip water variations

Rainwater $\delta^{18}\text{O}$ values at a specific location vary due to temperature changes, variation in the amount of rainfall, fluctuations in the source of the rainwater or cloud track (Rozanski et al., 1992). The rainwater $\delta^{18}\text{O}$ signal increases by a few per mille during the summer months, when air temperature is high (Fig. 3a and c). Six small drops are recorded in November 2012, December 2012, May 2013, July 2013, December 2013 and January 2014 and are numbered 1 through 6 in Fig. 3. These drops not correspond with air temperature decreases measured in the Han-sur-Lesse weather station. They however, systematically occur after a period marked by heavy rainfall (blue clouds in Fig. 3b). Therefore, the amount effect likely causes the smaller negative excursions in $\delta^{18}\text{O}$ values recorded in the rainwater. Negative $\delta^{18}\text{O}$ excursions appearing shortly after an increase in precipitation amount may be due to the bi-monthly collection of rainwater, which may explain a time lag of up to 15 days. A single larger drop in drip water $\delta^{18}\text{O}$ occurs in March 2013 and is indicated by a red arrow in Fig. 3. This large drop does not correspond with a temperature drop or an increase in rainfall. A modification in rainwater source could be a plausible explanation, but is not supported by changes in wind direction during that month based on RMI data. However, in March 2013, an unusually late snow layer covered the area for several weeks (RMI data). The observed decrease in $\delta^{18}\text{O}$ is then most probably related to the low $\delta^{18}\text{O}$ signal of the incorporate snow in the sampling bottle.

The $\delta^{18}\text{O}$ and δD compositions of the drip water display almost no variations throughout the year indicating that the epikarst-storage reservoir is relatively large and that the water residence time is sufficiently long to homogenize the isotopic composition of the drip water (Fig. 2j). In July and August, a small negative excursion in the drip water $\delta^{18}\text{O}$ of hardly 0.1 ‰ occurs. Although it is only of the size of the analytical uncertainty, we consider it meaningful because it is supported by two consecutive points as well as by trends of several points before and after. Since, meteoric water is by far the

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

major source of vadose/epikarst water and consequently of the drip water, the small negative $\delta^{18}\text{O}$ excursion in the drip water record is the subdued result of the $\delta^{18}\text{O}$ drop of several ‰ found in the meteoric water, as a result of snowfall in March 2013. The intense mixing of the percolating vadose water in the epikarst, deduced from the generally constant $\delta^{18}\text{O}$ in drip water record, would then reduce a $\delta^{18}\text{O}$ shift of about 8‰ in the meteoric to a hardly detectable one of about 0.1‰ in the drip water. This would also mean that the average residence time of the water in the epikarst must be estimated around five to six months (from March till August–September). Such a residence time is similar to that observed in the nearby Pere Noel cave, which formed in the same karst system as the Han-sur-Lesse cave, and where summer rainwaters only reach the cave in winter (Verheyden et al., 2008a). With an increase in drip rate within 24 h following a heavy rainfall event and a residence time of the water in the epikarst of 5 to 6 months, the epikarst hydrological system appears to act like a piston flow system with a delay of the order of half a year, long enough for the epikarst water to be well mixed before entering the cave.

5.2 Cave dynamics seasonal variations

Based on our observations, the temperatures of cave air, of drip water, and of outside air all follow the same seasonal evolution. However, the temperature range, which is around 20 °C outside the cave, is reduced to about 3 °C inside the cave with the drip water being about 1 °C colder than the cave air, an observation that is made in many caves (McDermott, 2004; Fairchild et al., 2006).

The Lesse River, which flows through the chamber, most probably causes the cave air to mix with that from neighboring chambers, including those openly connected to the outside. Compared to other Belgian caves where values up to 15 000 vppm are measured for cave air CO_2 (Verheyden, 2001), the CO_2 content in the Salle-du-Dôme chamber is very low, indicating that exchanges between cave air and external air must be relatively important. In the Salle-du-Dôme, CO_2 values fall close to outside air $p\text{CO}_2$, and values of 400 vppm are measured during much of the year (Fig. 2g). Only during

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rainfalls also affect the pH values of the drip water. In a piston-flow system, when high amounts of fresh water, – such as during a rainstorm –, are added to an epikarst that functions as a piston system, the total Ca^{2+} -ion concentration of the water diminishes and the water becomes more acid. This effect is illustrated during the very wet event of 30 May 2013 (“May wet event” in black box in the precipitation curve in Fig. 2a), where the discharge peaks to high values and where the pH (Fig. 2f) drops from 8.2 to 7.9 in one month (3 successive points), after a 2 month period of increase, during a much drier period.

To summarize: the cave system is subdivided into a “winter-mode” lasting from December to June and a “summer-mode” from June to December (Fig. 2). During the winter mode, cave air and drip water temperature are low. The plant coverage above the cave is inactive leading to a higher water discharge, low $p\text{CO}_2$ and high $\delta^{13}\text{C}$ values of the cave air CO_2 . Drip water pH is more basic due to the longer residence time of the summer rainwater in the epikarst. During the “summer-mode”, cave air and drip water temperatures increase. The plant coverage reactivates leading to lower discharge, higher $p\text{CO}_2$ and lower $\delta^{13}\text{C}$ values of the cave air CO_2 . The drip water pH is more acid due to its shorter residence time in the epikarst.

5.3 $\delta^{18}\text{O}$ composition of the precipitated calcite reflects temperature variations

The $\delta^{18}\text{O}$ composition of calcite deposited on the glass slabs varies seasonally with more negative values during summer (Fig. 2l). If the calcite is deposited in isotopic equilibrium with its drip water, these variations can be caused by changes in the $\delta^{18}\text{O}$ composition of the drip water and/or by changes in temperature that affect the fractionation factor between the drip water and the precipitating calcite (Lachniet, 2009). If not deposited in equilibrium, the seasonal $\delta^{18}\text{O}$ variations on the glass slabs are due to kinetic effects that require further investigation.

A first step in understanding the $\delta^{18}\text{O}$ transfer-function of the Salle-du-Dôme demands to determine whether the calcite is deposited in equilibrium or not. In speleothems, this is traditionally done by applying the Hendy-test (Hendy, 1971) that

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

compares the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values in the center of the stalagmite with those on the sides within a single growth layer. However, since we work on a drill core taken “in the middle” of a 1 to 2 m wide speleothem, the Hendy-test cannot be applied. As an alternative, we calculated equilibrium conditions of the deposited calcite using the calcite-water fractionation factor. Different authors have proposed fractionation factors based on three approaches.

The first approach, using in-lab experiments, has been tested in different studies each giving another value for the water-calcite fractionation factor. The most used in-lab established fractionation factors remain the ones by O’Neil et al. (1969) later modified by Friedman and O’Neil (1977), the relationship of Kim and O’Neil (1997) later modified by Kim et al. (2007), the results of Tarutani et al. (1969) and Jimenez-Lopez et al. (2001) (Table 2). A second category of fractionation factors is established by using theoretical models as the ones from Horita and Clayton (2007) and Chacko and Deines (2008). A third approach consists of using cave-monitoring data to make an average of the in-cave observed fractionation factors. Tremaine et al. (2011) established such a “cave calcite” relationship by doing a best fit through the data on a large number of modern caves at different latitudes, altitudes and temperatures.

Applying the different fractionation factors to our data shows that the measured $\delta^{18}\text{O}$ signals in the glass slabs calcite correspond within 1 ‰ with the calculated values suggesting that it is deposited close to equilibrium with the drip water (Table 3). However, variations occur between the results derived from different methods. Of the in-lab established relationships, that of O’Neil et al. (1969) modified by Friedmann and O’Neil (1977), corresponds best to our observations. This is also confirmed in other studies where this experimental fractionation factor is considered to give the best approximation for in-cave observations (McDermott et al., 2005; Riechelmann et al., 2013). The theoretical values (Horita and Clayton, 2007; Chacko and Deines, 2008) suggest a more negative $\delta^{18}\text{O}$ composition for the deposited calcites. This also confirms other studies where theoretical results seem to overestimate the fractionation factor (Tremaine et al., 2011). The optimal results are given by the Tremaine et al. (2011)

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- at values slightly more positive than the average $\delta^{18}\text{O}$ and δD composition of the rainwater indicating a well-mixed aquifer. A drop in isotopic composition of the drip water is seen 5 months after the March 2013 snowfall event indicating that the residence time of the water in the epikarst is 5 to 6 months. The drip water flow in the monitored Salle-du-Dôme reacts within 24 h on a heavy rainfall event indicating the piston flow character of the epikarst.
2. The records of the different measured factors in and outside the cave suggest that the physico-chemical behavior of the Han-sur-Lesse cave closely responds to the seasonally varying external climate. The cave interior climatology varies between a “summer-mode” lasting from June to December and a “winter-mode” from December to June. During the summer mode, the cave air and drip water temperatures are higher and the cave air $p\text{CO}_2$ increases due to an active vegetation cover above the cave. More negative plant derived CO_2 with a C3-signature is brought into the cave with the drip water decreasing the $\delta^{13}\text{C}$ signature of the cave air in summer. The drip flow decreases due to higher evaporation at the surface and because of higher water uptake by the vegetation. During the “winter-mode” the temperature cools down and the vegetation cover becomes less active leading to an opposite behavior of the measured factors compared to the “summer-mode”.
3. Calcite precipitated on glass slabs placed on top of the stalagmite and sampled throughout the year has more negative $\delta^{18}\text{O}$ and less negative $\delta^{13}\text{C}$ values when formed during summer. The opposite is observed in winter. After modeling of all possible mechanisms influencing these stable isotopic signals, we conclude that more negative $\delta^{18}\text{O}$ values of precipitated calcite in late summer are due to a temperature increase of the cave air and drip water. The less negative $\delta^{13}\text{C}$ values in summer are due to a decrease in water discharge.
4. The studied part of the Proserpine stalagmite displays seasonal layering with one dark and one white layer deposited every year. The opposite seasonal behavior

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Delvaux De Fenffe, D.: Geologie et tectonique du Parc de Lesse et Lomme au bord sud du Bassin de Dinant (Rochefort, Belgique), *Bulletins de la Societe Belge de Geologie*, 94, 81–95, 1985.

Dreybrodt, W.: Chemical kinetics, speleothem growth and climate, *Boreas*, 28, 347–356, 1999.

Dreybrodt, W.: Evolution of the isotopic composition of carbon in a calcite precipitating $\text{H}_2\text{O}-\text{CO}_2-\text{CaCO}_3$ solution and the related isotopic composition of calcite in stalagmites, *Geochim. Cosmochim. Ac.*, 72, 4712–4724, 2008.

Fairchild, I. J., Smith, C. L., Baker, A., Fuller, L., Spotl, C., Matthey, D., McDermott, F., and Eimp: Modification and preservation of environmental signals in speleothems, *Earth-Sci. Rev.*, 75, 105–153, 2006.

Friedman, I. and O’Neil, J. R.: Compilation of stable isotope fractionation factors of geochemical interest, *Geolog. Surv. Prof. Paper 440-KK*, 177, 1977.

Frisia, S., Fairchild, I. J., Fohlmeister, J., Miorandi, R., Spoetl, C., and Borsato, A.: Carbon mass-balance modelling and carbon isotope exchange processes in dynamic caves, *Geochim. Cosmochim. Ac.*, 75, 380–400, 2011.

Genty, D. and Deflandre, G.: Drip flow variations under a stalactite of the Pere Noel cave (Belgium), evidence of seasonal variations and air pressure constraints, *J. Hydrol.*, 211, 208–232, 1998.

Genty, D. and Quinif, Y.: Annually laminated sequences in the internal structure of some Belgian stalagmites – importance for paleoclimatology, *J. Sediment. Res.*, 66, 275–288, 1996.

Genty, D., Blamart, D., Ouahdi, R., Gilmour, M., Baker, A., Jouzel, J., and Van-Exter, S.: Precise dating of Dansgaard–Oeschger climate oscillations in western Europe from stalagmite data, *Nature*, 421, 833–837, 2003.

Hendy, C. H.: Isotopic geochemistry of speleothems: 1. Calculations of effects on different modes of formation on isotopic composition of speleothems and their applicability as paleoclimatic indicators, *Geochim. Cosmochim. Ac.*, 35, 801–824, 1971.

Horita, J. and Clayton, R. N.: Comment on the studies of oxygen isotope fractionation between calcium carbonates and water at low temperatures by Zhou and Zheng (2003; 2005), *Geochim. Cosmochim. Ac.*, 71, 3131–3135, 2007.

Jimenez-Lopez, C., Caballero, E., Huertas, F. J., and Romanek, C. S.: Chemical, mineralogical and isotope behavior, and phase transformation during the precipitation of calcium carbonate minerals from intermediate ionic solution at 25 °C, *Geochim. Cosmochim. Ac.*, 65, 3219–3231, 2001.

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Jochum, K. P., Scholz, D., Stoll, B., Weis, U., Wilson, S. A., Yang, Q., Schwalb, A., Boerner, N., Jacob, D. E., and Andreae, M. O.: Accurate trace element analysis of speleothems and biogenic calcium carbonates by LA-ICP-MS, *Chem. Geol.*, 318, 31–44, 2012.
- Kim, S. T. and O'Neil, J. R.: Equilibrium and nonequilibrium oxygen isotope effects in synthetic carbonates, *Geochim. Cosmochim. Ac.*, 61, 3461–3475, 1997.
- Kim, S.-T., Mucci, A., and Taylor, B. E.: Phosphoric acid fractionation factors for calcite and aragonite between 25 and 75 °C: revisited, *Chem. Geol.*, 246, 135–146, 2007.
- Lachniet, M. S.: Climatic and environmental controls on speleothem oxygen-isotope values, *Quaternary Sci. Rev.*, 28, 412–432, 2009.
- Mattey, D., Lowry, D., Duffet, J., Fisher, R., Hodge, E., and Frisia, S.: A 53 year seasonally resolved oxygen and carbon isotope record from a modern Gibraltar speleothem: reconstructed drip water and relationship to local precipitation, *Earth Planet. Sc. Lett.*, 269, 80–95, 2008.
- McDermott, F.: Palaeo-climate reconstruction from stable isotope variations in speleothems: a review, *Quaternary Sci. Rev.*, 23, 901–918, 2004.
- McDermott, F.: Centennial-scale Holocene climate variability revealed by a high-resolution speleothem $\delta^{18}\text{O}$ record from SW Ireland (9 November, pg 1328, 2001), *Science*, 309, 1816–1816, 2005.
- McDermott, F., Schwarcz, H. P., and Rowe, P. J.: Isotopes in speleothems, in: *Isotopes in Paleoenvironmental Research*, edited by: Leng, M. J., Springer Netherlands, 185–226, 2005.
- Mook, W. G.: Volume 1: introduction-theory, methods, review, in: *Environmental Isotopes in the Hydrological Cycle, Principles and Applications*, International Atomic Energy Agency and United Nations Educational, Scientific and Cultural Organization, 185 pp., 2000.
- Mühlhngaus, C., Scholz, D., and Mangini, A.: Modelling stalagmite growth and $\delta^{13}\text{C}$ as a function of drip interval and temperature, *Geochim. Cosmochim. Ac.*, 71, 2780–2790, 2007.
- Mühlhngaus, C., Scholz, D., and Mangini, A.: Modelling fractionation of stable isotopes in stalagmites, *Geochim. Cosmochim. Ac.*, 73, 7275–7289, doi:10.1016/j.gca.2009.09.010, 2009.
- O'Neil, J. R., Clayton, R. N., and Mayeda, T. K.: Oxygen isotope fractionation in divalent metal carbonates *J. Chem. Phys.*, 51, 5547–5558, 1969.
- Oster, J. L., Montanez, I. P., and Kelley, N. P.: Response of a modern cave system to large seasonal precipitation variability, *Geochim. Cosmochim. Ac.*, 91, 92–108, 2012.
- Polag, D., Scholz, D., Muehlinghaus, C., Spoetl, C., Schroeder-Ritzrau, A., Segl, M., and Mangini, A.: Stable isotope fractionation in speleothems: laboratory experiments, *Chem. Geol.*, 279, 31–39, 2010.

**Seasonal variations
recorded in cave
monitoring results**M. Van Rampelbergh
et al.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Quinif, Y.: Une nouvelle topographie de la Grotte de Han, Lapiaz, revue de l'équipe Spéléo du Centre de Mons (Hors-Série), 15–18, 1988.
- Riechelmann, D. F. C., Schroeder-Ritzrau, A., Scholz, D., Fohlmeister, J., Spoetl, C., Richter, D. K., and Mangini, A.: Monitoring Bunker Cave (NW Germany): a prerequisite to interpret geochemical proxy data of speleothems from this site, *J. Hydrol.*, 409, 682–695, 2011.
- Riechelmann, D. F. C., Deininger, M., Scholz, D., Riechelmann, S., Schroeder-Ritzrau, A., Spoetl, C., Richter, D. K., Mangini, A., and Immenhauser, A.: Disequilibrium carbon and oxygen isotope fractionation in recent cave calcite: comparison of cave precipitates and model data, *Geochim. Cosmochim. Ac.*, 103, 232–244, 2013.
- Romanek, C. S., Grossman, E. L., and Morse, J. W.: Carbon Isotopic Fractionation in synthetic aragonite and calcite-effects of temperature and precipitation rate, *Geochim. Cosmochim. Ac.*, 56, 419–430, 1992.
- Rozanski, K., Araguas-Araguas, L., and Gonfiantini, R.: Relationship between long-term trends of ^{18}O isotope composition of precipitation and climate, *Science*, 258, 981–985, 1992.
- Ruan, J. and Hu, C.: Seasonal variations and environmental controls on stalagmite calcite crystal growth in Heshang Cave, Central China, *Chinese Sci. Bull.*, 55, 3929–3935, 2010.
- Scholz, D., Muehlinghaus, C., and Mangini, A.: Modelling $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ in the solution layer on stalagmite surfaces, *Geochim. Cosmochim. Ac.*, 73, 2592–2602, 2009.
- Spotl, C. and Matthey, D.: Stable isotope microsampling of speleothems for palaeoenvironmental studies: a comparison of microdrill, micromill and laser ablation techniques, *Chem. Geol.*, 235, 48–58, 2006.
- Tarutani, T., Clayton, R. N., and Mayeda, T. K.: Effect of polymorphism and magnesium substitution on oxygen isotope fractionation between calcium carbonate and water, *Geochim. Cosmochim. Ac.*, 33, 987–996, 1969.
- Treble, P., Shelley, J. M. G., and Chappell, J.: Comparison of high resolution sub-annual records of trace elements in a modern (1911–1992) speleothem with instrumental climate data from southwest Australia, *Earth Planet. Sc. Lett.*, 216, 141–153, 2003.
- Tremaine, D. M., Froelich, P. N., and Wang, Y.: Speleothem calcite formed in situ: modern calibration of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ paleoclimate proxies in a continuously-monitored natural cave system, *Geochim. Cosmochim. Ac.*, 75, 4929–4950, 2011.
- Van Rampelbergh, M., Fleitmann, D., Verheyden, S., Cheng, H., Edwards, L., De Geest, P., De Vleeschouwer, D., Burns, S. J., Matter, A., Claeys, P., and Keppens, E.: Mid- to late Holocene

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Indian Ocean Monsoon variability recorded in four speleothems from Socotra Island, Yemen, Quaternary Sci. Rev., 65, 129–142, 2013.

Verheyden, S.: Speleothems as Palaeoclimatic Archives, Isotopic and Geochemical Study of the Cave Environment and its Late Quaternary Records, unpublished Ph.D- thesis, Vrije Universiteit Brussel, Belgium, 132 pp., 2001.

Verheyden, S., Baele, J.-M., Keppens, E., Genty, D., Cattani, O., Hai, C., Edwards, L., Hucai, Z., Van Strijdonck, M., and Quinif, Y.: The proserpine stalagmite (Han-sur-Lesse cave, Belgium): preliminary environmental interpretation of the last 1000 years as recorded in a layered speleothem, Geol. Belg., 9, 245–256, 2006.

Verheyden, S., Genty, D., Deflandre, G., Quinif, Y., and Keppens, E.: Monitoring climatological, hydrological and geochemical parameters in the Pere Noel cave (Belgium): implication for the interpretation of speleothem isotopic and geochemical time-series, Int. J. Speleol., 37, 221–234, 2008a.

Verheyden, S., Nader, F. H., Cheng, H. J., Edwards, L. R., and Swennen, R.: Paleoclimate reconstruction in the Levant region from the geochemistry of a Holocene stalagmite from the Jeita cave, Lebanon, Quaternary Res., 70, 368–381, 2008b.

Wang, Y., Cheng, H., Edwards, R. L., Kong, X., Shao, X., Chen, S., Wu, J., Jiang, X., Wang, X., and An, Z.: Millennial- and orbital-scale changes in the East Asian monsoon over the past 224 000 years, Nature, 451, 1090–1093, 2008.

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Table 1. (a) Comparison between samples measured immediately after sampling and samples measured one month after sampling to test the reproducibility of the “new” method used in this study to measure the $\delta^{13}\text{C}$ composition of the DIC in the drip water. Results show no difference with the samples measured one month after sampling. **(b)** A comparison of the “new” method used in this study for sampling and sample preparation for delta C analyses of DIC in waters with the method of Matthey et al. (2008). Both methods deliver similar results confirming that the new method used in this study delivers reproducible results.

(a) New method used in this study							
		immediate measurement	average	measured one month after sampling	average		
Sample 1	a	-11.15	-11.19	-11.18	-11.20		
	b	-11.21		-11.32			
	c	-11.20		-11.24			
	d	-11.21		-11.05			
(b) New method used in this study				Method by Matthey et al. (2008)			
		immediate measurement	average	immediate measurement	average		
Sample 2	a	-11.78	-11.77	Sample 2	a	-11.76	-11.80
	b	-11.73			b	-11.82	
	c	-11.79			c	-11.76	
					d	-11.82	
				e	-11.84		

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh
et al.

Table 3. Calculated $\delta^{18}\text{O}$ values for calcite deposited in equilibrium its drip water for different authors. Values were calculated using the average drip water composition of -7.65‰ for the lowest, highest and average measured temperatures at the Salle-du-Dôme. The second part of the table shows the difference between the calculated $\delta^{18}\text{O}$ values and the $\delta^{18}\text{O}$ values measured on the glass slabs. Best results are given by the “in-cave”-relationship of Tremaine et al. (2011).

Author	Predicted $d^{18}\text{O}$ (in ‰)			Difference with Glass slab values (in ‰)		
	T_{\min} 10.5 °C	T_{mid} 12 °C	T_{\max} 14.5 °C	$d^{18}\text{O}_{\min}$ -6.25	$d^{18}\text{O}_{\text{mid}}$ -6.55	$d^{18}\text{O}_{\max}$ -6.98
O'Neil et. al (1969)	-6.44	-6.80	-7.39	0.19	0.25	0.41
Kim and O'Neil (1997)	-5.71	-6.04	-6.59	-0.54	-0.51	-0.39
Horita and Clayton (2007)	-6.96	-7.30	-7.85	0.71	0.75	0.87
Chacko and Deines (2008)	-6.98	-7.32	-7.86	0.73	0.77	0.88
Demeny et al. (2010)	-6.30	-6.62	-7.15	0.05	0.07	0.17
Tremaine et al. (2011)	-6.26 ± 0.09	-6.55 ± 0.08	-7.04 ± 0.06	0.01 ± 0.09	0 ± 0.08	0.06 ± 0.06

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.




Fig. 1. The Han-sur-Lesse karst system with the Han-sur-Lesse cave being the northern cave system and the Pere Noel cave being the southern cave system. The Lesse-river enters the cave system at the “Gouffre de Belvaux” and exits 24 h later at the “Trou de Han”. The studied speleothem is located in the “Salle-du-Dôme” and grows on a pile of debris. Figure adapted after Quinif (1988).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

⏪

⏩

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.




Fig. 2. Cave monitoring results show that the cave conditions vary seasonally between a “summer-mode” lasting from June to December and a “winter-mode” lasting from December to June. The $\delta^{18}\text{O}$ signal of the glass slabs (i) is deposited in equilibrium with the drip water and reflects the seasonally varying cave temperatures (c). The $\delta^{13}\text{C}$ composition of the glass slabs (m) is not deposited in equilibrium and variations are affected by the seasonally varying discharge amounts (e).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.




Fig. 3. The $\delta^{18}\text{O}$ composition of the drip water averages the annual mean composition of the rainwater $\delta^{18}\text{O}$. 6 small drops occur in the rainwater composition in November 2012, December 2012, May 2013, July 2013, December 2013 and January 2014 (indicated in light blue) and correspond with increased rainfall periods. At the end of March a cold temperature peak and prolonged snowfall cause the rainwater $\delta^{18}\text{O}$ to display a large drop. A similar negative drop is seen in the drip water $\delta^{18}\text{O}$ composition in August indicating that this drop in $\delta^{18}\text{O}$ values in the rainwater can be used as spike and shows that the residence time of the water in the epikarst is 4 to 5 months.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




Fig. 4. Detail of the upper 16 mm of the Proserpine core drilled in 2011. 10 additional dark layers are counted compared to the core of Verheyden et al. (2006) confirming the seasonal character of the layering. Ages are based in on layer counting with one dark and one white layer deposited every year. The frame indicates the location of the monthly resolved $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ signal.

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 5. The $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ signals from the Proserpine stalagmite anticorrelate at monthly scale. At the end of a dark layer (dotted lines), the $\delta^{18}\text{O}$ values reach their minimum values while the $\delta^{13}\text{C}$ values reach their maximum. Dark layers are formed during summer when cave temperatures are high (leading the low $\delta^{18}\text{O}$) and discharge rates are low (leading to high $\delta^{13}\text{C}$). The clearer the lamination, the large the amplitude of the variations in the isotopic composition.

Seasonal variations recorded in cave monitoring results

M. Van Rampelbergh et al.

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[⏪](#) | [⏩](#)

[◀](#) | [▶](#)

[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

