



## Abstract

For this study two sediment cores from the Peruvian shelf covering the time period between the Little Ice Age (LIA) and present were examined for changes in productivity (biogenic opal concentrations (bSi)), nutrient utilisation (stable isotope compositions of silicon ( $\delta^{30}\text{Si}_{\text{opal}}$ ) and nitrogen ( $\delta^{15}\text{N}_{\text{sed}}$ )), as well as in ocean circulation and material transport (authigenic and detrital radiogenic neodymium ( $\varepsilon_{\text{Nd}}$ ) and strontium ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) isotopes).

For the LIA the proxies recorded weak primary productivity and nutrient utilisation reflected by low average bSi concentrations of  $\sim 10\%$ ,  $\delta^{15}\text{N}_{\text{sed}}$  values of  $\sim +5\text{‰}$  and intermediate  $\delta^{30}\text{Si}_{\text{opal}}$  values of  $\sim +0.9\text{‰}$ . At the same time the radiogenic isotope composition of the detrital sediment fraction indicates dominant local riverine input of lithogenic material due to higher rainfall in the Andean hinterland. These patterns were caused by permanent El Niño-like conditions characterized by a deeper nutricline, weak upwelling and low nutrient supply. At the end of the LIA,  $\delta^{30}\text{Si}_{\text{opal}}$  dropped to low values of  $+0.6\text{‰}$  and opal productivity reached its minimum of the past 650 years. During the following transitional period of time the intensity of upwelling, nutrient supply and productivity increased abruptly as marked by the highest bSi contents of up to 38%, by  $\delta^{15}\text{N}_{\text{sed}}$  of up to  $\sim +7\text{‰}$ , and by the highest degree of silicate utilisation with  $\delta^{30}\text{Si}_{\text{opal}}$  reaching values of  $+1.1\text{‰}$ . At the same time detrital  $\varepsilon_{\text{Nd}}$  and  $^{87}\text{Sr}/^{86}\text{Sr}$  signatures documented increased wind strength and supply of dust to the shelf due to drier conditions. Since about 1870, productivity has been high but nutrient utilisation has remained at levels similar to the LIA indicating significantly increased nutrient availability.

Comparison between the  $\delta^{30}\text{Si}_{\text{opal}}$  and  $\delta^{15}\text{N}_{\text{sed}}$  signatures suggests that during the past 650 years the  $\delta^{15}\text{N}_{\text{sed}}$  signature in the Peruvian Upwelling area has most likely primarily been controlled by surface water utilisation and not, as previously assumed, by subsurface nitrogen loss processes in the water column.

CPD

10, 3357–3396, 2014

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 1 Introduction

Global climate of the late Holocene was disrupted by major anomalies, the most recent of which being the Little Ice Age (LIA) between ca. 1400 and 1850 AD (Lamb, 1965; Grove, 2001). During that time a weakening of the Walker circulation (Conroy et al., 2008), reduced influence of the South Pacific subtropical high (SPSH) along the Peruvian margin (Sifeddine et al., 2008; Gutiérrez et al., 2009; Salvattecí et al., 2014), and a southward shift of the mean position of the Intertropical Convergence Zone (ITCZ) and the associated precipitation belt compared to today (Sachs et al., 2009) caused pronounced changes in rainfall patterns in the tropics. El Niño-like warmer conditions in the Eastern South Pacific were accompanied by an intensified South American summer monsoon (Bird et al., 2011) resulting in ~10% higher precipitation in northeast Peru (~5° S; Rabatel et al., 2008) and up to 20–30% higher precipitation in the Bolivian Andes (~16° S; Reuter et al., 2009). On the one hand this caused growth and extension of the Andean glaciers (Vuille et al., 2008) and on the other it enabled human settlements in the presently hyperarid southern Peruvian Andes (Unkel et al., 2007). In the upwelling areas off Peru and the western South American shelf regions the main consequence of these climatic conditions during the LIA was a deepening of the nutricline and a strongly diminished biological productivity (Vargas et al., 2007; Sifeddine et al., 2008; Valdés et al., 2008; Gutiérrez et al., 2009).

Sediment cores from the Peruvian shelf covering the period of time from the LIA until present indicate that the marine realm was characterised by an abrupt biogeochemical regime shift towards modern conditions at the end of the LIA due to the northward movement of the ITCZ and an expansion of the SPSH. While low productivity and a more oxygenated water column prevailed during the LIA, markedly increased biological productivity and pronounced oxygen depletion over wide areas of the shelf have characterised the system since the end of the LIA (Vargas et al., 2007; Sifeddine et al., 2008; Gutiérrez et al., 2009; Salvattecí et al., 2014).

### Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Nutrient utilisation  
and weathering  
inputs**

C. Ehlert et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In this study the stable silicon isotope composition of sedimentary diatoms ( $\delta^{30}\text{Si}_{\text{opal}}$ ) covering the period of time from the LIA to the present is analysed. The main goal is the reconstruction of the factors controlling the dynamics of nutrient cycling together with oxygen in the Peruvian upwelling, in particular a comparison between the  $\delta^{30}\text{Si}_{\text{opal}}$  and the stable nitrogen isotope composition ( $\delta^{15}\text{N}_{\text{sed}}$ ) of sedimentary organic matter. Both  $\delta^{30}\text{Si}_{\text{opal}}$  and  $\delta^{15}\text{N}_{\text{sed}}$  provide information about utilisation of silicic acid ( $\text{Si}(\text{OH})_4$ ) and nitrate ( $\text{NO}_3^-$ ) during primary productivity, e.g. during the formation of diatom frustules and associated organic matter, respectively (Altabet and Francois, 1994; De La Rocha et al., 1997). Diatoms preferentially incorporate the lighter isotopes from the dissolved  $\text{Si}(\text{OH})_4$  and  $\text{NO}_3^-$  pools leaving the residual dissolved nutrients enriched in the heavier isotopes (Wada and Hattori, 1978; Altabet et al., 1991; De La Rocha et al., 1997). For the  $\delta^{30}\text{Si}_{\text{opal}}$  this fractionation process is mainly controlled by the availability of  $\text{Si}(\text{OH})_4$  in surface waters (e.g., De La Rocha et al., 1998; Brzezinski et al., 2002; Egan et al., 2012). The  $\delta^{15}\text{N}$  of  $\text{NO}_3^-$  is partly controlled by  $\text{NO}_3^-$  utilisation of marine organisms but is also affected by N-loss processes in the water column (denitrification, anammox) (Codispoti et al., 2001; Dalsgaard et al., 2003) resulting in a marked enrichment of the upwelling source waters in the heavier  $^{15}\text{NO}_3^-$  (Liu and Kaplan, 1989; Lam et al., 2009; given that it is currently not possible to distinguish between different N-loss processes from the sediments we will use the term denitrification for simplicity). Consequently, sedimentary  $\delta^{15}\text{N}_{\text{sed}}$  records from areas dominated by oxygen-depleted waters such as the shelf region off Peru are usually interpreted to directly reflect changes in the intensity of subsurface  $\text{NO}_3^-$  loss and the extent and strength of oxygen depletion (e.g., De Pol-Holz et al., 2007, 2009; Agnihotri et al., 2008; Gutiérrez et al., 2009) whereas the effect of  $\text{NO}_3^-$  utilisation on the preserved  $\delta^{15}\text{N}_{\text{sed}}$  is often neglected. Comparison of both isotope systems can therefore provide information about the degree of utilisation of  $\text{NO}_3^-$  and  $\text{Si}(\text{OH})_4$  versus the influence of  $\text{NO}_3^-$  loss processes. Increasing nutrient utilisation should result in a consistent increase in both  $\delta^{30}\text{Si}_{\text{opal}}$  and  $\delta^{15}\text{N}_{\text{sed}}$ . In con-

trast, a change in  $\text{NO}_3^-$  reduction due to varying oxygen depletion in the water column would affect only the  $\delta^{15}\text{N}_{\text{sed}}$  leaving the  $\delta^{30}\text{Si}_{\text{opal}}$  unaffected.

The main forces driving surface productivity and subsurface oxygenation off Peru at centennial time scales during the past two millennia have been changes in the strength of the Walker circulation and in the expansion/contraction of the SPSH (Gutierrez et al., 2009; Salvattecchi et al., 2014). Therefore, the radiogenic isotope compositions of neodymium ( $\epsilon_{\text{Nd}}$ ) and strontium ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) of the authigenic ferromanganese (Fe-Mn) oxyhydroxide coatings of the sedimentary particles, which are expected to record the radiogenic isotope compositions of past bottom waters, as well as of the detrital fraction of the sediment were examined. These proxy data provide information about changes of (surface ocean) circulation and of transport processes, provenance of the sediments, and input mechanisms of terrigenous material as a function of changes in precipitation on land during the transition from wetter LIA-conditions to drier modern conditions. Weathering of continental source rocks delivers lithogenic particles of different origin and age to the shelf, which have distinct radiogenic isotope signatures ( $\epsilon_{\text{Nd}_{\text{detritus}}}$ ,  $^{87}\text{Sr}/^{86}\text{Sr}_{\text{detritus}}$ ) that can be used to trace their source areas (Goldstein et al., 1984). Central Peruvian Andean rocks have more radiogenic  $\epsilon_{\text{Nd}}$  signatures whereas southern Peruvian rocks are characterised by less radiogenic  $\epsilon_{\text{Nd}}$  signatures (Sarbas and Nohl, 2009), which is also reflected in the sediments along the shelf (Ehlert et al., 2013). Changes in detrital material input and transport pathways are generally closely related to climatic changes causing variations in the supply from the respective source areas (e.g. Grousset et al., 1988). It should therefore be possible to detect the transition from wetter LIA-conditions with higher local input from central Peru via rivers due to higher precipitation rates towards the drier presently prevailing conditions with an increased influence of eolian material transport from further south in the Atacama desert (Molina-Cruz, 1977) and deposition along the shelf after the LIA.

CPD

10, 3357–3396, 2014

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2 Material and methods

### 2.1 Core locations and age models

For the reconstruction of surface water  $\text{Si}(\text{OH})_4$  utilisation and terrestrial material input and transport for the period of time between the LIA and present two sediment cores with high sedimentation rates were analysed. Box core B0405-6 was recovered from the upper continental slope off Pisco at  $14^{\circ}07.9' \text{ S}$ ,  $76^{\circ}30.1' \text{ W}$  in a water depth of 299 m with the Peruvian R/V *José Olaya Balandra* in 2004 (Fig. 1) (Gutiérrez et al., 2006). The age model was previously published by Gutiérrez et al. (2009) and is based on downcore profiling of the activities of  $^{241}\text{Am}$ , excess  $^{210}\text{Pb}$ , and on radiocarbon ages obtained from bulk sedimentary organic carbon, which document that the core covers the past  $\sim 650$  years. The second core, multicorer M771-470, was taken at  $11^{\circ} \text{ S}$ ,  $77^{\circ}56.6' \text{ W}$  in 145 m water depth during cruise M77/1 with the German R/V Meteor in 2008 (Fig. 1). The age model was obtained by measuring excess  $^{210}\text{Pb}$  activities and modeling of the resulting profiles as described by Meysman et al. (2005) (for details see Supplement). Ages prior to  $\sim 1850$  AD were inferred using sedimentation rates from nearby core B0405-13 (Gutiérrez et al., 2009).

### 2.2 Methods

#### 2.2.1 Biogenic opal and silicon isotope analyses

The biogenic opal (bSi) contents in both cores were measured following the sequential leaching techniques described by DeMaster (1981) and Müller and Schneider (1993). Si isotope analyses were performed on the 11–32  $\mu\text{m}$  diatom-fraction that was extracted from the sediment applying the procedures described by Morley et al. (2004).

Approximately 300 mg of sediment were treated with 30 %  $\text{H}_2\text{O}_2$  and 35 % HCl to remove organic matter and carbonate. Afterwards the sediment was wet-sieved to separate the 11–32  $\mu\text{m}$  fraction. In a third step a heavy-liquid solution (sodium-

CPD

10, 3357–3396, 2014

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



polytungstate, 2.1–2.2 g mL<sup>-1</sup>) was applied in several steps to separate diatoms from the detrital lithogenic silicate material. All samples were screened under the microscope to verify their purity with respect to the detrital (clay) fraction.

The diatom samples were then transferred into Teflon vials and dissolved in 1 mL 0.1 M NaOH and diluted with MQ water according to Reynolds et al. (2008). More details are provided in Ehlert et al. (2012). Si concentrations of the dissolved diatom samples were measured colorimetrically using a photospectrometer (Hansen and Koroleff, 1999). Chromatographic separation and purification of the Si was achieved with 1 mL pre-cleaned AG50W-X8 cation exchange resin (mesh 200–400) (Georg et al., 2006; as modified by de Souza et al., 2012). Si isotope ratios were measured on a *NuPlasma HR* MC-ICPMS (Nu Instruments) at GEOMAR equipped with an adjustable source-defining slit, which can be set to medium resolution to ensure separation of the <sup>30</sup>Si peak from molecular interferences. The measurements were carried out applying standard-sample bracketing (Albarède et al., 2004). All solutions were measured at a Si concentration of 14–21 μmol kg<sup>-1</sup> of samples and standards depending on the performance of the instrument on the respective measurement day and were introduced into the plasma via a Cetac Aridus II desolvating nebulizer system equipped with a PFA nebulizer operated at a 60 to 80 μL min<sup>-1</sup> uptake rate. Si isotope compositions are reported in the δ<sup>30</sup>Si notation as deviations of the measured <sup>30</sup>Si/<sup>28</sup>Si from the NIST standard NBS28 in parts per thousand (‰). Repeated measurements of the reference materials IRMM018 and Big Batch gave average δ<sup>30</sup>Si values of -1.52 ± 0.18‰ (2σ<sub>(sd)</sub>) and -10.84 ± 0.18‰ (2σ<sub>(sd)</sub>), respectively, which are in good agreement with values obtained by other laboratories (Reynolds et al., 2007). Samples were measured three to five times within a one-day session and measurements were repeated on at least two separate days. The resulting uncertainties ranged between 0.04 and 0.23‰ (2σ<sub>(sd)</sub>) (Tables 1, 2). Replicate measurements of an in-house diatom matrix standard over longer periods of time gave an external reproducibility of 0.11‰ (2σ<sub>(sd)</sub>). Error bars provided in the figures correspond to that external reproducibility unless the uncertainties of the repeated sample measurements were higher.

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2.2 Neodymium and strontium isotope analyses

To obtain the radiogenic isotope composition of past bottom seawater at the sites of the sediment cores from the early diagenetic Fe-Mn coatings of the sediment particles, previously published methods were applied (Gutjahr et al., 2007; see Supplement for details). The residual detrital material was leached repeatedly to remove remaining coatings and was then treated with a mixture of concentrated HF-HNO<sub>3</sub>-HCl for total dissolution. The separation and purification of Nd and Sr in the leachates and in the completely dissolved detrital sediment fraction followed previously published procedures for Nd (Cohen et al., 1988) and Sr (Horwitz et al., 1992) applying ion exchange chromatography for separation of Rb/Sr from the rare earth elements (REEs) (0.8 mL AG50W-X12 resin, mesh 200–400) followed by separation of Sr from Rb (50 µL Sr-Spec resin, mesh 50–100), and separation of Nd from the other REEs (2 mL Eichrom Ln-Spec resin, mesh 50–100). All radiogenic isotope measurements were performed on the *NuPlasma HR MC-ICPMS* (Nu Instruments) at GEOMAR. Measured Nd isotope compositions were corrected for instrumental mass bias using a <sup>146</sup>Nd/<sup>144</sup>Nd of 0.7219 and were normalised to the accepted <sup>143</sup>Nd/<sup>144</sup>Nd literature value of 0.512115 of the JNdi-1 standard (Tanaka et al., 2000). All values are given as ε<sub>Nd</sub>, which corresponds to the measured <sup>143</sup>Nd/<sup>144</sup>Nd, normalised to the Chondritic Uniform Reservoir CHUR (0.512638), multiplied by 10 000. The external reproducibility was estimated by repeated measurements of the JNdi-1 standard and was always better than 20 ppm (2σ<sub>(sd)</sub> Tables 1, 2). Measured <sup>87</sup>Sr/<sup>86</sup>Sr ratios were corrected for instrumental mass bias using <sup>88</sup>Sr/<sup>86</sup>Sr = 8.3752 and were normalised to the accepted value for NIST SRM987 of 0.710245. The 2σ<sub>(sd)</sub> external reproducibility of repeated standard measurements was always better than 36 ppm (2σ<sub>(sd)</sub> Tables 1, 2). Procedural Nd and Sr blanks for leachates and total dissolutions of the detrital material were ≤ 83 pg and 2.1 ng, respectively, and thus negligible compared to the concentrations of the samples.

### Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







## 4 Discussion

After the end of the LIA around 1820 AD the mean position of the ITCZ shifted northward (Sachs et al., 2009) causing an intensification of alongshore winds and enhanced coastal upwelling off the Peruvian coast (Sifeddine et al., 2008; Gutiérrez et al., 2009), diminished coastal sea surface temperatures (Vargas et al., 2007), and a decrease in precipitation on land (Rabatel et al., 2008; Bird et al., 2011). Records of productivity and redox conditions based on  $\delta^{15}\text{N}_{\text{sed}}$  and the Mo and Cd content of the sediments indicate a rapid change of the biogeochemical composition of the source waters to higher nutrient concentrations causing higher biological productivity and lower subsurface oxygen, which have persisted until the present day (Sifeddine et al., 2008; Gutiérrez et al., 2009; Salvattecí et al., 2014). The shift after the end of the LIA constitutes a major anomaly of late Holocene climate in the Eastern Pacific, which was of the same order of magnitude as the changes in conditions off Chile during the Younger Dryas (De Pol-Holz et al., 2006). This study focuses on the reconstruction of the regime shift from the LIA and a transitional period towards modern conditions and its controlling factors including the evolution of nutrient utilisation and changes in the advection of water masses and material transport.

### 4.1 Changes in biological productivity and nutrient consumption

#### 4.1.1 Evolution of surface water productivity and nutrient utilisation

The pronounced change in the biogeochemical regime from low productivity during the LIA to higher productivity during the transitional and modern period thereafter is documented by several sedimentary records from the EEP region and has been dated at  $\sim 1820$  AD (Sifeddine et al., 2008; Gutierrez et al., 2009; Díaz-Ochoa et al., 2009, 2011; Salvattecí et al., 2014). Similarly, both cores M771-470 from  $11^\circ$  S and B0405-6 from  $14^\circ$  S off Pisco show the characteristic coeval pronounced increase in bSi and total N content (Fig. 2a, e) and  $\text{C}_{\text{org}}$  concentration (not shown here) after the end of the LIA

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and during the transition period. Therefore, three time periods that show distinct differences in productivity and nutrient utilisation have been identified from our records and will be discussed in the following: the LIA, the transition period from the LIA to modern conditions between  $\sim 1820$  and  $\sim 1870$  AD, and modern conditions after  $\sim 1870$  AD.

Both cores recorded a two- to threefold increase in bSi content from 10–12% prior to the end of the LIA to values of up to 27% in M771-470 and up to 38% in B0405-6 during the transition period (Fig. 2a, e). Afterwards the bSi contents decreased again but have remained at a level of  $\sim 20\%$  and thus significantly higher than prior to the end of the LIA. The increase in bSi content is also reflected by a marked increase in diatom accumulation rate in core B0405-6 (Fig. 2e) (Gutiérrez et al., 2009). Analyses of the downcore diatom assemblages have shown that the high diatom accumulation rates and bSi content in core B0405-6 during the transition period were associated with diatom layers dominated by *Skeletonema costatum* (Gutiérrez et al., 2009), a species that is today more abundant when upwelling is more intense during Austral winter/spring.

Both cores are characterised by a very high correlation between total N and  $C_{org}$  content ( $r^2 = 0.95$  for core M771-470 and 0.8 for core B0405-6, respectively) (Gutiérrez et al., 2009; this study). In contrast, bSi and total N contents do not co-vary throughout the records (Fig. 3a). Surface sediments from the Peruvian shelf region between the Equator and  $\sim 18^\circ$  S show a relatively weak but positive correlation between bSi and N contents ( $r^2 = 0.5$ , Fig. 3a) (Ehlert et al., 2012; Mollier-Vogel et al., 2012). Similar to the surface sediments, bSi and total N concentrations in core M771-470 are positively correlated, whereas they essentially do not correlate in core B0405-6. This is because the bSi maximum at the end of the transition period was more pronounced in core B0405-6 and higher than surface sediment bSi contents anywhere along the shelf region off Peru. At the same time only a rather gradual increase in total N content with some excursions to low values during the transition period occurred (Fig. 2a, e). The total N concentration also did not always co-vary with  $\delta^{15}N_{sed}$  (Fig. 3c). In particular,

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the samples from the late transition period show very low total N concentrations but high  $\delta^{15}\text{N}_{\text{sed}}$ , high  $\delta^{30}\text{Si}_{\text{opal}}$  and bSi content.

Sedimentary  $\delta^{15}\text{N}_{\text{sed}}$  data, which are only available for core B0405-6, show a shift from lower values around +4‰ to +5‰ during the LIA to higher values around +7‰ after the end of the LIA and have remained at that level since then (Fig. 2f) (Gutiérrez et al., 2009). The values in the younger part of the record are in good agreement with surface sediment  $\delta^{15}\text{N}_{\text{sed}}$  data measured in the main Peruvian upwelling region ranging from +6‰ to +9‰ (Mollier-Vogel et al., 2012). Bulk  $\delta^{15}\text{N}_{\text{sed}}$  signatures measured in core B0405-13 from 12° S (184 m water depth) close to the location of core M771-470 can be used for comparison and show very similar values, amplitude, and variability as core B0405-6 (Fig. 2i) (Gutiérrez et al., 2009). In contrast to  $\delta^{15}\text{N}_{\text{sed}}$ , the  $\delta^{30}\text{Si}_{\text{opal}}$  signatures, which mainly reflect changes in surface water nutrient utilisation, are not only characterised by a simple increase at the end of the LIA. Instead, both  $\delta^{30}\text{Si}_{\text{opal}}$  records closely follow the evolution of the bSi concentrations and show intermediate  $\delta^{30}\text{Si}_{\text{opal}}$  signatures between +0.8‰ and +0.9‰ during the LIA, a pronounced short-term decrease to +0.6‰ at the end of the LIA, which was followed by a marked increase to values around +1.1‰ during the transition period, and finally a return to intermediate values between +0.8‰ and +1.0‰ in the modern part of the records (Fig. 2b, f). The correspondence between bSi content and  $\delta^{30}\text{Si}_{\text{opal}}$  is more pronounced in core B0405-6 (Fig. 3b), which shows a higher variability and amplitude of bSi content. The difference in the  $\delta^{30}\text{Si}_{\text{opal}}$ ,  $\delta^{15}\text{N}_{\text{sed}}$ , bSi and total N content records during the transition from LIA to modern conditions reflects the different environmental factors controlling the proxies, which will be discussed in the following sections.

#### 4.1.2 Present day surface water utilisation versus subsurface nitrate loss

Diatoms are the dominant phytoplankton group of the Peruvian upwelling region (Estrada and Blasco, 1985; Bruland et al., 2005). The  $\delta^{30}\text{Si}_{\text{opal}}$  of these diatoms is primarily controlled by surface water diatom productivity and  $\text{Si}(\text{OH})_4$  utilisation (De La

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rocha et al., 1998; Brzezinski et al., 2002; Egan et al., 2012) but off Peru the  $\delta^{30}\text{Si}_{\text{opal}}$  has also been shown to be dependent on the isotopic signature of the advected surface and subsurface water masses (Ehlert et al., 2012). Similarly, the  $\delta^{15}\text{N}_{\text{sed}}$  of the organic matter is controlled by N isotope fractionation during  $\text{NO}_3^-$  uptake by phytoplankton, mostly diatoms. Off Peru, however, the  $\text{NO}_3^-$  supplied to the surface waters has previously been enriched in  $^{15}\text{NO}_3^-$  due to upwelling of oxygen-depleted subsurface waters, which had undergone significant  $\text{NO}_3^-$  – loss processes (mostly denitrification, but also anammox processes, associated with a high fractionation of up to 20–30‰) (Lam et al., 2009; Altabet et al., 2012). Bulk sediment  $\delta^{15}\text{N}_{\text{sed}}$  in areas with oxygen-depleted waters is therefore usually interpreted to reflect changes in the intensity of subsurface  $\text{NO}_3^-$  reduction and the extent and strength of the oxygen minimum zone (Altabet et al., 1999; De Pol-Holz et al., 2007; Agnihotri et al., 2008; Gutiérrez et al., 2009). The direct comparison of  $\delta^{30}\text{Si}_{\text{opal}}$ , reflecting mostly utilisation, and  $\delta^{15}\text{N}_{\text{sed}}$ , reflecting both utilisation and  $\text{NO}_3^-$  reduction, from core B0405-6 off Pisco will therefore provide insights into the strength of  $\text{NO}_3^-$  reduction in the past.

Subsurface water column  $\delta^{15}\text{N}_{\text{NO}_3^-}$  data from the present-day Peruvian shelf are isotopically very heavy, in particular along the southern shelf region between 10° S and 17° S, where values of up to +25‰ are reached due to the increasing oxygen deficit and intensification of  $\text{NO}_3^-$  – loss processes (Mollier-Vogel et al., 2012; Altabet et al., 2012). These isotopically enriched waters are upwelled along the shelf and represent the source for organic matter production in the surface waters. Therefore, it is expected, that the deposited sedimentary organic matter reflects these enriched subsurface water signatures. However, the latitudinal increase in surface sediment  $\delta^{15}\text{N}_{\text{sed}}$  from the same shelf region to maximum mean values around +9‰ is much lower than that measured in the water column (Mollier-Vogel et al., 2012). The reason for this observation is that the  $\delta^{15}\text{N}_{\text{sed}}$  signal in the southern shelf region (10–17° S) did not fully record the  $^{15}\text{NO}_3^-$  enrichment in the water column but is a combination of the isotopic effects associated with subsurface  $\text{NO}_3^-$  – loss and incomplete surface water  $\text{NO}_3^-$  utilisation



## Nutrient utilisation and weathering inputs

C. Ehlert et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



very close to a 1 : 1 ratio. In contrast, the samples from the end of the LIA and also the recent samples are shifted slightly towards the denitrification curve indicating a higher influence of  $\text{NO}_3^-$ -loss processes. This is particularly the case for the samples from the end of the LIA, which have the lowest  $\delta^{30}\text{Si}_{\text{opal}}$  but at the same time already show a strong increase in  $\delta^{15}\text{N}_{\text{sed}}$  to values of near +6‰. The most likely explanation is that upwelling was strongly increased during those brief periods resulting in high nutrient supply, high productivity, and either more complete  $\text{NO}_3^-$  utilisation (Gutiérrez et al., 2009) or increased  $\text{NO}_3^-$  – loss caused by enhanced subsurface oxygen depletion. Overall, however, the utilisation signal appears to have dominated both the Si and N isotope records.

If, however, the  $\delta^{15}\text{N}_{\text{sed}}$  is dominated by utilisation it is interesting that in the cores (both B0405-6 and -13)  $\delta^{15}\text{N}_{\text{sed}}$  and proxies for sediment redox conditions (e.g. molybdenum concentrations) are strongly coupled throughout the record (Sifeddine et al., 2008; Gutiérrez et al., 2009). One direct interpretation could be that the diatom blooms, and subsequently the degradation of the organic matter, strongly control the oxygen availability in the sediments after sedimentation and burial. Therefore, increased diatom productivity and higher  $\text{Si}(\text{OH})_4$  and  $\text{NO}_3^-$  utilisation would result in an increase in  $\delta^{15}\text{N}_{\text{sed}}$ . At the same time more oxygen is consumed during degradation of the organic matter in the sediments causing more reducing conditions in the sediments. Consequently, a change in the subsurface water column structure, e.g. enhanced re-supply of oxygen via ocean currents, may not be reflected in the  $\delta^{15}\text{N}_{\text{sed}}$  record.

### 4.1.4 Modelling the surface water utilisation

Following the above considerations we will try to quantify past utilisation based on our data. The theoretical relationship between the degree of surface water nutrient utilisation and the stable isotope composition of Si and N can be described assuming either Rayleigh-type (single input followed by no additional nutrients newly supplied to a particular parcel of water followed by fractional loss as a function of production and

export) or steady state (continuous supply and partial consumption of nutrients causing a dynamic equilibrium of the dissolved nutrient concentration and the product) fractionation behaviour (Fig. 5) (Mariotti et al., 1981). The lighter isotopes are preferentially incorporated into the diatom frustules and the organic matter, respectively, leaving the dissolved fraction enriched in the heavier isotopes (Wada and Hattori, 1978; Altabet et al., 1991; De La Rocha et al., 1997). The fractionation between  $\delta^{30}\text{Si}$  in seawater and  $\delta^{30}\text{Si}$  in the produced diatom opal has generally been assumed to be  $-1.1\text{‰}$  (De La Rocha et al., 1997) whereas between  $\delta^{15}\text{N}_{\text{NO}_3^-}$  of seawater and  $\delta^{15}\text{N}$  of the newly formed organic matter it is usually between  $-3\text{‰}$  to  $-6\text{‰}$  (DeNiro and Epstein, 1981; Minagawa and Wada, 1984). Here we adopted  $-5\text{‰}$ , which corresponds to present-day conditions along the central Peruvian shelf (Mollier-Vogel et al., 2012).

Along the Peruvian shelf region biological productivity in the euphotic zone is driven by upwelling of nutrients from subsurface waters. For the calculation of the utilisation of these nutrients, an initial  $\delta^{30}\text{Si}_{\text{Si(OH)}_4}$  of  $+1.5\text{‰}$  (Ehlert et al., 2012) and an initial  $\delta^{15}\text{N}_{\text{NO}_3^-}$  of  $+9\text{‰}$  (Mollier-Vogel et al., 2012) for the upwelled water masses at  $14^\circ\text{S}$  is assumed. The lower mean  $\delta^{15}\text{N}_{\text{sed}}$  of about  $+5\text{‰}$  and  $\delta^{30}\text{Si}_{\text{opal}}$  of  $+0.7\text{‰}$  signatures during the LIA in the southerly core B0405-6 correspond to a dissolved  $\delta^{15}\text{N}_{\text{NO}_3^-}$  and  $\delta^{30}\text{Si}_{\text{Si(OH)}_4}$  isotope signature of the surface waters of  $+10\text{‰}$  and  $+1.8\text{‰}$  and a calculated  $\text{NO}_3^-$  and  $\text{Si(OH)}_4$  utilisation of only 20–30 % for steady state-type fractionation (Fig. 5b) and 35–50 % for Rayleigh-type fractionation (Fig. 5a) behaviour, respectively. The highest mean values of  $+1.1\text{‰}$  for  $\delta^{30}\text{Si}_{\text{opal}}$  and  $+6.8\text{‰}$  for  $\delta^{15}\text{N}_{\text{sed}}$  for the transition period correspond to a much higher utilisation of  $\sim 60\%$  for steady state-type fractionation and  $\sim 80\%$  assuming Rayleigh-type fractionation. Consequently, the calculated utilisation of available  $\text{Si(OH)}_4$  and  $\text{NO}_3^-$  more than doubled, whereby bSi concentrations and diatom accumulation rates increased by about a factor of three (Fig. 2e).

The changes in  $\text{Si(OH)}_4$  and  $\text{NO}_3^-$  utilisation were of the same order of magnitude and reflect low nutrient utilisation during the LIA and much higher degree of utilisation

CPD

10, 3357–3396, 2014

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





mental conditions, e.g. source water signature and isotope enrichment during utilisation, and (2) a change in the dominating diatom assemblages has to be considered.

The assumed source water  $\delta^{30}\text{Si}_{\text{Si(OH)}_4}$  and  $\delta^{15}\text{N}_{\text{NO}_3^-}$  values of +1.5‰ and +9‰ (Figs. 5, 6), respectively, were measured in the present day subsurface waters under strong upwelling conditions during which high amounts of nutrients are supplied to the euphotic zone (Ehlert et al., 2012; Mollier-Vogel et al., 2012). Under strong upwelling conditions the bottom waters on the shallow shelf are today dominated by the southward directed high-nutrient Peru-Chile Undercurrent (PCUC) (Fig. 1) (Brink et al., 1983). Under LIA-conditions (prevailing El Niño-like conditions), however, atmospheric and oceanic circulation was different; the nutricline was deeper as a consequence of a weak Walker circulation and the upwelling favourable winds were weaker as a consequence of the SPSH contraction (Salvatteci et al., 2014). In fact, the pumped waters were likely nutrient-depleted, because the Ekman layer did not reach the subsurface nutrient-rich waters. Under these conditions, the subtropical and equatorial nutrient-depleted surface water masses may have occupied the entire surface layer in the coastal realm because they expanded both latitudinally and vertically in the water column (Montes et al., 2011). This may have changed the source water isotopic signatures and would therefore also change the calculated degrees of utilisation. If, for example, the assumed source water  $\delta^{15}\text{N}_{\text{NO}_3^-}$  was +6‰ instead of +9‰ (e.g. due to weaker subsurface  $\text{NO}_3^-$  – loss and weaker  $^{15}\text{N}$  enrichment during the LIA) the down-core  $\delta^{15}\text{N}_{\text{sed}}$  data of core B0405-6 would all plot closer to the denitrification curve (Fig. 4b). Redox proxies from the records, indeed, indicate a weaker OMZ (Gutiérrez et al., 2009), which would make a lower  $\delta^{15}\text{N}_{\text{NO}_3^-}$  in source waters likely. However, to date there is no reliable information if and how much the source water  $\delta^{30}\text{Si}_{\text{Si(OH)}_4}$  and  $\delta^{15}\text{N}_{\text{NO}_3^-}$  signatures changed over time.

Varying upwelling and nutrient supply conditions also cause changes in the dominating diatom assemblages. Recent results from culturing experiments suggest species-dependent enrichment factors for diatom- $\delta^{30}\text{Si}$  (–0.5‰ to –2.1‰; Sutton et al., 2013)

CPD

10, 3357–3396, 2014

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Nutrient utilisation and weathering inputs

C. Ehlert et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



and also diatom frustule-bound  $\delta^{15}\text{N}$  ( $-1.9\text{‰}$  to  $-11.2\text{‰}$ ; Horn et al., 2011). This raises the question whether a change in diatom assemblages may have been the cause for the observed downcore  $\delta^{30}\text{Si}_{\text{opal}}$  and, to a lesser extent, the bulk  $\delta^{15}\text{N}_{\text{sed}}$  variations. The quasi-monospecific diatom layers from the transition period 1820–1870 AD consist mainly of *Skeletonema costatum* (Gutiérrez et al., 2009), for which an enrichment factor  $\varepsilon$  of  $-1.0\text{‰}$  similar to the applied  $-1.1\text{‰}$  was determined (De la Rocha et al., 1997). The younger sediments are also dominated by upwelling-indicative species such as *Thalassionema nitzschioides* and *Chaetoceros* sp. (Abrantes et al., 2007), whereby *Chaetoceros brevis*, a species from the Southern Ocean, has been shown to have a much higher  $\varepsilon$  of  $-2.1\text{‰}$  (Sutton et al., 2013). That means, assuming the same surface water  $\delta^{30}\text{Si}_{\text{Si(OH)}_4}$  signatures, a dominance of *Chaetoceros* sp. in the sediments should result in a lower  $\delta^{30}\text{Si}_{\text{opal}}$  whereas assemblages dominated by *Skeletonema costatum* should be characterised by higher  $\delta^{30}\text{Si}_{\text{opal}}$  signatures, which is exactly what core B0405-6 shows. Consequently, the difference in  $\delta^{30}\text{Si}_{\text{opal}}$  over time could reflect the change in diatom assemblage and not a change in nutrient utilisation. On the other hand, *Chaetoceros brevis* is a polar species and it is not clear whether off Peru it undergoes the same high fractionation factor during frustule growth. Analysis have shown that the offset between modern surface water  $\delta^{30}\text{Si}_{\text{Si(OH)}_4}$  and surface sediment  $\delta^{30}\text{Si}_{\text{opal}}$  along the central Peruvian shelf is between  $-1.1\text{‰}$  and  $-1.3\text{‰}$  (Ehlert et al., 2012), which indicates that either the enrichment factor for the dominating *Chaetoceros* species off Peru does not deviate significantly from  $-1.1\text{‰}$  or that the mixing of different diatoms in the sediment samples overprints any isotopic excursions of single species caused by higher or lower fractionation factors. Given the paucity of data of the fractionation factors for the dominant diatom species off Peru, the importance of the role of downcore changes in the assemblage composition is hard to determine. However, in the following we investigate variations in past water mass circulation, upwelling conditions, as well as material input and transport to reconstruct the source water conditions.

## 4.2 Changes in detrital material input and transport

The radiogenic isotope composition of the lithogenic particles ( $\epsilon_{\text{Nd detritus}}$  and  $^{87}\text{Sr}/^{86}\text{Sr}_{\text{detritus}}$ ) of the sediments provides useful information about the source region of material and therefore about changes in material input and transport, either eolian or via ocean currents (e.g. Grousset et al., 1988). Surface sediments along the Peruvian shelf show highly variable signatures, which have overall more radiogenic  $\epsilon_{\text{Nd detritus}}$  values in the North and much less radiogenic  $\epsilon_{\text{Nd detritus}}$  values in the South off southern Peru and northern Chile (Ehlert et al., 2013). This north-south trend is a consequence of the southward increasing contributions of material input from the adjacent Andean hinterland rocks. The Andean rocks along the northwestern South American region display a wide range in  $\epsilon_{\text{Nd}}$  and  $^{87}\text{Sr}/^{86}\text{Sr}$  signatures (Fig. 6) (Sarbas and Nohl, 2009) varying from highly radiogenic  $\epsilon_{\text{Nd}}$  around 0 and unradiogenic  $^{87}\text{Sr}/^{86}\text{Sr}$  values around 0.704 in the equatorial region in northern Peru to much less radiogenic  $\epsilon_{\text{Nd}}$  mostly below  $-4$  and more radiogenic  $^{87}\text{Sr}/^{86}\text{Sr}$  mostly above 0.705 in southern Peru and northern Chile. The sedimentary  $\epsilon_{\text{Nd detritus}}$  and  $^{87}\text{Sr}/^{86}\text{Sr}_{\text{detritus}}$  records of the two cores off Callao at  $11^\circ\text{S}$  and off Pisco at  $14^\circ\text{S}$  show broad similarities, but also some differences. Both cores recorded a significant change in  $\epsilon_{\text{Nd detritus}}$  and  $^{87}\text{Sr}/^{86}\text{Sr}_{\text{detritus}}$ , and therefore a change in provenance, at the end of the LIA and during the transition period. Core M771-470, although being located further north, is overall characterised by less radiogenic  $\epsilon_{\text{Nd detritus}}$  values than core B0405-6 (Figs. 2, 6). The  $\epsilon_{\text{Nd detritus}}$  of core M771-470 recorded a trend from less radiogenic towards more radiogenic values prior to the end of the LIA, followed by a step of 1.5  $\epsilon_{\text{Nd}}$  units towards less radiogenic values, which afterwards remained at that level. In contrast, the  $\epsilon_{\text{Nd detritus}}$  record of core B0405-6 remained at a level around  $-3.6$  during the LIA and then slightly increased to maximum values of up to  $-2.5$  in the younger part. The  $^{87}\text{Sr}/^{86}\text{Sr}_{\text{detritus}}$  record in both cores is mainly characterised by a rapid shift towards more radiogenic values at the end of the LIA, whereby the change was much more pronounced in core M771-470 (Fig. 2d, h). The youngest samples of the cores are in good agreement with

measurements of surface sediments from the same area (Figs. 2, 6) (Ehlert et al., 2013). The variability in core M771-470 display the same magnitude as the complete glacial-interglacial variation in core SO147-106KL located at 12° S off Lima (Ehlert et al., 2013). All data of both M771-470 and B0405-6 plot within the provenance fields of southern Peru and northern Chile (Fig. 6).

Today, material input along the Peruvian shelf occurs mostly via riverine and minor eolian input (Molina-Cruz, 1977; Scheidegger and Krissek, 1982). The LIA was characterised by wetter conditions (Haug et al., 2001; Gutiérrez et al., 2009) due to the mean southward migration of the ITCZ and the associated precipitation belt, which also caused more intense rainfall in the central Andean hinterland (Rabatel et al., 2008; Reuter et al., 2009). Most terrigenous particles and weathering products such as clay minerals from the LIA show indications of increased riverine transport and discharge (Sifeddine et al., 2008; Salvattecchi et al., 2014). Consequently, material input during the LIA was dominated by local sources due to the higher river discharge. After the end of the LIA, when the climate became drier and alongshore winds became stronger, riverine input diminished and eolian dust input increased. The wind-blown dust has mainly originated from the Atacama Desert located in the southern Peruvian and northern Chilean Andes (Molina-Cruz, 1977). This material has less radiogenic  $\epsilon_{Nd}$  and much more radiogenic  $^{87}Sr/^{86}Sr$  values (Fig. 6) (Sarbas and Nohl, 2009). The record of core M771-470 is in agreement with this. The LIA-sediments indicate a local origin, probably via riverine input, whereas the younger sediments display characteristics from a more southerly origin and therefore increased eolian sources, possibly from the Atacama Desert. The signatures and overall small variations in core B0405-6 are much more difficult to interpret. There are fewer rivers in Southern Peru around Pisco compared to the Callao region. Therefore, riverine-derived material from northern and central Peru, which is transported via the PCUC, can get dispersed further south and can be deposited in the Pisco region. On the other hand, the influence of eolian deposition should be much higher at the southern core location. During the LIA river input increased in southern Peru as well, whereas eolian deposition was low. The invariable signature ob-

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



served might be the result of mixing of sediment from the different sources. Also, in comparison to core M771-470 core B0405-6 is located much closer to the coast, which most likely diminished the differences in material input and transport between the LIA and modern conditions.

In summary, our combined proxies demonstrate that during the LIA upwelling and nutrient supply, productivity and utilisation were low, because eolian wind forcing was low and the source waters of the upwelling carried less nutrients. In contrast, after the end of the LIA and especially in more recent times, the strong remineralisation of nutrients from subsurface waters fuelling enhanced productivity was only possible due to an increase in trade wind strength as also reflected by higher dust supply probably driven by an expansion of the SPSH, and a shoaling of the thermocline/nutricline due to a stronger Walker circulation.

## 5 Conclusions

Productivity, nutrient utilisation and material provenance proxies ( $\delta^{30}\text{Si}_{\text{opal}}$ ,  $\delta^{15}\text{N}_{\text{sed}}$ ,  $\epsilon_{\text{Nd}_{\text{detritus}}}$ , and  $^{87}\text{Sr}/^{86}\text{Sr}_{\text{detritus}}$ ) from two cores from the Peruvian shelf recorded significant changes in surface water  $\text{Si}(\text{OH})_4$  and  $\text{NO}_3^-$  concentration and utilisation due to changes in upwelling intensity and nutrient supply. During the LIA the overall nutrient content in the water column and in surface waters was low because the upwelling source waters contained less nutrients. Consequently, the Peruvian upwelling regime was characterised by persistent reduced primary productivity. The reasons for this were likely a contraction of the South Pacific Subtropical High and a weaker Walker circulation that produced a weakening of alongshore winds and a deepening of the nutricline.

During the LIA, the stronger rainfall associated with more moist conditions during prevailing El Niño-like conditions was recorded by the radiogenic isotope composition of the detrital material along the shelf, which was mainly transported via rivers from the Andean hinterland. At the end of the LIA, in accordance with a northward shift

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the ITCZ and an intensification of wind strength a higher dust transport of particles associated to drier conditions and eolian forcing is reflected by the radiogenic isotope composition of the detrital sediments. These conditions are also reflected in upwelling strength, with a rapid shoaling of the thermocline and nutricline, as well as enhanced nutrient supply and productivity to the surface waters. During a transition period a strong increase in diatom blooming events doubled the  $\text{Si}(\text{OH})_4$  and  $\text{NO}_3^-$  utilisation compared to the LIA, and was also higher than present day utilisation. After that transition period more persistent non-El Niño conditions favoured a high productivity accompanied with moderate utilisation of nutrients. The utilisation rates were on the same order as during the LIA, but productivity was much higher, which reflects the much higher concentration of nutrients in surface waters.

Most studies of past coastal upwelling regions have argued so far that the sedimentary  $\delta^{15}\text{N}_{\text{sed}}$  records were dominated by the large N isotope fractionation signature occurring during  $\text{NO}_3^-$  loss processes (denitrification or anammox) in oxygen-depleted subsurface waters upwelling at the coast. Comparison between  $\delta^{30}\text{Si}_{\text{opal}}$  and  $\delta^{15}\text{N}_{\text{sed}}$  in the same sediment samples of our study and assuming similar source water signatures as today, however, indicate that the  $\delta^{15}\text{N}_{\text{sed}}$  signatures to a large extent reflect expected utilisation signals, which has important implications for the reconstruction of variations in the intensity of oxygen depletion, the N cycle of the past and its controlling factors.

**The Supplement related to this article is available online at  
doi:10.5194/cpd-10-3357-2014-supplement.**

*Acknowledgements.* This work is a contribution of Sonderforschungsbereich 754 “Climate – Biogeochemistry Interactions in the Tropical Ocean” (www.sfb754.de), which is supported by the Deutsche Forschungsgemeinschaft. We acknowledge the help of Jutta Heinze in the laboratory of GEOMAR for the biogenic opal concentration measurements. We thank Ulrike Lomnitz and Klaus Wallmann for their help with the  $^{210}\text{Pb}$  dating and the establishment

of the age model of core M771-470.

The service charges for this open access publication  
have been covered by a Research Centre of the  
5 Helmholtz Association.

## References

Abrantes, F., Lopes, C., Mix, A. C., and Pisias, N. G.: Diatoms in Southeast Pacific surface sediments reflect environmental properties, *Quaternary. Sci. Rev.*, 26, 155–169, doi:10.1016/j.quascirev.2006.02.022, 2007.

10 Agnihotri, R., Altabet, M. A., Herbert, T. D., and Tierney, J. E.: Subdecadally resolved paleoceanography of the Peru margin during the last two millennia, *Geochem. Geophys. Geosy.*, 9, Q05013, doi:10.1029/2007GC001744, 2008.

Albarède, F., Telouk, P., Blichert-Toft, J., Boyet, M., Agranier, A., and Nelson, B.: Precise and accurate isotopic measurements using multiple-collector ICPMS, *Geochim. Cosmochim. Ac.*,  
15 68, 2725–2744, doi:10.1016/j.gca.2003.11.024, 2004.

Altabet, M. A., Deuser, W. G., Honjo, S., and Stienen, C.: Seasonal and depth-related changes in the source of sinking particles in the North Atlantic, *Nature*, 354, 136–139, 1991.

Altabet, M. A. and Francois, R.: Sedimentary nitrogen isotopic ratio as a recorder for surface ocean nitrate utilisation, *Global. Biogeochem. Cy.*, 8, 103–116, 1994.

20 Altabet, M. A., Pilskałn, C., Thunell, R. C., Pride, C. J., Sigman, D. M., Chavez, F. P., and Francois, R.: The nitrogen isotope biogeochemistry of sinking particles from the margin of the Eastern North Pacific, *Deep-Sea Res. I*, 46, 655–679, 1999.

Altabet, M. A., Ryabenko, E., Stramma, L., Wallace, D. W. R., Frank, M., Grasse, P., and Lavik, G.: An eddy-stimulated hotspot for fixed nitrogen-loss from the Peru oxygen minimum zone, *Biogeosciences*, 9, 4897–4908, doi:10.5194/bg-9-4897-2012, 2012.

25 Bird, B. W., Abbott, M. B., Vuille, M., Rodbell, D. T., Stansell, N. D., and Rosenmeier, M. F.: A 2300-year-long annually resolved record of the South American summer monsoon from the Peruvian Andes, *P. Natl. Acad. Sci.*, 108, 8583–8588, doi:10.1073/pnas.1003719108, 2011.

30 Brink, K. H., Halpern, D., Huyer, A., and Smith, R. L.: The Physical Environment of the Peruvian Upwelling System, *Prog. Oceanogr.*, 12, 285–305, 1983.

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Nutrient utilisation and weathering inputs

C. Ehlert et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Bruland, K. W., Rue, E. L., Smith, G. J., and DiTullio, G. R.: Iron, macronutrients and diatom blooms in the Peru upwelling regime: brown and blue waters of Peru, *Mar. Chem.*, 93, 81–103, doi:10.1016/j.marchem.2004.06.011, 2005.
- Brzezinski, M. A.: The Si:C:N ratio of marine diatoms: interspecific variability and the effect of some environmental variables, *J. Phycol.*, 21, 347–357, 1985.
- Brzezinski, M. A., Pride, C. J., Franck, V. M., Sigman, D. M., Sarmiento, J. L., Matsumoto, K., Gruber, N., Rau, G. H., and Coale, K. H.: A switch from Si(OH)<sub>4</sub> to NO<sub>3</sub><sup>-</sup> depletion in the glacial Southern Ocean, *Geophys. Res. Lett.*, 29, 3–6, doi:10.1029/2001GL014349, 2002.
- Codispoti, L. A., Brandes, J. A., Christensen, J. P., Devol, A. H., Naqvi, S. W. A., Paerl, H. W., and Yoshinory, T.: The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the anthropocene?, *Sci. Mar.*, 65, 85–105, 2001.
- Cohen, A. S., O’Nions, R. K., Siegenthaler, R., and Griffin, W. L.: Chronology of the pressure-temperature history recorded by a granulite terrain, *Contrib. Mineral. Petr.*, 98, 303–311, 1988.
- Conroy, J. L., Restrepo, A., Overpeck, J. T., Steinitz-Kannan, M., Cole, J. E., Bush, M. B., and Colinvaux, P. A.: Unprecedented recent warming of surface temperatures in the eastern tropical Pacific Ocean, *Nat. Geosci.*, 2, 46–50, doi:10.1038/ngeo390, 2008.
- Dalsgaard, T., Canfield, D. E., Petersen, J., Thamdrup, B., and Acuna-González, J.: N<sub>2</sub> production by the anammox reaction in the anoxic water column of Golfo Dulce, Costa Rica, *Nature*, 422, 606–608, 2003.
- De La Rocha, C. L., Brzezinski, M. A., and DeNiro, M. J.: Fractionation of silicon isotopes by marine diatoms during biogenic silica formation, *Geochim. Cosmochim. Ac.*, 61, 5051–5056, 1997.
- De La Rocha, C. L., Brzezinski, M. A., DeNiro, M. J., and Shemesh, A.: Silicon-isotope composition of diatoms as an indicator of past oceanic changes, *Nature*, 395, 680–683, 1998.
- De Pol-Holz, R., Robinson, R. S., Hebbeln, D., Sigman, D. M., and Ulloa, O.: Controls on sedimentary nitrogen isotopes along the Chile margin, *Deep-Sea Res. II*, 56, 1042–1054, doi:10.1016/j.dsr2.2008.09.014, 2009.
- De Pol-Holz, R., Ulloa, O., Dezileau, L., Kaiser, J., Lamy, F., and Hebbeln, D.: Melting of the Patagonian Ice Sheet and deglacial perturbations of the nitrogen cycle in the eastern South Pacific, *Geophys. Res. Lett.*, 33, L04704, doi:10.1029/2005GL024477, 2006.

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



De Pol-Holz, R., Ulloa, O., Lamy, F., Dezileau, L., Sabatier, P., and Hebbeln, D.: Late Quaternary variability of sedimentary nitrogen isotopes in the eastern South Pacific Ocean, *Paleoceanography*, 22, PA2207, doi:10.1029/2006PA001308, 2007.

De Souza, G. F., Reynolds, B. C., Rickli, J., Frank, M., Saito, M. A., Gerringa, L. J. A., and Bourdon, B.: Southern Ocean control of silicon stable isotope distribution in the deep Atlantic Ocean, *Global Biogeochem. Cy.*, 26, GB2035, doi:10.1029/2011GB004141, 2012.

DeMaster, D. J.: The supply and accumulation of silica in the marine environment, *Geochim. Cosmochim. Ac.*, 45, 1715–1732, 1981.

DeNiro, M. J. and Epstein, S.: Influence of diet on the distribution of nitrogen isotopes in animals, *Geochim. Cosmochim. Ac.*, 45, 341–351, 1981.

Díaz-Ochoa, J. A., Lange, C. B., Pantoja, S., De Lange, G. J., Gutiérrez, D., Munoz, P., and Salamanca, M.: Fish scales in sediments from off Callao, central Peru, *Deep-Sea Res. II*, 56, 1113–1124, doi:10.1016/j.dsr2.2008.09.015, 2009.

Díaz-Ochoa, J. A., Pantoja, S., De Lange, G. J., Lange, C. B., Sánchez, G. E., Acuña, V. R., Muñoz, P., and Vargas, G.: Oxygenation variability in Mejillones Bay, off northern Chile, during the last two centuries, *Biogeosciences*, 8, 137–146, doi:10.5194/bg-8-137-2011, 2011.

Egan, K. E., Rickaby, R. E. M., Leng, M. J., Hendry, K. R., Sloane, H. J., Bostock, H. C., and Halliday, A. N.: Diatom silicon isotopes as a proxy for silicic acid utilisation: A Southern Ocean core top calibration, *Geochim. Cosmochim. Ac.*, 96, 174–192, doi:10.1016/j.gca.2012.08.002, 2012.

Ehlert, C., Grasse, P., Mollier-Vogel, E., Bösch, T., Franz, J., De Souza, G. F., Reynolds, B. C., Stramma, L., and Frank, M.: Factors controlling the silicon isotope distribution in waters and surface sediments of the Peruvian coastal upwelling, *Geochim. Cosmochim. Ac.*, 99, 128–145, doi:10.1016/j.gca.2012.09.038, 2012.

Ehlert, C., Grasse, P., and Frank, M.: Changes in silicate utilisation and upwelling intensity off Peru since the Last Glacial Maximum – insights from silicon and neodymium isotopes, *Quaternary Sci. Rev.*, 72, 18–35, doi:10.1016/j.quascirev.2013.04.013, 2013.

Estrada, M. and Blasco, D.: Phytoplankton assemblages in coastal upwelling areas, in: *Simpósio Internacional Sobre Las Areas de Afloramiento Mas Importantes del Oeste Africano (Cabo Blanco y Benguela)*, edited by: Bas, C., Margalef, R., and Rubies, P., Barcelona: Instituto de Investigaciones Pesqueras, 379–402, 1985.

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Georg, R. B., Reynolds, B. C., Frank, M., and Halliday, A. N.: New sample preparation techniques for the determination of Si isotopic compositions using MC-ICPMS, *Chem. Geol.*, 235, 95–104, doi:10.1016/j.chemgeo.2006.06.006, 2006.

Goldstein, S. L., O’Nions, R. K., and Hamilton, P. J.: A Sm – N d isotopic study of atmospheric dusts and particulates from major river systems, *Earth Planet. Sc. Lett.*, 70, 221–236, 1984.

Grasse, P., Stichel, T., Stumpf, R., Stramma, L., and Frank, M.: The distribution of neodymium isotopes and concentrations in the Eastern Equatorial Pacific: Water mass advection versus particle exchange, *Earth Planet. Sc. Lett.*, 353–354, 198–207, doi:10.1016/j.epsl.2012.07.044, 2012.

Grousset, F. E., Biscaye, P. E., Zindler, A., Prospero, J., and Chester, R.: Neodymium isotopes as tracers in marine sediments and aerosols: North Atlantic, *Earth Planet. Sc. Lett.*, 87, 367–378, 1988.

Grove, M. J.: The initiation of the “Little Ice Age” in regions round the North Atlantic, *Climatic Change*, 48, 53–82, 2001.

Gutiérrez, D., Sifeddine, A., Field, D. B., Ortlieb, L., Vargas, G., Chávez, F. P., Velazco, F., Ferreira, V., Tapia, P., Salvattecchi, R., Boucher, H., Morales, M. C., Valdés, J., Reyss, J.-L., Campusano, A., Boussafir, M., Mandeng-Yogo, M., García, M., and Baumgartner, T.: Rapid reorganization in ocean biogeochemistry off Peru towards the end of the Little Ice Age, *Biogeosciences*, 6, 835–848, doi:10.5194/bg-6-835-2009, 2009.

Gutiérrez, D., Sifeddine, A., Reyss, J. L., Vargas, G., Velazco, F., Salvattecchi, R., Ferreira, V., Ortlieb, L., Field, D., Baumgartner, T., Boussafir, M., Boucher, H., Valdés, J., Marinovic, L., Soler, P., and Tapia, P.: Anoxic sediments off Central Peru record interannual to multidecadal changes of climate and upwelling ecosystem during the last two centuries, *Adv. Geosci.*, 6, 119–125, doi:10.5194/adgeo-6-119-2006, 2006.

Horn, M. G., Robinson, R. S., Rynearson, T. A., and Sigman, D. M.: Nitrogen isotopic relationship between diatom-bound and bulk organic matter of cultured polar diatoms, *Paleoceanography*, 26, PA3208, doi:10.1029/2010PA002080, 2011.

Horwitz, E. P., Chiarizia, R., and Dietz, M. L.: A Novel Strontium-Selective Extraction Chromatographic Resin, *Solvent Extr. Ion Exc.*, 10, 313–336, doi:10.1080/07366299208918107, 1992.

Hutchins, D. A., Hare, C. E., Weaver, R. S., Zhang, Y., Firme, G. F., DiTullio, G. R., Alm, M. B., Riseman, S. F., Maucher, J. M., Geesey, M. E., Trick, C. G., Smith, G. J., Rue, E. L.,

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Conn, J., and Bruland, K. W.: Phytoplankton iron limitation in the Humboldt Current and Peru Upwelling, *Limnol. Oceanogr.*, 47, 997–1011, 2002.

Kessler, W. S.: The circulation of the eastern tropical Pacific: A review, *Prog. Oceanogr.*, 69, 181–217, doi:10.1016/j.pocean.2006.03.009, 2006.

5 Lacan, F. and Jeandel, C.: Tracing Papua New Guinea imprint on the central Equatorial Pacific Ocean using neodymium isotopic compositions and Rare Earth Element patterns, *Earth Planet. Sc. Lett.*, 186, 497–512, 2001.

Lam, P., Lavik, G., Jensen, M. M., Van de Vossenberg, J., Schmid, M., Woebken, D., Gutiérrez, D., Amann, R., Jetten, M. S. M., and Kuypers, M. M. M.: Revising the nitrogen cycle in the  
10 Peruvian oxygen minimum zone, *P. Natl. Acad. Sci.*, 106, 4752–4757, 2009.

Lamb, H. H.: The early Medieval Warm Epoch and its sequel, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 1, 13–37, 1965.

Liu, K.-K. and Kaplan, I. R.: The eastern tropical Pacific as a source of  $^{15}\text{N}$ -enriched nitrate in seawater off southern California, *Limnol. Oceanogr.*, 34, 820–830, 1989.

15 Mariotti, A., Germon, J. C., Hubert, P., Kaiser, P., Letolle, R., Tardieux, A., and Tardieux, P.: Experimental determination of nitrogen kinetic isotope fractionation: some principles; illustration for the denitrification and nitrification processes, *Plant. Soil.*, 62, 413–430, 1981.

Meysman, F. J. R., Boudreau, B. P., and Middelburg, J. J.: Modeling reactive transport in sediments subject to bioturbation and compaction, *Geochim. Cosmochim. Ac.*, 69, 3601–3617, doi:10.1016/j.gca.2005.01.004, 2005.

20 Minagawa, M. and Wada, E.: Stepwise enrichment of  $^{15}\text{N}$  along food chains: Further evidence and the relation between  $\delta^{15}\text{N}$  and animal age, *Geochim. Cosmochim. Ac.*, 48, 1135–1140, 1984.

Molina-Cruz, A.: The Relation of the Southern Trade Winds to Upwelling Processes during the Last 75,000 Years, *Quaternary Res.*, 8, 324–338, 1977.

25 Mollier-Vogel, E., Ryabenko, E., Martinez, P., Wallace, D. W. R., Altabet, M. A., and Schneider, R. R.: Nitrogen isotope gradients off Peru and Ecuador related to upwelling, productivity, nutrient uptake and oxygen deficiency, *Deep-Sea Res. I*, 70, 14–25, doi:10.1016/j.dsr.2012.06.003, 2012.

30 Montes, I., Schneider, W., Colas, F., Blanke, B., and Echevin, V.: Subsurface connections in the eastern tropical Pacific during La Niña 1999–2001 and El Niño 2002–2003, *J. Geophys. Res.*, 116, C12022, doi:10.1029/2011JC007624, 2011.

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Morley, D. W., Leng, M. J., Mackay, A. W., Sloane, H. J., Rioual, P., and Battarbee, R. W.: Cleaning of lake sediment samples for diatom oxygen isotope analysis, *Journal of Paleolimnology*, 31, 391–401, 2004.
- Müller, P. J. and Schneider, R. R.: An automated leaching method for the determination of opal in sediments and particulate matter, *Deep-Sea Res. I*, 40, 425–444, 1993.
- Piepgas, D. J. and Wasserburg, G. J.: Isotopic Composition of Neodymium in Waters from the Drake Passage, *Science*, 217, 207–214, 1982.
- Rabatel, A., Francou, B., Jomelli, V., Naveau, P., and Grancher, D.: A chronology of the Little Ice Age in the tropical Andes of Bolivia (16°S) and its implications for climate reconstruction, *Quaternary Res.*, 70, 198–212, doi:10.1016/j.yqres.2008.02.012, 2008.
- Reuter, J., Stott, L., Khider, D., Sinha, A., Cheng, H., and Edwards, R. L.: A new perspective on the hydroclimate variability in northern South America during the Little Ice Age, *Geophys. Res. Lett.*, 36, L21706, doi:10.1029/2009GL041051, 2009.
- Reynolds, B. C., Aggarwal, J., André, L., Baxter, D. C., Beucher, C. P., Brzezinski, M. A., Engström, E., Georg, R. B., Land, M., Leng, M. J., Opfergelt, S., Rodushkin, I., Sloane, H. J., Van den Boorn, S. H. J. M., Vroon, P. Z., and Cardinal, D.: An inter-laboratory comparison of Si isotope reference materials, *J. Anal. Atom. Spectrom.*, 22, 561, doi:10.1039/b616755a, 2007.
- Reynolds, B. C., Frank, M., and Halliday, A. N.: Evidence for a major change in silicon cycling in the subarctic North Pacific at 2.73 Ma, *Paleoceanography*, 23, PA4219, doi:10.1029/2007PA001563, 2008.
- Sachs, J. P., Sachse, D., Smittenberg, R. H., Zhang, Z., Battisti, D. S., and Golubic, S.: Southward movement of the Pacific intertropical convergence zone AD 1400–1850, *Nat. Geosci.*, 2, 519–525, doi:10.1038/NGEO554, 2009.
- Salvatteci, R., Gutiérrez, D., Field, D., Sifeddine, A., Ortlieb, L., Bouloubassi, I., Boussafir, M., Boucher, H., and Cetin, F.: The response of the Peruvian Upwelling Ecosystem to centennial-scale global change during the last two millennia, *Clim. Past*, 10, 715–731, doi:10.5194/cp-10-715-2014, 2014.
- Sarbas, B. and Nohl, U.: The GEOROC database – A decade of “online geochemistry”, *Geochim. Cosmochim. Ac.*, (Goldschmidt Abstracts), A1158, 2009.
- Scheidegger, K. F. and Krissek, L. A.: Dispersal and deposition of eolian and fluvial sediments off Peru and northern Chile, *Geol. Soc. Am. Bull.*, 93, 150–162, doi:10.1130/0016-7606(1982)93<150:DADOEA>2.0.CO;2, 1982.

## Nutrient utilisation and weathering inputs

C. Ehlert et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




- Sifeddine, A., Gutiérrez, D., Ortlieb, L., Boucher, H., Velazco, F., Field, D. B., Vargas, G., Boussafir, M., Salvattecí, R., Ferreira-Bartrina, V., García, M., Valdés, J., Caquineau, S., Mandeng-Yogo, M., Cetin, F., Solis, J., Soler, P., and Baumgartner, T.: Laminated sedi-  
ments from the central Peruvian continental slope: A 500 year record of upwelling sys-  
tem productivity, terrestrial runoff and redox conditions, *Prog. Oceanogr.*, 79, 190–197,  
doi:10.1016/j.pocean.2008.10.024, 2008.
- Sutton, J. N., Varela, D. E., Brzezinski, M. A., and Beucher, C. P.: Species-dependent sil-  
icon isotope fractionation by marine diatoms, *Geochim. Cosmochim. Ac.*, 104, 300–309,  
doi:10.1016/j.gca.2012.10.057, 2013.
- Takeda, S.: Influence of iron availability on nutrient consumption ratio of diatoms in oceanic  
waters, *Nature*, 393, 774–777, 1998.
- Tanaka, T., Togashi, S., Kamioka, H., Amakawa, H., Kagami, H., Hamamoto, T., Yuhara, M., Ori-  
hashi, Y., Yoneda, S., Shimizu, H., Kunimaru, T., Takahashi, K., Yanagi, T., Nakano, T., Fuji-  
maki, H., Shinjo, R., Asahara, Y., Tanimizu, M., and Dragusanu, C.: JNdi-1: a neodymium iso-  
topic reference in consistency with LaJolla neodymium, *Chem. Geol.*, 168, 279–281, 2000.
- Unkel, I., Kadereit, A., Mächtle, B., Eitel, B., Kromer, B., Wagner, G., and Wackler, L.: Dating  
methods and geomorphic evidence of palaeoenvironmental changes at the eastern margin  
of the South Peruvian coastal desert (14°30' S) before and during the Little Ice Age, *Quatern.  
Int.*, 175, 3–28, doi:10.1016/j.quaint.2007.03.006, 2007.
- Valdés, J., Ortlieb, L., Gutiérrez, D., Marinovic, L., Vargas, G., and Sifeddine, A.: 250 years  
of sardine and anchovy scale deposition record in Mejillones Bay, northern Chile, *Prog.  
Oceanogr.*, 79, 198–207, doi:10.1016/j.pocean.2008.10.002, 2008.
- Vargas, G., Pantoja, S., Rutllant, J. A., Lange, C. B., and Ortlieb, L.: Enhancement of coastal  
upwelling and interdecadal ENSO-like variability in the Peru-Chile Current since late 19th  
century, *Geophys. Res. Lett.*, 34, L13607, doi:10.1029/2006GL028812, 2007.
- Vuille, M., Francou, B., Wagnon, P., Juen, I., Kaser, G., Mark, B. G., and Bradley, R. S.: Climate  
change and tropical Andean glaciers: Past, present and future, *Earth-Sci. Rev.*, 89, 79–96,  
doi:10.1016/j.earscirev.2008.04.002, 2008.
- Wada, E. and Hattori, A.: Nitrogen isotope effects in the assimilation of inor-  
ganic nitrogenous compounds by marine diatoms, *Geomicrobiol. J.*, 1, 85–101,  
doi:10.1080/01490457809377725, 1978.





Nutrient utilisation  
and weathering  
inputs

C. Ehlert et al.



**Figure 1.** Schematic circulation patterns in the Eastern Equatorial Pacific. Surface currents (solid lines): (n)SEC: (northern) South Equatorial Current, PCC: Peru-Chile Current, PCoastalC: Peru Coastal Current; subsurface currents (dashed lines): EUC Equatorial Undercurrent, PCUC Peru-Chile Undercurrent (after Brink, 1983; Kessler, 2006), the inset shows the detailed location of cores M771-470, B0405-6 and B0405-13 (grey dots). The bathymetry is given for 0 to 1000 m water depth in 100 m increments.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





Nutrient utilisation  
and weathering  
inputs

C. Ehlert et al.



**Figure 3.** Surface sediment (white triangles) and downcore data (core M771-470: black squares, B0405-6: grey diamonds, B0405-13: white circles) for **(a)** total N versus bSi concentrations (the dashed line marks the end of the LIA), **(b)**  $\delta^{30}\text{Si}_{\text{opal}}$  versus bSi concentration and **(c)**  $\delta^{15}\text{N}_{\text{sed}}$  versus total N concentrations. Error bars represent  $2\sigma_{(\text{sd})}$  external reproducibilities.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Figure 4.** Relationship between  $\delta^{15}\text{N}_{\text{sed}}$  versus  $\delta^{30}\text{Si}_{\text{opal}}$  for **(a)** surface sediments and **(b)** downcore data from core B0405-6. The crosses in **(a)** indicate  $\delta^{30}\text{Si}$  data obtained from hand-picked diatoms, which reflect a different growth season than bulk  $\delta^{30}\text{Si}_{\text{opal}}$  and which are influenced by stronger  $\text{Si}(\text{OH})_4$  limitation (higher  $\delta^{30}\text{Si}$ ) (Ehlert et al., 2012). The solid lines reflect theoretical utilisation (assuming 1 : 1 utilisation of  $\text{Si}(\text{OH})_4$  and  $\text{NO}_3^-$  by the diatoms) and the dashed lines mark the theoretically expected line for denitrification, which represent the expected signal preserved in the sediments, based on present-day measurements:  $\delta^{30}\text{Si}$  source signature and enrichment factor  $\varepsilon_{\text{diatom-Si}(\text{OH})_4}$  are always +1.5‰ (Ehlert et al., 2012) and -1.1‰ (De La Rocha et al., 1997), respectively.  $\delta^{15}\text{N}_{\text{sed}}$  source signature and  $\varepsilon_{\text{organic-NO}_3^-}$  vary with latitude (Mollier-Vogel et al., 2012), in the north at 3.6° S source signature and  $\varepsilon$  are +5.7‰ and -3.7‰ (red curves), along the central shelf at 13.7° S source signature and  $\varepsilon$  are +8.9‰ and -4.8‰ (green curves), and in the south at 17° S source signature and  $\varepsilon$  were measured to be +14.5‰ and -5.7‰ (blue curves), respectively. The samples are colour-coded according to their location on the shelf and relative to the  $\text{NO}_3^-$  utilisation/ $\text{NO}_3^-$ -loss that they experienced. Data points that plot above the utilisation curves reflect predominant  $\text{Si}(\text{OH})_4$  limitation whereas data points below record stronger  $\text{NO}_3^-$  limitation. The isotopic enrichment during denitrification was always set to be +20‰. For the downcore data **(b)** two different assumed source signatures are displayed: +9‰ (green lines, corresponding to the modern conditions along the central shelf region in **(a)**) and +6‰ (grey lines). Data points are colour-coded according to the respective time periods (black: LIA, white: transition period, grey: modern). Error bars represent  $2\sigma_{(\text{sd})}$  external reproducibilities.

**Nutrient utilisation and weathering inputs**


C. Ehlert et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Nutrient utilisation  
and weathering  
inputs


C. Ehlert et al.



**Figure 5.** Theoretical changes in  $\delta^{30}\text{Si}$  and  $\delta^{15}\text{N}$  values of seawater and the instantaneous and accumulated product as a function of  $f$  (remaining nutrients from the available pool =  $[\text{nutrient}_{\text{observed}}]/[\text{nutrient}_{\text{initial}}]$ ) with an initial  $\delta^{30}\text{Si}_{\text{Si(OH)}_4}$  value of +1.5 and  $\delta^{15}\text{N}_{\text{NO}_3^-}$  of +9‰. The formation of the product, e.g. diatom opal, follows either **(a)** Rayleigh-type fractionation or **(b)** steady state-type fractionation behaviour, with enrichment factors  $\epsilon$  of  $-1.1\text{‰}$  ( $\delta^{30}\text{Si}$ ) and  $-5\text{‰}$  ( $\delta^{15}\text{N}$ ) (corresponding to conditions along the modern central Peruvian shelf, see Fig. 4). The colour shadings mark the range of measured mean  $\delta^{30}\text{Si}_{\text{opal}}$  (both cores) and  $\delta^{15}\text{N}_{\text{sed}}$  (B0405-6 only) in the cores for the LIA (red), the transition period (blue) and modern sediments (green), from which the respective nutrient utilisation (%) can be deduced.

Nutrient utilisation  
and weathering  
inputs

C. Ehlert et al.



**Figure 6.**  $\epsilon_{\text{Nd,detritus}}$  versus  $^{87}\text{Sr}/^{86}\text{Sr}_{\text{detritus}}$  for core M771-470 (red diamonds) and B0405-6 (blue squares). Error bars represent  $2\sigma_{(\text{sd})}$  external reproducibilities. The green dots are data obtained from surface sediment samples at different sites on the Peruvian shelf. The grey shadings indicate potential sources and provenance endmembers of the detrital material.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

