

## **Response to the editor, Mart 27, 2015**

Dear Editor, In the final revised manuscript we have accounted for all of your corrections and suggestions.

Thank you very much.  
Dmitry Demezhko, Anastasia Gornostaeva

### **List of corrections**

#### **P1 L11:**

We use early obtained in the Middle Urals geothermal reconstruction of the ground surface temperature (GST) history to determine the surface heat flux (SHF) history over the past 35 kyr.

#### **is replaced by**

We use [geothermal reconstruction of the ground surface temperature \(GST\) history early obtained in the Middle Urals](#) to determine the surface heat flux (SHF) history over the past 35 kyr.

#### **P1 L16:**

...35-6 kyr BP (the linear correlation coefficient  $R = 0.99$ ) points to...

#### **is replaced by**

...35-6 kyr BP [with](#) the linear correlation coefficient  $R = 0.99$  points to...

#### **P2 L12-15:**

In the paper we first present the SHF history for the past 35 kyr obtained from GST early reconstructed on the basis of temperature-depth profile logged in the Urals superdeep borehole (Demezhko and Shchapov, 2001). The recently developed improved algorithm of GST-SHF transformation (Gornostaeva, 2014) was applied to estimate the SHF history.

#### **is replaced by**

[In the paper we use geothermal reconstruction of GST history early obtained in the Middle Urals \(Demezhko and Shchapov, 2001\) to determine the SHF history over the past 35 kyr. The recently developed improved algorithm of GST-SHF transformation \(Gornostaeva, 2014\) was applied to estimate the SHF history. Paleoclimatic interpretation of GST and SHF histories is based on comparison them with orbitally driven solar insolation and atmospheric carbon dioxide changes.](#)

**P2 L19:**

according to the Fourier's equation

**is replaced by**

according to Fourier's equation

**P2 L21-22**

where  $q$  is SHF,  $t$  is time,  $\lambda$  is thermal conductivity,  $T(z, t)$  is temperature anomaly at a depth  $z$ .

**is replaced by**

where  $q$  is SHF ( $\text{W/m}^2$ ),  $t$  is time,  $\lambda$  is thermal conductivity ( $\text{Wm}^{-1}\text{K}^{-1}$ ),  $T(z, t)$  is temperature anomaly ( $^{\circ}\text{C}$ ) at a depth  $z$ .

**P3 L1**

where  $a = \frac{\lambda}{\rho C}$  is thermal diffusivity,  $\rho$  is density,  $C$  is specific heat capacity

**is replaced by**

where  $a = \frac{\lambda}{\rho C}$  is thermal diffusivity ( $\text{m}^2/\text{s}$ ),  $\rho$  is density ( $\text{kg/m}^3$ ),  $C$  is specific heat capacity ( $\text{J}\cdot\text{kg}^{-1}\cdot^{\circ}\text{C}^{-1}$ ),

**P3 L5**

...represents the rock's thermal effusivity (thermal inertia) characterizing

**is replaced by**

...represents the rock's thermal effusivity ( $\text{J}\cdot\text{m}^{-2}\cdot\text{K}^{-1}\cdot\text{s}^{-1/2}$ ), or thermal inertia, characterizing

**P3 L9:**

...where  $i, j$  are positive integers related with the real time by the equations  $t = i\Delta t$ ,  $t = j\Delta t$ ,  $\Delta t$  is uniform time interval.

**is replaced by**

...where  $i, j$  are positive integers related to real time by  $t = i\Delta t$ ,  $t = j\Delta t$ , for a constant time interval  $\Delta t$ .

**P3 L12**

Using a recurrence equation

**is replaced by**

Using a recurrence [relation](#)

### **P3 L14**

one can estimate  $D_i$  for each interval of temperature curve and then by the equation

**is replaced by**

[the  \$D\_i\$  can be estimated for each interval of the temperature curve. The corresponding instantaneous heat flux values can then be calculated as](#)

### **P3 L16**

one can calculate heat flux instantaneous values at the end of interval. The SHF history reconstruction will be more accurate if we calculate

**is replaced by**

The [so obtained](#) SHF history [reconstruction](#) will be more accurate if we calculate

### **P4 L11:**

**we added**  $\pi/4\omega$

### **P4 L22**

We used the temperature history (Demezhko and Shchapov, 2001) early reconstructed from temperature-depth profile logged in the Urals superdeep borehole SG-4 (58° 24' N, 59° 44' E, Middle Urals, Russia) as initial data (Fig. 2).

**is replaced by**

[As an input to our SHF reconstruction, we used the GSTH obtained from the Urals superdeep borehole SG-4 \(58° 24' N, 59° 44' E, Middle Urals, Russia\) which was presented in an earlier contribution by Demezhko & Shchapov \(2001\) , Fig.2.](#)

### **P5 L26**

A minimal resolved interval of geothermal reconstruction is approximately  $2 \cdot t^*/3$  where  $t^*$  is time before present

**is replaced by**

[The minimum event length which can be resolved in the GSTH reconstruction has been estimated by Demezhko and Shchapov \(2001\) to be  \$2 \cdot t^\*/3\$  where  \$t^\*\$  is time before present.](#)

P7ff: Use numbers (enumerate in latex) or bullet points (itemize) for the list in the Discussion/conclusion

**we added bullet points**

**P7 L27:**

The procedure of averaging in uneven running windows was applied to modify the insolation curve to a form comparable with the geothermal one.

**is replaced by**

A procedure of running averages with changing window lengths was applied to the insolation in order to obtain a curve comparable to the geothermal one.

**P8 L1-5:** Maybe add a word on the particular rock types & a more general reference on the diffusivity values?

The reliability of the new time scale after synchronization with the orbital insolation also depends on how much the thermal diffusivity changed from the initial value. In our study the best coincidence of insolation and heat flux ( $R = 0.99$ ) in the most part of the reconstructed interval is achieved by varying the thermal diffusivity from  $1 \times 10^{-6} \text{ m}^2/\text{s}$  to  $0.71 \times 10^{-6} \text{ m}^2/\text{s}$ , i.e., within the range of its natural variability for the crystalline rocks of the Urals:  $(1 \pm 0.3) \times 10^{-6} \text{ m}^2/\text{s}$  (Demezhko, 2001).

**is replaced by**

- The reliability of the new time scale after synchronization with the orbital insolation also depends on how much the thermal diffusivity changed from the initial value  $1 \times 10^{-6} \text{ m}^2/\text{s}$  which is typical of common crustal rocks (e.g., Pasquale et al., 2014). The SG-4 borehole encountered 3.5 km of basalts, andesite–basalts and their tuffs (Shakhtorina, 1992). Thermal diffusivity of these rocks may decrease to  $(0.76-0.70) \times 10^{-6} \text{ m}^2/\text{s}$  (Durham et al., 1987, Demezhko and Solomina, 2009), and even to  $0.53 \times 10^{-6} \text{ m}^2/\text{s}$  (Eppelbaum et al., 2014). In our study the best coincidence of insolation and heat flux ( $R = 0.99$ ) in the most part of the reconstructed interval is achieved under the thermal diffusivity value  $0.71 \times 10^{-6} \text{ m}^2/\text{s}$ , which is quite realistic.

**We also added four references:**

Pasquale, V., Verdoya, M., and Chiozzi, P.: Geothermics. Heat Flow in the Lithosphere, Springer, Heidelberg, 119 pp, 2014

Shakhtorina, L.N.: Geological cross-section of borehole SG-4. In: The Ural Superdeep Borehole, GNPP Nedra, Yaroslavl, pp. 133–145, 1992 (in Russian).

Durham, W. B., Mirkovich, V. V., Heard, H. C.: (1987). Thermal diffusivity of igneous rocks at elevated pressure and temperature, *Journal of Geophysical Research: Solid Earth* (1978–2012), 92(B11), 11615-11634, 1987.

Eppelbaum L. V., Kutasov I., Pilchin A.: *Applied Geothermics*. Springer-Verlag, Berlin, Heidelberg, 750 pp, 2014.

**Marked-up manuscript version**

## **Late Pleistocene-Holocene ground surface heat flux changes reconstructed from borehole temperature data (the Urals, Russia)**

**D. Yu. Demezhko and A. A. Gornostaeva**

{Institute of Geophysics UB RAS, Yekaterinburg, Russia}

Correspondence to: D. Yu. Demezhko (ddem54@inbox.ru)

### **Abstract**

We use geothermal reconstruction of the ground surface temperature (GST) history early obtained in the Middle Urals to determine the surface heat flux (SHF) history over the past 35 kyr. A new algorithm of GST-SHF transformation was applied to solve this problem. The time scale of geothermal reconstructions has been corrected by comparing the estimated heat flux and annual insolation at the latitude of 60° N. The consistency of SHF and insolation changes on the interval 35-6 kyr BP with the linear correlation coefficient  $R = 0.99$  points to orbital factors as the main cause of climatic changes during the Pleistocene-Holocene transition. The amplitude of SHF variations is about 1.3% of the insolation changes amplitude. The increase of carbon dioxide concentrations lagged by 2-3 kyr from the SHF increase and occurred synchronously with GST changes.

## 1 Introduction

The role of orbital factors in Pleistocene climatic variations has been studied more than 100 years since Joseph Adhemar, James Croll and Milutin Milankovitch. A popular approach is comparing paleotemperatures reconstructed from proxy data (oxygen isotopes, palynological or others) with theoretically calculated insolation. Some investigators (Peixóto and Oort, 1984; Pielke, 2003; Douglass and Knox, 2012) criticized this approach. They noted that temperature field is not an optimal parameter for climate attribution, particularly for evaluation of climatic reaction on the external radiative forcing. There is a lag between external radiative flux and temperature changes, which is disappeared if we consider the heat content or the surface heat flux changes. The advantage of heat flux estimation over temperature one was not realized in full up to date. Wang and Bras (1999) proposed the integral relation to estimate surface heat flux (SHF) changes from ground surface temperature (GST) variations. A finite-difference approximation of the relation between the GST (represented by a piecewise linear function of temperature), and the SHF was proposed by Beltrami et al. (2002). SHF history reconstructions based on borehole temperature data were made in timescales from several centuries to millennium (Beltrami et al., 2002, 2006; Huang, 2006). Another approach was used in (Majorowicz et al., 2012). Subsurface temperatures were calculated from solar irradiance change using information about climate sensitivity.

In the paper we use geothermal reconstruction of GST history early obtained in the Middle Urals (Demezhko and Shchapov, 2001) to determine the SHF history over the past 35 kyr. The recently developed improved algorithm of GST-SHF transformation (Gornostaeva, 2014) was applied to estimate the SHF history. Paleoclimatic interpretation of GST and SHF histories is based on comparison them with orbitally driven solar insolation and atmospheric carbon dioxide changes.

## 2 The method

The GST-SHF transformation algorithm is based on the relation between surface heat flux and surface temperature changes according to the Fourier's equation in one dimension:

$$q(0,t) = -\lambda \left. \frac{\partial T(z,t)}{\partial z} \right|_{z=0} \quad (1)$$

where  $q$  is SHF ( $\text{W}\cdot\text{m}^{-2}$ ),  $t$  is time,  $\lambda$  is thermal conductivity ( $\text{W}\cdot\text{m}^{-1}\cdot\text{K}^{-1}$ ),  $T(z, t)$  is temperature anomaly ( $^{\circ}\text{C}$ ) at a depth  $z$ .

If GST is represented by an expression (Carslaw and Jaeger, 1959, Lachenbruch et al., 1982)

$$T(0,t) = D(t)^{\frac{n}{2}} \quad (2)$$

where  $D$  is a constant,  $n$  is positive integer (or 0) determining the shape of temperature changes, the transient temperature anomaly at any depth is

$$T(z,t) = 2^n \Gamma\left(\frac{1}{2}n + 1\right) i^n \operatorname{erfc} \frac{z}{\sqrt{4at}} T(0,t) \quad (3)$$

where  $a = \frac{\lambda}{\rho C}$  is thermal diffusivity ( $\text{m}^2/\text{s}$ ),  $\rho$  is density ( $\text{kg}/\text{m}^3$ ),  $C$  is specific heat capacity ( $\text{J}\cdot\text{kg}^{-1}\cdot\text{C}^{-1}$ ),  $i^n \text{erfc}(a)$  is the  $n$ -th repeated integral of the error function of  $a$  and  $\Gamma(\beta)$  is gamma-function of argument  $\beta$ . Differentiation of (3) yields SHF

$$q(0,t) = \frac{\Gamma\left(\frac{1}{2}n+1\right)}{\Gamma\left(\frac{1}{2}n+\frac{1}{2}\right)} \cdot \frac{\lambda}{\sqrt{at}} \cdot T(0,t) \quad (4)$$

Note that the ratio  $E = \lambda/(a)^{1/2}$  represents the rock's thermal effusivity ( $\text{J}\cdot\text{m}^{-2}\cdot\text{K}^{-1}\cdot\text{s}^{-1/2}$ ), or thermal inertia, characterizing the rate of heat exchange at the surface.

We approximate GST history by a sum of temperature changes corresponding to Eq. (2):

$$T_i = T_0 + \sum_{j=1}^i D_j (i-j+1)^{\frac{n}{2}} \quad (5)$$

where  $i, j$  are positive integers related to real time by  $t = i \cdot \Delta t$ ,  $t = j \cdot \Delta t$ , for a constant time interval  $\Delta t$ . For each addend of this sum

$$T_i = D_i i^{n/2}, \quad q_i = k_n \frac{E}{\sqrt{\Delta t}} D_i i^{\frac{n-1}{2}}, \quad k_n = \frac{\Gamma\left(\frac{1}{2}n+1\right)}{\Gamma\left(\frac{1}{2}n+\frac{1}{2}\right)}. \quad (6)$$

Using a recurrence relation

$$D_1 = T_1 - T_0$$

$$D_i = (T_i - T_0) - \sum_{j=1}^{i-1} D_j (i-j+1)^{\frac{n}{2}}, \quad i > 1 \quad (7)$$

the  $D_i$  can be estimated for each interval of the temperature curve. The corresponding instantaneous heat flux values can then be calculated as

$$q_i = k_n \frac{E}{\sqrt{\Delta t}} \sum_{j=1}^i D_j (i-j+1)^{\frac{n-1}{2}} \quad (8)$$

~~one can calculate heat flux instantaneous values at the end of interval.~~ The so obtained SHF history reconstruction will be more accurate if we calculate the average value of heat flux on the interval and refer it to the midpoint of the interval ( $i-0.5$ )

$$\overline{q}_{i-0.5} = q_{i-1} + \frac{2}{n+1} (q_i - q_{i-1}). \quad (9)$$

The GST-SHF transformation algorithm was tested by applying it to a harmonic function of surface temperature change with amplitude  $A$ , frequency  $\omega$ , and initial phase  $\varphi$ :

$$T(0,t) = A \sin(\omega t + \varphi), \quad (10)$$

the propagation of temperature waves in a homogeneous half-space with thermal diffusivity  $a$  is described by the expression

$$T(z,t) = Ae^{-kz} \sin(\omega t - kz + \varphi), \quad k = \sqrt{\omega/2a}. \quad (11)$$

Differentiating (11) with respect to  $z$ , we find the ground surface heat flux change  $q(0,t)$ :

$$q(0,t) = -\lambda \left. \frac{\partial T(z,t)}{\partial z} \right|_{z=0} = AE\sqrt{\omega} \sin(\omega t + \varphi + \frac{\pi}{4}) = E\sqrt{\omega} T(0, t + \frac{\pi}{4\omega}). \quad (12)$$

The relationship between the amplitudes of GST and SHF changes is determined by thermal effusivity  $E$  and frequency  $\omega$ . The heat flux changes are ahead of temperature changes by  $\pi/4\omega$ , i.e., one-eighth of the oscillation period.

The relative error of SHF estimation was calculated as the ratio of the standard error of the SHF estimation to the real amplitude of SHF variations. The test showed that approximation of temperature history by the Eq. (5) with  $n = 2, 3$  provides the most accurate results (Fig. 1). When GST discretization is 6 points per period we obtain the relative error of SHF history estimation equals to 3%, and given 10 points per period the relative error is less than 1% (Gornostaeva, 2014). For comparison, the algorithm proposed by Beltrami et al. (2002) under the same discretization conditions provides relative errors equals to 8% and 3.5% respectively.

### 3 GST data and SHF estimation

As an input to our SHF reconstruction, we used the GSTH obtained from the Urals superdeep borehole SG-4 (58° 24' N, 59° 44' E, Middle Urals, Russia) which was presented in an earlier contribution by Demezhko and Shchapov (2001), Fig.2. We analysed only the last 35 kyr of the GST history for the SHF reconstruction, while the paper mentioned above presents 80 kyr temperature history. Because of the decrease of the GSTH resolution with time the interval from 35 to 80 kyr BP does not contain any noticeable GST variations. The SHF may be considered as a constant on this time interval.

The reconstruction of the surface heat flux history was conducted using the algorithm described above with  $n = 3$  (see Fig. 2). GST and SHF curves are different in shape. The temperature increase started about 15 kyr BP and after a short break it continued to 1 kyr BP, while the heat flux increase began about 3 kyr earlier. The heat flux reached its maximum of 0.08 W/m<sup>2</sup> about 8 kyr BP and then it began to decline.

### 4 The comparison of the SHF with solar insolation

The reconstructed SHF changes are similar to the Northern Hemisphere solar insolation changes that are determined by the variations of the Earth's orbital parameters like eccentricity, inclination and the Earth's axis precession (Fig. 3). It is admissible to assume that insolation changes cause the surface heat flux changes. This assumption for the Middle Urals is also supported by the absence of Late Pleistocene ice sheets here (see Velichko et al., 1997, Svendsen et al, 2004 and references therein). However, there is some shift between insolation and SHF changes. The observed shift can be explained by several reasons. The first one is the influence of internal climatic factors and feedbacks translating the external heat flux on the Earth's surface with a certain delay and amplitude attenuation. The second reason is an overestimation of the effective thermal diffusivity that determines the rate of climatic signal propagation into the depth and therefore the time scale of geothermal reconstructions. The effective thermal diffusivity may be strongly affected by different factors like hydrogeological processes, thermophysical



inhomogeneities of rocks, thawing-freezing processes and so on. To synchronize SHF and insolation ( $\Delta I$ ) time series it is necessary to correct the initial value of thermal diffusivity (and time scale respectively) to maximize the correlation between them. The correlation maximum points to an optimal value of thermal diffusivity, which determines the degree of expansion/compression of SHF and GST time scale. Note that the direct comparison of these series is not so correct. The insolation temporal resolution is constant while SHF resolution power decreases back in time. [The minimum event length which can be resolved in the GSTH reconstruction has been estimated by Demezhko and Shchapov \(2001\) to be  \$2 \cdot t^\*/3\$  where  \$t^\*\$  is time before present.](#) The procedure of averaging in uneven running windows was proposed (Demezhko and Solomina, 2009) to modify the curve to a form comparable with the geothermal one. The insolation curve for the latitude of 60° N smoothed according to the resolution power of geothermal method is presented in Fig. 3b. A maximum correlation ( $R=0.99$ ) between SHF history and smoothed insolation is achieved by increasing SHF dates by 1.4 times. It corresponds to the thermal diffusivity decrease from initial value of  $a = 1.0 \cdot 10^{-6} \text{ m}^2/\text{s}$  to  $0.71 \cdot 10^{-6} \text{ m}^2/\text{s}$ . The interval of uncertainty for the optimal diffusivity value under  $R \geq 0.95$  is  $(0.71 \pm 0.06) \cdot 10^{-6} \text{ m}^2/\text{s}$ .

Linear regression analysis of  $q$  and  $\Delta I$  from 35 to 6 kyr BP showed that change of insolation on  $1 \text{ W/m}^2$  produces an additional surface heat flux equals to  $0.013 \text{ W/m}^2$ . At that only a small portion of insolation changes (about 1.3%) was spent to the increase of the lithosphere heat content. The ratio  $\Delta q/\Delta I$  may be considered as a dimensionless measure of climate sensitivity of the region under study to long-term orbital forcing variations.

Taking the climatically caused SHF before 35 kyr BP equals to  $0 \text{ W/m}^2$  and integrating it with respect to time we estimate changes in heat content. This value characterizes the additional amount of heat adsorbed in a rock column having a cross-sectional area of  $1 \text{ m}^2$  and limited by the depth of thermal anomaly penetration (i.e. by a few kilometers). Until 15 kyr BP a total heat balance was negative. A minimum value of heat content of  $-3.5 \text{ TJ/m}^2$  with respect to the reference value at 35 kyr BP was found about 20 kyr BP. From this moment the heat flux became positive. For the next 14 kyr (20-6 kyr BP) the heat content increased to  $22.0 \text{ TJ/m}^2$ . For comparison, during the period of modern warming (1765-2000), heat content of the continental lithosphere increased by  $0.1 \text{ TJ/m}^2$  (calculated using data from Beltrami, 2002).

## 5 The comparison of the SHF with CO<sub>2</sub> changes

Another source of the additional radiative forcing during the Pleistocene-Holocene transition could be greenhouse effect caused by the increase of carbon dioxide concentration in the atmosphere (see Shakun et al., 2012 and references therein). An additional downward heat flux necessarily would contribute to SHF changes. Figure 4 shows geothermal reconstructions of surface temperatures and heat fluxes from the borehole SG-4 (on the time scale corrected after SHF-insolation synchronization) and carbon dioxide concentration changes in Antarctic ice cores (Blunier et al., 1998; Indermühle et al., 1999; 19996; Smith, 1999; Barnola et al., 2003; Pedro et al., 2012). Despite the substantial dispersion of CO<sub>2</sub> estimations, a character and a chronology of CO<sub>2</sub> concentration changes are much closer to temperature changes rather than to heat flux variations. It may mean no significant contribution of CO<sub>2</sub> forcing to climatically caused heat flux and thus to the temperature increase during Pleistocene-Holocene warming.

About 10 kyr BP the increase of carbon dioxide concentration was replaced by its fall which ended about 8 kyr BP. This local minimum is not consistent with either GST or SHF histories. It

is possible that the CO<sub>2</sub> decrease was associated with a sharp increase of vegetation absorbing its excess.

## 6 Discussion and conclusions

The reconstruction of the surface heat flux history using data on the past surface temperature changes represents a new instrument for climate analysis. The reconstructed SHF variations and radiative forcing changes may be compared directly because they are expressed in the same units of energy flux (W/m<sup>2</sup>).

Time synchronization of heat flux and orbital insolation series is similar to the orbital tuning of glacial or marine sediments isotope records (Imbrie et al., 1984, Martinson et al, 1987; Waelbroeck et al., 1995, Shackleton, 2000; Bender et al., 2002; Parrenin et al, 2007). However, it has some special features.

- Since the concentration of  $\delta^{18}O$ ,  $\delta D$  in the ice cores or marine sediments associated with paleotemperature fluctuations, the time shift between the orbital insolation and temperature reaction can be estimated only from independent absolute markers. Because of the rarity of such markers it is generally considered that the shift is a constant (Parrenin et al, 2007). Unlike conventional approach we tune another paleoclimatic characteristics, the surface heat flux, which provides a physically reasonable shift. In (Waelbroeck et al., 1995) the phasing between the precession band of mid-June insolation at 65°N and  $\delta D$  was found about 3 kyr (with the uncertainty  $\pm 3$  kyr). A reliable estimation of the phase in the obliquity band was not obtained and therefore it was not accounted for. Considering the period of precession 23 kyr and using equation (12) we obtain the close estimate  $23/8 \approx 2.9$  kyr. For the obliquity band the phase shift is equal to  $41\text{kyr}/8 = 5.1$  kyr.
- For correct comparison with geothermal reconstruction the insolation curve must have the same resolution. [A procedure of running averages with changing window lengths was applied to the insolation in order to obtain a curve comparable to the geothermal one.](#) Such a procedure limits the tuning interval within the last cycle of precession.
- The reliability of the new time scale after synchronization with the orbital insolation also depends on how much the thermal diffusivity changed from the initial value  $1 \times 10^{-6} \text{ m}^2/\text{s}$  which is typical of common crustal rocks (e.g., Pasquale et al., 2014). The SG-4 borehole encountered 3.5 km of basalts, andesite–basalts and their tuffs (Shakhtorina, 1992). Thermal diffusivity of these rocks may decrease to  $(0.76-0.70) \times 10^{-6} \text{ m}^2/\text{s}$  (Durham et al., 1987, Demezhko and Solomina, 2009), and even to  $0.53 \times 10^{-6} \text{ m}^2/\text{s}$  (Eppelbaum et al., 2014). In our study the best coincidence of insolation and heat flux ( $R = 0.99$ ) in the most part of the reconstructed interval is achieved under the thermal diffusivity value  $0.71 \times 10^{-6} \text{ m}^2/\text{s}$ , which is quite realistic.
- Using the reconstructed surface heat flux instead of the surface temperature does not exclude the existence of residual time shift because the relation between insolation changes and the heat flux may be indirect. Such a shift can be caused by the climate delayed feedbacks. For example, orbital variations of insolation could change the extent of continental and sea ice cover in the Northern Hemisphere, albedo and North Atlantic warm currents. The secondary heat source distributed in the atmosphere arose, which could significantly affect spatial distribution of the SHF change. However, this is beyond the scope of our study.

Assuming the surface heat flux varies proportionally to the external forcing one can consider the ratio  $\Delta q/\Delta I$  as an alternative measure of the Earth's climatic sensitivity. The ratio of two heat

fluxes is a non-dimensional parameter, and additionally depends less on radiative forcing duration by contrast to traditional index of climatic sensitivity representing temperature reaction on the external radiative forcing ( $\Delta T/\Delta I$ ).

The reconstructed surface heat flux reflects impact of all possible sources of radiative forcing. In addition to solar insolation, greenhouse gases (such as CO<sub>2</sub>) can be a source of additional forcing. On the other hand the increase of carbon dioxide may be a consequence of temperature increasing. Comparing the chronology of surface flux, temperature and carbon dioxide concentration changes, we can draw some conclusions about the causes of climate changes.

The described algorithm of GST-SHF transformation is quite easy to realization and allows estimating of SHF history with high precision. Using this algorithm, we have first estimated long-term surface heat flux changes in the Urals for the past 35 kyr. The amplitude of heat flux variations was about 1.3 percent of the insolation changes range at the latitude of 60° N. The increase of carbon dioxide concentrations occurred 2-3 thousands of years later than the heat flux increase and synchronously with temperature response.

### **Acknowledgements**

We would like to thank Volker Rath, Jean-Claude Mareschal, Shaopeng Huang and anonymous reviewer for their valuable comments and suggestions in improving the paper.

This study was supported by the Russian Foundation for Basic Research (RFBR), projects № 13-05-00724-a and №14-05-31055 mol\_a.

## References

- Barnola J.-M., Raynaud D., Lorius C. and Barkov N.I.: Historical CO<sub>2</sub> record from the Vostok ice core, <http://cdiac.ornl.gov/trends/co2/vostok.html>, 2003
- Beltrami, H.: Climate from borehole data: energy fluxes and temperatures since 1500, *Geophys. Res. Lett.*, 29, 26-1–26-4, 2002
- Beltrami, H., Smerdon, J. E., Pollack, H. N., and Huang, S.: Continental heat gain in the global climate system, *Geophys. Res. Lett.*, 29, 8-1–8-3, 2002
- Beltrami, H., Bourlon, E., Kellman, L., and González-Rouco, J. F.: Spatial patterns of ground heat gain in the Northern Hemisphere, *Geophys. Res. Lett.*, 33, L06717, 2006
- Bender, M. L.: Orbital tuning chronology for the Vostok climate record supported by trapped gas composition, *Earth Planet. Sci. Lett.*, 204, 275–289, 2002.
- Berger A. and Loutre M.F.: Insolation values for the climate of the last 10 million of years, *Quat. Sci. Rev.*, 10 (4), 297–317, 1991
- Carslaw, H. S., and Jaeger, J. C.: *Conduction of heat in soils*. Clarendon Press, Oxford, 510 pp, 1958.
- Demezhko, D.Yu. and Shchapov, V.A.: 80,000 years ground surface temperature history inferred from the temperature-depth log measured in the superdeep hole SG-4 (the Urals, Russia), *Global and Planetary Change*, 29 (1-2), 219-230, 2001
- Demezhko, D. Y. *Geothermal Method for Paleoclimate Reconstruction (Examples From the Urals, Russia)*, Russ. Acad. of Sci., Urals Branch, Ekaterinburg, 143 pp. 2001 (in Russian)
- Demezhko, D. Yu., and Solomina, O. N.: Ground surface temperature variations on Kunashir Island in the last 400 years inferred from borehole temperature data and tree-ring records, *Doklady Earth Sciences*, 426 (1), 628-631, 2009
- Douglass D. H., and Knox R. S.: Ocean heat content and Earth's radiation imbalance. II. Relation to climate shifts, *Phys. Lett. A.*, 376, 1226-1229, 2012
- Durham, W. B., Mirkovich, V. V., Heard, H. C.: (1987). Thermal diffusivity of igneous rocks at elevated pressure and temperature, *Journal of Geophysical Research: Solid Earth (1978–2012)*, 92(B11), 11615-11634, 1987
- Eppelbaum L. V., Kutasov I., Pilchin A.: *Applied Geothermics*. Springer-Verlag, Berlin, Heidelberg, 750 pp, 2014
- Gornostaeva A.A.: The calculation algorithm of ground surface heat flux changes from ground temperature changes, *Urals geophysical herald*, 1, 30-39, 2014 (In Russian).
- Huang S.: 1851–2004 annual heat budget of the continental landmasses *Geophys. Res. Lett.*, 33, L04707, 2006
- Imbrie, J., Hays, J.D., Martinson, D.G., McIntyre, A., Mix, A.C., Morley, J.J., Pisias, N.G., Prell, W.L., and Shackleton, N.J.: The orbital theory of Pleistocene climate: support from a revised chronology of the marine  $\delta^{18}O$  record. In: Berger AL et al. (eds) *Milankovitch and climate*, part 1. D. Riedel, Hingham, MA, USA, 269–305, 1984

- Indermühle, A., Monnin, E., Stauffer, B., Stocker, T. F., and Wahlen, M.: Atmospheric CO<sub>2</sub> concentration from 60 to 20 kyr BP from the Taylor Dome ice core, Antarctica, *Geophys. Res. Lett.*, *27*, 735–738, 1999
- Indermühle, A., Stocker, T. F., Joos F., Fischer H., Smith H.J., Wahlen M., Deck B., Mastroianni D., Tschumi J., Blunier T., Meyer R. and Stauffer B.: Holocene carbon-cycle dynamics based on CO<sub>2</sub> trapped in ice at Taylor Dome, Antarctica, *Nature*, *398*, 121–126, 1999b (<ftp://ftp.ncdc.noaa.gov/pub/data/paleo/icecore/antarctica/taylor/>)
- Lachenbruch A., Sass J.H., Marshall B.V., and Mases Jr. T.H.: Permafrost, heat flow, and the geothermal regime at Prudhoe Bay, Alaska, *J. Geophys. Res.*, *87*, 9301-9316, 1982
- Martinson, D.G., Pisias, N.G., Hays, J.D., Imbrie, J., Moore, T.C., Shackleton, N.J.: Age dating and the orbital theory of the Ice Ages: development of a high-resolution 0–300000 years chronostratigraphy. *Quart. Res.*, *27*, 1–30, 1987.
- Majorowicz J., Scinner W. and Safanda J.: Western Canadian Sedimentary Basin temperature–depth transients from repeated well logs: evidence of recent decade subsurface heat gain due to climatic warming, *J. Geophys. Eng.*, *9*, 127-137, 2012
- [Pasquale, V., Verdoya, M., and Chiozzi, P.: Geothermics. Heat Flow in the Lithosphere, Springer, Heidelberg, 119 pp, 2014](#)
- Parrenin, F., Barnola, J. M., Beer, J., Blunier, T., Castellano, E., Chappellaz, J., and Wolff, E.: The EDC3 chronology for the EPICA Dome C ice core. *Climate of the Past* *3.3*, 485-497, 2007.
- Peixóto J. P., and Oort A. H.: Physics of climate, *Rev. Mod. Phys.*, *56* (3), 365-429, 1984
- Pedro J.B., Rasmussen S.O. and Van Ommen T.D.: Tightened constraints on the time-lag between Antarctic temperature and CO<sub>2</sub> during the last deglaciation, *Clim. Past*, *8*, 1213–1221, 2012
- Pielke Sr R. A.: Heat storage within the Earth system, *Bull. Am. Meteorol. Soc.*, *84* (3), 331-335, 2003
- Shackleton, N. J.: The 100 000-year ice-age cycle identified and found to lag temperature, carbon dioxide, and orbital eccentricity, *Science*, *289*(5486), 1897–1902, 2000.
- [Shakhtorina, L.N.: Geological cross-section of borehole SG-4. In: The Ural Superdeep Borehole, GNPP Nedra, Yaroslavl, pp. 133–145, 1992 \(in Russian\).](#)
- Shakun J.D, Clark P.U, He F, Marcott S.A, Mix A.C, Liu Z, Otto-Bliesner B., Schmittner A., and Bard E.: Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation, *Nature*, *484* (7392), 49-54, 2012
- Smith H. J., Fischer H., Mastroianni D., Deck B. and Wahlen M.: Dual modes of the carbon cycle since the Last Glacial Maximum, *Nature*, *400*, 248–250, 1999 (<ftp://ftp.ncdc.noaa.gov/pub/data/paleo/icecore/antarctica/taylor/>)
- Svendsen, J. I., Alexanderson, H., Astakhov, V. I., Demidov, I., Dowdeswell, J. A., Funder, S., Gataullin, V., Henriksen, M., Houmark-Nielsen, V., Hubberten, H.W., Ingolfsson, O., Jakobsson, M., Kjaer K.H., Larsen, E., Lokrantz, H., Lunkka, J.P., Lysa, A., Mangerud, J., Matiouchkov, A., Murray, A., Moller, P., Niessen, F., Nikolskaya, O., Polyak, L., Saarnisto, M., Siegert, C., Siegert, M.J., Spielhagen, R.F., and Stein, R.: Late Quaternary ice sheet history of northern Eurasia. *Quaternary Science Reviews*, *23.11*, 1229-1271, 2004.

Velichko, A. A., Yu M. Kononov, and M. A. Faustova.: The last glaciation of Earth: size and volume of ice-sheets. *Quaternary International*, 41, 43-51, 1997.

Wang J., and Bras R.L.: Ground heat flux estimated from surface soil temperature, *J.Hydrol.*, 216 (3-4), 214-226, 1999.

Waelbroeck, C., Jouzel, J., Labeyrie, L., Lorius, C., Labracherie, M., Stievenard, M., and Barkov, N. I.: A comparison of the Vostok ice deuterium record and series from Southern Ocean core MD 88-770 over the last two glacial-interglacial cycles. *Climate Dynamics*, 12(2), 113-123, 1995.




Figure 1. Testing the algorithms of GST-SHF transformation by applying it to a harmonic function of GST change. Relative error of SHF estimation (the ratio of the standard error of the SHF estimation to the real amplitude of SHF variations) versus the GST discretization frequency (points per period).




Figure 2. Initial data and surface heat flux a) Temperature-depth profile from the borehole SG-4 (Demezhko and Shchapov, 2001, brown line), b) GST history  $T(t)$  ( $a = 1.0 \cdot 10^{-6} \text{ m}^2/\text{s}$ , Demezhko and Shchapov, 2001, brown line) and calculated according to Eqs. (1-5) SHF history  $q(t)$  ( $E = 2500 \text{ J} \cdot \text{m}^{-2} \cdot \text{K}^{-1} \cdot \text{s}^{-1/2}$ , blue line).




Figure 3. The comparison of SHF history with solar insolation changes in the Northern Hemisphere caused by changes in Earth's orbital parameters and time scale correcting. a) Annual insolation changes  $\Delta I(t)$  at the latitudes of 40-70° N (Berger, Loutre, 1991); b) annual solar insolation at the latitude of 60° N smoothed in uneven running windows (green line), SHF history in the initial timescale ( $a = 1.0 \cdot 10^{-6} \text{ m}^2/\text{s}$ , blue dashed line) and SHF history in the corrected timescale ( $a = 0.71 \cdot 10^{-6} \text{ m}^2/\text{s}$ , blue solid line).






Figure 4. The comparison of GST history  $T(t)$  (brown line), SHF history  $q(t)$  (blue line) and CO<sub>2</sub> concentration in the Antarctic ice cores (multicolored markers).