## Thermal evolution of the western South Atlantic and the adjacent continent during Termination 1

3

### 4 C. M. Chiessi<sup>1</sup>, S. Mulitza<sup>2</sup>, G. Mollenhauer<sup>2,3</sup>, J. B. Silva<sup>4</sup>, J. Groeneveld<sup>2</sup>, M. 5 Prange<sup>2</sup>

6 [1]{School of Arts, Sciences and Humanities, University of São Paulo, São Paulo, Brazil}

7 [2]{MARUM – Center for Marine Environmental Sciences, University of Bremen, Bremen,
8 Germany}

9 [3]{Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany}

10 [4] {Institute of Geosciences, University of São Paulo, São Paulo, Brazil}

11 Correspondence to: C. M. Chiessi (chiessi@usp.br)

12

#### 13 Abstract

14 During Termination 1, millennial-scale weakening events of the Atlantic meridional 15 overturning circulation (AMOC) supposedly produced major changes in sea surface 16 temperatures (SST) of the western South Atlantic, and in mean air temperatures (MAT) over 17 southeastern South America. It was suggested, for instance, that the Brazil Current (BC) 18 would strengthen (weaken) and the North Brazil Current (NBC) would weaken (strengthen) 19 during slowdown (speed-up) events of the AMOC. This anti-phase pattern was claimed to be 20 a necessary response to the decreased North Atlantic heat piracy during periods of weak 21 AMOC. However, the thermal evolution of the western South Atlantic and the adjacent 22 continent is so far largely unknown. Here we address this issue, presenting high temporal 23 resolution SST and MAT records from the BC and southeastern South America, respectively. 24 We identify a warming in the western South Atlantic during Heinrich Stadial 1 (HS1), which 25 is followed first by a drop and then by increasing temperatures during the Bølling-Allerød, in-26 phase with an existing SST record from the NBC. Additionally, a similar SST evolution is 27 shown by a southernmost eastern South Atlantic record, suggesting a South Atlantic-wide 28 pattern in SST evolution during most of Termination 1. Over southeastern South America, our 29 MAT record shows a two-step increase during Termination 1, synchronous with atmospheric CO<sub>2</sub> rise (i.e., during the second half of HS1 and during the Younger Dryas), and lagging
abrupt SST changes by several thousand years. This delay corroborates the notion that the
long duration of HS1 was fundamental to drive the Earth out of the last glacial.

4

#### 5 **1** Introduction

The thermal bipolar seesaw describes the warming occurring in the Southern Hemisphere due 6 7 to diminished northward heat transport within the Atlantic Ocean when the Atlantic 8 meridional overturning circulation (AMOC) is weakened (Mix et al., 1986; Stocker, 1998). 9 This mechanism is particularly efficient for perturbations of the AMOC through positive 10 anomalous freshwater fluxes in the high latitudes of the North Atlantic (Crowley, 1992; 11 Manabe and Stouffer, 1988). Heinrich Stadial 1 (HS1) is probably the best example for a 12 freshwater-forced AMOC reduction (McManus et al., 2004). It has been suggested that the 13 southward flowing Brazil Current (BC) might redirect the excess heat to the South Atlantic 14 during times of AMOC slowdown (Crowley, 2011; Maier-Reimer et al., 1990). Yet, little is 15 known about the thermal evolution of the western South Atlantic and the adjacent continent during Termination 1. The few available oceanic (e.g., Carlson et al., 2008) and continental 16 17 (e.g., Bush et al., 2004) records do not show the necessary temporal resolution to 18 appropriately resolve HS1. The lack of high temporal resolution records from the BC (Clark 19 et al., 2012), for instance, hinders the evaluation of the previously hypothesized anti-phase 20 behavior between the BC and the North Brazil Current (NBC) during periods of a stalled AMOC (Arz et al., 1999; Schmidt et al., 2012; Chiang et al., 2008). 21

Here we address this issue using an oceanic and a continental temperature record based on Mg/Ca analyses in planktonic foraminifera and lipid analyses in continentally-derived organic matter, respectively. Our records come from a single marine sediment core collected off southeastern South America under the influence of the BC and spanning Termination 1 with high temporal resolution. Our data provide evidence for millennial-scale fluctuations in the oceanic temperature record associated to changes in AMOC strength, and a two-step increase in the continental temperature record associated to changes in atmospheric CO<sub>2</sub>.

29

#### 1 2 Regional setting

#### 2 2.1 Western South Atlantic

3 Upper level circulation in the subtropical western South Atlantic is dominated by the southward-flowing BC (Fig. 1A) (Peterson and Stramma, 1991; Stramma and England, 1999). 4 5 The BC originates between 10 and 15°S from the bifurcation of the Southern South Equatorial Current (SSEC). At the bifurcation, the SSEC feeds both the BC and the northward flowing 6 7 NBC (also termed the North Brazil Undercurrent (Stramma et al., 1995) between the 8 bifurcation and ca. 5°S). Around 37°S the BC converges with the northward-flowing Malvinas 9 Current (Olson et al., 1988), where both currents turn southeastward and flow offshore as the 10 South Atlantic Current and the northern branch of the Antarctic Circumpolar Current, respectively. The position of the Brazil-Malvinas Confluence varies seasonally between ca. 11 12 34 and 40°S, with a northward penetration of the Malvinas Current during austral winter and early spring and a southward shift of the BC during austral summer and early autumn (Olson 13 14 et al., 1988). In its uppermost 100 m, the BC transports Tropical Water (>20 °C and >36 psu) in the mixed layer, and from ca. 100 until 600 m the BC transports South Atlantic Central 15 16 Water (6–20°C and 34.6–36 psu) in the permanent thermocline (Locarnini et al., 2010; Antonov et al., 2010). 17

18 The deficit in the southward BC transport relative to what would be expected from the wind 19 field is a consequence of the northward-directed upper branch of the thermohaline circulation 20 (Stommel, 1957; Peterson and Stramma, 1991). The formation of North Atlantic Deep Water 21 in the high latitudes of the North Atlantic requires a net transfer of thermocline water from the 22 South Atlantic to the North Atlantic together with net northward fluxes of intermediate and 23 bottom waters (Rintoul, 1991; Peterson and Stramma, 1991). Thus, under modern conditions 24 the NBC receives the larger portion (ca. 12 Sv) of the SSEC volume transport if compared to the BC (ca. 4 Sv) (e.g., Stramma et al., 1990). 25

#### 26 2.2 Southeastern South America

Throughout the year atmospheric circulation over southeastern South America is dominated by northerly winds (Fig. 1B) (Kalnay et al., 1996). During Southern Hemisphere summer, the South Atlantic Convergence Zone, a northwest-southeast-oriented convective band along the northeastern boundary of the La Plata River drainage basin (LPRDB), and the South

American Low-Level Jet, a northwesterly low-level flow that transports moisture from the 1 2 western Amazon to the LPRDB, are key features of the South American summer monsoon (Carvalho et al., 2004; Zhou and Lau, 1998). During Southern Hemisphere winter, 3 equatorward incursions of mid-latitude cold dry air result in cyclonic storms (Vera et al., 4 5 2002). Precipitation in the LPRDB is dominated by Southern Hemisphere summer rainfall associated to the South American summer monsoon (Fig. 2) (Zhou and Lau, 1998; Xie and 6 7 Arkin, 1997). Correspondingly, maximum La Plata River discharge occurs in late Southern 8 Hemisphere summer (Fig. 2). Winter precipitation triggered by occasional northward 9 migration of extratropical cyclones results in less pronounced rainfall (Vera et al., 2002). Histograms of long-term mean average monthly precipitation display a strong Southern 10 11 Hemisphere winter minimum (Fig. 2), particularly in the north-western sector of the LPRDB, 12 which supplies most of the particulate load of the La Plata River (Depetris and Kempe, 1993; 13 Depetris et al., 2003).

Air temperatures at low atmospheric levels over South America are dominated by the equatorto-pole thermal gradient (Fig. 1B) (Garreaud et al., 2009). The meridional temperature profile is rather flat between the equator and 20°S, centered around 20°C. To the south of 20°S, temperatures gradually decrease to 0°C over the southern tip of the continent. Zonal departures from this meridional gradient are relatively small to the east of the Andes, as is the case for the LPRDB.

20

#### 21 **3** Material and methods

#### 22 **3.1 Marine sediment core**

We investigated sediment core GeoB6211-2 (Schulz et al., 2001; Wefer et al., 2001) collected from the continental slope off southeastern South America (32.51°S / 50.24°W / 657 m water depth / 774 cm long) (Figs. 1A, 2). The gravity core was raised at the Rio Grande Cone, a major sedimentary feature in the western Argentine Basin (Schulz et al., 2001). Because our focus here is Termination 1 we analyzed the section from 86 until 583 cm core depth that corresponds to the period from 10.2 until 19.3 cal ka BP (see Section 4.1 below).

29 One meter long sections of core GeoB6211-2 were longitudinally split and described onboard,

30 and then stored at 4°C. Visual inspection of core GeoB6211-2 does not provide evidence for

31 depositional or erosive disturbance (Wefer et al., 2001). Onshore, the last deglaciation section

of the core was sampled at 1 cm intervals. Samples for radiocarbon, Mg/Ca and stable oxygen isotope ( $\delta^{18}$ O) analyses were wet sieved, oven-dried at 50°C, and the residues from the 150 µm size sieve were stored in glass vials. Hand-picking of foraminiferal tests was performed under a binocular microscope. Samples for lipid analyses were stored at 4°C until processing.

#### 5 **3.2** Radiocarbon analyses and age model

The age model of core GeoB6211-2 is based on nine accelerator mass spectrometry 6 7 radiocarbon ages (Table 1, Fig. 3). Five ages are based on tests of shallow dwelling planktonic foraminifera Globigerinoides ruber (pink and white) and Globigerinoides 8 9 sacculifer, while the remaining four ages are based either on mixed planktonic foraminifera (i.e., two ages) or epibenthic bivalve shells (i.e., two ages). Apart from the age obtained at 10 11 315 cm core depth, all ages were previously published by Chiessi et al. (2008) and Razik et al. 12 (2013). For each sample, we collected around 10 mg of CaCO<sub>3</sub> from the sediment fraction 13 larger than 150 µm. One of the samples was measured at the National Ocean Sciences Accelerator Mass Spectrometry Facility at Woods Hole (USA), while the other eight were 14 15 measured at the Leibniz-Laboratory for Radiometric Dating and Stable Isotope Research at 16 Kiel (Germany). All radiocarbon ages were calibrated with the calibration curve Marine13 17 (Reimer et al., 2013) with the software Calib 7.0 (Stuiver and Reimer, 1993). Following the 18 arguments from Chiessi et al. (2008) we decided not to use a specific reservoir age to the 19 radiocarbon ages based on epibenthic bivalve shells. Also, no additional marine reservoir 20 correction was applied because our core site is located far from upwelling zones and 21 significantly to the north of the Brazil-Malvinas Confluence, both being places where 22 corrections are typically necessary (Reimer and Reimer, 2001). All ages are indicated as 23 calibrated years before present (cal a BP; present is 1950 AD), except where noted otherwise. To construct the age model, we linearly interpolated the calibrated ages. For each dated depth 24 25 we used in the interpolation the mean value from the  $1\sigma$  range of the calibrated age.

#### 26 **3.3 Mg/Ca analyses and sea surface temperatures**

Around 40 tests of *G. ruber* (white, sensu stricto according to Wang (2000)) within the size range 250–350 µm were used for Mg/Ca analyses. Analyses were performed at approximately every cm between 86 and 123 cm core depth, and at approximately every four cm below 123 cm core depth. Different spacing was applied to compensate for the lower sedimentation rates in the section 86-123 cm core depth as compared to the section 123-583 cm core depth (see

Section 4.1 below). After gently crushing the tests, shell fragments were cleaned according to 1 2 the standard cleaning protocol for foraminiferal Mg/Ca analyses suggested by Barker et al. (2003) and slightly modified by Groeneveld and Chiessi (2011). Before dilution, samples 3 4 were centrifuged for 10 min (6000 rpm) to exclude any remaining insoluble particles from the analyses. Samples were diluted with Seralpur water before analysis with an inductively 5 coupled plasma - optical emission spectrometer (ICP-OES) (Agilent Technologies, 700 6 7 Series with autosampler (ASX-520 Cetac) and micro-nebulizer) at the MARUM – Center for 8 Marine Environmental Sciences, University of Bremen, Germany. Instrumental precision of 9 the ICP- OES was monitored by analysis of an in- house standard solution with a Mg/Ca of 2.93 mmol mol<sup>-1</sup> after every five samples (long-term standard deviation of 0.026 mmol mol<sup>-1</sup> 10 or 0.91 %). To allow inter-laboratory comparison we analyzed an international limestone 11 12 standard (ECRM752–1) with a reported Mg/Ca of 3.75 mmol mol<sup>-1</sup> (Greaves et al., 2008). 13 The long-term average of the ECRM752-1 standard, which was routinely analyzed twice before each batch of 50 samples in every session, is 3.78 mmol mol<sup>-1</sup> ( $1\sigma = 0.066$  mmol mol<sup>-1</sup> 14 <sup>1</sup>). Analytical error based on three replicate measurements of each sample for G. ruber was 15 0.14 % ( $1\sigma = 0.004 \text{ mmol mol}^{-1}$ ) for Mg/Ca. To convert Mg/Ca into sea surface temperatures 16 (SST) we used the calibration equation of Anand et al. (2003) for G. ruber (white) in the size 17 range 250–350 µm with no pre-assumed exponential constant: 18

19 
$$Mg/Ca = 0.34 \exp(0.102 * SST)$$

20 The propagation of uncertainties typically results in  $1\sigma$  error of about 1°C for SST (Mohtadi et al., 2014).

According to Hönisch et al. (2013), the small sensitivity of *G. ruber* Mg/Ca to changes in salinity (i.e.,  $3.3 \pm - 1.7$  % per salinity unit) supports the use of this paleotemperature proxy given the range of salinity change in our study area (see Section 4.3 below).

25 We measured Mg/Ca in tests of G. ruber (white) because it dwells in the uppermost water 26 column and reflects mixed layer conditions (Chiessi et al., 2007). Moreover, G. ruber (white) 27 records austral hemisphere summer conditions at our core site (Fraile et al., 2009a; Lombard 28 et al., 2011), with no significant change in seasonal preference during the Last Glacial Maximum (LGM) (Fraile et al., 2009b). Furthermore, the mean Mg/Ca based SST (i.e., 29 30 23.1°C) obtained for the uppermost two samples of multicore GeoB6211-1 (collected in the 31 same site as gravity core GeoB6211-2) compares favorably with the modern mean summer 32 SST in the top 20 m of the local water column (i.e., 24.1°C) and differs considerably from

(1)

1 modern mean winter SST (i.e., 17.8°C), corroborating the austral hemisphere summer signal

2 recorded by *G. ruber* (white) (Chiessi et al., 2014).

#### 3 3.4 Stable oxygen isotope analyses and sea surface salinities

4 Ten hand-picked tests of G. ruber (white, sensu stricto according to Wang (2000)) within the 5 size-range 250-350 µm from approximately every cm of core GeoB6211-2 were used for  $\delta^{18}$ O analyses. Results between 448 and 123 cm core depth were previously published by 6 7 Chiessi et al. (2009). Stable oxygen isotope analyses were performed on a Finnigan MAT 252 8 mass spectrometer equipped with an automatic carbonate preparation device at the MARUM -9 Center for Marine Environmental Sciences, University of Bremen, Germany. Isotopic results were calibrated relative to the Vienna Peedee belemnite (VPDB) using the NBS19 standard. 10 11 The standard deviation of the laboratory standard was lower than 0.07 ‰ for the measuring 12 period.

13 To calculate the  $\delta^{18}$ O of continental-ice-volume-corrected surface sea water ( $\delta^{18}$ O<sub>ivc-ssw</sub>), a 14 proxy for relative sea surface salinity, we used: (i) our *G. ruber* Mg/Ca SST and  $\delta^{18}$ O; (ii) the 15 paleotemperature equation from Mulitza et al. (2003) for *G. ruber* (white):

16 
$$SST(^{\circ}C) = -4.44 * \left(\delta^{18}O_{G.ruber} - \delta^{18}O_{sw}\right) + 14.20$$
 (2)

17 (iii) the VPDB to Vienna Standard Mean Ocean Water conversion factor from Hut (1987); 18 (iv) the sea level curve from Lambeck and Chappell (2001); and (v) the global average change 19 in  $\delta^{18}O_{sw}$  since the LGM from Schrag et al. (2002). The sea level curve from Lambeck and 20 Chappell (2001) is consistent with the timing of meltwater pulse 1A reported by Deschamps 21 et al. (2012) (14.5 and 14.6 cal ka BP, respectively). The propagation of uncertainties 22 typically results in 1 $\sigma$  error of about 0.3 ‰ for  $\delta^{18}O_{ivc-ssw}$  (Mohtadi et al., 2014).

#### 23 **3.5** Lipid analyses and continental mean air temperatures

Lipid analyses were performed at approximately every 6 cm. Lipid extraction of freeze-dried powdered samples was performed by the use of ultrasonic probes. Extracts were saponified and further separated on Bond-Elut SiO<sub>2</sub> columns. Polar fractions containing glycerol dialkyl glycerol tetraethers (GDGTs) were eluted with 2 mL MeOH. Prior to analysis by high performance liquid chromatography / atmospheric pressure chemical ionization – mass spectrometry (HPLC/APCI-MS), samples were filtered through a 4  $\mu$ m pore size PTFE filter and dissolved in hexane/isopropanol (99:1; v/v). An Agilent 1200 series HPLC/APCI-MS
system equipped with a Grace Prevail Cyano column (150 mm x 2.1 mm; 3 µm) was used,
and separation was achieved in normal phase using the method described by Hopmans et al.
(2004).

5 Mean air temperature (MAT) was estimated according to Peterse et al. (2012). GDGTs with 6 the following protonated molecular ion masses were quantified: 1022 (Ia), 1020 (Ib), 1018 7 (Ic); 1036 (IIa), 1034 (IIb), 1032 (IIc); 1050 (IIIa). Ratios of peak areas were used to calculate 8 the Methylation Branched Tetraether (MBT<sup>^</sup>) and Cyclisation Branched Tetraether (CBT) 9 indices as follow:

10 
$$MBT' = \frac{(Ia + Ib + Ic)}{(Ia + Ib + Ic + IIa + IIb + IIc + IIIa)}$$
(3)

11 
$$CBT = -\log\left[\left(\frac{Ib + IIb}{Ia + IIa}\right)\right]$$
 (4)

Index values calculated using equations (3) and (4) were subsequently converted to MATestimates according to:

14 
$$MAT(^{o}C) = 0.81 - 5.67 * CBT + 31.0 * MBT'$$
 (5)

The production of GDGTs by some uncharacterized microbial community in marine sediments has gained recent attention (e.g., Zhu et al., 2011). For in situ production in the marine realm, the authors consistently describe an increase in the relative abundance of those GDGTs containing cyclopentane moieties (e.g., GDGT Ic and GDGT IIc) as well as a decrease in the relative abundance of the compounds GDGT Ia and GDGT IIa. We carefully screened our results for a similar behavior.

21 Temperature estimates are thought to reflect mean annual air temperature (Peterse et al., 22 2012). During Termination 1, our core site received terrigenous material discharged from the 23 La Plata River drainage basin (LPRDB) as attested by Nd isotopes (Lantzsch et al., 2014). 24 Within the LPRDB, most of the suspended load (Depetris et al., 2003) and particulate organic 25 matter (Depetris and Kempe, 1993) originates from the Bermejo River sub-basin, located in the northwest domain. The amount of river suspended load corresponds to the discharge 26 27 (Depetris et al., 2003), and most of the particulate organic matter is soil-derived (Depetris and 28 Kempe, 1993). Thus, we expect our MAT record to represent a LPRDB-integrated signal with 29 a predominant contribution from its north-western domain.

1

#### 2 4 Results

#### 3 4.1 Radiocarbon analyses and age model

The investigated section (i.e., 86-583 cm core depth) of core GeoB6211-2 recorded the period between 10.2 and 19.3 cal ka BP (Table 1, Fig. 3). The Marine13 calibration curve produced very similar ages (i.e., difference smaller than 0.2 kyr) if compared to the previously published values (Chiessi et al., 2008; Razik et al., 2013) calibrated with Marine04 (Hughen et al., 2004) and Marine09 (Reimer et al., 2009). Thus, the age model used here is very similar to the age models published by Chiessi et al. (2008) between 19.3 and 14.1 cal ka BP, and by Razik et al. (2013) between 14.1 and 10.2 cal ka BP.

Sedimentation rates of the investigated section of core GeoB6211-2 show a two-step decrease
from the LGM to the early Holocene (Fig. 3). Mean values decrease from ca. 160 to 80 cm
kyr<sup>-1</sup> at 18.45 cal ka BP, and from ca. 80 to 10 cm kyr<sup>-1</sup> at 14.1 cal ka BP.

14 Considering the sampling strategy and the sedimentation rates for core GeoB6211-2, the 15 mean temporal resolution is ca. 30 yr for Mg/Ca analyses, ca. 10 yr for  $\delta^{18}$ O analyses, and ca. 16 80 yr for lipid analyses for the period before 18.45 cal ka BP, ca. 60 yr for Mg/Ca analyses, 17 ca. 15 yr for  $\delta^{18}$ O analyses, and ca. 70 yr for lipid analyses for the period between 18.45 and 14.1 cal ka BP, and ca. 120 yr for Mg/Ca analyses, ca. 105 yr for  $\delta^{18}$ O analyses, and ca. 555 19 yr for lipid analyses for the period after 14.1 cal ka BP.

#### 20 4.2 Mg/Ca analyses and sea surface temperatures

Mg/Ca from G. ruber range from 2.50 to 3.60 mmol mol<sup>-1</sup> and are equivalent to 19.5 and 23.1 21 22 °C, respectively (Fig. 4B). Reconstructed SST increase since the LGM (averaging 20.6°C) 23 until ca. 18 cal ka BP, remaining roughly constant (averaging 21.9°C) until ca. 16 cal ka BP. 24 A marked SST drop reaching minimum value (20.4°C) at ca. 15.5 cal ka BP ends the period 25 of relatively stable SST. A double-peak structure culminating at ca. 15 cal ka BP (23.0°C) and ca. 13 cal ka BP (22.9°C) was followed by low temperatures (averaging 21.7°C) until ca. 11.9 26 27 cal ka BP. After that, the record is characterized by oscillating SST (averaging 22.2°C) SST. 28 Thus, the deglacial SST rise is ca. 1.6°C.

#### **4.3** Stable oxygen isotope analyses and sea surface salinities

Values of *G. ruber* δ<sup>18</sup>O show a stepwise decrease from 0.75 ‰ during the LGM to -0.06 ‰
during the early Holocene (Fig. 4C). There are three major steps and they occurred at ca. 15.5,
13.5 and 11.5 cal ka BP.

5 Ice-volume corrected  $\delta^{18}O_{ssw}$  values range from 0.88 to 2.15 ‰ (Fig. 4D). From the LGM 6 until ca. 14 cal ka BP, temporal changes in  $\delta^{18}O_{ivc-ssw}$  are similar to the changes described for 7 SST. After that, the record is marked by roughly constant values (averaging 1.65 ‰) until 8 11.5 cal ka BP and a rather large variability around 1.47 ‰ during the early Holocene.

#### 9 4.4 Lipid analyses and continental mean air temperatures

We first examined our data set for indications of marine in-situ production of GDGTs as 10 11 described in section 3.5, which was not the case (Chiessi et al., 2015b). Then, we calculated continental MAT values that range from 11.5°C at 18.0 cal ka BP to 14.9°C at 11.5 cal ka BP 12 13 (Fig. 4E). Reconstructed MAT show a small gradual increase from the base of the record until ca. 16.5 cal ka BP when a sharp increase of ca. 1.1°C takes place. Temperatures remain 14 15 relatively stable until ca. 12.5 cal ka BP when an increase of ca. 1.0°C within ca. 1 kyr was recorded. After that, stable MAT characterize the record until the early Holocene. Although 16 17 our MAT record does not cover the LGM, the deglacial MAT rise calculated using the averaged value for the oldest and youngest 500 yr values of our time series is 2.5°C. The 18 19 marked decrease in the mean temporal resolution of our MAT record after 14.1 cal ka BP that 20 shifts from ca. 70 years to ca. 555 years is worthy of note. This has to be considered while 21 interpreting the MAT trends described for the period after 14.1 cal ka BP.

22

#### 23 **5 Discussion**

The two major decreases in sedimentation rates found in GeoB6211-2 are remarkably synchronous (within age model uncertainties) with outstanding events of sea-level rise related to meltwater pulses that occurred at ca. 19 and 14.6 cal ka BP (Deschamps et al., 2012; Yokoyama et al., 2000). The sea-level drop preceding the LGM shifted the coastline towards our core site. With the resulting narrow continental shelf, the large sediment supply of the La Plata River was directed to the Rio Grande Cone via the La Plata paleo-valley, that was responsible for the high sedimentation rates typical for the lowermost section of core GeoB6211-2 (Fig. 3) (Chiessi et al., 2008; Lantzsch et al., 2014). The stepwise rise in sealevel following the LGM caused abrupt displacements of the coastline towards the continent
trapping a large amount of the La Plata River sediment supply on the shelf and controlling the
stepwise decrease in sedimentation rate at our site (Chiessi et al., 2008; Lantzsch et al., 2014).
Because of the high sedimentation rates (i.e., ca. 100 cm kyr<sup>-1</sup>) found between the LGM and
14.1 cal ka BP, core GeoB6211-2 is particularly well suited to investigate HS1.

### 5.1 Sea surface temperatures and salinities of the western South Atlantic during Termination 1

9 The high SST reconstructed for our western South Atlantic site between 18 and 16 cal ka BP 10 as well as the peak in SST at ca. 15 cal ka BP (Fig. 4B) fall within HS1, as defined by 11 Sarnthein et al. (2001). It has been suggested that during HS1 a strong slowdown of the 12 AMOC (Fig. 5B) (McManus et al., 2004) produced by a positive anomalous freshwater 13 discharge into the high latitudes of the North Atlantic (Bond et al., 1992) would have been 14 responsible for a decreased cross equatorial heat transport in the Atlantic (Fig. 5A) (Bard et al., 2000). Under a sluggish AMOC, the residual heat not transported to the North Atlantic 15 16 would be trapped in the Southern Hemisphere (Broecker, 1998; Crowley, 1992). Many water 17 hosing model experiments that show a strong decrease in AMOC strength suggested that the Southern Hemisphere warming should have affected the surface layer of the BC (Kageyama 18 19 et al., 2013; Stouffer et al., 2006). This warming has been suggested for experiments under both LGM and pre-industrial boundary conditions. Here we show the first record that 20 21 corroborates this suggestion (Fig. 4B). We propose that the surface layer of the BC acted as a 22 conduit and storage volume for part of the heat not transported to the North Atlantic during 23 HS1 that was eventually shunted towards higher latitudes in the South Atlantic (Barker et al., 24 2009; Anderson et al., 2009). Moreover, the marked drop in our SST record reaching 25 minimum values at ca. 15.5 cal ka BP could be related to an intervening warm spell registered 26 within HS1 in the North Atlantic mid-latitudes (Martrat et al., 2014). We hypothesize that not 27 only millennial-scale changes associated to Termination 1 (e.g., HS1) affected the BC, but also centennial-scale fluctuations (e.g., internal structure of HS1) were registered. However, 28 29 we primarily discuss the millennial-scale events because of age model uncertainties.

Interestingly, the other high temporal resolution Mg/Ca based SST record from the western
South Atlantic covering Termination 1 shows similar changes in SST during HS1 (Figs. 1A,
5C) (Weldeab et al., 2006). This core (i.e., GeoB3129-1/3911-3) was collected off NE Brazil

at 4.61°S, thus under the influence of the NBC. The similarity in SST between both western
South Atlantic records goes beyond HS1, and is also valid for the SST drop with minimum
values at ca. 14 cal ka BP, and peak SST at ca. 13 cal ka BP, during the Bølling-Allerød (BA).
Thus, our SST record (from the BC) together with the SST record from Weldeab et al. (2006)
(from the NBC) suggest a generally in-phase behavior of the BC and the NBC regions during
HS1 and the BA. This contradicts the BC-NBC anti-phase relationship suggested by Arz et al.
(1999), at least concerning SST since we have no proxy to assess current strength.

It is worthy of note that Arz et al. (1999) based their suggestion exclusively on  $\delta^{18}$ O records 8 of planktonic foraminifera. The more negative excursion in foraminiferal  $\delta^{18}$ O that those 9 10 authors reported during HS1 for the cores collected under the influence of the BC (i.e., 11 GeoB3229-2, GeoB3202-1) if compared to the less negative excursion reported for the cores 12 under the influence of the NBC (i.e., GeoB3104-1, GeoB3117-1, GeoB3129-1/3911-3, 13 GeoB3176-1) supported the notion that the sluggish AMOC would have triggered a 14 weakening in the NBC and a strengthening in the BC (Fig. 1A). This would have been responsible for the low HS1 meridional gradient in the  $\delta^{18}$ O records published by Arz et al. 15 16 (1999).

Based on absolute SST and  $\delta^{18}O_{ivc-ssw}$  values from the NBC (Weldeab et al., 2006) and the 17 BC (this study) we are now able to show that the HS1-LGM SST ( $\delta^{18}O_{iyc-ssw}$ ) anomaly at the 18 NBC site amounts to ca. 2.5°C and 0.5 ‰ respectively, while at our BC site it is limited to ca. 19 1.3°C and 0.3 % respectively. Thus, the NBC showed larger SST and  $\delta^{18}O_{ivc-ssw}$  increases if 20 compared to the BC during HS1. Since temperature and  $\delta^{18}O_{sw}$  influence for a forminiferal  $\delta^{18}O_{sw}$ 21 22 in opposite directions, the signal of the stronger warming at the NBC was dampened by the larger increase in  $\delta^{18}O_{ivc-ssw}$ , preventing the  $\delta^{18}O$  signal in G. ruber to change (assuming a 0.2) 23 ‰ °C<sup>-1</sup>; (Mulitza et al., 2003)). So far, this stands for no BC-NBC anomaly in foraminiferal 24  $\delta^{18}$ O during HS1. Nevertheless, our BC site is located ca. 12° downstream the sites 25 investigated by Arz et al. (1999) in the BC. Because the N-S SST gradient in the western 26 South Atlantic was larger than the one for  $\delta^{18}O_{ivc-ssw}$  during HS1, it is expected that a larger 27 warming at the southern sites studied by Arz et al. (1999) overprinted the  $\delta^{18}O_{ivc-ssw}$  effect, 28 and produced the reported negative excursion in foraminiferal  $\delta^{18}$ O. 29

30 Together with the NBC record, our SST reconstruction provides evidence that the western
31 South Atlantic was indeed affected by Northern Hemisphere rapid climate change during

Termination 1. However, the thermal response of the surface layer of the western South
 Atlantic cannot be described as an anti-phase in SST between the BC and the NBC regions
 (Arz et al., 1999), but rather as a widespread and in-phase increase in SST.

4 The low SST from our record during the Younger Dryas (YD) do not agree with the high 5 temperatures reported by Weldeab et al. (2006) for the same event (Fig. 5C, D). The 6 inconsistency of the YD SST signal in the western South Atlantic may be due to: (i) the 7 smaller amplitude of the AMOC slowdown that characterized the YD if compared to HS1 8 (McManus et al., 2004; Ritz et al., 2013); (ii) the shorter duration of the YD if compared to 9 HS1 (EPICA Community Members, 2006; Rasmussen et al., 2006; Sarnthein et al., 2001) 10 related to the time needed for the South Atlantic to equilibrate after an anomalous freshwater 11 pulse in the high latitudes of the North Atlantic; and (iii) the different boundary conditions of 12 the YD if compared to those present during HS1 (Clark et al., 2012). Numerical model experiments provide key insights to these three non-exclusive possibilities. First, water hosing 13 14 model experiments that retain an active and relatively strong AMOC indeed showed a much 15 weaker expression of the bipolar seesaw, if compared to simulations in which the AMOC strongly decreases (Kageyama et al., 2013; Otto-Bliesner and Brady, 2010). Second, the 16 17 reduction of the AMOC intensity due to freshwater perturbation increases with increasing 18 duration and amount of the freshwater perturbation (Rind et al., 2001; Prange et al., 2002). 19 Third, freshwater discharge to different geographic regions in the North Atlantic has been 20 shown to trigger different responses in the AMOC (Roche et al., 2009; Otto-Bliesner and 21 Brady, 2010). Thus, all three possibilities may have acted together or independently 22 producing a different response of the western South Atlantic during the YD and HS1.

23 In addition to the bipolar seesaw, another mechanism that acts to cool the western South 24 Atlantic during specific slowdown events of the AMOC seems to exist. This mechanism may be related to a change in the wind field, more precisely to a weakening of the subtropical high 25 pressure cell (Prange and Schulz, 2004). Based on climate model results and proxy records 26 from the Benguela upwelling region, Prange and Schulz (2004) suggested a weakening of the 27 South Atlantic subtropical anticyclone in response to a reduced cross-equatorial Atlantic 28 29 Ocean heat transport. This would result in a weakening of the BC and its associated heat 30 transport from the tropics and hence a cooling at our core site. Which mechanism dominates 31 (i.e., bipolar seesaw or wind field) may depend on boundary conditions and freshwater 32 forcing function.

A similar thermal evolution spanning most of Termination 1 (i.e., HS1 and BA) is not only a 1 2 pervasive feature of the western South Atlantic (this study; Weldeab et al., 2006), but also includes the southernmost eastern South Atlantic, as reconstructed from a core raised at 3 41.10°S / 7.80°E / 4981 m water depth (Fig. 5E) (Barker et al., 2009). The high temporal 4 5 resolution Mg/Ca based SST record from the southernmost eastern South Atlantic also presents high SST during HS1 that is followed by a marked drop at ca. 14 cal ka BP and 6 7 increasing temperatures towards the onset of the YD (Barker et al., 2009). The striking 8 similarity of the three high temporal resolution (i.e., 150 yr or less between adjacent samples) 9 Mg/Ca based SST records from the South Atlantic (this study; Weldeab et al., 2006; Barker et 10 al., 2009) not influenced by continental margin upwelling (Farmer et al., 2005) or continental 11 freshwater discharge (Weldeab et al., 2007) suggest an emerging South Atlantic-wide pattern 12 in SST evolution during most of Termination 1. Still, the view of the YD as a replicate of HS1 13 seems not to hold for the western South Atlantic.

14 Recently, Weber et al. (2014) suggested that Antarctic meltwater pulses may have cooled the 15 upper water column of the South Atlantic. Indeed, our sea surface temperature record shows minor (i.e., ca. 0.5°C) decreases around the two most prominent events of increased flux of 16 17 iceberg-rafted debris at the Scotia Sea (a proxy for Antarctic meltwater pulses) recorded 18 during Termination 1 (i.e., Antarctic Ice Sheet discharge (AID) event 7 between 16.91 and 19 15.75 cal ka BP, and event AID6 between 14.86 and 13.94 cal ka BP) (Chiessi et al., 2015a). 20 Thus, Antarctic meltwater pulses may have contributed to the variability of SST from the 21 subtropical domain of the Brazil Current on top of the mechanisms already described.

### 5.2 Continental mean air temperatures over southeastern South America during Termination 1

24 Most of the warming in our step-like structured MAT record takes place during the second 25 half of HS1 and during the YD, but due to the marked decrease in temporal resolution of our 26 MAT record after 14.1 cal ka BP we raise a note of caution while interpreting the temperature 27 rise during the YD (Fig. 4E) (Sarnthein et al., 2001; Rasmussen et al., 2006). Our MAT 28 record is remarkably similar to deglacial rise in atmospheric CO<sub>2</sub> and Antarctic temperatures (Fig. 6C, D, E) (EPICA Community Members, 2006; Monnin et al., 2004). Not only the 29 30 timing of the two pulses of MAT increase in our record (i.e., ca. 16.5 and 12.5 cal ka BP) is 31 synchronous with intervals of marked increases in global atmospheric CO<sub>2</sub> and Antarctic temperatures, but also the periods of relatively stable MAT are contemporaneous with periods 32

1 of a small rate of global atmospheric  $CO_2$  and Antarctic temperature increase. The two pulses 2 of sharp increase in deglacial atmospheric CO<sub>2</sub> also occurred simultaneously with increased upwelling in the Southern Ocean (Anderson et al., 2009). As suggested by Toggweiler et al. 3 (2006), warming around Antarctica may have increased upwelling through a poleward shift in 4 5 the Southern Westerlies and a corresponding increase in northward Ekman transport of surface waters. Still, the trigger for the changes in upwelling in the Southern Ocean probably 6 7 resided in the high latitudes of the North Atlantic (i.e., HS1 and the YD) and was transmitted 8 to the Southern Hemisphere via changes in atmospheric (Lee et al., 2011) or oceanic (Knutti 9 et al., 2004) circulation.

10 Since atmospheric circulation in the LPRDB is dominated by northerly winds (Fig. 1B) 11 (Kalnay et al., 1996), deglacial evolution of continental surface air temperature in the region 12 is also expected to follow the mean warming trend of low latitude regions in South America. High temporal resolution and continuous MAT records from tropical South America to the 13 14 east of the Andes spanning most of Termination 1 are, to our knowledge, still absent (Shakun 15 et al., 2012). Nevertheless, the comparison of our MAT record to other continental 16 temperature records with relatively lower temporal resolution allows evaluating the spatial 17 variability of MAT. Our MAT record, for instance, bears some resemblance with a pollen-18 derived temperature record from eastern Peru (Lake Consuelo) collected at 13.95°S (Fig. 6B, 19 C) (Bush et al., 2004). The linkage of MAT in the LPRDB to low latitude temperature 20 evolution is supported by modern observations of a relatively flat MAT profile in tropical to 21 subtropical South America between 10°N and 20°S (Fig. 1B) (Legates and Willmott, 1990).

The general agreement between our MAT and eastern Peru temperatures (Bush et al., 2004) taken together with the rise in atmospheric CO<sub>2</sub> content (Monnin et al., 2004) (Fig. 6B, C, D) suggests that greenhouse gas concentrations exerted a strong control on South American MAT during Termination 1.

Additionally, Antarctic meltwater pulses may have decreased MAT over southeastern South America (Weber et al., 2014). The modeled cooling amounts to ca. 1.0°C over the southernmost portion of the LPRDB. Given age model uncertainties, the two most prominent events of increased flux of iceberg-rafted debris at the Scotia Sea (i.e., AID7 and AID6; Weber et al., 2014) partially correlate with negative anomalies in our MAT record (Chiessi et al., 2015a). Thus, Antarctic meltwater pulses may have contributed to the variability of MAT over the LPRDB in addition to changes in atmospheric CO<sub>2</sub> concentration.

# 5.3 Combining sea surface temperatures in the western South Atlantic and continental mean air temperatures on the adjacent continent during Termination 1

Our SST record suggests that the South Atlantic, and the BC more specifically, was of 4 5 paramount importance for the southward propagation of the thermal bipolar seesaw signal of 6 HS1 (Fig. 5). Indeed, the western South Atlantic was more sensitive to AMOC forcing than 7 lowland South America, which appears to be more susceptible to atmospheric CO<sub>2</sub> changes 8 (Figs. 5 and 6). Thus, our SST and MAT records sum up to other lines of evidence supporting 9 the notion that global continental air temperature closely tracked the increase in atmospheric 10 CO<sub>2</sub> concentration during Termination 1, and that variations in the AMOC caused a 11 seesawing of heat between the hemispheres mainly impacting the oceans in the Southern 12 Hemisphere (Shakun et al., 2012).

13 Assuming no significant delay in the transport of the continental temperature signal to our 14 core site (Weijers et al., 2007; Schefuss et al., 2011), our records allow establishing a phase 15 relationship between changes in AMOC strength related to the onset of HS1 and the rise in 16 atmospheric CO<sub>2</sub>. According to our records, the decrease in AMOC strength already impacted 17 SST in the western South Atlantic as early as ca. 19 cal ka BP, while the rise in atmospheric 18 CO<sub>2</sub> only affected MAT at ca. 16.5 cal ka BP. As suggested by Denton et al. (2010), the long 19 duration of last deglaciation stadials was of fundamental importance to produce the necessary 20 large oceanic  $CO_2$  release via the Southern Ocean (Anderson et al., 2009). Thus, increasing 21 Northern Hemisphere summer insolation alone was insufficient to terminate the last 22 glaciation, and the impact of rising atmospheric  $CO_2$  was a key factor to complete the last 23 deglaciation (Denton et al., 2010; Shakun et al., 2012).

Interestingly, our SST and MAT records present different amplitudes in deglacial temperature rise. Similarly to the oceanic and continental temperature records reported by Weijers et al. (2007), our oceanic temperatures (i.e.,  $1.6^{\circ}$ C) showed a smaller amplitude if compared to our continental temperatures (i.e.,  $2.5^{\circ}$ C) (Fig. 4B, E). The deglacial amplitude of our SST record is very similar to global compilations (i.e.,  $1-2^{\circ}$ C) (e.g., MARGO Project Members, 2009) and regional reconstructions (i.e.,  $1-2^{\circ}$ C) (Carlson et al., 2008; Toledo et al., 2007), even considering the lower temporal resolution of those reconstructions if compared to our record. On the other hand, the deglacial amplitude of our MAT record is remarkably smaller than the amplitude of the few other available continental records for tropical South America, namely 5-7°C from Behling (2002) and 5-9°C from Bush et al. (2004). Nevertheless, pollen-based tropical and subtropical temperature reconstructions should be interpreted with caution since changes in moisture availability may also impact the recorded signal.

6 The difference between oceanic and continental warming during Termination 1 reported in 7 this study agrees with climate model simulations that suggest an average continental deglacial 8 warming in the tropics ca. 1.5 times higher than the deglacial warming of the tropical oceans 9 (Otto-Bliesner et al., 2006; Braconnot et al., 2012). The land/sea warming ratio is usually 10 explained through differences in evaporation between land and ocean, and through land-11 surface feedbacks (Braconnot et al., 2012).

12

#### 13 6 Conclusions

14 Our SST record from the BC in the western South Atlantic shows a marked positive anomaly during HS1. This is the first record that corroborates model suggestions that the surface layer 15 of the BC acted as an important conduit and storage volume for part of the heat not 16 transported to the North Atlantic under a sluggish AMOC. Thus, the BC was of paramount 17 18 importance in propagating southwards the thermal bipolar seesaw signal of HS1. Moreover, 19 the marked similarity to a SST record from the NBC suggests an in-phase thermal evolution 20 of the BC and the NBC during HS1 (and the BA), contradicting previous assumptions of a 21 BC-NBC anti-phase. Similar changes in SST are not only a pervasive feature of the western 22 South Atlantic but also include the southernmost eastern South Atlantic, suggesting a South 23 Atlantic-wide pattern in SST evolution during most of Termination 1. Over southeastern 24 South America, our MAT record shows that most of the deglacial warming occurred during 25 the second half of HS1 and during the YD. Changes in MAT are remarkably synchronous 26 with atmospheric CO<sub>2</sub> rise, suggesting that greenhouse gas concentrations exerted a strong 27 control on South American surface temperatures during Termination 1. The ca. 2.5 kyr lag of 28 MAT rise if compared to SST rise after the LGM corroborates the notion that the long 29 duration of HS1 was fundamental to drive the Earth out of the last glacial.

30

#### 31 Acknowledgements

1 We thank M. Segl and S. Pape for their help with the isotope and trace element analyses, 2 respectively. We thank W. Duleba, S.L. Ho and U. Merkel for discussion, and L. Barbara 3 (reviewer), one anonymous reviewer and J.P. Bernal (editor) for constructive comments. 4 Logistic and technical assistance was provided by the Captain and Crew of the R/V Meteor. 5 CMC acknowledges the financial support from FAPESP (grant 2012/17517-3). This study was partially performed during a stay of CMC at the Hanse Institute for Advanced Study in 6 Delmenhorst, Germany. SM was funded through the DFG Research Center/Cluster of 7 8 Excellence "The Ocean in the Earth System". Sample material has been provided by the 9 GeoB Core Repository at the MARUM - Center for Marine Environmental Sciences, 10 University of Bremen, Germany. The data reported in this paper will be archived in Pangaea 11 (www.pangaea.de).

12

#### 1 References

- Anand, P., Elderfield, H., and Conte, M. H.: Calibration of Mg/Ca thermometry in planktonic
   foraminifera from a sediment trap time series, Paleoceanography, 18, 1050,
   10.1029/2002PA000846, 2003.
- Anderson, R. F., Ali, S., Bradtmiller, L. I., Nielsen, S. H., Fleisher, M. Q., Anderson, B. E., and
   Burckle, L. H.: Wind-driven upwelling in the Southern Ocean and the deglacial rise in
   atmospheric CO2, Science, 323, 1443-1448, 10.1126/science.1167441, 2009.
- Antonov, J. I., Seidov, D., Boyer, T. P., Locarnini, R. A., Mishonov, A. V., Garcia, H. E.,
  Baranova, O. K., Zweng, M. M., and Johnson, D. R.: World Ocean Atlas 2009 Volume
  2: Salinity, edited by: Levitus, S., NOAA Atlas NESDIS 69, U.S. Gov. Printing Office,
  Washington, D. C., 184, 2010.
- Arz, H. W., Pätzold, J., and Wefer, G.: The deglacial history of the western tropical Atlantic as
   inferred from high resolution stable isotope records off northeastern Brazil, Earth
   and Planetary Science Letters, 167, 105-117, 10.1016/S0012-821X(99)00025-4,
   1999.
- Bard, E., Rostek, F., Turon, J. –L., and Gendreau, S.: Hydrological Impact of Heinrich Events in
   the Subtropical Northeast Atlantic, Science, 289, 1321-1324,
   10.1126/science.289.5483.1321, 2000.
- Barker, S., Greaves, M., and Elderfield, H.: A study of cleaning procedures used for
   foraminiferal Mg/Ca paleothermometry, Geochemistry, Geophysics, Geosystems, 4,
   8407, 10.1029/2003GC000559, 2003.
- Barker, S., Diz, P., Vautravers, M. J., Pike, J., Knorr, G., Hall, I. R., and Broecker, W. S.:
   Interhemispheric Atlantic seesaw response during the last deglaciation, Nature, 457,
   1097-1102, 10.1038/nature07770, 2009.
- Behling, H.: South and southeast Brazilian grasslands during Late Quaternary times: a
   synthesis, Palaeogeography, Palaeoclimatology, Palaeoecology, 177, 19-27,
   10.1016/S0031-0182(01)00349-2, 2002.
- Berbery, E. H., and Barros, V. R.: The Hydrologic Cycle of the La Plata Basin in South America,
  Journal of Hydrometeorology, 3, 630-645, 10.1175/15257541(2002)003<0630:THCOTL>2.0.CO;2, 2002.
- Bond, G., Heinrich, H., Broecker, W., Labeyrie, L., McManus, J., Andrews, J., Huon, S.,
   Jantschik, R., Clasen, S., Simet, C., Tedesco, K., Klas, M., Bonani, G., and Ivy, S.:
   Evidence for massive discharges of icebergs into the North Atlantic ocean during the
   last glacial period, Nature, 360, 245-249, 1992.
- Braconnot, P., Harrison, S. P., Kageyama, M., Bartlein, P. J., Masson-Delmotte, V., Abe-Ouchi,
   A., Otto-Bliesner, B., and Zhao, Y.: Evaluation of climate models using palaeoclimatic
   data, Nature Climate Change, 2, 417-424, 10.1038/nclimate1456, 2012.
- Broecker, W.: Paleocean circulation during the last deglaciation: A bipolar seesaw?,
   Paleoceanography, 13, 119-121, 1998.

- Bush, M. B., Silman, M. R., and Urrego, D. H.: 48,000 years of climate and forest change in a
   biodiversity hot spot, Science, 303, 827-829, 10.1126/science.1090795, 2004.
- Carlson, A. E., Oppo, D. W., Came, R. E., LeGrande, A. N., Keigwin, L. D., and Curry, W. B.:
   Subtropical Atlantic salinity variability and Atlantic meridional circulation during the
   last deglaciation, Geology, 36, 991-994, 10.1130/g25080a.1, 2008.
- Carvalho, L. M. V., Jones, C., and Liebmann, B.: The South Atlantic Convergence Zone:
   intensity, form, persistence, and relationships with intraseasonal to interannual
   activity and extreme rainfall, Journal of Climate, 17, 88-109, 2004.
- 9 Chiang, J. C. H., Cheng, W., and Bitz, C. M.: Fast teleconnections to the tropical Atlantic
   10 sector from Atlantic thermohaline adjustment, Geophysical Research Letters, 35,
   11 L07704, 10.1029/2008GL033292, 2008.
- Chiessi, C. M., Mulitza, S., Groeneveld, J., Silva, J. B., Campos, M. C., and Gurgel, M. H. C.:
   Variability of the Brazil Current during the late Holocene, Palaeogeography,
   Palaeoclimatology, Palaeoecology, 415, 28-36, 10.1016/j.palaeo.2013.12.005, 2014.
- Chiessi, C. M., Mulitza, S., Mollenhauer, G., Silva, J. B., Groeneveld, J., Prange, M.: Interactive
   comment on "Thermal evolution of the western South Atlantic and the adjacent
   continent during Termination 1" by C. M. Chiessi et al., Climate of the Past
   Discussions, 10, C2489-C2489, 2015a
- Chiessi, C. M., Mulitza, S., Mollenhauer, G., Silva, J. B., Groeneveld, J., Prange, M.: Interactive
   comment on "Thermal evolution of the western South Atlantic and the adjacent
   continent during Termination 1" by C. M. Chiessi et al., Climate of the Past
   Discussions, 10, C2490-C2490, 2015b
- Chiessi, C. M., Mulitza, S., Pätzold, J., Wefer, G., and Marengo, J. A.: Possible impact of the
   Atlantic Multidecadal Oscillation on the South American summer monsoon,
   Geophysical Research Letters, 36, 10.1029/2009gl039914, 2009.
- Chiessi, C. M., Mulitza, S., Paul, A., Pätzold, J., Groeneveld, J., and Wefer, G.: South Atlantic
   interocean exchange as the trigger for the Bølling warm event, Geology, 36, 919 922, 10.1130/g24979a.1, 2008.
- Chiessi, C. M., Ulrich, S., Mulitza, S., Pätzold, J., and Wefer, G.: Signature of the Brazil Malvinas Confluence (Argentine Basin) in the isotopic composition of planktonic
   foraminifera from surface sediments, Marine Micropaleontology, 64, 52-66,
   10.1016/j.marmicro.2007.02.002, 2007.
- Clark, P. U., Shakun, J. D., Baker, P. A., Bartlein, P. J., Brewer, S., Brook, E., Carlson, A. E.,
  Cheng, H., Kaufman, D. S., Liu, Z., Marchitto, T. M., Mix, A. C., Morrill, C., OttoBliesner, B. L., Pahnke, K., Russell, J. M., Whitlock, C., Adkins, J. F., Blois, J. L., Clark,
  J., Colman, S. M., Curry, W. B., Flower, B. P., He, F., Johnson, T. C., Lynch-Stieglitz, J.,
  Markgraf, V., McManus, J., Mitrovica, J. X., Moreno, P. I., and Williams, J. W.: Global
  climate evolution during the last deglaciation, Proceedings of the National Academy
  of Sciences, 109, E1134-E1142, 10.1073/pnas.1116619109, 2012.
- 40 Crowley, T. J.: North Atlantic Deep Water cools the southern hemisphere, 41 Paleoceanography, 7, 489-497, 10.1029/92PA01058, 1992.

- Crowley, T. J.: Proximal trigger for late glacial Antarctic circulation and CO<sub>2</sub> changes, PAGES
   news, 19, 2, 2011.
- Denton, G. H., Anderson, R. F., Toggweiler, J. R., Edwards, R. L., Schaefer, J. M., and Putnam,
   A. E.: The last glacial termination, Science, 328, 1652-1656,
   10.1126/science.1184119, 2010.
- Depetris, P. J., and Kempe, S.: Carbon dynamics and sources in the Paraná River, Limnology
   and Oceanography, 38, 382-395, 1993.
- Bepetris, P. J., Probst, J.-L., Pasquini, A. I., and Gaiero, D. M.: The geochemical characteristics
   of the Paraná River suspended sediment load: an initial assessment, Hydrological
   Processes, 17, 1267-1277, 10.1002/hyp.1283, 2003.
- Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., Henderson, G.
   M., Okuno, J., and Yokoyama, Y.: Ice-sheet collapse and sea-level rise at the Bolling
   warming 14,600 years ago, Nature, 483, 559-564, 10.1038/nature10902, 2012.
- EPICA Community Members: One-to-one coupling of glacial climate variability in Greenland
   and Antarctica, Nature, 7116, 195-198, 2006.
- Farmer, E. C., deMenocal, P. B., and Marchitto, T. M.: Holocene and deglacial ocean
   temperature variability in the Benguela upwelling region: Implications for low latitude atmospheric circulation, Paleoceanography, 20, PA2018,
   10.1029/2004pa001049, 2005.
- Fraile, I., Mulitza, S., and Schulz, M.: Modeling planktonic foraminiferal seasonality:
   Implications for sea-surface temperature reconstructions, Marine
   Micropaleontology, 72, 1-9, 10.1016/j.marmicro.2009.01.003, 2009a.
- Fraile, I., Schulz, M., Mulitza, S., Merkel, U., Prange, M., and Paul, A.: Modeling the seasonal
   distribution of planktonic foraminifera during the Last Glacial Maximum,
   Paleoceanography, 24, PA2216, 10.1029/2008PA001686, 2009b.
- Garreaud, R. D., Vuille, M., Compagnucci, R., and Marengo, J.: Present-day South American
   climate, Palaeogeography, Palaeoclimatology, Palaeoecology, 281, 180-195,
   10.1016/j.palaeo.2007.10.032, 2009.
- 29 Greaves, M., Caillon, N., Rebaubier, H., Bartoli, G., Bohaty, S., Cacho, I., Clarke, L., Cooper, 30 M., Daunt, C., Delaney, M., deMenocal, P., Dutton, A., Eggins, S., Elderfield, H., 31 Garbe-Schoenberg, D., Goddard, E., Green, D., Groeneveld, J., Hastings, D., 32 Hathorne, E., Kimoto, K., Klinkhammer, G., Labeyrie, L., Lea, D. W., Marchitto, T., 33 Martínez-Botí, M. A., Mortyn, P. G., Ni, Y., Nuernberg, D., Paradis, G., Pena, L., 34 Quinn, T., Rosenthal, Y., Russell, A., Sagawa, T., Sosdian, S., Stott, L., Tachikawa, K., 35 Tappa, E., Thunell, R., and Wilson, P. A.: Interlaboratory comparison study of 36 calibration standards for foraminiferal Mg/Ca thermometry, Geochemistry, 37 Geophysics, Geosystems, 9, Q08010, 10.1029/2008gc001974, 2008.
- Groeneveld, J., and Chiessi, C. M.: Mg/Ca of Globorotalia inflata as a recorder of permanent
   thermocline temperatures in the South Atlantic, Paleoceanography, 26, PA2203,
   10.1029/2010pa001940, 2011.

- 1Hönisch, B., Allen, K. A., Lea, D. W., Spero, H. J., Eggins, S. M., Arbuszewski, J., deMenocal, P.,2Rosenthal, Y., Russell, A. D., and Elderfield, H.: The influence of salinity on Mg/Ca in3planktic foraminifers Evidence from cultures, core-top sediments and4complementary  $\delta^{18}$ O, Geochimica et Cosmochimica Acta, 121, 196-213,510.1016/j.gca.2013.07.028, 2013.
- Hopmans, E. C., Weijers, J. W. H., Schefuß, E., Herfort, L., Sinninghe Damsté, J. S., and
  Schouten, S.: A novel proxy for terrestrial organic matter in sediments based on
  branched and isoprenoid tetraether lipids, Earth and Planetary Science Letters, 224,
  107-116, 2004.
- Hughen, K. A., Baillie, M. G. L., Bard, E., Beck, J. W., Bertrand, C. J. H., Blackwell, P. G., Buck,
  C. E., Burr, G. S., Cutler, K. B., Damon, P. E., Edwards, R. L., Fairbanks, R. G.,
  Friedrich, M., Guilderson, T. P., Kromer, B., McCormac, G., Manning, S., Ramsey, C.
  B., Reimer, P. J., Reimer, R. W., Remmele, S., Southon, J. R., Stuiver, M., Talamo, S.,
  Taylor, F. W., Van der Plicht, J., and Weyhenmeyer, C. E.: Marine04 marine
  radiocarbon age calibration, 0–26 cal kyr BP, Radiocarbon, 46, 1059-1086, 2004.
- Hut, G.: Consultants group meeting on stable isotope reference samples for geochemical and
   hydrological investigations, International Atomic Energy Agency, Vienna, 42, 1987.
- Kageyama, M., Merkel, U., Otto-Bliesner, B., Prange, M., Abe-Ouchi, A., Lohmann, G.,
   Ohgaito, R., Roche, D. M., Singarayer, J., Swingedouw, D., and Zhang, X.: Climatic
   impacts of fresh water hosing under Last Glacial Maximum conditions: a multi model study, Climate of the Past, 9, 935-953, 10.5194/cp-9-935-2013, 2013.
- 22 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., 23 White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., 24 Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, 25 D.: The NCEP/NCAR 40-Year Reanalysis Project, Bulletin of the American 26 Meteorological Society, 437-471, 10.1175/1520-77, 27 0477(1996)077<0437:TNYRP>2.0.CO;2, 1996.
- Knutti, R., Fluckiger, J., Stocker, T. F., and Timmermann, A.: Strong hemispheric coupling of
   glacial climate through freshwater discharge and ocean circulation, Nature, 430,
   851-856, 10.1038/nature02786, 2004.
- Lambeck, K., and Chappell, J.: Sea level change through the last glacial cycle, Science, 292,
   679-686, 10.1126/science.1059549, 2001.
- Lantzsch, H., Hanebuth, T. J. J., Chiessi, C. M., Schwenk, T., and Violante, R. A.: The high supply, current-dominated continental margin of southeastern South America
   during the late Quaternary, Quaternary Research, 81, 339-354,
   10.1016/j.yqres.2014.01.003, 2014.
- Lee, S.-Y., Chiang, J. C. H., Matsumoto, K., and Tokos, K. S.: Southern Ocean wind response to
   North Atlantic cooling and the rise in atmospheric CO<sub>2</sub>: Modeling perspective and
   paleoceanographic implications, Paleoceanography, 26, PA1214,
   10.1029/2010pa002004, 2011.

- Legates, D. R., and Willmott, C. J.: Mean seasonal and spatial variability in global surface air
   temperature, Theoretical and Applied Climatology, 41, 11-21, 10.1007/BF00866198,
   1990.
- Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K.,
  Zweng, M. M., and Johnson, D. R.: World Ocean Atlas 2009, Volume 1: Temperature,
  edited by: Levitus, S., NOAA Atlas NESDIS 68, U.S. Government Printing Office,
  Washington, D. C., 184, 2010.
- Lombard, F., Labeyrie, L., Michel, E., Bopp, L., Cortijo, E., Retailleau, S., Howa, H., and
   Jorissen, F.: Modelling planktic foraminifer growth and distribution using an
   ecophysiological multi-species approach, Biogeosciences, 8, 853-873, 10.5194/bg-8 853-2011, 2011.
- Maier-Reimer, E., Mikolajewicz, U., and Crowley, T.: Ocean GCM sensitivity experiment with
   an open Central American Isthmus, Paleoceanography, 5, 349-366, 1990.
- 14Manabe, S., and Stouffer, R. J.: Two Stable Equilibria of a Coupled Ocean-Atmosphere Model,15JournalofClimate,1,841-866,10.1175/1520-160442(1988)001<0841:TSEOAC>2.0.CO;2, 1988.
- MARGO Project Members: Constraints on the magnitude and patterns of ocean cooling at
   the Last Glacial Maximum, Nature Geoscience, 2, 127-132, 10.1038/ngeo411, 2009.
- Martrat, B., Jimenez-Amat, P., Zahn, R., and Grimalt, J. O.: Similarities and dissimilarities
   between the last two deglaciations and interglaciations in the North Atlantic region,
   Quaternary Science Reviews, 99, 122-134, 10.1016/j.quascirev.2014.06.016, 2014.
- McManus, J. F., Francois, R., Gherardi, J. M., Keigwin, L. D., and Brown-Leger, S.: Collapse
   and rapid resumption of Atlantic meridional circulation linked to deglacial climate
   changes, Nature, 428, 834-837, 10.1038/nature02494, 2004.
- Mix, A., Ruddiman, W. F., and McIntyre, A.: Late Quaternary paleoceanography of the
   tropical Atlantic, 1: Spatial variability of annual mean sea-surface temperatures, 0 20,000 years B.P., Paleoceanography, 1, 43-66, 1986.
- Monnin, E., Steig, E. J., Siegenthaler, U., Kawamura, K., Schwander, J., Stauffer, B., Stocker, T.
   F., Morse, D. L., Barnola, J.-M., Bellier, B., Raynaud, D., and Fischer, H.: Evidence for
   substantial accumulation rate variability in Antarctica during the Holocene, through
   synchronization of CO<sub>2</sub> in the Taylor Dome, Dome C and DML ice cores, Earth and
   Planetary Science Letters, 224, 45-54, 10.1016/j.epsl.2004.05.007, 2004.
- Mulitza, S., Boltovskoy, D., Donner, B., Meggers, H., Paul, A., and Wefer, G.:
   Temperature:δ<sup>18</sup>O relationships of planktonic foraminifera collected from surface
   waters, Palaeogeography, Palaeoclimatology, Palaeoecology, 202, 143-152,
   10.1016/s0031-0182(03)00633-3, 2003.
- 37 NGRIP members: High-resolution record of Northern Hemisphere climate extending into the
   38 last interglacial period, Nature, 431, 147-151, 10.1038/nature02805, 2004.
- 39Olson, D. B., Podestá, G. P., Evans, R. H., and Brown, O. B.: Temporal variations in the40separation of Brazil and Malvinas Currents, Deep Sea Research Part A.

- 1Oceanographic Research Papers, 35, 1971-1990, 10.1016/0198-0149(88)90120-3,21988.
- 3 Otto-Bliesner, B., Brady, E., Clauzet, G., Tomas, R., Levis, S., and Kothavala, Z.: Last Glacial 4 Maximum and Holocene Climate in CCSM3, Journal of Climate, 19, 2529-2547, 2006.
- Otto-Bliesner, B. L., and Brady, E. C.: The sensitivity of the climate response to the magnitude
   and location of freshwater forcing: last glacial maximum experiments, Quaternary
   Science Reviews, 29, 56-73, 10.1016/j.quascirev.2009.07.004, 2010.
- Peterse, F., van der Meer, J., Schouten, S., Weijers, J. W. H., Fierer, N., Jackson, R. B., Kim, J.H., and Sinninghe Damsté, J. S.: Revised calibration of the MBT–CBT
  paleotemperature proxy based on branched tetraether membrane lipids in surface
  soils, Geochimica et Cosmochimica Acta, 96, 215-229, 10.1016/j.gca.2012.08.011,
  2012.
- Peterson, R. G., and Stramma, L.: Upper-level circulation in the South Atlantic Ocean,
   Progress in Oceanography, 26, 1-73, 10.1016/0079-6611(91)90006-8, 1991.
- Prange, M., Romanova, V., and Lohmann, G.: The glacial thermohaline circulation: Stable or
   unstable?, Geophysical Research Letters, 29, 2028, 10.1029/2002GL015337, 2002.
- Prange, M., and Schulz, M.: A coastal upwelling seesaw in the Atlantic Ocean as a result of
   the closure of the Central American Seaway, Geophysical Research Letters, 31,
   L17207, 10.1029/2004gl020073, 2004.
- Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen,
   H. B., Siggaard-Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler,
   M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., and Ruth, U.: A
   new Greenland ice core chronology for the last glacial termination, Journal of
   Geophysical Research, 111, D06102, 10.1029/2005jd006079, 2006.
- Razik, S., Chiessi, C. M., Romero, O. E., and von Dobeneck, T.: Interaction of the South
   American Monsoon System and the Southern Westerly Wind Belt during the last 14
   kyr, Palaeogeography, Palaeoclimatology, Palaeoecology, 374, 28-40,
   10.1016/j.palaeo.2012.12.022, 2013.
- Reimer, P. J., and Reimer, R. W.: A marine reservoir correction database and on-line
   interface, Radiocarbon, 34, 461-463, 2001.

Reimer, P. J., Baillie, M. G. L., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Ramsey, C. B.,
Buck, C. E., Burr, G. S., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P.,
Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B.,
McCormac, F. G., Manning, S. W., Reimer, R. W., Richards, D. A., Southon, J. R.,
Talamo, S., Turney, C. S. M., Van der Plicht, J., and Weyhenmeyer, C. E.: IntCal09 and
Marine09 radiocarbon age calibration curves, 0–50,000 years cal BP, Radiocarbon,
51, 1111-1150, 2009.

Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Ramsey, C. B., Buck, C. E.,
Cheng, H., Lawrence Edwards, R., Friedrich, M., Grootes, P. M., Guilderson, T. P.,
Haflidason, H., Hajdas, I., Hatte, C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G.,
Hughen, K. A., Kaiser, K. F., Kromer, B., Manning, S. W., Niu, M., Reimer, R. W.,
Richards, D. A., Scott, E. M., Southon, J. R., Staff, R. A., Turney, C. S. M., and Van der

- Plicht, J.: IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years
   cal BP, Radiocarbon, 55, 1869-1887, 10.2458/azu\_js\_rc.55.16947, 2013.
- Rind, D., Russell, G., Schmidt, G., Sheth, S., Collins, D., deMenocal, P., and Teller, J.: Effects of
   glacial meltwater in the GISS coupled atmosphere-ocean model: 2. A bipolar seesaw
   in Atlantic Deep Water production, Journal of Geophysical Research: Atmospheres,
   106, 27355-27365, 10.1029/2001JD000954, 2001.
- Rintoul, S. R.: South Atlantic interbasin exchange, Journal of Geophysical Research, 96, 2675,
   10.1029/90jc02422, 1991.
- Ritz, S. P., Stocker, T. F., Grimalt, J. O., Menviel, L., and Timmermann, A.: Estimated strength
   of the Atlantic overturning circulation during the last deglaciation, Nature
   Geoscience, 6, 208-212, 10.1038/ngeo1723, 2013.
- Roche, D. M., Wiersma, A. P., and Renssen, H.: A systematic study of the impact of
   freshwater pulses with respect to different geographical locations, Climate
   Dynamics, 34, 997-1013, 10.1007/s00382-009-0578-8, 2009.
- Sarnthein, M., Stattegger, K., Dreger, D., Erlenkeuser, H., Grootes, P., Haupt, b., Jung, S.,
  Kiefer, T., Kuhnt, W., Pflaumann, U., Schäfer-Neth, C., Schulz, H., Schulz, M., Seidov,
  D., Simstich, J., van Kreveld, S., Vogelsang, E., Völker, A., and Weinelt, M.:
  Fundamental Modes and Abrupt Changes in North Atlantic Circulation and Climate
  over the last 60 ky Concepts, Reconstruction and Numerical Modeling, in: The
  Northern North Atlantic: A Changing Environment, edited by: Schäfer, P., Ritzrau,
  W., Schlüter, M., and Thiede, J., Springer, Berlin, 365-410, 2001.
- Schefuss, E., Kuhlmann, H., Mollenhauer, G., Prange, M., and Patzold, J.: Forcing of wet
   phases in southeast Africa over the past 17,000 years, Nature, 480, 509-512,
   10.1038/nature10685, 2011.
- Schlitzer, R.: Ocean Data View, available at: http://odv.awi.de (last access: 31 October 2014),
   2014.
- Schmidt, M. W., Chang, P., Hertzberg, J. E., Them, T. R., Ji, L., and Otto-Bliesner, B. L.: Impact
   of abrupt deglacial climate change on tropical Atlantic subsurface temperatures,
   Proceedings of the National Academy of Sciences, 109, 14348-14352,
   10.1073/pnas.1207806109, 2012.
- Schrag, D. P., Adkins, J. F., McIntyre, K., Alexander, J. L., Hodell, D. A., Charles, C. D., and
   McManus, J. F.: The oxygen isotopic composition of seawater during the Last Glacial
   Maximum, Quaternary Science Reviews, 21, 331-342, 10.1016/S0277 3791(01)00110-X, 2002.
- 35 Schulz, H. D., Ayres Neto, A., Boetius, A., Enneking, K., Fabian, K., Feseker, T., Funk, J., Gorke, 36 M., Heidersdorf, F., Hensen, C., Heuer, V., Hill, H. G., Hinrichs, S., Kasten, S., Klann, 37 M., Lacerda de Souza, C., Martinez Briao, A., Meyer, S., Mulitza, S., Niebler, H.-S., 38 Ochsenhirt, W.-T., Panteleit, B., Pfeifer, K., Schewe, F., Schwenk, T., Senorans, J. L., 39 Siemer, S., Steinmetz, E., and Wenzhöfer, F.: Report and preliminary results of 40 Meteor Cruise M46/2, Recife (Brazil) - Montevideo (Uruguay), December 2 -41 December 29, 1999, Fachbereich Geowissenschaften, Universität Bremen, Bremen, 42 107, 2001.

- Shakun, J. D., Clark, P. U., He, F., Marcott, S. A., Mix, A. C., Liu, Z., Otto-Bliesner, B.,
   Schmittner, A., and Bard, E.: Global warming preceded by increasing carbon dioxide
   concentrations during the last deglaciation, Nature, 484, 49-54,
   10.1038/nature10915, 2012.
- 5 Stocker, T. F.: The seasaw effect, Science, 282, 61-62, 10.1126/science 6 10.1126/science.282.5386.61.282.5386.61, 1998.
- 7 Stommel, H.: A survey of ocean current theory, Deep Sea Research, 4, 149-184, 1957.
- Stouffer, R. J., Yin, J., Gregory, J. M., Dixon, K. W., Spelman, M. J., Hurlin, W., Weaver, A. J.,
  Eby, M., Flato, G. M., Hasumi, H., Hu, A., Jungclaus, J. H., Kamenkovich, I. V.,
  Levermann, A., Montoya, M., Murakami, S., Nawrath, S., Oka, A., Peltier, W. R.,
  Robitaille, D. Y., Sokolov, A., Vettoretti, G., and Weber, S. L.: Investigating the
  Causes of the Response of the Thermohaline Circulation to Past and Future Climate
  Changes, Journal of Climate, 19, 1365-1387, 10.1175/JCLI3689.1, 2006.
- Stramma, L., Ikeda, Y., and Peterson, R. G.: Geostrophic transport in the Brazil Current region
   north of 20°S Deep Sea Research, 37, 1875-1886, 1990.
- Stramma, L., Fischer, J., and Reppin, J.: The North Brazil Undercurrent, Deep-Sea Research I,
   42, 773-795, 1995.
- Stramma, L., and England, M.: On the water masses and mean circulation of the South
   Atlantic Ocean, Journal of Geophysical Research: Oceans, 104, 20863-20883,
   10.1029/1999JC900139, 1999.
- 21 Stuiver, M., and Reimer, P. J.: Extended <sup>14</sup>C data base and revised CALIB 3.0 <sup>14</sup>C age 22 calibration program, Radiocarbon, 35, 215-230, 1993.
- Toggweiler, J. R., Russell, J. L., and Carson, S. R.: Midlatitude westerlies, atmospheric CO<sub>2</sub>,
   and climate change during the ice ages, Paleoceanography, 21, PA2005,
   10.1029/2005pa001154, 2006.
- Toledo, F. A. L., Costa, K. B., and Pivel, M. A. G.: Salinity changes in the western tropical
   South Atlantic during the last 30 kyr, Global and Planetary Change, 57, 383-395,
   10.1016/j.gloplacha.2007.01.001, 2007.
- Vera, C. S., Vigliarolo, P. K., and Berbery, E. H.: Cold Season Synoptic-Scale Waves over
   Subtropical South America, Monthly Weather Review, 130, 684-699, 10.1175/1520 0493(2002)130<0684:CSSSWO>2.0.CO;2, 2002.
- Wang, L.: Isotopic signals in two morphotypes of *Globigerinoides ruber* (white) from the
   South China Sea: implications for monsoon climate change during the last glacial
   cycle, Palaeogeography, Palaeoclimatology, Palaeoecology, 161, 381-394,
   10.1016/S0031-0182(00)00094-8, 2000.
- Weber, M. E., Clark, P. U., Kuhn, G., Timmermann, A., Sprenk, D., Gladstone, R., Zhang, X.,
   Lohmann, G., Menviel, L., Chikamoto, M. O., Friedrich, T., and Ohlwein, C.:
   Millennial-scale variability in Antarctic ice-sheet discharge during the last
   deglaciation, Nature, 510, 134-138, 10.1038/nature13397, 2014.
- Wefer, G., Adegbie, A. T., Böckel, B., Brauner, R., Büttner, U., Diekamp, V., Dillon, M.,
   Dobeneck, T. v., Donner, B., Eberwein, A., Heinecke-Herzog, M., Kim, J. H., Meier, S.,

- Mollenhauer, G., Olguin, H., Romero, O., Schewe, F., Schlemm, V., Schmieder, F.,
   Schünemann, H., Streng, M., and Volbers, A. N. A.: Report and preliminary results of
   Meteor Cruise M46/4, Mar del Plata (Argentina) Salvador da Bahia (Brazil),
   February 10 March 13, 2000. With partial results of Meteor Cruise M46/2.,
   Fachbereich Geowissenschaften, Universität Bremen, Bremen, 136, 2001.
- Weijers, J. W., Schefuss, E., Schouten, S., and Sinninghe Damste, J. S.: Coupled thermal and
   hydrological evolution of tropical Africa over the last deglaciation, Science, 315,
   1701-1704, 10.1126/science.1138131, 2007.
- Weldeab, S., Schneider, R. R., and Kölling, M.: Deglacial sea surface temperature and salinity
   increase in the western tropical Atlantic in synchrony with high latitude climate
   instabilities, Earth and Planetary Science Letters, 241, 699-706,
   10.1016/j.epsl.2005.11.012, 2006.
- Weldeab, S., Lea, D. W., Schneider, R. R., and Andersen, N.: 155,000 years of West African
   monsoon and ocean thermal evolution, Science, 316, 1303-1307,
   10.1126/science.1140461, 2007.
- Xie, P., and Arkin, P. A.: Global Precipitation: A 17-Year Monthly Analysis Based on Gauge
   Observations, Satellite Estimates, and Numerical Model Outputs, Bulletin of the
   American Meteorological Society, 78, 2539-2558, 10.1175/1520 0477(1997)078<2539:GPAYMA>2.0.CO;2, 1997.
- Yokoyama, Y., Lambeck, K., De Deckker, P., Johnston, P., and Fifield, L. K.: Timing of the Last
   Glacial Maximum from observed sea-level minima, Nature, 406, 713-716,
   10.1038/35021035, 2000.
- Zhu, C., Weijers, J. W. H., Wagner, T., Pan, J. -M., Chen, J. -F., and Pancost, R. D.: Sources and
   distributions of tetraether lipids in surface sediments across a large river-dominated
   continental margin, Organic Geochemistry, 42, 376-386,
   10.1016/j.orggeochem.2011.02.002, 2011.
- Zhou, J., and Lau, K. M.: Does a Monsoon Climate Exist over South America?, Journal of
   Climate, 11, 1020-1040, 10.1175/1520-0442(1998)011<1020:DAMCEO>2.0.CO;2,
   1998.
- 30

Lab ID	Core depth (cm)	Species	Radiocarbon age $\pm 1 \sigma$ error (yr BP)	1 σ calibrated age range (cal ka BP)	Calibrated age (cal ka BP)	Additional age used in the age model (cal ka BP)	References
NOSAMS75186	86	<i>G. ruber</i> and <i>G. sacculifer</i>	$9370\pm40$	10.234 - 10.168	10.2		Razik et al. (2013)
KIA35163	95	<i>G. ruber</i> and <i>G. sacculifer</i>	9920 ± 70	10.997 - 10.762	10.9		Razik et al. (2013)
	98					10.85 <sup>a</sup>	Razik et al. (2013)
KIA35162	101	<i>G. ruber</i> and <i>G. sacculifer</i>	9810 ± 110	10.891 - 10.582	10.75		Razik et al. (2013)
KIA30526	123	<i>G. ruber</i> and <i>G. sacculifer</i>	$12600\pm70$	14.180 - 13.985	14.1		Chiessi et al. (2008)
KIA30525	218	<i>G. ruber</i> and <i>G. sacculifer</i>	$13340\pm80$	15.599 - 15.306	15.45		Chiessi et al. (2008)
KIA35159	315	Mixed planktonic foraminifera <sup>b</sup>	$14520\pm30$	17.388 - 16.985	17.2		This study
KIA30524	358	Mixed planktonic foraminifera <sup>b</sup>	$14860\pm90$	17.750 - 17.484	17.6		Chiessi et al. (2008)
KIA30531	448	Yoldia riograndensis	$15590 \pm 100$	18.576 - 18.333	18.45		Chiessi et al. (2008)
KIA30530	583	Yoldia riograndensis	$16400 \pm 120$	19.479 - 19.143	19.3		Chiessi et al. (2008)

1 Table 1. Accelerator mass spectrometer radiocarbon ages and calibrated ages used to

2 construct the age model of core GeoB6211-2.

<sup>3</sup> <sup>a</sup>Interpolated value between the calibrated radiocarbon ages at 95 and 101 cm depth.

4 <sup>b</sup>Mixed planktonic foraminifera contained G. ruber (pink and white), G. sacculifer, G. bulloides, G. siphonifera, T.

5 quinqueloba, G. glutinata, G. uvula, G. conglobatus, and G. falconensis.

6

1 Figure 1. Location of the marine sediment core investigated in this study and other archives 2 discussed in the text. (A) Annual mean sea surface temperatures (color shading, °C) 3 (Locarnini et al., 2010), and main annual mean surface ocean currents in the western South 4 Atlantic (black arrows) (Peterson and Stramma, 1991; Stramma and England, 1999). Thin 5 white lines depict the main tributaries of the La Plata River. Numbers indicate the locations of 6 the following archives: (1) GeoB3104-1 and GeoB3117-1 (Arz et al., 1999); (2) GeoB3129-7 1/3911-3 (Arz et al., 1999; Weldeab et al., 2006); (3) Lake Consuelo (Bush et al., 2004); (4) 8 GeoB3202-1 and GeoB3229-2 (Arz et al., 1999); (5) mean location of the sites described by 9 Behling (2002); (6) SAN76 (Toledo et al., 2007); (7) KNR159-5-36GGC (Carlson et al., 10 2008); (8) GeoB6211-2 (this study). ACC: Antarctic Circumpolar Current; BC: Brazil 11 Current; MC: Malvinas Current; NBC: North Brazil Current; SAC: South Atlantic Current; 12 SSEC: Southern South Equatorial Current. (B) Annual mean air temperature (color shading, 13 °C) (Legates and Willmott, 1990), and annual mean low-level atmospheric circulation (black 14 arrows) (Kalnay et al., 1996). Dashed black line depicts the La Plata River drainage basin. 15 This figure was partly produced with Ocean Data View (Schlitzer, 2014).

16

**Figure 2.** Histograms of the mean annual cycle of precipitation at selected stations in the La Plata River drainage basin (Xie and Arkin, 1997), and mean annual cycle of the La Plata River discharge on the lower right panel (Berbery and Barros, 2002). The color shading in the central panel depicts annual mean sea surface temperatures (°C) (Locarnini et al., 2010). The number "1" in the central panel indicates the location of marine sediment core GeoB6211-2 (this study). This figure was partly produced with Ocean Data View (Schlitzer, 2014).

23

Figure 3. Age model and sedimentation rates for marine sediment core GeoB6211-2.

25

Figure 4. Proxy records for the western South Atlantic and southeastern South America spanning Termination 1 based on marine sediment core GeoB6211-2 together with ice core temperature records. (A) North Greenland Ice Core Project (NGRIP members, 2004)  $\delta^{18}$ O plotted versus the Greenland Ice Core Chronology 2005 (GICC05) (Rasmussen et al., 2006). (B) GeoB6211-2 *Globigerinoides ruber* (white) Mg/Ca and Mg/Ca based sea surface temperatures. (C) GeoB6211-2 *Globigerinoides ruber* (white) stable oxygen isotope ( $\delta^{18}$ O)

(partially from Chiessi et al. (2009)). (D) GeoB6211-2 continental-ice-volume-corrected 1 oxygen isotopic composition of surface sea water ( $\delta^{18}O_{ivc-ssw}$ ), a proxy for salinity. (E) 2 GeoB6211-2 Methylation Branched Tetraether (MBT') and Cyclisation Branched Tetraether 3 4 (CBT) based mean air temperature (MAT). (F) EPICA Dronning Maud Land (EPICA Community Members, 2006)  $\delta^{18}$ O plotted versus its original chronology. Black symbols at 5 6 the bottom of the panel depict calibrated radiocarbon ages used to produce the age model for 7 GeoB6211-2. Grey vertical bars depict Heinrich Stadial 1 (HS1) (Sarnthein et al., 2001) and 8 the Younger Dryas (YD) (Rasmussen et al., 2006). BA: Bølling-Allerød; EH: early Holocene.

9

10 Figure 5. Millennial-scale variability of sea surface temperatures of the Brazil Current 11 spanning Termination 1 compared to available circum-Atlantic records. (A) SU8118  $U^{K'_{37}}$ based sea surface temperatures (Bard, 2000). (B) OCE326-GGC5 <sup>231</sup>Pa/<sup>230</sup>Th based record of 12 13 the strength of the Atlantic meridional overturning circulation (McManus et al., 2004). (C) 14 GeoB3129-1/3911-3 Globigerinoides ruber (white) Mg/Ca based sea surface temperatures 15 (Weldeab et al., 2006). (D) GeoB6211-2 Globigerinoides ruber (white) Mg/Ca based sea 16 surface temperatures. (E) TN057-21 Globigerina bulloides Mg/Ca based sea surface 17 temperatures (Barker et al., 2009). (F) Southern Hemisphere proxy temperature stack (Shakun 18 et al., 2012). Grey vertical bars depict Heinrich Stadial 1 (HS1) (Sarnthein et al., 2001) and 19 the Younger Dryas (YD) (Rasmussen et al., 2006). BA: Bølling-Allerød; EH: early Holocene.

20

21 Figure 6. Millennial-scale variability of the mean air temperature of southeastern South 22 America spanning Termination 1 compared to selected available records. (A) Southern 23 Hemisphere proxy temperature stack (Shakun et al., 2012). (B) Consuelo Lake pollen based 24 temperature anomalies relative to modern (Bush et al., 2004). (C) GeoB6211-2 Methylation 25 Branched Tetraether (MBT') and Cyclisation Branched Tetraether (CBT) based mean air 26 temperature (MAT). (D) EPICA Dome C (Monnin et al., 2004) atmospheric CO<sub>2</sub> plotted 27 versus its original chronology. (E) EPICA Dronning Maud Land (EPICA Community 28 Members, 2006)  $\delta^{18}$ O plotted versus its original chronology. Grey vertical bars depict 29 Heinrich Stadial 1 (HS1) (Sarnthein et al., 2001) and the Younger Dryas (YD) (Rasmussen et 30 al., 2006). BA: Bølling-Allerød; EH: early Holocene.











