

1 **Multi-century lake area changes in the Southern Altiplano: A tree-ring based**
2 **reconstruction**

3

4 M.S. Morales¹, J. Carilla², H. R. Grau², R. Villalba¹

5

6 ¹Instituto Argentino de Nivología, Glaciología y Cs. Ambientales, CONICET CCT

7 Mendoza, Argentina (mmorales@mendoza-conicet.gob.ar)

8 ²Instituto de Ecología Regional, Universidad Nacional de Tucumán, Argentina.

9 CONICET

10

11 *Correspondence to:* M.S. Morales (mmorales@mendoza-conicet.gob.ar)

1 **Abstract.** Size fluctuations in endorheic lakes in northwestern Argentina (NWA) and
2 southwestern Bolivia (SWB) are very sensitive to basin hydrological balances, and
3 consequently, very vulnerable to deleterious effects from climatic changes. The
4 management of these water resources and their biodiversity requires a comprehensive
5 knowledge of their natural variability over multiple time scales. In this study, we
6 present a multi-century reconstruction of past lake-area fluctuations in NWA and SWB.
7 The evidence used to develop and validate this reconstruction includes satellite images
8 and a century-long tree-ring record from *P. tarapacana*. Inter-annual fluctuations in
9 lake area of nine lakes were quantified based on Landsat satellite images over the
10 interval 1975 to 2009. A regional *P. tarapacana* tree-ring chronology, composite from
11 two sampling sites, was used as predictors in a regression model to reconstruct the mean
12 annual (January-December) lake area from the nine lakes. The reconstruction model
13 captures 62% of the total variance in lake-area fluctuations and shows adequate levels
14 of cross-validation. This high-resolution reconstruction covers the past 601 yr and
15 characterized the occurrence of annual to multi-decadal lake area fluctuations and its
16 main oscillation modes of variability. Our reconstruction points out that the late twenty
17 century decrease in lake area was exceptional over the period 1407–2007; a persistent
18 negative trend in lake area is clear in the reconstruction and consistent with glacier
19 retreat and other climate proxies from the Altiplano and the tropical Andes. Since the
20 mid 1970s, the Vilama-Coruto lake system recorded an accelerated decrease in area
21 consistent with an increasing recurrence of extremely small lake-area events.
22 Throughout the 601 yr, the reconstruction provides valuable information about spatial
23 and temporal stabilities of the relationships between changes in lake area, ENSO, and
24 PDO, highlighting the Pacific influence over most modes of lake area variability.
25 Global and regional climate models for the Altiplano project a marked reduction in

1 precipitation to the end of the 21st century, exacerbating presently dry conditions. These
2 results provide a baseline for the historical range of variability in lake fluctuations and
3 thus should be considered for the management of biodiversity and water resources in the
4 Central Andes during the next decades.

5

6 **1. Introduction**

7

8 The southern Altiplano or dry Puna, is located in the southern subtropical Andes. A
9 prominent feature of the landscape between 21.5° and 24°S is the shallow salty lakes in
10 the lowest point of endorheic watersheds at very high elevation (above 4400 masl).

11 Productivity in the dry Puna is low and mainly concentrated in the wetlands (lakes and
12 peat bogs), which consequently play a critical role in sustaining a unique diversity of
13 rare and endemic biota (Squeo et al., 2006). Bird and large vertebrate species depend
14 upon the wetlands for grazing, nesting and water. This is an extreme environment with a
15 prolonged dry season, strong winds, high daily temperature amplitude with frequent
16 frosts, intense solar radiation, and hypoxia. These conditions prevent agriculture
17 development and permanent grazing, resulting in the absence of human settlement
18 above 4000 m. During summer, however, the zone is used as grazing area for livestock,
19 which is an important component of the local indigenous economy (Nielsen, 2003). The
20 ecological dynamics of these wetlands are sensitive to the watershed's hydrological
21 balance, related mainly to precipitation and temperature (Carilla et al., 2013); and lake
22 ecosystems act as direct and indirect indicators of climate variability.

23 During the twentieth century, the tropical and subtropical Andes have
24 experienced a persistent warming trend (Vuille and Bradley, 2000). Climate scenarios
25 for the 21st century predict a decrease in precipitation and an increase in temperature

1 (Urrutia and Vuille, 2009), posing a major threat for water resources in this arid region.
2 Changes in lake area, reflecting hydrological dynamics and vegetation productivity
3 (Carilla et al., 2013), allow the assessment of climate change effects in areas and
4 periods with poor or non-existing instrumental records. Lake hydrological dynamics are
5 usually measured as fluctuations in lake levels recorded at gauging stations but this
6 region lacks such instrumental data. However, inter-annual changes in lake area
7 between 1985 and 2009 were quantified based on Landsat satellite images and their
8 association with tree-ring and instrumental climatic records (adjacent to our study area)
9 was analyzed (Carilla et al., 2013).

10 *P. tarapacana* are small trees that grow between 4200 and 5200 masl on the
11 slopes of the volcanoes near the lacustrine areas studied. This extremely moisture
12 sensitive woody species has allowed the development of a network of multi-century
13 tree-ring width chronologies along the South American Altiplano from 16 to 23°S
14 (Morales et al., 2012). As the yearly growth of *Polylepis* trees varies depending on
15 hydrological balance (Soliz et al., 2009; Christie et al., 2009), annual rings of this
16 species may therefore be used to reconstruct past lake area fluctuations.

17 The main goal of our study was to use *P. tarapacana* ring-width records and
18 Landsat images to develop a high-resolution multi-century reconstruction of past
19 fluctuations in lake area from the Vilama-Coruto region in Northwest Argentina (NWA)
20 and Southwest Bolivia (SWB). We describe temporal fluctuations of lake area and put
21 recent changes and trends in the context of the past six centuries. We compare this
22 reconstruction with palaeo-hydroclimatic records from the subtropical and tropical
23 Andes. In addition, we analyzed the recurrence of extreme drought events, the presence
24 of persistent periodicities and dominant oscillation modes of variation at different time
25 scales. Finally, in order to identify the major climatic factors influencing variations in

1 lake area, we compared the Vilama-Coruto lake reconstruction with indices of El Niño-
2 Southern Oscillation (ENSO) and the Pacific Decadal Oscillation (PDO).

3

4 **2. Methods**

5

6 **2.1. Study system**

7

8 The study area is located on the border between Argentina (Rinconada, Jujuy)
9 and Bolivia (Sud Lipez, Potosí) from 22° 10' to 22° 45'S and from 66° 25'to 67° 13'W
10 (Fig. 1). This area shows the influence of a prolonged volcanic activity during the
11 Upper Cenozoic (Coira et al., 1996) with the presence of a large concentration of
12 shallow lakes that form an interrupted chain of endorheic watersheds above 4400 m.
13 The study area belongs to the broader ecological region called dry Puna, located in the
14 high-elevation plateaus of the southern subtropical Andes (Cabrera et al., 1976).
15 Vegetation cover does not exceed 20% and is dominated by Poaceae (*Festuca*,
16 *Deyeuxia*), several small shrubby species (*Parastrephia*, *Adesmia*, *Acantholipia*), and
17 cushion plants (*Azorella*, *Pycnophyllum*) (Cabrera et al., 1976). *P. tarapacana* is a
18 small tree, ca 2-3m tall, and is the largest woody species that grows on the slope of the
19 high volcanoes between 4200-5200 m. Extreme aridity (<150 mm annual rainfall) and
20 low temperatures (mean annual < 5°C) characterize this region
21 (<http://www.worldclim.org>). Precipitation is of convective type and mostly concentrated
22 during summer. More than 80% of the total annual rainfall falls between November and
23 March (Vuille et al., 2003; Vuille and Keimig, 2004). Summer precipitation variability
24 is related to local orography and with changes of the upper troposphere zonal wind,
25 which in turn favor or block the ingression of wet easterly flow transporting humid air

1 masses from the Amazon basin (Vuille et al., 2000; Garreaud and Aceituno, 2001;
2 Bradley et al., 2003; Garreaud et al., 2003).

3

4 **2.2. Lake area records**

5

6 Between 1975 and 2009, inter-annual fluctuations in lake area from nine lakes were
7 quantified using Landsat images. The nine lakes are located between 4400 and 4600 m
8 in the Vilama-Coruto region from NWA-SWB (Fig. 1). Lake area variations were
9 derived from Landsat MSS (80 m pixel resolution, path 249 row 076) for the period
10 1975-1982, and Landsat TM images (30 m pixel resolution path 232 and row 076) for
11 the period 1984-2009. We calculated lake area based on the number of pixels and their
12 sizes. Pixel size for Landsat MSS is 80 x 80m (0.0064 km²) and for Landsat TM is 30 x
13 30m (0.0009km²). For example in November 1978 Cerro Negro lake encompassed 316
14 pixels of Landsat MSS corresponding to 2.02 km² (316 * 0.0064km²). In October 1988
15 the same lake, encompassed 2551 pixels of Landsat TM image, which is the same as
16 saying 2.3 km² (2551 * 0.0009 km²). The area of the lakes in each particular image was
17 determined using the non-parametric method Support Vector Machine (SVM; Hsu et
18 al., 2007). Good quality available images for the different years of the period 1975-
19 2009, ranged from one to ten (dates of images are in the supplement, S1). Despite of the
20 variable acquisition dates and number of the images, relative low intra-annual area
21 variability was recorded for the nine lakes. A comparison between the coefficients of
22 variation for each lake showed that inter-annual variability represented more than the
23 double of the intra-annual variability (see the supplement, S2.1-9). Therefore, we
24 assume that the monthly or seasonal variation was negligible in relation to the higher
25 inter-annual lake area variation, and an area value from a particular image could well

1 represent the mean lake area condition of the year. To further validate this assumption,
2 autocorrelation analyses were performed to determine persistence in the monthly and
3 annual time series averaged for the nine lakes from 1975 to 2009 period (see the
4 supplement, S3). This analysis showed strong 19-month persistence in the monthly
5 series, while two-year persistence was found in the annual series. AnClim program
6 (Štěpánek, 2008) was used to perform this analysis.

7 The annual (January-December) lake area record for each of the nine lakes was
8 estimated by averaging every available image (from one to ten) for a particular year.

9 Based on the nine individual records, we developed a regional record of annual lake
10 area fluctuations. To minimize the influences of lakes with different sizes on the
11 regional mean, annual lake area records were standardized as Z-score. Finally, the
12 standardized lake area records from the nine lakes were averaged to develop a regional
13 mean covering the interval 1975-2009. There was no image available in the entire
14 region for the year 1983. The model used for the calibration and verification of the past
15 lake area reconstruction requires continuous time series without missing years.
16 Therefore, a linear regression model between the regional lake area and a regional
17 precipitation index from the Altiplano was applied to estimate the 1983 lake area value.
18 A list of the meteorological stations used for this analysis is reported in the supplement
19 (S5).

20

21 **2.3. Tree-ring data collection and standardization**

22

23 Our study area encompasses the southern latitudinal limit of *P. tarapacana* distribution
24 and tree stands are mostly scattered on wetter north and east slopes of high volcanoes
25 but not on the south and west slopes. Sample sites include the *P. tarapacana*

1 populations located in relatively close proximity to the study lakes. Cross-sections and
2 wedges were obtained from dead and living trees. From a total of seven sampled sites,
3 we selected those that were highly replicated (> 60 tree-ring series), contain at least 300
4 years of record, and were high correlated with lake area fluctuations ($r > 0.5$; Table 1).
5 Two sites met these criteria: the tree-ring records from the Granada and Uturunco
6 volcanoes (Table 1). The Schulman's convention (1956) for the Southern Hemisphere
7 was used to assign to each ring the date of the calendar year in which radial growth
8 starts. We measured total ring width to ± 0.001 mm and cross-dated samples using
9 standard procedures (Stokes and Smiley, 1968; Fritts, 1976). The program COFECHA
10 (Holmes et al., 1983) was used to detect measurement and cross-dating errors. The ring-
11 width chronologies from Granada and Uturunco shared large percentage of common
12 signal in tree-ring variations ($r > 0.62$; 1620-2006 common period), therefore, 155 tree-
13 ring series were merged into a regional chronology. Ring-width measurements were
14 standardized to remove the age-related growth trends and minimize the growth
15 variations not related to climate (Fritts, 1976). A negative exponential curve or straight
16 line was used to standardized ring-width series. When this standardized growth trend is
17 removed, some of the variance related to climatic forcing signal can also be removed,
18 leading to a trend distortion in resulting index series (Melvin, 2004). To prevent trend
19 distortion the signal free detrending procedure (Melvin, 2004; Melvin and Briffa, 2008)
20 was applied to the tree-ring measurements. The regional tree-ring chronology was
21 produced with the RCSigFree program (Tree Ring Lab-LDEO, Columbia University
22 <http://www.ldeo.columbia.edu/tree-ring-laboratory/resources/software>).

23 The quality of the tree-ring chronology was tested based on standard chronology
24 diagnostics such as the RBar (Briffa, 1995) and the Expressed Population Signal (EPS;
25 Wigley et al., 1984; Table 1; S7). The RBar is the mean correlation coefficient from all

1 possible pairings among individual cores in the chronology (Briffa, 1995). EPS is a
2 measurement of the strength of the common signal in the chronology used to identify
3 the most reliable interval in the chronology valid for climate reconstructions. To
4 calculate the RBar and EPS, we used a 50-year window with an overlap of 25-year
5 between adjacent windows.

6

7 **2.4. Lake area reconstruction**

8

9 First, we identified the best relationship between variations of regional lake area and
10 inter-annual tree growth. For that purpose, we computed correlation coefficients
11 between the regional standard *P. tarapacana* chronology and annual mean lake area
12 records. The highest correlation coefficient with tree-ring variations was found using for
13 comparison the regional mean annual (January-December) lake area ($r = 0.70$; $p <$
14 0.0001). We reconstructed, using a linear regression, the mean January-December lake
15 area from the Vilama-Coruto lacustrine region over the period 1407-2007. To capture
16 possible climate-related persistence in the tree-ring series and enhance the common
17 climate signal, we included as predictors in the regression the regional chronology in
18 the temporal lags significantly correlated ($p < 0.05$) to annual lake area during the
19 calibration period 1975–2007. The chronology at year t and a lag ($t+1$) were considered
20 as candidate predictors of annual lake area in a stepwise multiple regression (Weisberg,
21 1985). The “leave-one-out” cross-validation procedure (Michaelsen, 1987; Meko and
22 Baisan, 2001) was used to validate the model (Fig. 2). This method allowed the use of
23 all available data to calibrate and validate the model, and it is very useful when
24 observed records are short temporal series, such as our lake area records. In this
25 approach, the reconstruction model is calibrated using the full available overlap period.

1 Then, the method sequentially withheld one observation and calibrates additional
2 models with the remaining observations. Each model is used to estimate the withheld
3 observation. Finally, all these estimated values are assembled in a time series used as
4 the validation data set. The quality of the model during the calibration period was
5 measured by the proportion of variance explained by the regression (R^2 adj). The model
6 accuracy was evaluated using the reduction of error (RE) statistic, which measures the
7 model skill in estimating values of the predictand in the cross-validation process
8 (Wilkes, 1995; Fritts, 1976). We also computed the root-mean-square error (RMSE)
9 statistic as a measure of inherent uncertainties in the reconstruction (Weisberg, 1985).

10

11 **2.5. Identification of extreme high/low events in the lake area reconstruction**

12

13 Significant shifts in the mean state conditions of the lake area reconstruction were
14 identified using a simple regime shift detection technique (Rodionov, 2004) with a cut-
15 off segment length $l = 25$ years, a target probability level $p = 0.1$ and a Hubert's weight
16 parameter $h = 1$ (Fig. 3). In order to determine the intensity and duration of drought
17 (pluvial) events in the lake area from the Vilama-Coruto region, we computed the
18 lowest (highest) reconstructed n -year running means for $n = 1, 5, 25$ and 50 yr (Table
19 2). In addition, we analyzed the occurrence rate of extreme reduction in lake area within
20 the reconstruction (Fig. 4). For that purpose, we employed a kernel estimation technique
21 (Mudelsee et al., 2004; Girardin et al., 2006). This method allows the detection of non-
22 monotonic trends and imposes no parametric restrictions. A Gaussian kernel function
23 was applied to weigh reconstructed years of extreme decrease in lake area and calculate
24 the occurrence rate (probability per time unit). A Kernel bandwidth of 50 years was
25 used and pseudo-data generation rule 'reflection' to reduce boundary bias (Mudelsee et

1 al., 2004). Confidence bands at the 90% level were obtained by using 2000 bootstrap
2 simulations (Cowling et al., 1996; Mudelsee et al., 2004). The dates of extreme events
3 were define by a percentile threshold of 20%.

4

5 **2.6. Spatial and spectral analyses of the reconstructed lake area and climate** 6 **forcing**

7

8 The influence of Pacific sea surface temperatures on lake area fluctuations in the
9 Vilama-Coruto area was estimated by examining the spatial correlation coefficient
10 pattern between the reconstructed annual (January–December) lake area and the annual
11 averaged (January-December; 2.5×2.5 gridded cell) sea surface temperatures from the
12 NCEP reanalysis global dataset (Kistler et al., 2001:
13 <http://www.cpc.ncep.noaa.gov/data/indices/>; Fig. 5). Additionally, power-spectral and
14 cross-spectral analyses (Jenkins and Watts, 1968) were used to determine the frequency
15 domains of the reconstructed lake area variations in the Vilama-Coruto region and the
16 annual averaged Pacific sea surface temperatures from the N3.4 Pacific sector (N3.4
17 SST; Fig. 6).

18 Blackman-Tukey (BT) spectral analysis (Jenkins and Watts, 1968) was
19 computed for the periods 1407–2007 and 1872–2007 for the reconstruction and the
20 Pacific N3.4-SST, respectively. The number of lags used for the auto- and cross-
21 covariance functions was equal to 30% of the series length. Spectral functions were
22 smoothed with the Hamming filter. The 95% confidence level of the spectrum was
23 estimated from a “red noise” first-order Markov null continuum based on the lag-1
24 autocorrelation of the time series (Mitchell et al., 1966). A coherency spectral analysis
25 was computed over the common period 1872–2007 between the reconstructed lake area
26 and N3.4-SST records. The coherency spectrum measures the variance in common

1 between reconstructed lake area and N3.4-SST as a function of frequency. Both, BT and
2 coherency analysis were performed using the AnClim program (Štěpánek, 2008).

3 To detect the dominant, non-stationary oscillatory modes of variability in the
4 601-yr series of reconstructed lake area we used Singular Spectral Analysis (SSA;
5 Vautard and Ghil, 1989; Fig.7). SSA is a non-parametric method related to empirical
6 orthogonal function analysis that samples lagged copies of time series at equal time
7 intervals and calculates eigenvalues and eigenvectors of the auto-covariance matrix
8 (Vautard and Ghil, 1989; Vautard, 1995). SSA enables the user to evaluate the changing
9 periodic behaviour in a single time series by extracting ‘reconstructed components’ or
10 ‘waveforms’, which represent the dominant periodic modes of the time series. This
11 method detects changes in amplitude and phasing of the dominant cycles over time
12 because waveforms retain their temporal periodic behaviour (Vautard and Ghil, 1989;
13 Vautard, 1995; Speer et al., 2001). In addition, SSA was computed for the N3.4-SST
14 and PDO indices to assess the relationship in the time frequency domain between the
15 reconstructed lake area and Pacific climate forcing. The SSA analysis was performed by
16 using the SSA program (Boninsegna and Holmes, unpublished operating manual [on
17 file at Laboratory of Tree-Ring Research, University of Arizona]).

18 19 **3. Results**

20 21 **3.1. Tree-ring chronology and reconstruction model**

22
23 In this study, 155 tree-ring samples from dead and living trees of *P. tarapacana* were
24 combined to develop a robust composite tree-ring chronology (S6) for the Vilama-
25 Coruto region in Northwest Argentina (NWA) and Southwest Bolivia (SWB). The
26 regional chronology covers the period AD 1242-2008, but is best replicated since AD

1 1407 (N > 10 ring-width series). As previously demonstrated (Morales et al., 2004,
2 2012; Carilla et al., 2013), *P. tarapacana* trees are drought sensitive and show relative-
3 high variability in tree-ring widths (mean sensitivity = 0.30). The RBar and EPS
4 statistics indicate that the chronology is of good quality with a strong common signal in
5 ring width between individuals (Table 1). The chronology statistics are similar to those
6 previously documented for *P. tarapacana* tree-ring records from the Bolivian Altiplano
7 and the Argentine Puna (Morales et al., 2004; Solíz et al., 2009; Christie et al., 2009;
8 Morales et al., 2012; Carilla et al., 2013). The EPS remains above the threshold of 0.85
9 over the period 1399-2008. The mean EPS for the period 1407-2008 is 0.96 (Table 1).
10 The “signal free” standardization method, using negative exponential curves, allows to
11 capture appreciable amounts of both low- (multi-decadal) and high-frequency (inter-
12 annual) variability in lake area fluctuations.

13 We present here a 601-yr high-resolution tree-ring reconstruction of annual area
14 fluctuations from nine lakes in the Vilama-Coruto region. Our reconstruction, calibrated
15 and validated on satellite image January-December lake area (1975–2007; Fig. 2),
16 accounts for 62% of the total variability in lake area changes (R_{adj}^2 adjusted for loss of
17 degrees-of-freedom; Draper and Smith, 1981). The consistence between the observed
18 and estimated lake area indicates good quality in the calibration model for reproducing
19 past lake area fluctuations. The high predictive power of the calibration model is
20 consistent with a highly significant F value (F=25.4; P < 0.001) and a largely positive
21 Reduction of Error (RE = 0.54). Analyses of regression residuals indicated no violations
22 of regression assumptions. The residuals of the regression models are normally
23 distributed, not significantly autocorrelated and without a significant trend according to
24 Durbin-Watson tests (Fig. 2b).

25

1 **3.2. Temporal evolution of the lake area reconstruction**

2

3 The lake area reconstruction for the Vilama-Coruto region is characterized by inter-
4 annual variations embedded within decadal to multi-decadal fluctuations (Fig. 3).
5 Significant persistent changes in the mean state of the reconstruction were detected with
6 the Rodionov analysis (Rodionov, 2004; Fig. 3). This analysis shows that lake-area
7 conditions around the long-term mean for the period 1407-1440 were followed by a lake
8 area increase lasting to the beginning of the 16th century (1503). The 15th century high
9 lake level stand was characterized by high inter-annual variance, with extreme high- and
10 low-lake area years. The longest period of comparatively low-lake area spans most of
11 the 16th century (1504-1595). This long-term dry period was interrupted by two decade-
12 scale periods showing contrasting high and low lake areas centered on the years 1600
13 and 1620, respectively. After that, significant high lake area conditions prevailed for
14 more than 160-yr (1627-1790). Two of the five years with the largest lake areas were
15 recorded during this period (1734 and 1744; Table 2). The 35-yr smoothing spline
16 shows a period of gradual reduction in lake area between 1750 and 1790). However, no
17 significant shifts in mean lake-area extent were detected over this interval. For the
18 period 1997-1818 low lake area conditions were identified. A marked shift from low- to
19 high-lake areas was identified around 1820. The period 1820-1859 encompasses the
20 highest lake areas reconstructed during the past 601 years. After this long-term increase
21 in lake area, a gradual decrease in lake areas was reconstructed to the first decades of
22 the 20th century. After that, a steady negative trend in lake area begins in the 1930s and
23 persists to the present. The most pronounced negative decline for the entire
24 reconstruction is recorded for the period 1976 to 2007. During this 30 year period 1987

1 was the only year to exceed the long-term historical mean area. Three of the five years
2 with the lowest lake areas occurred during the last three decades (Table 2).

3 A different picture of the long-term history of changes in lake area periods is
4 given by averaging non-overlapping intervals of 5-, 25-, and 50-years (Table 2). Based
5 on a 5-year moving average, three of the five periods of lowest lake area occurred since
6 1930 (i.e. 1938-1942, 1998-2002, 2003-2007), whereas the highest lake-area periods in
7 the reconstruction were centered around 1740s and 1840s (Table 2). In terms of 25-year
8 moving averages, the period 1983-2007 is has the smallest 25 year mean and is
9 substantially lower than the other four lowest events. The standardized anomalies of
10 lake area during this interval were substantially lower than during 1608–1632, which
11 ranks second among the 25-year long lowest periods (Table 2). The marked lake area
12 increase from 1833 to 1857 is also a prominent feature revealed by the 25-year moving
13 averages. The intervals 1958–2007 and 1808–1857 are similarly the lowest and highest
14 50 year periods in this 600 year record (Table 2). These results indicate that lake-area
15 during the second half of 20th century has been comparatively the lowest, whereas
16 during the 19th century have been the higher in the context of the past 601 years.

17 Following these results, the reconstruction suggests an increase in the occurrence
18 rate of extreme events in lake-area reduction during the 20th century (Fig. 4). To assess
19 changes in the occurrence rate of extreme lake-area along the reconstruction, we
20 employed the recurrence analysis by Mudelsee et al. (2004). The recurrence interval of
21 extreme events (< 20% of the mean area) was between 4-7 years in the 15th to 18th
22 centuries and between 7-15 years in the 19th century. It has steadily increased from ca 6
23 in the 1930s to 2 in recent years.

24

25 **3.3. Spectral properties and Pacific influence on lake area reconstruction**

1

2 The spatial correlation pattern between the reconstructed lake area and the global
3 gridded sea surface temperatures (SSTs) were determined for the interval 1948–2006
4 (Fig. 5). The annual lake area is significantly negatively correlated with SSTs from the
5 tropics to the mid-latitude Pacific coasts of the Americas, resembling an El Niño
6 Southern Oscillation (ENSO) – Pacific Decadal Oscillation (PDO) like pattern (Fig. 5).

7 Figure 6 shows the BT spectra for the annual lake-area reconstruction over the
8 Vilama-Coruto region and the N3.4 SSTs. Peaks that exceed the 95% confidence limit
9 in the lake area reconstruction are prominent at 12.9, 5.9-6.3, 3.2 and 3 years. Except for
10 the decadal peak of 12.9, all significant peaks coincided with the N3.4 SSTs peaks
11 recorded in Figure 6b (5.9, 3.6-3.7). At an inter-annual time scale, the lake
12 reconstruction contained similar periodicities to N3.4 SSTs, falling within the preferred
13 frequency band of ENSO (2-7 years).

14 The coherency spectrum was significant at 2.8, 3.4-3.9 year cycles, revealing
15 that the lake area reconstruction captures the inter-annual variability in SST records
16 (Fig. 6c). These results suggest that part of the variance in the lake-area reconstruction
17 from the Vilama-Coruto region reproduces the timing and duration of high frequency
18 SST anomalies.

19 We used singular spectral analysis (SSA), to isolate the main oscillatory modes
20 of the lake area reconstruction variability (Fig. 7). Several important waveforms were
21 recorded at multi-decadal (26-33 yrs.), decadal (11-15 yrs.) and inter-annual (5-8 and 2-
22 4yrs.) time scales (Fig. 7). The multi-decadal waveform amplitudes around 1600s were
23 the largest in the series at a time when two severe decade-scale reductions in lake area
24 (1583-1595; 1614-1627) were interrupted by a decade of increased lake area (1596-
25 1613). The decadal oscillation mode showed high amplitude variability during the

1 fifteenth, eighteenth and the beginning of the nineteenth centuries, and was reduced
2 during the sixteenth and the second half of nineteenth centuries. The reconstruction
3 contains an important 5-8 year cycle waveform that explains a high percentage (27%) of
4 the lake-area reconstruction variability. The amplitude of this oscillatory mode was
5 relatively large over the entire reconstruction but decreased from the 1920s to present.
6 The inter-annual oscillation mode (2-4 yrs.) shows high amplitudes during the
7 fifteenth, first half of sixteenth, second half of nineteenth and in particular during the
8 second half of twentieth centuries.

9 A comparative analysis between the dominant oscillatory modes in the
10 reconstruction with those in the Pacific Decadal Oscillation index (PDO) and in the
11 N3.4 SSTs, highlighted the occurrence of common waveforms between them at
12 different frequencies (Fig. 8). The lake-area reconstruction shares similar oscillations
13 with PDO at multi-decadal modes of variability, while by decadal and inter-annual
14 modes, the reconstruction waveforms are closely related to those recorded at N3.4
15 SSTs. High correspondence in the inter-annual modes are evident in the amplitudes of
16 the lake reconstruction and N3.4 SSTs waveforms showing that both records have large
17 and small amplitudes during 1870-1890/1960-2008 and 1930-1960, respectively; (Fig.
18 8c).

19

20 **4. Discussion and concluding remarks**

21

22 Our study represents the first high-resolution, tree-ring based reconstruction of
23 variations in lake area over the past six centuries in the Andes. Through the analysis of
24 601 years of water balances of the dry Puna lakes, this study provides a novel insight
25 into the climate variability of the subtropical Andes. The evidence used to develop and

1 validate this reconstruction includes satellite images and century-long tree-ring series of
2 *P. tarapacana*, a climate-sensitive species from the Altiplano. Consistent with the
3 results obtained by Carilla et al. (2013), the use of satellite-derived lake area records as
4 a measured of water balance, resulted a useful approach to explore hydrological trends
5 in this region devoid of local meteorological records.

6 The regression model used to develop the reconstruction explains 60% of the
7 total variance in the January-December lake area over the 1975–2007 calibration period.
8 This value is comparatively higher to that previously reported for a tree-ring based
9 precipitation reconstruction of the Southern Altiplano ($R^2_{\text{adj}}=0.54$; Morales et al., 2012).
10 At the sampling sites, precipitation explains high percentage of the total radial growth
11 variance, but also summer temperatures which increase evapotranspiration and reduced
12 soil water supply, affect tree growth variability (Morales et al., 2004). In the same way,
13 the areal extent and depth of water in lakes respond to changes in precipitation and
14 evaporation integrated over the lakes and their catchment basins (Mason et al., 1994).
15 Both, tree-growth and lake area integrate changes in climatic parameters including
16 precipitation, temperature, radiation and wind speed. As a consequence, a closer
17 association is expected between tree growth and lake extents than tree growth and
18 singular parameters such as precipitation or temperature.

19 Extended periods of small lake areas in our reconstruction, such those recorded
20 in 1504-1595, 1614-1626, 1790-1818 and 1976-2007, highlight the occurrence of
21 decadal to multi-decadal droughts in the region. Similar, relatively long-term humid
22 conditions were reconstructed during the intervals 1441-1503, 1596-1613, 1627-1796,
23 1819-1934. Concurrent dry and wet periods to those registered in our reconstruction
24 have been documented in a tree-ring based precipitation reconstruction for the South

1 American Altiplano (Morales et al., 2012) and in a gridded multiproxy PDSI (Palmer
2 Drought Severity Index) for the South American subtropics (Boucher et al., 2011).

3 Wet and cold conditions in NWA-SWB, associated with large areas of the lakes
4 during the period 1627-1796, have also been documented for other climate proxies from
5 northern locations in tropical Andes such as the $\delta^{18}\text{O}$ isotope records in the ice core
6 from the Quelccaya (Thomson et al., 2006) and in the sediment core from Pumacocha
7 (Bird et al., 2011). However, wet and cold conditions in these records occurred during
8 almost 300 years (1500-1800), whereas in our tree-ring reconstruction, these conditions
9 last 169 years from 1627 to 1796. In contrast to the Quelccaya and Pumacocha records,
10 our reconstruction shows persistent low lake areas in the 16th century. Differences
11 between these records may reflect distinct climate conditions between sites separate by
12 more than 1000 km in north-south direction along the Andes. Discrepancies between
13 proxies are also related to the individualistic nature of records (tree-rings, ice cores and
14 lake sediments) capturing environmental changes at different frequency and time lags.
15 In addition, small lake areas were recorded in this relatively wet time interval (1627-
16 1796) for the years 1661-62, 1724, 1740, and 1753-55. These dry years were also
17 identified in other high resolution precipitation reconstruction based on historical
18 documents from the silver mines from Potosi (southern Bolivia) across the period 1585-
19 1807 (Gioda and Prieto, 1999). The annual dry events of the historical series and lake
20 area reductions were consistently related at inter-annual variations during the common
21 period (see statistical analysis in the supplement; S7).

22 19th century showed the highest lake area records of the reconstruction,
23 consistent with the most important wet period identified in the precipitation
24 reconstruction from the Altiplano (Morales et al., 2012). Also, others low resolution
25 environmental proxies such as the peat-accumulating wetland from C° Tuzgle in the dry

1 Puna, Argentina (Schitteck et al., 2015) and packrat middens in the Andean precordillera
2 from the northern Atacama Desert, Chile (Mujica et al., 2014), highlight the large
3 pluvial period of the 19th century.

4 Analyses of satellite-derived lake area and vegetation index in the Vilama-
5 Coruto region by Carilla et al. (2013), revealed sustained negative trends in water
6 balance and ecosystem productivity for the periods 1985-2009 and 2000-2010,
7 respectively. These environmental changes have been associated with recent climate
8 variability in NWA-SWB (Carilla et al., 2013). Our lake-area reconstruction indicates
9 that the recorded dry condition of the late 20th century is exceptional over the 1407-
10 2007 period. Three of the five 5-yr periods with smallest lake areas occurred from 1930
11 to present, and the driest 25 and 50 year intervals occur in the second half of 20th
12 century.

13 Recurrence of extremely small lake-area events has been a common feature in
14 the Vilama-Coruto region over the past six centuries. However, the 20th century shows
15 the highest rate of extreme events, that changes from a recurrence interval of 15-yr in
16 the second half of 19th century to every 2-yr in the late 20th century.

17 Since the mid 1970s, the Vilama-Coruto lake system recorded an accelerated
18 decrease in area consistent with the driest condition of the past six centuries. This shift
19 to arid condition has also been identified in $\delta^{18}\text{O}$ isotope records in the ice core from
20 Quelccaya (Thompson et al., 2006) and sediment core from Pumacocha (Bird et al.,
21 2011). In both records, this pervasive dry period was also unprecedented in the context
22 of the last 600 yrs. The dry condition seem to be a regional pattern affecting the
23 southern tropical Andes, and a major cause of the steady shrinking of glaciers from
24 1930 to 2005 in southern Perú and Bolivia (Ramirez et al., 2001; Francou et al., 2003;
25 Vuille et al., 2008). Glacier retreat is also related to the increasing temperatures during

1 the last decades across the region (Vuille and Bradley, 2000; Urrutia and Vuille, 2009,
2 Rabatel et al., 2013). Annual temperatures deviations averaged over the tropical Andes
3 (1°N-23°S) indicate a significant warming trend for the period 1939-1998. However,
4 this trend has intensified since the mid 1970s (Vuille and Bradley, 2000) and was highly
5 consistent with the faster rate of global temperatures increase (Jones et al., 1999).

6 Significant changes in precipitation patterns in tropical and subtropical South
7 America have been identified in instrumental records during the 20th century. A trend
8 analysis (1950-1998) of atmospheric vertical motion and winds across a north-south
9 transect at 65°W in tropical-subtropical South America shows a significant increase in
10 precipitation and cloudiness in the inner tropics whereas the southern tropics and
11 subtropics become drier and less cloudy (Vuille et al., 2008). These changes in
12 precipitation and cloudiness have been associated with an intensification of the regional
13 Hadley circulation, with a more vigorous vertical ascent, favorable for convective
14 activity, in the tropics, balanced by enhanced subsidence and less cloudiness in the
15 subtropics (see figure 6, in Vuille et al., 2008).

16 The main moisture source for the Altiplano is the easterly influx from the
17 Amazon Basin (Lenters and Cook, 1997; Garreaud et al., 2009). However, wet or dry
18 episodes are related to changes in the mean zonal wind, largely modulated by Pacific
19 Ocean surface temperature (Vuille et al., 2000; Garreaud and Aceituno, 2001; Bradley
20 et al., 2003). Pacific SSTs properties, such as amplitude, frequency and associated
21 teleconnection patterns, showed considerable natural variability from inter-annual to
22 centennial time scales (Li et al., 2013). In this study spectral analyses reveal
23 periodicities and dominant waveforms in the lake–area reconstruction at multi-decadal
24 (26-33 yr), decadal (11-15 yr) and inter-annual (5-8 yr and 2-4 yr) modes of variability.
25 These oscillations were consistent with the spectral properties from the N3.4 SSTs and

1 PDO indices. The amplitudes of the different modes change throughout the
2 reconstruction. A dominant multi-decadal mode shows large-amplitude oscillations
3 around 1600, whereas the decadal mode was more important in the 1500s and from
4 1700s to the beginning of 1800s. The inter-annual 5-8 yr bandwidth was dominant for
5 almost the entire period analysed, except after 1950, where amplitudes decreased. The
6 inter-annual 2-4 yr bandwidth was strong during the second half of nineteenth and
7 twentieth centuries. The shift from a dominant decadal to an inter-annual mode was also
8 recorded in a PDO index reconstruction based on western North America and southern
9 Central Andes tree-ring chronologies (D'Arrigo et al., 2001; Villalba et al., 2001).
10 These authors present evidence of a more dominant decadal mode from 1700 to middle
11 1800s with a shift towards more inter-annual-dominated forcing around that time
12 (D'Arrigo et al., 2001; Villalba et al., 2001).

13 The main oscillatory modes between the lake-area reconstruction and the
14 instrumental PDO and N3.4 SSTs indices are similar though negatively related over the
15 1900-2007 and 1872-2007 periods, respectively. Decrease in reconstructed lake areas
16 and Pacific-SSTs amplitudes at inter-annual modes was recorded from 1930 to 1960,
17 which was coherent with a reported low ENSO activity during the same interval
18 (Aceituno and Montecinos, 1993; Torrence and Webster, 1999; Sutton and Hodson,
19 2003). These results were highly consistent with the reported relationships between the
20 main oscillation modes of the precipitation reconstruction from the Altiplano and the
21 N3.4 SSTs (Morales et al., 2012).

22 The BT spectral density analysis suggests that the lake-area reconstruction and
23 N3.4 SSTs contain similar inter-annual periodicities within the preferred frequency
24 band of ENSO (2-7 years). In particular, the strong spectral coherency between both
25 series within the 2-4 yr bandwidth indicates that the timing and magnitude of high-

1 frequency anomalies in Pacific SSTs modulate the interannual lake-area variability in
2 the Vilama-Coruto system. During the past 600 years, 16% of the variance in lake area
3 is associated with the 2-4 yr bandwidth (Fig. 7d); however, the largest contribution to
4 the total variance (27%) arises from cycles at the 5-8 yr bandwidths (Fig. 7c). The
5 amplitude of the 5-8 yr waveforms decreased from the 1930s to present. In contrast, an
6 increase in amplitudes of the 2-4 yr oscillation is recorded over the same interval.

7 In contrast to high-frequency oscillations associated with ENSO, decadal to
8 multi-decadal modes of variability in the lake-area reconstruction appear to be
9 associated with changes in the PDO index. Previous studies have documented a long-
10 lived pattern of Pacific climate variability in palaeoclimate records from Western North
11 America and the Central Andes (D'Arrigo et al., 2001; Villalba et al., 2001, Masiokas et
12 al., 2010). The persistent trend to arid conditions recorded from mid 1970s to present in
13 the Vilama-Coruto lake-area reconstruction was coincident with a documented shift
14 from the negative to the positive phase of the PDO in 1976 (Mantua et al., 1997). This
15 PDO positive phase, which lasted to the end of the 20th century, coincided with a
16 marked increases in global mean surface temperatures (Trenberth et al., 2013, 2014) and
17 has significantly impacted climate and ecosystems across the western coasts of North
18 and South America (D'Arrigo et al., 2001; Villalba et al., 2001; Masiokas et al., 2010).
19 During this interval of positive PDO phase, an increased in ENSO was recorded in both
20 instrumental and reconstructed data (Li et al., 2013).

21 Interactions between PDO and ENSO have been documented for both North and
22 South America (Andreoli and Kayano, 2005). More intense El Niño (La Niña) rainfall
23 anomalies occur when both ENSO and PDO share positive (negative) phases. In
24 contrast weaken anomalies occur when they are in opposite phases. It has been
25 suggested that the PDO switched to a negative phase (La Niña-like pattern) around

1 1999, coincident with a pause in the sustained increase in global mean surface
2 temperatures (Trenberth et al., 2014). However, we note that the downward trend in
3 Coruto-Vilama lake areas has persisted since 1999. This suggests that, in addition to
4 PDO, different forcings may have contributed to the persistence of dry conditions in our
5 study region for the past decades. However, we note that the downward trend in Coruto-
6 Vilama lake areas have persisted since the year 1999. This suggests that, in addition to
7 PDO, different forcings may have contributed to the persistence of dry conditions in our
8 study region for the past decade. A dominance of the upper-tropospheric Westerlies
9 during second half of 20th and early 21st centuries, leading to decreased precipitation by
10 blocking the moisture easterly flow from the Amazon basin (Neukom et al. 2015),
11 together with a continuous warming trend of the upper troposphere over the Central
12 Andes (Vuille et al. 2015) may thus explain much of the recent dramatic lake area
13 decrease. Furthermore, an anomalous ENSO activity in the late 20th century (Li et al.
14 2013) would even strengthen the negative relationship between Westerlies and
15 precipitation, with the consequent intensification of drier conditions.

16 In the present study, common cycles and periodicities were identified between
17 lake area and ENSO (at inter-annual/decadal oscillation modes) and PDO (at multi-
18 decadal oscillation mode). These results allow to identified ENSO and PDO frequency,
19 amplitudes and associated teleconnection patterns. These findings were consistent to
20 those recorded for the ENSO related precipitation reconstruction from the Altiplano
21 (Morales et al. 2012), while no previous analyzes were made about the PDO influence
22 on the reconstructed precipitation. Since most palaeo-reconstructions of ENSO devoid
23 of tropical/subtropical records (Li et al., 2013), our lake-area reconstruction provides
24 valuable information about ENSO and also PDO evolution during the past 600 years
25 and should be considered as a key input in further high-resolution reconstructions of

1 these hemispheric forcing. Given that ENSO and PDO variability provided major
2 factors affecting hydrological patterns in the high-elevation lake systems from NWA-
3 SWB, the longer time frames provide here may also complement those studies
4 discussing the regional manifestations and long-term variability of these large-scale
5 ocean-atmosphere features (Stahle et al., 1998; Evans et al., 2001; D'Arrigo et al., 2001;
6 Villalba et al., 2001; Cobb et al., 2003; D'Arrigo et al., 2005; Gergis and Fowler, 2009;
7 Li et al., 2013).

8 At local and regional scales, the lake-area reconstruction should be used as input
9 in water system models to refine the reliability of water-balance assessments and
10 ecosystem productivity in this arid region with persistent hydrological deficits. This
11 study is especially useful in understanding the temporal variability of water availability
12 in the Vilama-Coruto lake system, where the water resource is fundamental for
13 biodiversity conservation and socio-economic activities such as grazing, mining and
14 tourism. Global and regional climate models for the Altiplano and Puna (Urrutia and
15 Vuille, 2009; Minvielle and Garreaud, 2011; IPCC, 2013; Neukom et al. 2015) project a
16 marked reduction in precipitation to the end of the 21st century, exacerbating presently
17 dry conditions. If the negative trend in precipitation documented in recent decades
18 continues, it is likely that the ranges of natural variability of these high altitude
19 ecosystems are exceeded. Therefore, quantitative information on past hydrological
20 changes is required to provide reliable information to better adaptation to future drier
21 conditions in the Altiplano. Further improvements in the existing lake area and
22 precipitation reconstructions could be achieved by expanding the temporal and spatial
23 coverage of the tree-ring network in the Puna and Altiplano from Argentina, Bolivia,
24 Chile and Peru. Similar reconstruction exercises could be performed for neighboring
25 high altitude lake systems from the Altiplano and northernmost tropical Andes.

1
2 *Acknowledgements.* This work was carried out with the aid of grants from the
3 Argentinean Research Council (PIP 0968), National Geographic (NGS 8681-09), the
4 Argentinean Agency for Promotion of Science (PICT 07-246; PICT 2012-1565; PICT
5 2013-1880), National University of Tucumán (PIUNT-G516), and the Inter-American
6 Institute for Global Change Research (IAI) CRN II # 2047 supported by the US
7 National Science Foundation (GEO-0452325), We are grateful to Alvaro Gonzalez for
8 his help on drought events recurrence analysis. To Alberto Ripalta for his help during
9 fieldwork, and Juan Carlos Gómez for tree-ring samples preparation. We acknowledge
10 David Meko for providing the “leave one out” routine on Matlab, and the national water
11 agencies DGA-Chile and SENHAMI-Bolivia for providing the instrumental
12 precipitation records. The N3.4-SSTs data were obtained from UCAR-NCAR
13 (www.cgd.ucar.edu/cas/catalog/ climind/ TNI_N34/index.html#Sec5) website. Satellite
14 Image data were downloaded from U.S. Geological Survey, Department of the
15 Interior/USGS. The manuscript was greatly benefitted from comments and corrections
16 by J. Martinez (editor), B. Luckman and D. Urrego (reviewers).

17

18

19 **Reference**

20

21 Aceituno, P. and Montecinos, A.: Circulation anomalies associated with dry and wet
22 periods in the South American Altiplano, in: Preprints 4th Int. Conf. on Southern
23 Hemisphere Meteorology and Oceanography, Hobart, Australia, 29 March–2 April,
24 330–331, 1993.

- 1 Andreoli, R. V. and Kayano, M. T.: ENSO-related rainfall anomalies in South America
2 and associated circulation features during warm and cold Pacific decadal oscillation
3 regimes, *Int. J. Climatol.*, 25, 2017–2030, 2005.
- 4 Bird, W. B., Abbott, M. B., Vuille M., Rodbell, D. T., Stansella, N. D., and Rosenmeier,
5 M. F.: A 2,300-year-long annually resolved record of the South American summer
6 monsoon from the Peruvian Andes, *Proc. Natl. Acad. Sci. USA*, 108, 8583–8588,
7 doi:10.1073/pnas.1003719108, 2011.
- 8 Boucher, E., Guiot, J., and Chapron, E.: A millennial multi-proxy reconstruction of
9 summer PDSI for Southern South America, *Clim. Past*, 7, 957–974, doi:10.5194/cp-
10 7-957-2011, 2011.
- 11 Bradley, R. S., Vuille, M., Hardy, D., and Thompson, L. G.: Low latitude ice cores
12 record Pacific sea surface temperatures. *Geophys. Res. Lett.*, 30, 1174,
13 doi:10.1029/2002GL016546, 2003.
- 14 Briffa, K. R.: Interpreting high-resolution proxy climate data. The example of
15 dendroclimatology, in: *Analysis of climate variability, applications of statistical
16 techniques*, edited by: von Storch, H. and Navarra, A., Springer Verlag, Berlin, 77-
17 94, 1995.
- 18 Cabrera, A.: *Regiones fitogeográficas argentinas*. Enciclopedia Argentina de
19 *Agronomía y Jardinería*, editorial Acme, Buenos Aires, 1976.
- 20 Carilla, J., Grau, H. R., Paolini, L., and Morales, M. S.: Lake fluctuations, plant
21 productivity, and long-term variability in high-elevation tropical andean ecosystems,
22 *Arct. Antarct. Alp. Res.*, 45: 179–189. 2013.
- 23 Christie, D. A., Lara, A., Barichivich, J., Villalba, R., Morales, M. S., and Cuq, E.: El
24 Niño-Southern Oscillation signal in the world's high-elevation tree-ring

- 1 chronologies from the Altiplano, Central Andes, *Palaeogeogr. Palaeocl. Paleoecol.*,
2 281, 309–319, 2009.
- 3 Cobb, K. M., Charles, C. D., Cheng, H., and Edwards, R. L.: El Niño/Southern
4 Oscillation and tropical Pacific climate during the last millennium, *Nature*, 424,
5 271–276, 2003.
- 6 Coira, B., Caffè, P., Díaz, A., and Ramírez, A.: Sistema de Calderas Anidadas del
7 Cenozoico Superior: Vilama-Coruto, Puna (Argentina) – Altiplano (Bolivia).
8 *Memorias del XII Congreso Geológico de Bolivia*, pp. 991-1002, Tarija, 1996.
- 9 Cowling, A., Hall, P., and Phillips, M. J.: Bootstrap confidence regions for the intensity
10 of a Poisson point process, *J. Am. Stat. Assoc.*, 91: 1516–1524, 1996.
- 11 D'Arrigo, R. and Villalba, R., Wiles, G.: Tree-ring estimates of Pacific decadal climate
12 variability. *Climate Dynamics*, 18, 219-224, 2001.
- 13 D'Arrigo, R., Wilson, R., Deser, C., Wiles, G., Cook, E., Villalba, R., Tudhope, S., Cole,
14 J., and Linsley, B.: Tropical-north Pacific climate linkages over the past four
15 centuries. *J. Clim.*, 18, 5253–5265, 2005.
- 16 Draper, N. R. and Smith, H.: *Applied Regression Analysis*, John Wiley & Son, New
17 York, 1981.
- 18 Evans, M. N., Cane, M. A., Schrag, D. P., Kaplan, A., Linsley, B. K., Villalba, R., and
19 Wellington, G. M.: Support for tropically-driven Pacific decadal variability based on
20 paleoproxy evidence, *Geophys. Res. Lett.*, 28, 3869–3692, 2001.
- 21 Francou, B., Vuille, M., Wagnon, P., Mendoza, J., and Sicart, J. E.: Tropical climate
22 change recorded by a glacier in the central Andes during the last decades of the 20th
23 century: Chacaltaya, Bolivia, 16°S, *J. Geophys. Res.*, 108, 4154,
24 doi:10.1029/2002JD002959, 2003.
- 25 Fritts, H. C.: *Tree rings and climate*, Academic Press, London, 1976.

- 1 Garreaud, R. and Aceituno, P.: Interannual rainfall variability over the South American
2 Altiplano, *J. Climate*, 14, 2779–2789, 2001.
- 3 Garreaud, R., Vuille, M., and Clement, C.: The climate of the Altiplano: Observed
4 current conditions and mechanisms of past changes, *Palaeogeogr. Palaeoclimatol.*, 194, 5–
5 22, 2003.
- 6 Garreaud, R. D., Vuille, M., Compagnucci, R., and Marengo, J.: Present day South
7 American climate, *Palaeogeogr. Palaeoclimatol.*, 281, 180–195, 2009.
- 8 Gergis, J. L. and Fowler, A. M.: A history of ENSO events since A.D. 1525:
9 implications for future climate change, *Climatic Change*, 92, 343–387, 2009.
- 10 Gioda, A., Prieto, M. R.: Histoire des sécheresses andines: Potosi El Niño et le Petit
11 Age Glaciaire, *La Météorologie*, 8, 33–42, 1999.
- 12 Girardin, M. P., Bergeron, Y., Tardif, J. C., Flannigan, M. D., Gauthier, S., and
13 Mudelsee, M.: A 229-year dendroclimatic-inferred record of forest fire activity for
14 the Boreal Shield of Canada, *Int J Wildland Fire*, 15, 375–388, 2006.
- 15 Holmes, R. L.: Computer-assisted quality control in tree-ring dating and measurements,
16 *Tree Ring Bull.*, 43, 69–75, 1983.
- 17 Hsu, C. W., Chang, C. C., and Lin, C. J.: A practical guide to support vector
18 classification, National Taiwan University
19 http://ntu.csie.org/_cjlin/papers/guide/guide.pdf, 2007.
- 20 IPCC: Long-term Climate Change: Projections, Commitments and Irreversibility, in:
21 *Climate Change 2013: The Physical Science Basis*, Contribution of Working Group
22 I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change,
23 edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K.,
24 Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M., Cambridge
25 University Press, Cambridge, United Kingdom and New York, 2013.

- 1 Jenkins, G. M. and Watts, D. G.: Spectral analysis and its applications, Holden-Day,
2 San Francisco, 1968.
- 3 Jones, P. D., New, M., Parker, D. E., Martin, S. and Rigor, I. G.: Surface air
4 temperature and its changes over the past 150 years, *Rev. Geophys.*, 37, 173-199,
5 1999.
- 6 Kistler, R., Kalnay, E., Collins, W., Saha, S., White, G., Woollen, J., Chelliah, M.,
7 Ebisuzaki, W., Kanamitsu, M., Kousky, V., Van den Dool, H., Jenne, R., and
8 Fiorino, M.: The NCEP-NCAR 50-year reanalysis: monthly means CD-ROM and
9 documentation, *B. Am. Meteorol. Soc.*, 82, 247–267, 2001.
- 10 Lenters, J. D. and Cook, K. H.: On the origin of the Bolivian High and related
11 circulation features of the South American climate, *J. Atmos. Sci.*, 54, 656–677,
12 1997.
- 13 Li, J., Xie, S.-P., Cook, E. R., Morales, M. S., Christie, D., Chen, F., D'Arrigo, R.,
14 Johnson, N. C., Fowler, A. M., Gou, X., and Fang, K.: El Niño modulations during
15 the past seven centuries. *Nature Clim. Change*, 3, 822-826, 2013.
- 16 Mantua, N. J., Hare S.R., Zhang Y., Wallace J. M., and Francis R. C. A.: Pacific
17 interdecadal climate oscillation with impacts on salmon production. *Bull. Amer.*
18 *Meteor. Soc.*, 78, 1069–1079. 1997.
- 19 Masiokas, M., Villalba, R., Luckman, B., and Mauget, S.: Intra- to Multidecadal
20 Variations of Snowpack and Streamflow Records in the Andes of Chile and
21 Argentina between 30° and 37°S. *J. Hydrometeorol.*, 11, 822-831, 2010.
- 22 Mason, I. M., Guzkowska M. A. J., Rapley C. G., and Street-Perrot F. A.: The Response
23 of Lake Levels and Areas, *Climatic Change*, 27, 161-197, 1994.

- 1 Meko, D. M. and Baisan, C. H.: Pilot study of latewood-width of conifers as an
2 indicator of variability of summer rainfall in the North American monsoon region,
3 *Int. J. Climatol.*, 21: 697-708, 2001.
- 4 Melvin, T. M.: Historical growth rates and changing climatic sensitivity of boreal
5 conifers, Ph.D. thesis, University of East Anglia, Norwich, United Kingdom, 2004.
- 6 Melvin, T. and Briffa K.: A 'signal-free' approach to dendroclimatic standardization,
7 *Dendrochronologia*, 26, 71–86, 2008.
- 8 Michaelsen, J.: Cross-validation in statistical climate forecast models, *J. Clim. Appl.*
9 *Meteorol.*, 26, 1589–1600, 1987.
- 10 Minvielle, M. and Garreaud, R.: Projecting rainfall changes over the South American
11 Altiplano, *J. Climate*, 24, 4577–4583, 2011.
- 12 Mitchell, Jr., J. M., Dzerdseevskii, B., Flohn, H., Hofmeyr, W. L., Lamb, H. H., Rao,
13 N., and Wallen, C. C.: *Climatic change*. World Meteorological Organization,
14 Geneva, 1966.
- 15 Morales, M. S., Villalba, R., Grau, H. R., and Paolini, L.: Rainfall controlled tree
16 growth in high elevation subtropical treelines, *Ecology*, 85, 3080–3089, 2004.
- 17 Morales, M. S., Christie, D. A., Villalba, R., Argollo, J., Pacajes, J., Silva, J. S.,
18 Alvarez, C. A., Llancabure, J. C., and Soliz, C.: Precipitation changes in the South
19 American Altiplano since 1300 AD reconstructed by tree-rings, *Clim Past*, 8: 653–
20 666, 2012.
- 21 Mudelsee, M., Börngen, M., Tetzlaff, G., and Grünewald, U.: Extreme floods in central
22 Europe over the past 500 years: Role of cyclone pathway “Zugstrasse Vb”, *J.*
23 *Geophys. Res.* 109:D23101. doi: 10.1029/2004JD005034, 2004.
- 24 Mujica, M. I., Latorre, C., Maldonado, A., González-Silvestre, L., Pinto, R., de Pol-
25 Holz, R., and Santoro, C. M.: Late Quaternary climate change, relict populations

- 1 and present-day refugia in the northern Atacama Desert: A case study from
2 Quebrada La Higuera (18° S), *J. Biogeogr.* doi:10.1111/jbi.12383, 2014.
- 3 Nielsen, A. E.: Ocupaciones prehispanicas de la etapa Agropastoril en la laguna de
4 Vilama (Jujuy, Argentina), *Cuadernos FHyCS-UNJu*, 20: 81-108, 2003.
- 5 Neukom, R., Rohrer, M., Calanca, P., Salzmann, N., Huggel, C., Acuña, D., Christie, D.
6 A., and Morales, M. S.: Facing unprecedented drying of the Central Andes?
7 Precipitation variability over the period AD 1000-2100. *Environ. Res. Lett.* 10,
8 084017, doi:10.1088/1748-9326/10/8/084017, 2015.
- 9 Rabatel, A., Francou, B., Soruco, A., Gomez, J., Cáceres, B., Ceballos, J. L., Basantes,
10 R., Vuille, M., Sicart, J.E., Huggel, C., Scheel, M., Lejuene, Y., Arnaud, Y., Collet,
11 M., Condom, T., Consoli, G., Favier, V., Jomelli, V., Galarraga, R., Ginot, P.,
12 Maisincho, L., Mendoza, J., Ménégos, M., Ramírez, M., Ribstein, P., Suarez, W.,
13 Willacis, M., Wagnon, P.: Current state of glaciers in the tropical Andes: A multi-
14 century perspective on glacial evolution and climate change, *Cryosphere*, 7, 81–102,
15 2013.
- 16 Ramirez, E., Francou, B., Ribstein, P., Descloitres, M., Guérin, R., Mendoza, J.,
17 Gallaire, R., Pouyaud, B., and Jordan, E.: Small glaciers disappearing in the tropical
18 Andes. A case study in Bolivia: Glacier Chacaltaya (16° S), *J. Glaciol.*, 47, 187–194,
19 2001.
- 20 Rodionov, S. N.: A sequential algorithm for testing climate regime shifts, *Geophys.*
21 *Res. Lett.*, 31, L09204, doi:10.1029/2004GL019448, 2004.
- 22 Schitteck, K., Kock, S. T., Lücke, a., Ohlendorf, C., Kulemeyer, J. J., Lupo, L. C., and
23 Schäbitz, F.: Environmental and climatic history in the NW Argentine Andes (24°
24 S) over the last 2100 years inferred from a high-altitude peatland record. *Clim. Past*
25 *Discuss.* 11, 2037–2076, 2015.

- 1 Schulman, E.: Dendroclimatic changes in semiarid America, Tucson, University of
2 Arizona Press, 1956.
- 3 Squeo, F. A., Warner, B. G., Aravena, R., and Spinoza, D.: Bofedales: high altitude
4 peatlands of central Andes, *Rev. Chil. Hist. Nat.*, 79: 245–255, 2006.
- 5 Solíz, C., Villalba, R., Argollo, J., Morales, M. S., Christie, D. A., Moya, J., and
6 Pacajes, J.: Spatio-temporal variations in *Polylepis tarapacana* radial growth across
7 the Bolivian Altiplano during the 20th century, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*,
8 281, 296–330, 2009.
- 9 Speer, J. H., Swetnam, T. W., Wickman, B. E., Youngblood, A.: Changes in pandora
10 moth outbreak dynamics during the past 622 years, *Ecology*, 82, 679–697, 2001.
- 11 Stahle, D. W., D'Arrigo, R. D., Krusic, P. J., Cleaveland, M. K., Cook, E. R., Allan, R.
12 J., Cole, J. E., Dunbar, R. B., Therrell, M. D., Gay, D. A., Moore, M. D., Stokes, M.
13 A., Burns, B. T., Villanueva-Diaz, J., and Thompson, L. G.: Experimental
14 dendroclimatic reconstruction of the Southern Oscillation. *B. Am. Meteorol. Soc.*,
15 79, 2137–2152, 1998.
- 16 Štěpánek, P.: AnClim - software for time series analysis, Dept. of Geography, Fac. of
17 Natural Sciences, Munich, <http://www.climahom.eu/AnClim.html>, 2008.
- 18 Stokes, M. A. and Smiley, T. L.: An introduction to tree-ring dating, University of
19 Chicago Press, Chicago, 1968.
- 20 Sutton, R. T. and Hodson, D. L. R.: Influence of the ocean on North Atlantic climate
21 variability 1871–1999, *J. Climate*, 16, 3296–3313, 2003.
- 22 Thompson, L. G., Mosley-Thompson, E., Brecher, H., Davis, M., León, B., Les, D.,
23 Lin, P. N., Masiotta, T., and Mountain, K.: Abrupt tropical climate change: Past
24 and present, *P. Natl. Acad. Sci.*, 103, 10536–10543, 2006.

- 1 Torrence, C. and Webster, P.: Interdecadal changes in the ENSO Monsoon system, *J.*
2 *Climate*, 12, 2679–2690, 1999.
- 3 Trenberth, K. E. and Fasullo, J. T.: An apparent hiatus in global warming?, *Earth's*
4 *Future*, 1: 19-32, 2013.
- 5 Trenberth K. E., Fasullo, J. T., Branstator G., and Phillips A. S.: Seasonal aspects of the
6 recent pause in surface warming, *Nature Clim. Change*, 4: 911-916, 2014.
- 7 Urrutia, R. and Vuille, M.: Climate change projections for the tropical Andes using a
8 regional climate model: Temperature and precipitation simulations for the end of the
9 21st century, *J. Geophys. Res.*, 114, D02108, doi:10.1029/2008JD011021, 2009.
- 10 Vautard, R. and Ghil, M.: Singular spectrum analysis in nonlinear dynamics, with
11 applications to paleoclimatic time series, *Physica D.*, 35, 395–424, 1989.
- 12 Vautard, R.: Patterns in Time: SSA and MSSA, in: *Analysis of climate variability.*
13 *Applications of statistical techniques*, edited by: von Storch, H., Navarra, A.,
14 Springer, Berlin, 259–287, 1995.
- 15 Villalba, R., D'Arrigo, R. D., Cook, E. R., Wiles, G., and Jacoby, G. C.: Decadal-scale
16 climatic variability along the extra-tropical western coast of the Americas over past
17 centuries inferred from tree-ring records, in: *Interhemispheric Climate Linkages*,
18 edited by: Markgraf V., Cambridge University Press, Cambridge, UK, 2001.
- 19 Vuille, M. and Bradley, R.: Mean annual temperature trends and their vertical structure
20 in the tropical Andes, *Geophys. Res. Lett.*, 27, 3885–3888, 2000.
- 21 Vuille, M., Bradley, R., Werner, M., and Keimig, F.: 20th century climate change in the
22 tropical Andes: observations and model results, *Climatic Change*, 59, 75–99, 2003.
- 23 Vuille, M. and Keimig, F.: Interannual variability of summertime convective cloudiness
24 and precipitation in the central Andes derived from ISCCP-B3 data, *J. Climate*, 17,
25 3334–3348, 2004.

- 1 Vuille, M., Bradley, R. S., and Keimig, F.: Interannual climate variability in the Central
2 Andes and its relation to tropical Pacific and Atlantic forcing, *J. Geophys. Res.*, 105,
3 12447–12460, 2000.
- 4 Vuille, M., Francou, B., Wagnon, P., Juen, I., Kaser, G., Mark, B., and Bradley, R.:
5 Climate change and tropical Andean glaciers: Past, present and future, *Earth-Sci*
6 *Rev.*, 89, 79–96, 2008.
- 7 Vuille, M., Franquist, E., Garreaud, R., Sven, W., Casimiro, L., and Cáceres, B.: Impact
8 of the global warming hiatus on Andean temperature, *J. Geophys. Res.*, 120,
9 doi:10.1002/2015JD023126, 2015.
- 10 Weisberg, S.: *Applied Linear Regression*, John Wiley & Son, New York, 1985.
- 11 Wigley, T. M. L., Briffa, K. R., and Jones, P. D.: On the average value of correlated
12 time series, with applications in dendroclimatology and hydrometeorology, *J. Clim.*
13 *Appl. Meteorol.*, 23, 201–213, 1984.
- 14 Wilkes, D. S.: *Statistical methods in the atmospheric sciences*, Academic Press, Boston,
15 1995.

1 **Tables**

2

3 **Table 1.** Geographical location and statistics of the *Polylepis tarapacana* ring width
 4 chronologies. Mean RBar and mean EPS were averaged over chronology periods
 5 including > 10 tree-ring series.

Chronology	Location (°S/°W)	Period (years)	No. of series	Mean Inter- correlation	Mean sensitivity	Mean RBar	Mean EPS	<i>r</i> with region al lakes
Uturunco	22 16/67 10	1242- 2006	81	0.59	0.28	0.39	0.95	0.59
Granadas	22 35/66 33	1620- 2008	74	0.63	0.31	0.40	0.95	0.58
Regional		1407- 2008	155	0.58	0.30	0.37	0.96	0.69

6

7

8

1

2

3 **Table 2.** Lowest and highest non-overlapping averages of the reconstructed (1407–
 4 2007) lake area fluctuations for the Vilama-Coruto lake system. Annual lake area
 5 expressed as Z-score of the 1975–2007 lake area mean. Ranks 1–5 correspond to the
 6 five most extreme reconstructed lake area years or set of consecutive years. Rank 1
 7 represents the most important minimum (maximum) lake area record.

8

Rank	Minimum Lake Area			
	1 yr	5 yr	25 yr	50 yr
1	-0.803 (2007)	-0.425 (2003-2007)	0.040 (1983-2007)	0.398 (1958-2007)
2	-0.727 (1998)	-0.225 (1618-1622)	0.661 (1608-1632)	0.775 (1508-1557)
3	-0.661 (1505)	-0.148 (1998-2002)	0.741 (1533-1557)	0.915 (1608-1657)
4	-0.610 (1621)	-0.091 (1753-1757)	0.754 (1783-1807)	0.924 (1758-1807)
5	-0.553 (1983)	-0.127 (1938-1942)	0.757 (1958-1982)	0.926 (1558-1607)

Rank	Maximum Lake Area			
	1 yr	5 yr	25 yr	50 yr
1	3.277 (1456)	2.352 (1838-1842)	1.727 (1833-1857)	1.399 (1808-1857)
2	2.941 (1734)	2.216 (1733-1737)	1.358 (1858-1882)	1.345 (1858-1908)
3	2.844 (1840)	1.962 (1743-1747)	1.335 (1883-1907)	1.196 (1458-1507)
4	2.826 (1744)	1.936 (1453-1457)	1.295 (1458-1482)	1.188 (1658-1708)
5	2.802 (1431)	1.912 (1843-1847)	1.281 (1908-1933)	1.164 (1908-1957)

9

10

1 **Figure captions**

2

3 **Fig. 2. (a)** Observed (green line) and tree-ring predicted (blue line) annual lake area
4 (January–December) variations in the NWA-SWB. Annual lake area expressed as a Z-
5 score of the satellite-derived 1975–2007 lake area mean. Calibration and verification
6 statistics: the Pearson correlation coefficient (r) between observed and reconstructed
7 values, the coefficient of determination adjusted for the degrees of freedom in the model
8 (R^2 adj) over the calibration period, the F-value of regression (the ratio between the
9 explained and the unexplained variability in the model), and the reduction of error (RE).
10 **(b)** The linear trend of regression residuals (slope: blue line) and the Durbin-Watson (D-
11 W) statistics used to test for first-order autocorrelation of the regression residuals are
12 shown.

13

14 **Fig. 3.** Annual (January-December) Vilama-Coruto lake area reconstruction for the
15 period AD 1407–2007. Annual lake area expressed as Z-score of the 1975–2007 lake
16 area mean area. The 35-yr smoothing-spline curve highlights the multi-decadal
17 variability. Significant (95% c.l.) regime shifts (blue arrows) and the mean of periods
18 (orange horizontal line) detected by the Rodionov (2004) method (window length=25
19 yr). Dates of the regime shifts are shown in the figure. Uncertainties of the
20 reconstruction are shown by the light green band (± 1 RMSE).

21

22 **Fig. 4.** Changing probability of extreme low values ($< 20^{\text{th}}$ percentile; blue line) of the
23 Vilama-Coruto lake area reconstruction from 1407 to 2007. A Kernel smoothing

1 method was used with a bandwidth of 50 years. The shaded represent 95% confidence
2 interval based on 1000 bootstrap simulations.

3

4 **Fig. 5.** Spatial correlation patterns over the interval 1948–2006 between the 2.5 x 2.5
5 gridded monthly averaged January-December sea surface temperatures (SSTs) and the
6 reconstructed January-December lake area for the Vilama-Coruto region. Spatial
7 correlations were obtained from the National Oceanic and Atmospheric Administration
8 website (<http://www.esrl.noaa.gov/psd/data/correlation/>). The reconstructed lake area
9 region is indicated by the red square.

10

11 **Fig. 6.** Blackman-Tukey power spectra analysis of the (a) reconstructed lake area
12 (1407–2007) and (b) the sea surface temperatures from the N3.4 Pacific sector (N3.4
13 SST; 1872-2007). (c) Coherency spectrum of N3.4 SST and reconstructed lake area in
14 the Vilama-Coruto region, estimated over the common period 1872–2007. Short dashed
15 and dotted lines represent the 0.05 probability levels and the red noise band,
16 respectively. Dominant periodicities are indicated in each panels.

17

18 **Fig. 7.** (a) Multi-decadal, (b) decadal and (c,d) inter-annual main oscillation modes of
19 variability extracted by Singular Spectral Analysis (SSA) of reconstructed lake area
20 from 1407 to 2007 interval. The frequencies for each SSA are indicated in years with
21 the corresponding explained variance in percentage (%).

22


23 **Fig. 8.** Comparisons between the waveforms of the lake area reconstruction (red line),
24 PDO (blue line) and the N3.4 SST (green line) extracted by Singular Spectrum Analysis

1 (SSA) over the (a) 1900-2007 (PDO) and (b,c) 1872-2007 (N3.4 SST) intervals. The
 2 periodicities and percentage of variance explained by each frequency are indicated in
 3 parentheses. The correlation coefficients between the two series are shown at the right
 4 corner. The PDO and N3.4 SST waveforms are shown inverse to facilitate the
 5 comparison between records (right axis).

6

7 **Figures**

8



9

10 **(Fig. 1)**

1

2

3

4


5

6


7

8

9

**(Fig. 2)**


1



2

3 (Fig. 3)


1



2

3 (Fig. 4)


1



3 (Fig. 5)

4


1



2

3 (Fig. 6)


1



2

3 (Fig. 7)

1



2

3 (Fig. 8)

4