

This discussion paper is/has been under review for the journal Climate of the Past (CP).  
Please refer to the corresponding final paper in CP if available.

# A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney<sup>1</sup>, R. Jones<sup>2</sup>, C. Fogwill<sup>1</sup>, J. Hatton<sup>2</sup>, A. N. Williams<sup>3,4</sup>, A. Hogg<sup>5</sup>,  
Z. Thomas<sup>1</sup>, J. Palmer<sup>1</sup>, and S. Mooney<sup>1</sup>

<sup>1</sup>School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, Australia

<sup>2</sup>Department of Geography, Exeter University, Devon, EX4 4RJ, UK

<sup>3</sup>Fenner School of Environment and Society, The Australian National University, Canberra, ACT 0200, Australia

<sup>4</sup>Archaeological and Heritage Management Solutions Pty Ltd, 2/729 Elizabeth Street, Waterloo, NSW 2017, Australia

<sup>5</sup>Waikato Radiocarbon Laboratory, University of Waikato, Private Bag 3105, Hamilton, New Zealand

Received: 24 April 2015 – Accepted: 8 May 2015 – Published: 3 June 2015

Correspondence to: C. Turney (c.turney@unsw.edu.au)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**A 250 year periodicity  
in Southern  
Hemisphere westerly  
winds over the last  
2600 years**

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

Southern Hemisphere westerly airflow has a significant influence on the ocean–atmosphere system of the mid- to high-latitudes with potentially global climate implications. Unfortunately historic observations only extend back to the late nineteenth century, limiting our understanding of multi-decadal to centennial change. Here we present a highly resolved (30 yr) record of past westerly air strength from a Falkland Islands peat sequence spanning the last 2600 years. Situated under the core latitude of Southern Hemisphere westerly airflow, we identify highly variable changes in exotic pollen derived from South America which can be used to inform on past westerly air strength and location. The results indicate enhanced airflow over the Falklands between 2000 and 1000 cal. yr BP, and associated with increased burning, most probably as a result of higher temperatures and/or reduced precipitation, comparable to records in South America. Spectral analysis of the charcoal record identifies a 250 year periodicity within the data, suggesting solar variability has a modulating influence on Southern Hemisphere westerly airflow with potentially important implications for understanding global climate change through the late Holocene.

## 1 Introduction

A major limitation for quantifying the magnitude and impact of change across the Southern Ocean is the relatively short duration or low resolution of ocean–atmosphere records. This is particularly significant with regards the Southern Hemisphere westerly storm belt, which since the mid-1970s, has undergone a significant intensification and southward shift (Gillett et al., 2008; Messié and Chavez, 2011). One measure of this change in atmospheric circulation is the Southern Annular Mode (SAM), described as the pressure difference between Antarctica (65° S) and the latitude band at around 40° S (Karpechko et al., 2009; Marshall, 2003). Since the mid-1970s SAM appears to have undergone a positive shift in the troposphere, which has been associated with

CPD

11, 2159–2180, 2015

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

hemispheric-wide changes in the atmosphere–ocean-ice domains, including the delivery of precipitation and significant surface and subsurface warming (Cook et al., 2010; Delworth and Zeng, 2014; Domack et al., 2005; Gille, 2008, 2014; Thompson et al., 2011). This trend is projected to continue during the 21st century as a result of both ongoing greenhouse gas emissions and a persistence of the Antarctic ozone hole (Liu and Curry, 2010; Thompson et al., 2011; Yin, 2005), potentially resulting in reduced Southern Ocean uptake of anthropogenic CO<sub>2</sub> (Ito et al., 2010; Le Quéré et al., 2009; Lenton et al., 2013; Marshall, 2003; Marshall and Speer, 2012).

While no observational records for SAM extend beyond the late nineteenth century (Fogt et al., 2009; Marshall, 2003; Visbeck, 2009), proxy records of past westerly airflow have been generated on annual to centennial timescales through the Holocene (Abram et al., 2014; Björck et al., 2012; Lamy et al., 2010; Strother et al., 2015; Villalba et al., 2012). Crucially the association between proxies and changes in westerly wind strength and/or latitude is often implied but few provide a direct measure of past airflow or directly test their interpretation through time. One possibility is the identification of exotic airborne pollen preserved in sedimentary sequences. Ideally, the peat or lake record should be close enough to the source vegetation to have a relatively high input of pollen but not so close that the influx is constant over time. Whilst numerous studies have been undertaken in the Arctic (Fredskild, 1984; Jessen et al., 2011) and the high-latitudes of the Indian and Pacific oceans (McGlone et al., 2000; Scott and van Zinderen Barker, 1985), few have been reported from the south Atlantic. Recent work on a lake core taken from Annekov Island, South Georgia (Strother et al., 2015) demonstrates the considerable potential of this approach but the relatively large distance from the nearest source in South America (Fig. 1) (approximately 2100 km) limits the delivery of pollen.

Here we report a new high-resolution record of westerly airflow over the past 2600 years from the Falkland Islands. The Falkland Islands (52° S) lie west of Annekov Island and 500 to 730 km east of Argentina within the main latitudinal belt of Southern Hemisphere westerly airflow. The close proximity to South America means that these



## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Macphail and Cantrill, 2006) and the pollen type slide collection at Exeter University. Past fire activity was assessed using micro-charcoal counts of fragments identified on the pollen slides e.g. (Whitlock and Larsen, 2001). Counts were undertaken at each level until a fixed total of 50 lycopodium spores was counted and the total expressed as a concentration (fragments per  $\text{cm}^3$ ). The results were sub-divided into two size classes: < 50 and 50–150  $\mu\text{m}$ .

Terrestrial plant macrofossils (fruits and leaves) were extracted from the peat sequence and given an acid-base-acid (ABA) pretreatment and then combusted and graphitized in the University of Waikato AMS laboratory, with  $^{14}\text{C}/^{12}\text{C}$  measurement by the University of California at Irvine (UCI) on a NEC compact (1.5SDH) AMS system. The pretreated samples were converted to  $\text{CO}_2$  by combustion in sealed pre-baked quartz tubes, containing Cu and Ag wire. The  $\text{CO}_2$  was then converted to graphite using  $\text{H}_2$  and an Fe catalyst, and loaded into aluminum target holders for measurement at UCI. This was supplemented by  $^{137}\text{Cs}$  measurements down the profile to detect the onset of nuclear tests.  $^{137}\text{Cs}$  analysis was undertaken following standard techniques (Appleby, 2001) with measurements made using an ORTEC high-resolution, low-background coaxial germanium detectors. The first detectable measurements were obtained between 8.5 and 9.5 cm and assigned an age of CE 1963, the time of peak radionuclide fallout (Hancock et al., 2011).

The radiocarbon and  $^{137}\text{Cs}$  ages were used to develop an age model using a P<sub>sequence</sub> deposition model in OxCal 4.2 (Ramsey, 2008). The  $^{14}\text{C}$  ages were calibrated against the Southern Hemisphere calibration (SHCal13) dataset (Hogg et al., 2013). Using Bayes theorem, the algorithms employed sample possible solutions with a probability that is the product of the prior and likelihood probabilities. Taking into account the deposition model and the actual age measurements, the posterior probability densities quantify the most likeliest age distributions. All ages returned an *agreement index* of > 60 % indicating they had a large overlap with the likelihood probability distribution ( $A_{\text{model}} = 67.2$ ;  $A_{\text{overall}} = 66.0$ ). Calibrated ages are reported here as thousands

of calendar years BP or cal. BP (Table 1 and Fig. 2). The pollen sequence reported here spans the last 2600 yr with an average 30 yr resolution.

To investigate the periodicities preserved in the palaeoenvironmental proxies utilised herein, we undertook Multi-Taper Method (MTM) analysis using a narrowband signal and red noise significance (with a resolution of 2 and number of tapers, 3) (Thomson, 1982). To test the robustness of the obtained periodicities, we also undertook single spectrum analysis (SSA) which applies an empirical orthogonal function (EOF) analysis to the autocovariance matrix on the chronologies (Burg, 1978). Both analyses were undertaken using the software *kspectra* version 3.4.3 (3.4.5). Wavelet analysis was undertaken on the 30 year averaged charcoal data using the R package “biwavelet” (Gouhier, 2013). The Morlet continuous wavelet transform was applied, and the data were padded with zeros at each end to reduce wraparound effects (Torrence and Webster, 1999).

### 3 Results and discussion

Only a limited number of Holocene pollen records come from the Falkland Islands (Barrow, 1978). The pollen record in the uppermost 90 cm at Port Stanley Airport is dominated by *Poaceae* and *Empetrum*, consistent with previous work and today’s vegetation (Barrow, 1978; Broughton and McAdam, 2003; Clark et al., 1998). The most significant change in the pollen taxa is a pronounced shift to increased representation of *Asteroidae* (accompanied by a relative decline in *Poaceae*) centered on 47 cm (equivalent to 1250 cal. BP) (Fig. 2). Although undifferentiated in the counts, the *Asteroidae* are most likely *Chilliotrichum diffusum*, common on the island across a range of habitats including *Empetrum* heath e.g. (Broughton and McAdam, 2003). The shift in the pollen diagram therefore most likely reflects the replacement of upland grasslands by *Empetrum* heath. Highly variable charcoal counts were obtained through the sequence, with the same peaks identified in both the < 50 and > 50  $\mu\text{m}$  fractions (Fig. 2), suggesting sustained periods of fire across the island and possibly as far afield as South

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

America. Importantly, the sequence preserves a record of continuous exotic pollen delivery into the site, with *Nothofagus*, *Podocarp*, *Ephedra fragilis* and *Anacardium*-type, all originating from South America. Crucially all samples contain measurable quantities of *Nothofagus*, a taxa not known to have grown on the Falkland Islands since the Middle Miocene/Early Pliocene (Macphail and Cantrill, 2006) but detected as pollen in Lateglacial (Clark et al., 1998) and Holocene (Barrow, 1978) sequences. The youngest arboreal macrofossils of the other exotic taxa is dated to late Tertiary deposits on West Point Island, West Falkland (Birnie and Roberts, 1986).

The above taxa were converted to frequency and concentration values to explore changing input into the site over the last 2600 yr (Fig. 3). Crucially, the *Nothofagus* demonstrate a comparable trend to charcoal preserved in the sequence. While the *Nothofagus* pollen concentration is relatively variable, consistent with more distal studies (Strother et al., 2015), the counts are sufficiently high to recognize identical features in the charcoal record, with several periods of low fire frequency associated with implied weaker westerly airflow. Periods with prominent peaks in *Nothofagus* and charcoal are recognized at approximately 2450, 2100, 1800, 1500, 1200, 600 and 150 cal. years BP (Fig. 3).

In contrast to previous work at Annenkov Island, which suggested enhanced westerly airflow is associated with wetter conditions (Strother et al., 2015), we appear to observe the reverse, with stronger westerly airflow linked to warmer/drier conditions, leading to greater fire frequency. However, modern comparisons between the SAM (as a measure of westerly airflow) and air temperature suggest a positive correlation (Abram et al., 2014). Comparing historic observations of SAM (Marshall, 2003) with ERA79 Interim reanalysis (Dee et al., 2011), we observe a highly significant relationship between enhanced westerly airflow and 2 m air temperature across much of South America, the Antarctic Peninsula and the Falkland Islands (Fig. 4), supporting our observations. The contrasting moisture interpretation is most probably a result of the rain shadow effect of the Andes on the Falklands with the South Atlantic playing a more significant role on South Georgia.

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The MTM analysis identifies two different periodicities in the charcoal (< 50  $\mu\text{m}$ ) record from Port Stanley Airport significant above 95 %: 146 and 244 yr (Fig. 5a) with the latter exhibiting a broad multi-decadal peak. To test whether the MTM spectral peaks are robust, we undertook SSA on the sequence chronologies. A Monte Carlo significance test identified two significant periodicities (above 95 %) at 233 and 361 yr (Fig. 5b). The 233–244 yr signal is significant in both analyses, suggesting this periodicity is pervasive through the record and the more robust.

The existence of a 200–250 yr periodicity has been identified in numerous Holocene records globally (Galloway et al., 2013; Poore et al., 2004), including Southern Ocean productivity as recorded in Palmer Deep (Domack et al., 2001; Leventer et al., 1996) and dust deposition over Antarctica (Delmonte et al., 2005). Importantly, a similar periodicity has been observed in records of atmospheric  $^{14}\text{C}$  (Stuiver and Braziunas, 1993; Turney et al., 2005), suggesting solar variability may play a role (Leventer et al., 1996).

The role changing solar output may have on westerly airflow is not immediately apparent. One possibility is that the 200–250 yr periodicity may change salinity in the North Atlantic (Stuiver and Braziunas, 1993), driving changes in the Meridional Overturning Circulation that are transmitted globally. However, the existence of the same periodicity in the delivery of dust on to the East Antarctic Ice Sheet (Delmonte et al., 2005) does imply a direct atmospheric link, either through changing sea ice extent or sea surface temperatures, or via the westerlies themselves. A recent modeling study suggested centennial-duration periods of lower solar activity may lead to an equatorward shift in the mean westerly latitude (Varma et al., 2011). Future work in this area is required to understand the driving mechanism(s) behind the 200–250 yr periodicity on global climate.

Importantly, the 250 yr periodicity identified here varies in amplitude over the last 2600 yr (Fig. 6). A Gaussian filtered curve and wavelet plot shows this periodicity is most strongly expressed between 2000 and 1000 cal. BP (Fig. 6b and c). Our results imply that the central Southern Hemisphere westerlies were particularly strong during this period and/or lay close to the latitude of the Falkland Islands, at least within the





## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

der the present core of Southern Hemisphere westerly airflow (the so-called “furious fifties”) and spanning the last 2600 years. We observe peaks in taxa from South America (particularly *Nothofagus*) that appear to be linked to warm, dry conditions, leading to fire across the island. Spectral analysis identifies a robust 250 yr periodicity, with evidence of stronger westerly airflow between 2000 and 1000 calendar yrs. Along with other records, this periodicity strongly suggests solar forcing plays a significant role in modulating the strength of the Southern Hemisphere westerlies, something hitherto not recognised, and will form the focus of future research.

*Acknowledgements.* CSMT and CF acknowledge the support of the Australian Research Council (FL100100195, FT120100004 and DP130104156). We thank the Falkland Islands Government for permission to undertake sampling on the island (permit number: R07/2011) and Darren Christie for assisting with the fieldwork.

The authors declare no competing financial interests.

## References

- Abram, N. J., Mulvaney, R., Vimeux, F., Phipps, S. J., Turner, J., and England, M. H.: Evolution of the Southern Annular Mode during the past millennium, *Nat. Clim. Change*, 4, 564–569, 2014.
- Appleby, P. G.: Chronostratigraphic techniques in recent sediments, in: *Tracking Environmental Change using Lake Sediments. Volume 1: Basin Analysis, Coring, and Chronological Techniques*, Kluwer Academic, Dordrecht, the Netherlands, 2001.
- Barrow, C.: Postglacial pollen diagrams from south Georgia (sub-Antarctic) and West Falkland island (South Atlantic), *J. Biogeogr.*, 5, 251–274, 1978.
- Birnie, J. F. and Roberts, D. E.: Evidence of Tertiary forest in the Falkland Islands (Ilas Malvinas), *Palaeogeogr. Palaeoclimatol.*, 55, 45–53, 1986.
- Björck, S., Rundgren, M., Ljung, K., Unkel, I., and Wallin, Å.: Multi-proxy analyses of a peat bog on Isla de los Estados, easternmost Tierra del Fuego: a unique record of the variable Southern Hemisphere Westerlies since the last deglaciation, *Quaternary Sci. Rev.*, 42, 1–14, 2012.

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Broughton, D. A. and McAdam, J. H.: The current status and distribution of the Falkland Islands pteridophyte flora, *Fern Gazette*, 17, 21–38, 2003.
- Burg, J. P.: A new analysis technique for time series data, in: *Modern Spectrum Analysis*, edited by: Childers, D. G., IEEE Press, New York, 42–48, 1978.
- 5 Clark, R., Huber, U. M., and Wilson, P.: Late Pleistocene sediments and environmental change at Plaza Creek, Falkland Islands, South Atlantic, *J. Quaternary Sci.*, 13, 95–105, 1998.
- Cook, A. J., Poncet, S., Cooper, A. P. R., Herbert, D. J., and Christie, D.: Glacier retreat on South Georgia and implications for the spread of rats, *Antarct. Sci.*, 22, 255–263, 2010.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,  
10 Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and  
15 performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, 2011.
- Delmonte, B., Petit, J., Krinner, G., Maggi, V., Jouzel, J., and Udisti, R.: Ice core evidence for secular variability and 200 year dipolar oscillations in atmospheric circulation over East Antarctica during the Holocene, *Clim. Dynam.*, 24, 641–654, 2005.
- Delworth, T. L. and Zeng, F.: Regional rainfall decline in Australia attributed to anthropogenic greenhouse gases and ozone levels, *Nat. Geosci.*, 7, 583–587, 2014.
- 20 Domack, E., Leventer, A., Dunbar, R., Taylor, F., Brachfeld, S., and Sjunneskog, C.: Chronology of the Palmer Deep site, Antarctic Peninsula: a Holocene palaeoenvironmental reference for the circum-Antarctic, *Holocene*, 11, 1–9, 2001.
- Domack, E., Duran, D., Leventer, A., Ishman, S., Doane, S., McCallum, S., Amblas, D., Ring, J., Gilbert, R., and Prentice, M.: Stability of the Larsen B ice shelf on the Antarctic Peninsula during the Holocene epoch, *Nature*, 436, 681–685, 2005.
- 25 Faegri, K. and Iversen, J.: *Textbook of Pollen Analysis*, Blackwell, Oxford, 1975.
- Fogt, R. L., Perlwitz, J., Monaghan, A. J., Bromwich, D. H., Jones, J. M., and Marshall, G. J.: Historical SAM variability. Part II: Twentieth-century variability and trends from reconstructions, observations, and the IPCC AR4 models, *J. Climate*, 22, 5346–5365, 2009.
- 30 Fredskild, B.: Holocene palaeo-winds and climatic changes in West Greenland as indicated by long-distance transported and local pollen in lake sediments, in: *Climatic Changes on*



## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

- Lamy, F., Kilian, R., Arz, H. W., Francois, J.-P., Kaiser, J., Prange, M., and Steinke, T.: Holocene changes in the position and intensity of the southern westerly wind belt, *Nat. Geosci.*, 3, 695–699, 2010.
- Le Quére, C., Raupach, M. R., Canadell, J. G., and Marland, G. E. A.: Trends in the sources and sinks of carbon dioxide, *Nat. Geosci.*, 2009, 831–836, doi:10.1038/ngeo1689, 2009.
- Lenton, A., Tilbrook, B., Law, R. M., Bakker, D., Doney, S. C., Gruber, N., Ishii, M., Hoppema, M., Lovenduski, N. S., Matear, R. J., McNeil, B. I., Metzl, N., Mikaloff Fletcher, S. E., Monteiro, P. M. S., Rödenbeck, C., Sweeney, C., and Takahashi, T.: Sea–air CO<sub>2</sub> fluxes in the Southern Ocean for the period 1990–2009, *Biogeosciences*, 10, 4037–4054, doi:10.5194/bg-10-4037-2013, 2013.
- Leventer, A., Domack, E. W., Ishman, S. E., Brachfeld, S., McClennen, C. E., and Manley, P.: Productivity cycles of 200–300 years in the Antarctic Peninsula region: understanding linkages among the sun, atmosphere, oceans, sea ice, and biota, *Geol. Soc. Am. Bull.*, 108, 1626–1644, 1996.
- Lister, D. H. and Jones, P. D.: Long-term temperature and precipitation records from the Falkland Islands, *Int. J. Climatol.*, doi:10.1002/joc.4049, 2014.
- Liu, J. and Curry, J. A.: Accelerated warming of the Southern Ocean and its impacts on the hydrological cycle and sea ice, *P. Natl. Acad. Sci. USA*, 107, 14987–14992, 2010.
- Macphail, M. and Cantrill, D. J.: Age and implications of the Forest Bed, Falkland Islands, southwest Atlantic Ocean: evidence from fossil pollen and spores, *Palaeogeogr. Palaeoclimatol.*, 240, 602–629, 2006.
- Marshall, G.: Trends in the Southern Annular Mode from observations and reanalyses, *J. Climate*, 16, 4134–4143, 2003.
- Marshall, J. and Speer, K.: Closure of the meridional overturning circulation through Southern Ocean upwelling, *Nat. Geosci.*, 5, 171–180, 2012.
- McGlone, M., Wilmshurst, J. M., and Wiser, S. K.: Lateglacial and Holocene vegetation and climatic change on Auckland Island, Subantarctic New Zealand, *Holocene*, 10, 719–728, 2000.
- Messié, M. and Chavez, F.: Global modes of sea surface temperature variability in relation to regional climate indices, *J. Climate*, 24, 4314–4331, 2011.
- Poore, R. Z., Quinn, T. M., and Verardo, S.: Century-scale movement of the Atlantic Intertropical Convergence Zone linked to solar variability, *Geophys. Res. Lett.*, 31, L12214, doi:10.1029/2004GL019940, 2004.

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

- Ramsey, C. B.: Radiocarbon dating: revolutions in understanding, *Archaeometry*, 50, 249–275, 2008.
- Roig, F. A., Le-Quesne, C., Boninsegna, J. A., Briffa, K. R., Lara, A., Grudd, H., Jones, P. D., and Villagrán, C.: Climate variability 50,000 years ago in mid-latitude Chile as reconstructed from tree rings, *Nature*, 410, 567–570, 2001.
- Scott, L. and van Zinderen Barker, E. M.: Exotic pollen and long-distance wind dispersal at a Sub-Antarctic island, *Grana*, 24, 45–54, 1985.
- Strother, S. L., Salzmann, U., Roberts, S. J., Hodgson, D. A., Woodward, J., Van Nieuwenhuyze, W., Verleyen, E., Vyverman, W., and Moreton, S. G.: Changes in Holocene climate and the intensity of Southern Hemisphere Westerly Winds based on a high-resolution palynological record from sub-Antarctic South Georgia, *Holocene*, 25, 263–279, 2015.
- Stuiver, M. and Braziunas, T. F.: Sun, ocean, climate and atmospheric  $^{14}\text{CO}_2$ : an evaluation of causal and spectral relationships, *Holocene*, 3, 289–305, 1993.
- Thompson, D. W. J., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M., and Karoly, D. J.: Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change, *Nat. Geosci.*, 4, 741–749, 2011.
- Thomson, D. J.: Spectrum estimation and harmonic analysis, *Proc. IEEE*, 70, 1055–1096, 1982.
- Torrence, C. and Webster, P. J.: Interdecadal changes in the ENSO–Monsoon system, *J. Climate*, 12, 2679–2690, 1999.
- Turney, C., Baillie, M., Clemens, S., Brown, D., Palmer, J., Pilcher, J., Reimer, P., and Leuschner, H. H.: Testing solar forcing of pervasive Holocene climate cycles, *J. Quaternary Sci.*, 20, 511–518, 2005.
- Upton, J. and Shaw, C. J.: An overview of the oceanography and meteorology of the Falkland Islands, *Aquat. Conserv.*, 12, 15–25, 2002.
- van Oldenborgh, G. J. and Burgers, G.: Searching for decadal variations in ENSO precipitation teleconnections, *Geophys. Res. Lett.*, 32, L15701, doi:10.1029/2005GL023110, 2005.
- Varma, V., Prange, M., Lamy, F., Merkel, U., and Schulz, M.: Solar-forced shifts of the Southern Hemisphere Westerlies during the Holocene, *Clim. Past*, 7, 339–347, doi:10.5194/cp-7-339-2011, 2011.
- Villalba, R., Lara, A., Masiokas, M. H., Urrutia, R., Luckman, B. H., Marshall, G. J., Mundo, I. A., Christie, D. A., Cook, E. R., Neukom, R., Allen, K., Fenwick, P., Boninsegna, J. A., Srur, A. M., Morales, M. S., Araneo, D., Palmer, J. G., Cuq, E., Aravena, J. C., Holz, A., and LeQuesne, C.:

Unusual Southern Hemisphere tree growth patterns induced by changes in the Southern Annular Mode, *Nat. Geosci.*, 5, 793–798, 2012.

Visbeck, M.: A station-based Southern Annular Mode Index from 1884 to 2005, *J. Climate*, 22, 940–950, 2009.

5 Whitlock, C. and Larsen, C.: Charcoal as a fire proxy, in: *Tracking Environmental Change using Lake Sediments*, Springer Dordrecht, the Netherlands, 75–97, 2001.

Yin, J. H.: A consistent poleward shift of the storm tracks in simulations of 21st century climate, *Geophys. Res. Lett.*, 32, doi:10.1029/2005GL023684, 2005.

# CPD

11, 2159–2180, 2015

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

**Table 1.** Radiocarbon ages and modelled calibrated age ranges using SHCal13 (Hogg et al., 2013) and Bomb04SH (Hua and Barbetti, 2004) using the P\_sequence option in OxCal (Ramsey, 2008).


Depth, cm	Wk lab number	Material	$^{14}\text{C} \pm 1\sigma$	Cal. yr BP $\pm 1\sigma$
8–9	34 598	Fruits and leaves	$117.0 \pm 0.4\% \text{M}$	$-22 \pm 13$
11–12	32 994	Fruits and leaves	$107.8 \pm 0.4\% \text{M}$	Failed age model
18–19	37 007	Fruits and leaves	$107.3 \pm 0.3\% \text{M}$	$-7 \pm 2$
25–26	35 146	Fruits and leaves	$95 \pm 25$	$85 \pm 61$
35–36	37 008	Fruits and leaves	$647 \pm 25$	$603 \pm 29$
39–40	33 445	Fruits and leaves	$761 \pm 25$	$660 \pm 27$
57–58	32 996	Fruits and leaves	$1818 \pm 25$	$1672 \pm 50$
70–71	32 350	Fruits and leaves	$2235 \pm 25$	$2204 \pm 66$
97–98	32 997	Fruits and leaves	$2749 \pm 25$	$2802 \pm 31$

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.



**Figure 1.** Location of the Falkland Islands in the South Atlantic Ocean with mean locations of the Polar and Southern Boundary fronts (dashed lines), the continental shelf (grey areas) and prevailing westerly airflow (solid arrows) **(a)**; and Port Stanley Airport site in the east Falkland Islands **(b)**. **(a)** was modified from (Strother et al., 2015) and **(b)** was obtained from Google Earth.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 2.** Pollen concentration diagram (%TLP) and charcoal counts plotted with age–depth relationship for the Port Stanley Airport peat sequence (blue envelope denotes 68 % confidence limits).

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years**

C. Turney et al.



**Figure 3.** Pollen frequency and concentration diagram from Port Stanley Airport site plotted against calendar age.

[Title Page](#)

[Abstract](#)   [Introduction](#)

[Conclusions](#)   [References](#)

[Tables](#)   [Figures](#)

[◀](#)   [▶](#)

[◀](#)   [▶](#)

[Back](#)   [Close](#)


[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years


C. Turney et al.



**Figure 4.** Correlation (a) and significance (b) of relationship between the hemispherically-averaged Southern Annular Mode (SAM) index (Marshall, 2003) with 2 m air temperature in the ERA-79 Interim reanalysis (Dee et al., 2011) (July–June 1979–2013). Analyses were made with KNMI Climate Explorer (van Oldenborgh and Burgers, 2005).

## A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years

C. Turney et al.



**Figure 5.** Multi-Taper Method (MTM) **(a)** and Monte-Carlo Single Spectrum Analysis (SSA) analyses **(b)** of charcoal from the Port Stanley Airport sequence. Error bars denote 95 % confidence.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 6.** Charcoal concentration ( $< 50 \mu\text{m}$ ) (a), charcoal Gaussian filtered for the periodicity  $244 \pm 49$  years (b) and wavelet analysis of charcoal concentration (c) from Port Stanley Airport ( $52^\circ \text{S}$ ). Solid black line in wavelet denotes 95% confidence in periodicity; white dashed line denotes cone of influence. (d) shows biogenic carbonate accumulation rate (AR) from Palm2, Chile ( $53^\circ \text{S}$ ) (Lamy et al., 2010).

**A 250 year periodicity in Southern Hemisphere westerly winds over the last 2600 years**

C. Turney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion