We thank both reviewers for their positive comments about the changes that have been made following their recommendations. We tried to do our best to address their latest suggestions with a point-by-point response.

#### **Reviewer 1**

The authors have responded well to earlier comments and I think the paper as it stands is an interesting and novel contribution which can be published. I have some suggestions for some further small revisions specified below which could further improve the paper.

1. The conclusions that the FIS "survives" (losing only 10% of its mass) as the LIS is raised is an important one which could be more strongly and clearly highlighted, for example with an extra bullet point in Section 6 and with a specific comment in the abstract.

We followed the suggestions to clearly highlight this result. We therefore added a new sentence towards the end of the abstract and modified the conclusion section with a new bullet point (see point 5 in section 6).

Added in the abstract: "The most striking result is that the Eurasian ice sheet is maintained under full-LGM North American ice-sheet conditions, but loses ~10 % of its mass compared to the case in which the North American ice sheet is flat"

Added in the conclusion: "Within the context of glacial-interglacial cycles, the FIS inception under full-LGM LIS conditions appears to be unrealistic. Therefore, we performed additional simulations with a high FIS and flat and full-LGM LIS. The results of these new sensitivity experiments show a similar response to the first series of experiments: in the presence of a full LIS, the FIS undergoes a significant mass loss. This confirms that a large LIS counteracts the growth of the FIS although it is still maintained, such as it does within more realistic contexts, and highlights therefore the self-consistency of the overall study."

2. Further to the above, an interesting point which is not explicitly made in the paper is that the decrease in FIS ice volume with increasing LIS is consistent with the results of Kleman et al (2013) who show in their Fig 3 that FIS and LIS grow together in the early part of the last glacial, while in the later part the LIS continues to grow while the FIS diminishes. This point could be made explicit in p. 17, for example.

We acknowledge that we did not insist enough on the agreement with the results of Kleman et al. (2013). We therefore modified the text accordingly in section 5.1:

"More recently, Kleman et al. (2013) combined geological evidences and numerical modelling to reconstruct the paleo-topography of past Northern hemisphere ice sheets during MIS5b (~86 kyr), and MIS4 (~64 kyr) and 2 (LGM). They found that the LIS and the FIS grow in concert in the early part of the last glacial cycle (i.e. until MIS5b), while after MIS4, their evolution is radically different with a further growth of the LIS and a significant shrinkage of the FIS due to its southwestward retreat towards the main precipitation sources. [....]. The large positive SMB values over the SBK sector when the LIS is low combined with the westward displacement of the Eurasian ice sheet as the LIS gets higher are fully in line

with the ice-sheet paleo-topography reconstructions provided by Kleman et al. (2013) for the last glacial cycle".

3. I insist on my earlier comment that Lofverstrom et al (2014) show warm anomalies over Central Asia and Siberia mostly due to the effect of the FIS; it's true that the LIS gives a small warm anomaly in northwestern Siberia also, but this is much smaller than that due to the FIS. The interpretation in Lofverstrom et al is that the FIS is largely self-limiting as regards its eastward expansion.

The text has been modified in the discussion Section (see section 5.1):

"Löfverström et al. (2014) propose to attribute the westward shift of the FIS and the decrease of the Kara ice sheet to warm temperatures over the southeastern part of the Fennoscandian region, induced by the FIS itself, along with the upslope precipitation effect proposed by Sanberg and Oerlemans (1983). In their study, the LIS exerts only a small influence on the retreat of the Kara ice sheet.

#### Reviewer 2

This study well explores the impact of different LIS configurations on FIS mass balance by employing experiments in AGCM and ISM. The questions mentioned in last review process are solved properly in this version. A minor flaw of this study, as also mentioned by authors in the text, is its implication to realistic glaciations of Eurasian ice sheet (EIS). In discussion part 5.1, the authors discussed their results against previous modeling studies regarding FIS evolution through last glacial cycle.

According to the model setup, the land-sea mask, especially in the area covered by FIS, is prescribed to the LGM. This thus provides an ad hoc sea level drop (land area) for FIS build-up as evaluating changes in FIS mass balance in response to LIS. How will it be if Barents-Kara sea still exists? Will it influence on the present conclusion? To clarify this issue, authors are suggested either providing extra experiments or proper discussion in the revised manuscript.

We fully agree with the reviewer. The choice of an LGM land-sea mask provides an ad-hoc sea level drop and as a result the Barents-Kara (BK) sector is considered as a continental area. In this study, we focused on large-scale atmospheric circulation changes induced by the LIS albedo and topography effects. Therefore, huge differences in these large-scale changes caused by a modification of the land-sea mask in the BK sector are unlikely. However, having land in the BK area may have local consequences. As mentioned in section 3.1.1, this favors snow accumulation and therefore glacial inception through the positive albedo feedback. As a consequence, in the presence of a marine BK sector, glaciation would likely to be reduced or at least slowed down. As it is now explained in the revised manuscript, our experimental setup uses LGM SST and sea ice boundary conditions provided by the coupled atmosphere-ocean IPSL model (Kageyama et al. 2013). In this run, the BK sector was covered by the FIS, and it is thus impossible to investigate the conditions that would have prevailed in case of a marine BK sector. This is why further studies using a fully coupled atmosphere-ocean will be necessary in the future to investigate further the relationships between the LIS and the FIS

within more realistic contexts. Nevertheless, we tried to clearly present the limitations induced by our experimental setup and we added a new paragraph at the beginning of section 5.2:

"Our experimental setup has been designed to investigate how the atmospheric circulation changes in response to the LIS albedo and topography effects may impact the climate over the Eurasian region and therefore the evolution of the surface mass balance of the Fennoscandian ice sheet. We have therefore neglected the potential impact of the ocean.

First of all, all experiments presented in this study have used the LGM land-sea mask and therefore the Barents-Kara region is considered as a continental area. The consequences on the simulated large-scale atmospheric circulation of this assumption are likely to be negligible. However, accounting for a marine Barents-Kara area would decrease the positive albedo feedback in this region, and thus the FIS inception (see section 3.1.1). Our experimental setup uses LGM SST and sea ice boundary conditions provided by the coupled atmosphere-ocean IPSL model (Kageyama et al. 2013). This prevents us from studying the conditions that could have prevailed with a marine Barents-Kara area, because this sector was covered by the FIS in this run, and future studies will be necessary to investigate further the relationships between the LIS and the FIS within more realistic contexts".

In addition, it would be very meaningful if authors can provide their viewpoints regarding lead-lag relationship of glaciations among different ice sheets in Northern Hemisphere based on their study.

In the absence of any transient simulation, it is not easy to discuss the potential lead-lag relationships of the Northern hemisphere ice sheets. Nevertheless, based on our results we proposed a plausible glaciation scenario presented in the last paragraph of the conclusion section. We did not provide any new idea, but with this last paragraph we intended to better highlight the implications of our study.

"It is worth noting that without a transient simulation of the last glacial period, it is not possible to accurately elaborate the lead-lag relationships of the Northern hemisphere ice sheets during the last glacial period. However, based on our results, a plausible glaciation scenario can be proposed with an early phase of FIS glacial inception favored by the areal expansion of the LIS, and followed by a limited growth of the FIS with a progressive westward retreat of the FIS itself as the LIS altitude gradually increased. Such proposed relationships between ice-sheet build-up demonstrate that this study also provides new insights in the understanding of the various Northern hemisphere ice sheet configurations during the different glacial-interglacial cycles."

# How might the North American ice sheet influence the Northwestern Eurasian climate?

D	Daahin <sup>1</sup> C	Charle:41	C Dumas <sup>1</sup>	M Vacas	yama <sup>1</sup> , C. Ritz <sup>2</sup>
Р.	Begnin, S	s. Charbit .	, C. Dumas ,	, M. Kage <sup>,</sup>	yama , C. Kitz

 $Corresponding\ author: sylvie.charbit@lsce.ipsl.fr$ 

(Revised manuscript)

<sup>&</sup>lt;sup>1</sup>Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS, Gif-sur-Yvette, France

<sup>&</sup>lt;sup>2</sup>Laboratoire de Glaciologie et de Géophysique de l'Environnement, CNRS, Saint Martin d'Hères, France

#### **Abstract**

1

2 It is now widely acknowledged that past Northern hemisphere ice sheets covering Canada and 3 northern Europe at the last glacial maximum (LGM) exerted a strong influence on climate by 4 causing changes in atmospheric and oceanic circulations. In turn, these changes may have 5 impacted the development of the ice sheets themselves through a combination of different 6 7 feedback mechanisms. The present study is designed to investigate the potential impact of the 8 North American ice sheet on the surface mass balance (SMB) of the Eurasian ice sheet driven by simulated changes in the past glacial atmospheric circulation. Using the LMDZ5 9 atmospheric circulation model, we carried out twelve experiments under constant LGM 10 conditions for insolation, greenhouse gases and ocean. In these experiments, the Eurasian ice 11 sheet is removed. The twelve experiments differ in the North American ice-sheet topography, 12 ranging from a white and flat (present-day topography) ice sheet to a full-size LGM ice sheet. 13 This experimental design allows the albedo and the topographic impacts of the North American 14 15 ice sheet onto the climate to be disentangled. The results are compared to our baseline 16 experiment where both the North American and the Eurasian ice sheets have been removed. In 17 summer, the sole albedo effect of the American ice sheet modifies the pattern of planetary waves with respect to the no-ice sheet case, resulting in a cooling of the northwestern Eurasian 18 region. By contrast, the atmospheric circulation changes induced by the topography of the 19 North American ice sheet lead to a strong decrease of this cooling. In winter, the Scandinavian 20 and the Barents-Kara regions respond differently to the American ice sheet albedo effect: in 21 22 response to atmospheric circulation changes, Scandinavia warms and total precipitation is more abundant whereas the Barents-Kara area cools with decrease of convective processes causing a 23 decrease of total precipitation. The gradual increase of the altitude of the American ice sheet 24 leads to less total precipitation and snowfall and to colder temperatures over both the 25 Scandinavian and the Barents-Kara Seas sectors. We then compute the resulting annual surface 26 mass balance over the Fennoscandian region from the simulated temperature and precipitation 27 fields used to force an ice-sheet model. It clearly appears that the SMB is dominated by the 28 ablation signal. In response to the summer cooling induced by the American ice-sheet albedo, 29 high positive SMB values are obtained over the Eurasian region, leading thus to the growth of 30 an ice sheet. On the contrary, the gradual increase of the American ice sheet altitude induces 31 32 more ablation over the Eurasian sector, hence limiting the growth of Fennoscandia. To test the robustness of our results w.r.t the Eurasian ice sheet state, we carried out two additional LMDZ 33 34 experiments with new boundary conditions involving both the American (flat or full-LGM) and a high Eurasian ice sheets. The most striking result is that the Eurasian ice sheet is maintained 35 under full-LGM North American ice-sheet conditions, but loses ~10 % of its mass compared to 36 the case in which the North American ice sheet is flat. These new findings qualitatively confirm 37 the conclusions from our first series of experiments and suggest that the development of the 38 Eurasian ice sheet may have been slowed down by the growth of the American ice sheet, 39 offering thereby a new understanding of the evolution of Northern hemisphere ice sheets 40 throughout glacial-interglacial cycles. 41

#### 1. Introduction

42

The last million years is characterized by alternating glacial and interglacial phases. During 43 glacial periods, large ice sheets covering present-day Canada and northwestern Eurasia (Dyke 44 45 and Prest 1987; Clark et al. 1993; Peltier 2004; Svendsen et al. 2004; Lambeck et al. 2006; Tarasov et al. 2012) exerted a strong influence on climate. Several studies highlighted the 46 47 importance the climatic changes induced by continental-scale ice sheets (e.g. Clark 1999). It has been recognized that the ice-sheet topography is likely to be the main factor altering the 48 49 atmospheric circulation in the Northern hemisphere (Broccoli and Manabe 1987; Pausata et al. 2011). As an example, the pioneering study carried out by Manabe and Broccoli (1985) with an 50 51 atmospheric general circulation model shows that the North American ice sheet caused a split 52 of the westerlies. The authors also highlight a larger amplitude of the planetary waves due to 53 the presence of ice sheets. Recently, Ullman et al. (2014) tested the influence of different North American ice-sheet reconstructions on the climate. They showed that a higher American ice 54 sheet leads to a more zonal Atlantic jet, thereby confirming the key role of ice-sheet topography 55 on atmospheric circulation. Using model outputs from the Paleo Modelling Intercomparison 56 Project 2 (PMIP2) (Braconnot et al. 2007), Laîné et al. (2009) and Rivière et al. (2010) 57 highlight a strengthening and an equatorward displacement of the sub-tropical jet-stream during 58 the Last Glacial Maximum (LGM) w.r.t. the pre-industrial period. Changes in the position and 59 the strength of the North Atlantic jet stream induce changes in the storm tracks, and therefore in 60 precipitation (Hall et al. 1996; Kageyama and Valdes 2000; Laîné et al. 2009; Rivière et al. 61 2010). These changes also have an influence on the energy transport and can therefore modify 62 63 the temperature. The climatic changes induced by large-scale ice sheets exert an influence on both temperature 64 and precipitation that drive the ice-sheet surface mass balance (SMB), defined as the sum of 65 snow accumulation and snow and ice ablation. Using a simple ice-sheet model based on an 66 idealized geometry coupled to a stationary-wave model, Roe and Lindzen (2001a,b) underline 67 the importance of accounting for the feedbacks between ice sheets and temperatures induced by 68 69 changes in stationary waves to properly simulate the evolution of an ice sheet. They show that the self-induced temperature anomaly due to an ice sheet leads to a warming over the ice-sheet 70 71 western part. This may explain the absence of ice over Alaska at the LGM. They also suggest 72 that the stationary waves excited by the North American ice sheet may have contributed to a warming over Europe, influencing the development of the Eurasian ice sheet. In the same way, 73 with a three-dimensional stationary wave model, Liakka et al. (2011) showed that the position 74

of the southern margin of ice sheets strongly depends on the temperature anomalies due to stationary waves, which are modified by the ice sheet itself. More recently, with the use of the CAM3 atmospheric model run under four different climatic contexts (last interglacial, marine isotope stages 5b and 4 (MIS5b and MIS4), and LGM periods), Löfverström et al. (2014) showed how the atmospheric circulation changes induced by the ice sheets could have influenced the growth of the ice sheets themselves. Similarly to Roe and Lindzen (2001a), they show that the summer atmospheric circulation change due to the presence of ice sheets may cause a sufficient warming over Siberia and Alaska to inhibit ice growth. They also perform two experiments under MIS4 conditions to test the influence of one ice sheet on the other. They conclude that the summer temperature anomaly induced by the presence of the MIS4 American ice sheet is too weak to explain the small size of the European ice sheet at that period, but may have contributed to the westward shift of the ice-sheet mass centre. Although several studies have been devoted to the mutual influences of the ice sheets (Roe and Lindzen 2001a; Beghin et al. 2014; Löfverström et al. 2014), no study has specifically investigated the mechanisms through which the American ice sheet may have influenced the European climate, and therefore the European ice-sheet surface mass balance.

The aim of this paper is to investigate the atmospheric response to changes in the North American ice sheet boundary conditions and to examine how the related atmospheric circulation changes impact the climate over the Scandinavian and Barents-Kara Seas regions. To achieve this goal, we use the LMDZ5 atmospheric general circulation model run with different thicknesses of the North American ice sheet taken as boundary conditions, with no ice at all over Eurasia. We investigate the mechanisms by which the American ice sheet may change the surface mass balance of the Eurasian ice sheet. Besides gradually increasing the altitude of the North American ice sheet, the originality of the present study also relies on ice-sheet model simulations. We use climatic fields simulated by the LMDZ5 model as inputs to a three-dimensional ice-sheet model to compute the surface mass balance of the Eurasian ice sheet. The description of the climate and ice-sheet models is given in section 2 as well as the experimental design. The model results, presented in sections 3 and 4 are followed by a discussion (section 5). Section 6 summarizes the main findings of our study.

## 2. Model and experiment

107

108

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

#### 2.1 The atmospheric model

109 The LMDZ5tmospheric model used in this study belongs to the LMDZ model family developed at Laboratoire de Météorologie Dynamique (Sadourny and Laval 1984; Le Treut et al. 1994; Le 110 Treut et al. 1998, Hourdin et al., 2006). LMDZ is the atmospheric component of the IPSL-111 CM5A coupled ocean-atmosphere model (Dufresne et al. 2013). The dynamical equations are 112 113 discretized on a longitude-latitude-staggered Arakawa C-grid (Kasahara 1977). The model ensures the conservation of both enstrophy (square of wind rotational) for baroclinic flows 114 (Sadourny 1975b; Sadourny 1975a) and the axi-symmetric component of the angular 115 momentum. The model version used in this study has 39 vertical levels and runs on a 96 x 95 116 model grid resolution (3.75° x 1.9°). A complete description of the model can be found in 117 (Hourdin et al. 2013). 118

#### 2.2 The ice-sheet model

The GRISLI ice-sheet model is a three-dimensional thermo-mechanical model which simulates the evolution of ice-sheet geometry (extension and thickness) and the coupled temperaturevelocity fields in response to climate forcing. A comprehensive description of the model can be found in (Ritz et al. 2001) and (Peyaud et al. 2007). Here, we only summarize the main characteristics of this model. The equations are solved on a cartesian grid (40 km x 40 km). Over the grounded part of the ice sheet, the ice flow resulting from internal deformation is governed by the shallow-ice approximation (Hutter 1983; Morland et al. 1984). The model deals with ice flow through ice shelves using the shallow-shelf approximation (MacAyeal 1989). It also predicts the large-scale characteristics of the ice streams using criteria based on the effective pressure and hydraulic load. At each time step, the velocity and vertical profiles of temperature in the ice are computed as well as the new geometry of the ice sheet. The isostatic adjustment of bedrock in response to ice load is governed by the flow of the asthenosphere, with a characteristic time constant of 3000 years, and by the rigidity of the lithosphere. The temperature field is computed both in the ice and in the bedrock by solving a time-dependent heat equation. The surface mass balance is defined as the sum between accumulation and ablation computed by the empirical positive degree-day (PDD) method (Reeh 1991; Fausto et al. 2009). This method assumes that melt rates of snow and ice are linearly related to the number of PDD through degree-day factors for snow and ice materials (Braithwaite 1984; 1995).

# 2.3 Experimental set-up

139

In order to investigate the albedo and topography effects induced by the North American ice 140 sheet on the Eurasian climate at the LGM, we carried out 12 simulations under LGM conditions 141 142 for greenhouse gases (GHG), insolation, sea-surface temperatures and sea-ice). In this series of simulations, the altitude of the North American ice sheet surface ranges from that of the 143 144 present-day surface to 100 % of that used in the PMIP3 LGM experiments. The PMIP3 LGM ice sheets result from a combination of three reconstructions, namely ICE-6G v2.0 (Toscano et 145 146 al. 2011), GLAC-1 (Tarasov et al. 2012) and ANU (Lambeck 2001). The way this new reconstruction has been obtained is explained in detail on the PMIP3 website 147 148 (https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:design:pi:final:icesheet) and in Abe-Ouchi et al. (2015). In our baseline experiment (noIS), the land-ice mask is modified (w.r.t. PMIP3) to 149 150 remove both the European and the American ice sheets and the altitude is that of the presentday (Fig. 1). In the other simulations, we only remove the European ice-sheet mask. These 151 simulations are referred to as xxdhL, where dhL represents the surface height difference 152 between the PMIP3 LGM Laurentide ice sheet and the present-day surface, and xx represents 153 the percentage of dhL taken into account. To simplify the writing, the American and Eurasian 154 ice sheets are respectively referred to as "Laurentide ice sheet" (LIS) and the "Fennoscandian 155 ice sheet" (FIS) in the following. 156 157 The topography of the 00dhL experiment is therefore the same as today, but the land-ice mask is set to the LGM one (Figs. 1 and 2). Greenland and Antarctic ice sheets are the same as in the 158 PMIP3 experimental set-up. Both insolation and GHG boundary conditions are similar to those 159 160 defined in the PMIP3 protocol: the orbital forcing is taken at 21 kyr BP from (Berger 1978), while atmospheric GHG concentrations are those recorded in Antarctic ice cores ( $CO_2 = 185$ 161 162 ppm, (Lüthi et al. 2008);  $CH_4 = 350$  ppb, (Loulergue et al. 2008);  $N_2O = 200$  ppb, (Spahni et al. 2005)). The LGM land-sea mask is also taken into account, with closed Bering Strait, and land 163 in Hudson Bay and Barents Sea (Fig. 1). The sea-surface temperatures (SST) and sea-ice cover 164 come from 50 consecutive years of the IPSL PMIP3 LGM run outputs (Kageyama et al. 2013). 165 These oceanic surface conditions are depicted in Figure 1. In the North Atlantic area, the 166 temperature meridional gradient is steep in the western part of the basin and weakens in its 167 168 eastern part, as for the pre-industrial period. The sea-ice edge is located South of Iceland in 169 winter and retreats to the northern limits of the Nordic Seas in summer. These rather warm North Atlantic / Nordic Seas conditions are due to a strong Atlantic meridional overturning 170 circulation (Kageyama et al. 2013). Using these boundary conditions, the LMDZ5 simulations 171

are run for 50 years, and we study the last 40 years to be at equilibrium. Two-sided Student t-172

tests with a p-value = 0.05 (von Storch and Zwiers, 2003) are performed on key diagnostics. 173

#### 3. AGCM results

174

176

177

179

180

196

197

In this section, we focus on how the LIS influences the climatic fields which can have an 175

impact on the surface mass balance of an ice sheet. The key climatic variables used to compute

the amount of ablation with the PDD method are the monthly temperatures and the monthly

178 total (solid plus liquid) precipitation.

#### 3.1 The northwestern Eurasian temperature

# 3.1.1 Impact of the Laurentide ice sheet

Using the comparison between the 00dhL and the noIS experiments, we first consider the 181 impact of the LIS albedo on the 2-meter summer air temperature over the Scandinavian-182 Barents-Kara (SBK) region (Fig. 3, 00dhL experiment, left column). The most striking feature 183 is a cooling over the whole Eurasian continent with a maximum (more than 10 °C) centered 184 over the Barents-Kara Seas and a moderate cooling (2 to 4°C) over Scandinavia. The area 185 experiencing the largest cooling also corresponds to the area with the largest albedo (orange 186 contours). This is a direct consequence of the LGM land-sea mask used in this study for which 187 188 the Barents-Kara Seas are represented as continental areas. This favors snow accumulation and therefore higher albedo values in this area, causing in turn a positive feedback on temperatures. 189 190 Figure 3 also displays the 2m mean summer temperature over the SBK region for different 191 altitudes of the LIS. Here, we only give results of selected experiments to simplify the presentation of our results. It appears that the SBK cooling gets smaller when the LIS gets 192 193 higher. Overall, the simulated summer surface air temperature evolves together with the LIS

altitude, with a progressive reduction of the zonal and meridional extent of the largest cooling 194 195

zone. When the LIS reaches its full LGM size (100dhL- noIS), the summer cooling over

Barents-Kara Seas is reduced to 4-6°C and to 2°C at most over Scandinavia. The topography of

the LIS has therefore a warming effect on the SBK region for the summer season.

In winter, the response of the temperature to the LIS albedo and its evolution with the LIS 198

199 altitude are different from the summer response (Fig. 3, right panels). First, the LIS albedo

induces warmer temperatures (+3 °C) over South Scandinavia, the British Isles and present-day 200

North and Baltic Seas, and cooler temperatures over an area in the Barents Sea expanding from 201

Svalbard to Novaya Zemlya (3 °C). Surprisingly, in the 30dhL experiment, the amplitude of 202

- these temperature changes (w.r.t. noIS) is reduced, showing that the response of winter temperatures to changes in LIS altitude is far from being linear. Beyond 60dhL, the cooling zone progressively expands, and spreads across the entire British Isles and Scandinavia-Barents-Kara area when the LIS has its full LGM size. In contrast to the climatic response simulated for the summer season, the LIS topography leads to colder surface air temperatures in winter.
- To understand the origin of these contrasted responses, we investigate the processes through which the changes in the altitude of the Laurentide ice sheet modify the atmospheric circulation.

# 3.1.2 Atmospheric circulation processes

212

224

225

226

227

228

229

230

231

- The comparison of the zonal anomaly (00dhL vs. noIS) of the 500 hPa summer geopotential 213 height shows that the sole albedo effect of the LIS is sufficient to drastically change the 214 atmospheric circulation (Fig. 4, 00dhL and noIS, left column). The ridge over the Rockies, 215 clearly visible in noIS, disappears in 00dhL, and the trough over the Labrador sector is more 216 217 extensive. Ridges over the North Atlantic, the Greenland ice sheet and northern Europe are more developed in 00dhL (w.r.t. noIS). By contrast, the troughs over Iceland, the Norvegian 218 219 and Barents Seas appearing in noIS are weaker or even vanish in 00dhL. In noIS, the trough over Svalbard implies southerlies over the Barents Sea (Fig. 4, right column) and therefore, 220 221 warm temperatures in this area (Fig. 3). The weakening of this trough in 00dhL as well as the stronger ridge over Greenland induces northerlies over Svalbard and the Barents Sea (Fig. 4) 222 and therefore colder temperatures (Fig. 3). 223
  - The Labrador trough and the North Atlantic ridge become stronger when the LIS gets higher, until it reaches 60 % of its full size. Beyond 60dhL, the Labrador trough and the Atlantic ridge keep more or less the same amplitude. When the LIS gets higher, the Greenland ridge gets weaker, the European ridge remains almost unchanged (Fig. 4), and there is a return of the Iceland-Svalbard-Barents trough. This new trough, centered between Greenland and Svalbard, along with the weakening of the Greenland ridge, brings southerlies again over the northern Barents Sea. The Barents winds shift from northerlies to southerlies when the LIS is 60 % of its full size. The southerlies bring warmer temperatures, explaining the SBK warming with the increase of the LIS height (Figs 3 and 4).

In winter (Fig. 5), the sole albedo effect of the LIS does not drastically change the geopotential 233 anomaly. However, the meridional wind field shows a slight decrease of the southerlies over 234 the Barents Sea, explaining the slight cooling over this region. There is also a slight decrease of 235 236 the northerlies simulated over the British Isles, the eastern North Atlantic Ocean and southern Scandinavia, consistent with the moderate warming simulated over this region. In the 30dhL 237 experiment, a strengthening of the southerlies is observed (Fig. 5) inducing a warming (w.r.t. 238 00dhL) over the SBK region (Fig. 3) and explaining the non-linear behavior on winter 239 temperatures. When the altitude of the LIS further increases, there is a progressive emergence 240 241 of a trough over the Scandinavian-Barents sector. This trough comes with northerlies over the Norvegian Sea and weaker southerlies over the Barents sea (Fig. 5, right column), consistently 242 243 with colder temperatures over this area, as seen in Fig. 3.

244 This analysis explains why opposite sign temperature responses are obtained for the summer and winter seasons. Since ablation is rather sensitive to the summer season, more ice is 245 expected over the Fennoscandian area when the sole albedo effect of the LIS is operating (i.e. 246 00dhL experiment). However, the development of an ice sheet in this area also depends on the 247 248 amount of snow accumulation. Therefore, we next examine hereafter the LIS impact on the 249 SBK total precipitation and snowfall.

#### 3.2 Total and solid precipitation over northwestern Eurasia

#### 3.2.1 Impact of the Laurentide ice sheet

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

In this section, we investigate the impact of the LIS on the mean annual precipitation and snowfall over northwestern Eurasia (Fig. 6). In response to the LIS albedo effect (00dhL-noIS panel) a slight excess of precipitation is simulated over the northeastern North Atlantic and the southern and western parts of Scandinavia, but less precipitation is simulated over the Barents-Kara Sea area. However, over the whole SBK region, the snowfall anomaly is positive. This increase in snowfall is due to the summer signal (not shown) related to the cooling observed over this region (Fig. 3). This cooling is strong enough to allow for an increase of the fraction of solid precipitation and to maintain the snow cover.

When the LIS reaches its full LGM altitude (100dhL-noIS), the positive anomaly of total precipitation over Scandinavia simulated in the 00dhL experiment shifts southward, reaching even the French and the Iberian Peninsula Atlantic coasts. This results in dry conditions over the western flank of the Scandinavian region, both in terms of total and solid precipitation. Concomitantly, the negative anomaly of total precipitation over the Barents-Kara Sea region

simulated in 00dhL expands westward leading to a drier climate and to a decrease of snowfall over Scandinavia. These results suggest that the Barents-Kara Seas and the Scandinavian regions are sensitive to different characteristics of the LIS: whereas the precipitation over Scandinavia appears to be sensitive to the LIS height, the precipitation anomaly over the Barents-Kara Seas mainly appears to result from the LIS albedo effect and is rather insensitive to height.

#### 3.2.2 Large-scale vs convective precipitation

To disentangle the mechanisms responsible for precipitation changes in these regions we split up the total precipitation into its large-scale and convective components (Fig. 7). The large-scale component behaves similarly to the total precipitation shown in Figure 6 with a positive anomaly in the 00dhL experiment simulated over Scandinavia and the eastern North Atlantic sector and the southward shift of this pattern simulated when the LIS gets higher. Hence the drier signal over the western flank of Scandinavia obtained with the full-LGM LIS (100dhL-noIS) can also be found in the large-scale component of the total precipitation. This strongly suggests that the Scandinavian precipitation, as expected, is driven by large-scale processes.

The negative anomaly of precipitation appearing in the Arctic Ocean north of the Barents-Kara area (Fig. 6) is also clearly visible in the large-scale component (Fig. 7); by contrast, the large negative precipitation anomaly south of the Novaya Zemlya clearly arises from convective processes. Over the Barents-Kara area, the origin of the precipitation signal is less clear but seems to be a combination of both convective and large-scale processes. The decrease of convective precipitation negative anomaly (w.r.t noIS) over the entire SBK sector when the LIS is higher can be easily explained by the weakening of the summer cooling discussed in section 3.1.

To further investigate the mechanisms governing precipitation over Scandinavia and the Barents-Kara area, we examine the seasonality of precipitation. We first compute the average precipitation above Scandinavia (55-70°N: 0-20°E) (Fig. 8a) and the Barents-Kara region (65-90°N: 20-100°E) (Fig 8b), then we compare summer and winter precipitation anomalies with the annual precipitation anomaly. Results displayed in Figure 8 show that over Scandinavia winter precipitation changes are clearly correlated with annual precipitation (r = 0.99). This means that the annual signal observed over Scandinavia (Figs 6 and 7) is mainly due to winter precipitation. Conversely, the Barents-Kara sector is rather governed by summer precipitation

(at least for LIS altitudes up to 60% of the PMIP3 one), as shown by the positive correlation (r = 0.98) obtained for the summer season (Fig. 8b).

# 3.2.3 Precipitation changes over Scandinavia: implication of the jet stream shift

- We therefore look for a mechanism prevailing in winter to explain the Scandinavian
- precipitation differences. Using the PMIP3 model outputs for the LGM and the PI time slices,
- Beghin et al. (accepted) found a correlation between the southward shift of the North Atlantic
- jet stream from the pre-industrial to the LGM periods and the winter precipitation changes over
- the Iberian Peninsula. Following the same approach, we examine whether a similar mechanism
- 304 to the one found in (Beghin et al. accepted) may explain the precipitation changes over
- 305 Scandinavia when the LIS characteristics (albedo and topography) are modified.
- Similarly to Chavaillaz et al. (2013), we use the 850 hPa zonal wind to define the jet stream.
- The position of the jet displayed in Fig. 9 corresponds to the position of the 850 hPa meridional
- wind maximum. The LIS albedo effect induces a slight northward shift (w.r.t. noIS) of the
- North Atlantic jet stream (00dhL-noIS panel). As the LIS becomes higher, the jet moves
- progressively southward and shifts southward of its noIS position between 30dhL and 60dhL.
- 311 From 60dhL and beyond, the Scandinavian winter precipitation anomaly becomes more and
- more negative. This suggests a relationship between the shift of the North Atlantic jet stream
- and the precipitation over the Scandinavian region.

- To confirm this assumption, we plot the shift of the jet  $(\Delta \phi)$  as a function of the precipitation
- 315 difference between the noIS and the xxdhL experiments over Scandinavia (Fig. 10). Δφ is
- defined as the difference of latitudes between noIS and xxdhL experiments where the 850 hPa
- zonal wind is at its highest value. The latitude of this maximum has been found by computing
- 318 the zonal mean of the zonal wind over the Atlantic basin  $(50^{\circ}W 10^{\circ}W)$ . To minimize the
- 319 uncertainties due to model discretization, we used a parabolic interpolation around the grid
- point where the mean zonal wind is largest. The maximum of the corresponding parabola yields
- a better approximation of the exact latitude of the maximum jet wind. This method allows to
- find more accurately the latitude of the jet maximum (Chavaillaz et al. 2013; Beghin et al.
- accepted). We find a good linear correlation between the shift of the North Atlantic jet and the
- 324 Scandinavian precipitation anomaly during the winter season (Fig. 10). The quantification of
- 325 the range of the precipitation anomalies and of the jet shift is obtained by bootstrapping: a
- sample of thousand values is obtained by calculating the average of 1000 randomly picked
- samples of 40 year-duration in the noIS and the xxdhL original samples. Linear regressions of

- the 5th and 95th percentiles of the bootstrapping sample confirm the close link between the shift of the North Atlantic jet stream and the winter precipitation changes over Scandinavia due to the effect of LIS.
- The 500 hPa geopotential height (z500) zonal anomaly over the East Laurentide sector (90°W-60°W) does not change significantly when a flat LIS is imposed (Figs 5, 11). As the LIS gets
- higher, the Labrador trough expands southward: figure 11 shows that the zero isoline is located
- at ~43°N in the noIS and 00dhL experiments and progressively reaches 37°N for a full-LGM
- 335 LIS. Over the Atlantic sector (50°W-10°W) the Icelandic low / Azores high pattern remain
- similar in all experiments up to 60 dhL and only show a significant change in the full LIS
- experiment (Figs 5, 11). Indeed, in 100dhL, the Icelandic low's amplitude weakens but
- simultaneously expands southward (Fig. 11). Consistently with these changes in the z500 zonal
- anomaly, the 500hPa westerlies shift southward over the whole Atlantic sector (Fig; 11,
- bottom). Again, the largest southward shift for this variable is between the 60dhL and 100dhL
- experiments, showing the non-linearity of the atmospheric circulation to the LIS height.

# 3.2.4 Precipitation changes over the Barents-Kara area

- 343 As previously mentioned in section 3.2.2, the mean annual precipitation changes over the
- Barents-Kara area are dominated by the summer signal when the LIS remains at a relatively
- low level (i.e. up to 60dhL). This result is confirmed by the comparison of the patterns of
- annual (Fig. 6), summer and winter (Fig 9) precipitation anomalies. The amount of summer
- precipitation is at its lowest level when the sole albedo effect of the LIS is accounted for. It
- increases with the LIS altitude in response to warmer temperatures (Fig; 3). Conversely, the
- summer snowfall amount decreases from 00dhL to 100dhL due to the progressive lowering of
- 350 the fraction of solid vs liquid precipitation.

#### 3.3 Summary

342

- 352 To summarize, the LIS albedo effect induces colder summer temperatures over the entire
- northwestern Eurasian region (SBK area) w.r.t. the noIS experiment. Over South Scandinavia,
- 354 the winter temperatures (Fig. 3) are warmer due to weaker northerlies (Fig. 5). More
- precipitation is also simulated in winter in response to the poleward shift of the North Atlantic
- jet stream (Fig. 6). As the Barents-Kara sector is dominated by the summer signal, less
- precipitation is simulated in response to colder summer temperatures (w.r.t. noIS), while a
- 358 larger snowfall amount is obtained.

When the LIS gets higher the negative anomaly of summer temperatures over the Barents-Kara region becomes weaker due to changes in the atmospheric circulation. In addition, the higher the LIS, the smaller the amounts of annual precipitation and snowfall over the Scandinavian and the Barents-Kara areas, in response to the equatorward shift of the jet.

# 4. What is the influence of the LIS on the growth of the FIS?

#### 4.1 In case of no initial Fennoscandian ice sheet

What are the impacts of the LIS on the surface mass balance of both Scandinavian and Barents-Kara areas? To investigate this question, we use the LMDZ5 monthly temperatures and precipitation fields of each experiment to force the ice-sheet model GRISLI. LMDZ5 climatic fields are downscaled over the GRISLI grid using a bilinear interpolation. Due to the difference of resolution between the atmospheric and the ice-sheet models, temperature is vertically corrected using a linear vertical gradient of 6 K/km. Precipitation is also vertically corrected using an exponential function of the temperature (Charbit et al. 2002; Charbit et al. 2007). Snowfall is recalculated using the downscaled precipitation and temperature because the more detailed topography of GRISLI allows snowfall when LMDZ5 provides only liquid precipitation. 

Figure 12 displays the snow accumulation and the ablation fields computed by GRISLI for the five selected experiments. As expected from the LMDZ5 results of annual precipitation and snowfall, snow accumulation over Scandinavia is larger when the LIS is flat and decreases as the LIS is higher. A similar observation can be made over the Barents-Kara region. Consistently with the simulated summer temperatures (Fig. 3), when the LIS is flat, the ablation is weaker over the entire SBK region, and increases as the LIS gets higher. Nevertheless, even with the full-LGM LIS, the ablation remains weaker than in the noIS experiment.

The resulting surface mass balance is shown in Fig. 13a. The similarities between the ablation (Fig. 12b) and the surface mass balance patterns indicate that the surface mass balance is dominated by the ablation. In the absence of LIS, the surface mass balance is positive over only a small part of the SBK area, namely over Svalbard. Under these conditions, no ice sheet grows (Fig. 13b). When the LIS is flat, the surface mass balance is positive over the Barents-Kara Seas and over the northern part of Scandinavia, allowing the growth of ice. The resulting FIS is less extended than those provided by the ICE-6G (Toscano et al. 2011) and the ANU (Lambeck 2001) reconstructions. This is likely due to the absence of the albedo feedback, since our approach is not based on a full coupling method and does not allow to account for the impact of

ice-sheet changes onto the climate. As the LIS gets higher, the limit of the positive surface mass balance shifts westward and northward, excluding the Kara Sea and Scandinavia from the positive surface-mass balance area. When the LIS has its full LGM size, the surface mass balance is positive only over the Svalbard. As a consequence, the simulated FIS is smaller.

# 4.2 What if the Fennoscandian ice sheet is initially high?

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

The aim of this study is not to provide a fully realistic scenario of the evolution of the ice sheets throughout an entire glacial cycle. Rather, based on idealized cases, it is to investigate the impact of the Laurentide ice sheet on the Eurasian ice sheet surface mass balance. In the first part of this study, we have shown that the growth of the LIS prevents the development of the FIS. However, back to a more realistic context in which both ice sheets co-evolved, the relevance of this conclusion may be questioned and the issue of the FIS inception under full LGM LIS conditions may appear somehow artificial. Thus, to make our previous conclusions more robust, we perform two additional LMDZ experiments with new ice-sheet boundary conditions. In both simulations, the FIS has the elevation computed by the ice-sheet model for the 00dhL run (i.e. the largest FIS elevation of all the GRISLI experiments), while the LIS is either "flat" (00dhLFIS) or has its full LGM altitude (100dhLFIS). Here, our objective is to test the self-consistency of the overall study, by checking that the growth of the FIS is slowed down when the LIS gets higher. We first examine how a high FIS modifies the northwestern Eurasian climate under 00dhL and 100dhL conditions (i.e. with flat or full LGM LIS ice sheet), in terms of the temperature and precipitation fields used in our modelling approach to compute the icesheet surface mass balance. We then use the climatic outputs to force the ice-sheet model.

#### **4.2.1** Response of the atmosphere

Figure 14 displays the summer surface air temperature, which is the main driver of the ablation, for the 00dhL (flat LIS, no FIS), 00dhLFIS (flat LIS, high FIS), 100dhL (high LIS, noIS) and 100dhLFIS (high LIS, high FIS) experiments. The comparison between Figs 14a and 14b and Figs 14c and 14d illustrate the impact of the Fennoscandian ice sheet on the 2m summer temperature. As expected, the presence of Fennoscandia induces a large cooling over the ice sheet itself, due to the temperature-altitude relationship and to the albedo effect. In previous sections, we have shown that the cooling observed over the Barents-Kara region is weakened when the LIS gets higher due to a modification of the large-scale atmospheric circulation (Figs 3-4). A similar mechanism is observed when the Fennoscandian ice sheet is high (100dhLFIS experiment): the cooling over the ice sheet is reduced and the warming along the southern

margin is amplified (Fig. 14d vs Fig. 14b). These features result from an increase of warm air advection in the Barents-Kara sector and to a decrease of northerly heat flux in the south. The presence of the FIS also induces a split of the North Atlantic jet stream in two branches resulting in northward deflected heat fluxes along the western margin of the ice sheet and the Barents Sea sector (Figs 15b,d and 16b,d). The heat advection modulates the ice-sheet altitude and albedo effects but is not sufficient to cancel out the cooling observed over the ice sheet.

Figure 17 displays maps of snow accumulation differences (after downscaling on the GRISLI grid) between the four experiments computed by GRISLI after downscaling. The presence of the Fennoscandian ice sheet results in an increase of precipitation and snow accumulation over its western flank, along the Norwegian Sea coast (Figs 17a,c). This is due to the barrier effect of the ice sheet and to the subsequent increase of the vertical wind. This effect is less pronounced when the LIS is high (Fig. 17c) as a result of the southward displacement of the North Atlantic jet stream (Figs 6,11). A second striking feature is the increase of snow accumulation in the southern part of the Eurasian ice sheet with respect to the no FIS situation. This is mainly due to colder temperatures over the ice sheet, increasing the fraction of total precipitation which falls as snow. In case of a flat LIS, this accumulation zone is confined to the southern margin and drier conditions (w.r.t no FIS case) in the interior of the Eurasian ice sheet are observed, mainly located over the Kara Sea (Fig. 17a), and likely due to the strong barrier effect. By contrast, when the LIS is high (Fig. 17c), differences between FIS and no FIS experiments are larger and expand over more than the southern half of the FIS. This coincides with a larger temperature difference over the FIS area when the LIS is high (Figs 14c,d) compared to the low LIS situation (Figs 14a,b).

As mentioned above (section 3.2.1 and Fig. 6), in the absence of FIS, the total precipitation is driven by large-scale processes over the Scandinavian region and by convection over the Barents-Kara sector. When the LIS is high, the southward shift of the winter jet stream (Fig. 9) and the increase of summer temperature lead to an overall decrease of snowfall over the Eurasian region. When the FIS is present, the situation is rather different and convective precipitation is negligible (not shown) due to the cold temperatures induced by the presence of the FIS. The differences between the 00dhLFIS and 100dhLFIS experiments are mainly located over the western and southeastern parts of the ice sheet with respectively larger and smaller snowfall amounts simulated in the 00dhLFIS experiment (w.r.t 100dhLFIS). The excess of precipitation (total and solid) in the flat LIS/high FIS case is a direct consequence of the

increase of both zonal and meridional humidity transports in winter (not shown) favoured by the enhanced strength of the 850 hPa horizontal winds (Fig 18).

#### 4.2.2 Consequences on the simulated Eurasian ice sheets

The climatic outputs simulated in the 00dhLFIS and 100dhLFIS experiments have been used to compute the surface mass balance of the Eurasian ice sheet at the GRISLI resolution and to force the ice-sheet model with the downscaling procedure described in section 4.1. Starting from the initial state provided by the 00dhL GRISLI run with flat (00dhLFIS) or full LGM LIS (100dhLFIS), GRISLI has been integrated during 100 kyr. At the end of the simulation, the simulated ice volume and ice covered area simulated in the 00dhLFIS experiment are respectively 10 and 11 % greater than those obtained with the 100dhLFIS simulation. The differences in simulated ice thickness between both simulations are displayed in figure 19 (00dhLFIS - 100dhLFIS). In most parts of the simulated ice covered areas, ice thickness differences are fully consistent with the snow accumulation signal (Fig. 17). In particular, large positive anomalies (> 500 m) are found in the southwestern sector, that are greater than 1000 m near the ice sheet margin. In the northern part, lower ice thicknesses obtained in the 00dhLFIS experiment correspond to the area of lower snow accumulation (w.r.t 100dhLFIS), as ablation is very small due to summer temperatures largely below 0°C. Surprisingly, another large positive difference is found in the eastern part which is characterized by a lower amount of snow accumulation in case of flat LIS. (w.r.t full LIS). This pattern results from warmer temperatures simulated in the southern and eastern parts of the 100dhLFIS Eurasian ice sheet in response to the decrease of cold air advection (see section 4.2.1). Warmer temperatures found in the south and east of the FIS when the LIS has its full LGM size favour the westward and the northward retreats of the eastern and southern margins of the ice sheet. Since our approach does not account for the feedback of the simulated ice sheets on the climate, the amplitude of the different responses are likely to be underestimated due to an underestimation of the albedo effect. However, this analysis confirms that the growth of the FIS is slowed down when the LIS is high and demonstrates the self-consistency of the overall study.

#### 5. Discussion

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

#### 5.1. Glaciation scenario

Geomorphological reconstructions revealed that the Fennoscandian ice sheet eastern limit reached the Putorana Mountains in Siberia during the Early Weichselian (90-80 kyr BP) (Mangerud et al. 1998; Svendsen et al. 2004). According to Svendsen et al. (2004), this eastern

487 limit moved back to the Kara sea around 60 kyr BP, along with the southward extension of the Scandinavian ice sheet (Andersen and Mangerud 1989; Lundqvist 1992; Mangerud et al. 1998; 488 489 Houmark-Nielsen 1999). At the LGM, the ice sheet finally reached the British Isles, and the 490 Kara ice shelf did not cover the entire Kara sea anymore (Landvik et al. 1998; Ehlers et al. 491 2004; Svendsen et al. 2004). More recently, Kleman et al. (2013) combined geological 492 evidences and numerical modelling to reconstruct the paleo-topography of past Northern hemisphere ice sheets during MIS5b (~ 86 kyr), and MIS4 (~64 kyr) and 2 (LGM). They found 493 that the LIS and the FIS grow in concert in the early part of the last glacial cycle (i.e. until 494 495 MIS5b), while after MIS4, their evolution is radically different with a further growth of the LIS and a significant shrinkage of the FIS due to its southwestward retreat towards the main 496 497 precipitation sources.

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

Löfverström et al. (2014) propose to attribute the westward shift of the FIS and the decrease of the Kara ice sheet to warm temperatures over the southeastern part of the Fennoscandian region, induced by the FIS itself, along with the upslope precipitation effect proposed by Sanberg and Oerlemans (1983). In their study, the LIS exerts only a small influence on the retreat of the Kara ice sheet. While Löfverström et al. (2014) used realistic LIS topographies provided by the Kleman's et al. (2013) reconstructions, we used idealized geometries to study separately the impact of the LIS albedo and topography on climate. The results of our first series of xxdhL experiments clearly show that a large but low LIS favors a positive SMB over the SBK sector (Fig. 13, left panels) and therefore the FIS maintenance (Fig.13, right panels). By contrast, as the LIS gets higher, ablation increases leading to a smaller FIS. Back to a more realistic glacial context in which both North American and Eurasian ice sheets exist (xxdhLFIS experiments), we also simulate a net decrease of snow accumulation under full LGM LIS conditions (w.r.t flat LIS) over the Kara sector combined with an increase over the western part of FIS (Fig. 17d), consistently with the upslope precipitation effect proposed by Sandberg and Oerlemans (1983). The large positive SMB values over the SBK sector when the LIS is low combined with the westward displacement of the Eurasian ice sheet as the LIS gets higher are fully in line with the ice-sheet paleo-topography reconstructions provided by Kleman et al. (2013) for the last glacial cycle. The remote interactions highlighted in the present study between the LIS and the FIS systems also suggest that the ablation increase over the Barents-Kara Seas when the LIS has its LGM size may have contributed to the LGM FIS small size. Conversely, the larger size of the FIS (Svendsen et al. 2004) during the late Saalian (around 140 ka) w.r.t. the LGM may have been related to a smaller LIS.

- The remote interactions highlighted in the present study between the LIS and the FIS systems
- also suggest that the larger size of the FIS (Svendsen et al. 2004) during the late Saalian
- (around 140 ka) w.r.t. the LGM may have been related to a smaller LIS.

#### 5.2 Oceanic effects

- Our experimental setup has been designed to investigate how the atmospheric circulation
- changes in response to the LIS albedo and topography effects may impact the climate over the
- 526 Eurasian region and therefore the evolution of the surface mass balance of the Fennoscandian
- ice sheet. We have therefore neglected the potential impact of the ocean.
- First of all, all experiments presented in this study have used the LGM land-sea mask and
- therefore the Barents-Kara region is considered as a continental area. The consequences of this
- assumption on the simulated large-scale atmospheric circulation are likely to be negligible.
- However, accounting for a marine Barents-Kara area would decrease the positive albedo
- feedback in this region, and thus the FIS inception (see section 3.1.1). Our experimental setup
- uses LGM SST and sea ice boundary conditions provided by the coupled atmosphere-ocean
- 534 IPSL model (Kageyama et al. 2013). This prevents us from studying the conditions that could
- have prevailed with a marine Barents-Kara area, because this sector was covered by the FIS in
- this run, and future studies will be necessary to investigate further the relationships between the
- 537 LIS and the FIS within more realistic contexts.
- Secondly, the potential impacts of oceanic circulation changes have also been ignored. Yet,
- changes in atmospheric circulation related to the presence of continental ice sheets may induce
- large modifications of the oceanic circulation. For example, Wunsch (2006) proposed a
- scenario in which phase shifts of the stationary waves caused by the development of ice sheets
- 542 modify the wind direction and strength leading to abrupt changes in temperature and
- 543 precipitation fields (and thus to climatic events having similar features as Dansgaard-Oeschger
- events). These changes eventually produce large-scale modifications of the oceanic circulation.
- In turn, any change in the large-scale oceanic circulation may influence the evolution of the
- Northern hemisphere ice sheets as well as their mutual interactions through the atmosphere-
- 547 ocean coupling.
- Recently, using the Earth system model COSMOS, Zhang et al. (2014) found that gradual
- increase of the Northern hemisphere ice sheet height is able of causing abrupt transitions of the
- 550 AMOC through a positive atmosphere-ocean-sea-ice feedback resulting in an adjustment of
- heat distribution and sea ice coverage in the North Atlantic ocean. In this study, abrupt AMOC
- 552 transitions from a weak to a strong mode, occurring for intermediate ice sheet sizes, are

accompanied by a noticeable increase of surface air temperatures in the northeastern North Atlantic. The warming effect simulated in our study over the Eurasian region when the LIS gets higher could be therefore amplified by the North Atlantic warmer temperatures. However, the mechanism highlighted in Zhang et al. (2014) is primarily driven by the simulated northward shift of the westerlies when the Northern hemisphere ice sheet altitude gradually increases. On the contrary, we simulate a southward displacement of the jet stream when the height of the Laurentide ice sheet increases. Also, using the IPSL-CM4 model, Pausata et al (2011) showed that the position of the North Atlantic jet stream is not substantially influenced by sea-surface temperatures or sea-ice coverage. In the same way, using the CAM3 atmospheric model coupled to a slab ocean, Löfverström et al. (2014) did not find any significant changes in the simulated temperature patterns over Alaska, Asia and the North Atlantic sector when two different oceanic heat transports corresponding to interglacial and LGM conditions are used. The diversity of these models'responses to changes in the LIS characteristics show that they are highly model-dependent. This underlines the need for investigating more in-depth the links between large-scale oceanic and atmospheric circulations with different models run under climatic conditions rather different from those of the present-day periods. In fact, the recent inter-comparison study of the North Atlantic jet stream in LGM vs pre-industrial conditions in the PMIP3 models (Braconnot et al. 2012) carried out by Beghin et al. (accepted) clearly shows that the response of the jet stream to LGM conditions is highly model-dependent. The range of model results found in this latter study is confirmed by the range of various results found in the recent literature. Using the outputs of four comprehensive atmosphere-ocean models, Laîné et al. (2009) found a southward displacement of the North Atlantic jet stream under LGM conditions. A similar conclusion can be drawn from the study of Ullman et al. (2014) carried out with the GISS model and two different LIS topographies as LGM ice-sheet boundary conditions: the difference of the simulated jet speed obtained with these different LIS reconstructions (see their figure 2) shows a more southward position of the jet when the LIS is higher. Similarly to our study, Löfverström et al. (2014) found an equatorward position of the winter zonal winds and a more zonal jet under LGM conditions (w.r.t. MIS5b and MIS4). On the contrary, Roberts et al. (2014) found that a more zonal structure is simulated with the FAMOUS model after the occurrence of a Heinrich event, and thus after a decrease of the Laurentide ice thickness. This indicates a southward shift of the North Atlantic jet stream when the LIS altitude is lower, and, by contrast, a northward displacement when it is higher. All these contrasted responses call for further studies to determine why the results are so modeldependent.

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

#### 6. Conclusions

- The goal of this study was to investigate the atmospheric processes through which the LIS can
- influence the northwestern Eurasian climate, and therefore how it can influence the FIS surface
- 590 mass balance. We performed twelve simulations under LGM conditions, but with different ice-
- sheet configurations. In our baseline experiment, the LIS and the FIS are removed. In the first
- set of simulations, we only removed the FIS, and the LIS has different heights. In the final
- series of experiments, we considered the impact of the LIS height on a high FIS. We therefore
- identified the effect of the LIS albedo and topography on the northwestern European climate.
- We focused on the fields having an impact on ice-sheet surface mass balance, i.e. the
- temperature and the precipitation. We showed that:
- 597 1/ The LIS albedo acts to decrease the summer temperatures over the Fennoscandian sector.
- 598 This decrease is amplified by positive snow albedo feedback. This positive feedback is
- somewhat related to our experimental setup, which imposes land over Barents-Kara instead of
- water, allowing snow maintenance. The temperature decrease is weaker when the LIS is higher,
- due to atmospheric circulation changes.
- 602 2/ In winter, the LIS albedo impact decreases the Barents-Kara temperature, but increases the
- Scandinavian temperature. The higher the LIS, the colder the temperature over the whole
- 604 Fennoscandian sector.
- 605 3/ The effect of LIS albedo also tends to shift the North Atlantic jet stream poleward, and to
- bring more precipitation and snowfall over Scandinavia. When the LIS is higher, the Atlantic
- jet shifts equatorward, bringing less precipitation over northern Europe. As a consequence,
- precipitation and snowfall decrease over Scandinavia and Barents-Kara sectors when the LIS is
- 609 higher.
- 610 4/ The analysis of the GRISLI simulations forced by the LMDZ precipitation and temperature
- fields shows that the LIS albedo favors the growth of the FIS, essentially because of weak
- ablation. As the LIS gets higher, the FIS gets smaller due to more ablation, in accordance with
- 613 the LMDZ5 summer temperature response. When the LIS reaches its full-LGM size, ice
- 614 remains over Svalbard only.
- 615 5/ Within the context of glacial-interglacial cycles, the FIS inception under full-LGM LIS
- conditions appears to be unrealistic. Therefore, we performed additional simulations with a
- 617 high FIS and flat and full-LGM LIS. The results of these new sensitivity experiments show a

similar response to the first series of experiments: in the presence of a full LIS, the FIS undergoes a significant mass loss. This confirms that a large LIS counteracts the growth of the FIS although it is still maintained, such as it does within more realistic contexts, and highlights therefore the self-consistency of the overall study.

This study highlights the mechanisms by which the LIS can influence the FIS surface mass balance. The use of the ice-sheet model allows to quantify the impact of climatic changes due to the LIS presence on the FIS. It shows that the albedo of the LIS favors the FIS growth whereas the LIS topography acts against the FIS development. This conclusion is reinforced by the results obtained with a high FIS. It is worth noting that without a transient simulation of the last glacial period, it is not possible to accurately elaborate the lead-lag relationships of the Northern hemisphere ice sheets during the last glacial period. However, based on our results, a plausible glaciation scenario can be proposed with an early phase of FIS glacial inception favored by the areal expansion of the LIS, and followed by a limited growth of the FIS with a progressive westward retreat of the FIS itself as the LIS altitude gradually increased. Such proposed relationships between ice-sheet build-up demonstrate that this study also provides new insights in the understanding of the various Northern hemisphere ice sheet configurations during the different glacial-interglacial cycles.

# **Acknowledgments:**

The authors would like to thank Daniel Lunt, Francis Codron, David Salas-y-Mélia and Hubert Gallée for fruitful discussions about this work. They are also very grateful to two anonymous reviewers for their constructive comments and suggestions that helped to improve the quality and the clarity of this manuscript. Pauline Beghin received a grant from CEA (Commissariat à l'Energie Atomique et aux Energies Alternatives) and this work was supported by the Université Versailles-Saint-Quentin. This work also benefited from the HPC resources of CCRT made available by GENCI (Grand Equipement National de Calcul Intensif), CEA and CNRS (Centre National de la Recherche Scientifique). 

#### References

- Abe-Ouchi A., Saito F., Kageyama M., Braconnot P., Harrison S. P., Lambeck K., Otto-Bliesner B. L.,
  Peltier W. R., Tarasov L., Peterschmitt J.-Y. and Takahashi K., Ice-sheet configuration in the
  CMIP5/PMIP3 Last Glacial Maximum experiments, Geoscience Model Development
  Discussions, 8, 4293-4336, doi: 10.5194/gmdd-8-4293-2015, 2015.
- Andersen B. G. and Mangerud J.: The last interglacial-glacial cycle in Fennoscandia, Quaternary International, 3, 21-29, 1989.

- Beghin P., Charbit S., Dumas C., Kageyama M., Roche D. M. and Ritz C.: Interdependence of the growth of the Northern Hemisphere ice sheets during the last glaciation: the role of atmospheric circulation, Climate of the Past, 10, 345–358, 2014.
- Beghin P., Charbit S., Kageyama M., Combourieu-Nebout N., Hatte C., Dumas C. and Peterschmitt J.-Y.: What drives LGM precipitation over the Western Mediterranean? A study focused on the Iberian peninsula and northern Moroco, Climate Dynamics, in press.
- Berger A.: Long-term variations of daily insolation and quaternary climatic changes, Journal of Atmospheric Sciences, 35, 2362-2367, 1978.
- Braconnot P., Otto-Bliesner B., Harrison S., Joussaume S., Peterchmitt J.-Y., Abe-Ouchi A., Crucifix M., Driesschaert E., Fichefet T., Hewitt C. D., Kageyama M., Kitoh A., Laîné A., Loutre M.-F., Marti O., Merkel U., Ramstein G., Valdes P., Weber S. L., Yu Y. and Zhao Y.: Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum. Part 1: experiments and large-scale features, Climate of the Past, 3, 261–277, doi: 10.5194/cp-3-261-2007, 2007.
- Braithwaite R. J.: Calculation of degree-days for glacier-climate research, Zeitschrift für Gletscherkunde und Glazialgeologie, 20, 1-18, 1984.
- Braithwaite R. J.: Positive degree-day factors for ablation on the Greenland ice sheet studies by energy-balance modelling, Journal of Glaciology, 41(137), 153-160, 1995.
- Broccoli A. J. and Manabe S.: The influence of continental ice, atmospheric CO<sub>2</sub>, and land albedo on the climate of the last glacial maximum, Climate Dynamics, 1, 87-99, 1987.
- 671 Charbit S., Ritz C., Philippon G., Peyaud V. and Kageyama M.: Numerical reconstuctions of the 672 Northern Hemisphere ice sheets through the last glacial-interglacial cycle, Climate of the Past, 3, 673 15-37, 2007.
- 674 Charbit S., Ritz C. and Ramstein G.: Simulations of Northern Hemisphere ice-sheet retreat: sensitivity to 675 physical mechanisms involved during the Last Deglaciation., Quaternary Science Reviews, 21, 676 243-265, 2002.
- Chavaillaz Y., Codron F. and Kageyama M.: Southern westerlies in LGM and future (RCP4.5) climates,
   Climate of the Past, 9, 517-524, doi: 10.5194/cp-9-517-2013, 2013.
- 679 Clark P. U.: Northern Hemisphere Ice-Sheet Influences on Global Climate Change, Science, 286(5442), 680 1104-1111, 1999.
- Clark P. U., Clague J. J., Curry B. B., Dreimanis A., Hicock S. R., Miller G. H., Berger G. W., Eyles N.,
   Lamothe M., Miller B. B., Mott R. J., Oldale R. N., Stea R. R., Szabo J. P., Thorleifson L. H. and
   Vincent J.-S.: Initiation and developement of the Laurentide and Cordilleran ice sheets following
   the last interglaciation, Quaternary Science Reviews, 12, 79-114, 1993.
- 685 Dufresne J.L., Foujols M.A., Denvil S., Caubel A., Marti O., Aumont O., Balkanski Y., Bekki S., Bellenger H., Benshila R., Bony S., Bopp L., Braconnot P., Brockmann P., Cadule P., Cheruy F., 686 Codron F., Cozic A., Cugnet D., De Noblet N., Duvel J.P., Ethé C., Fairhead L., Fichefet T., 687 Flavoni S., Friedlingstein P., Grandpeix J.Y., Guez L., Guilyardi E., Hauglustaine D., Hourdin F., 688 Idelkadi A., Ghattas J., Joussaume S., Kageyama M., Krinner G., Labetoulle S., Lahellec A., 689 Lefebvre M.P., Lefevre F., Levy C., Li Z.X., Lloyd J., Lott F., Madec G., Mancip M., Marchand 690 M., Masson S., Meurdesoif Y., Mignot J., Musat I., Parouty S., Polcher J., Rio C., Schulz M., 691 Swingedouw D., Szopa S., Talandier C., Terray P., Viovy N., Vuichard N.: Climate change 692 projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5. Climate 693 Dynamics. 40, 2123-2165, 2013. 694
- Dyke A. S. and Prest V. K. Late Wisconsinan and Holocene retreat of the Laurentide ice sheet. Geological Survey of Canada. Ottawa, Ontario Map 1702A, scale 1701:500000, 1987.
- Ehlers J., Eissmann L., Lippstreu L., Stephan H.-J. and Wansa S.: Pleistocene glaciations of north Germany, Developments in Quaternary Sciences, 2, 135-146, 2004.

- Fausto R. S., Ahlström A. P., Van As D., Johnsen S., Langen P. L. and Steffen K.: Improving surface boundary conditions with focus on coupling snow densification and meltwater retention in large-scale ice-sheet models of Greenland, Journal of Glaciology, 55(193), 869-878, 2009.
- Hall N. M. J., Valdes P. J. and Dong B.: The maintenance of the last great ice sheets: A UGAMP GCM study, Journal of Climate, 9, 1004-1019, 1996.
- Houmark-Nielsen M.: A lithostratigraphy of Weichselian glacial and interstadial deposits in Denmark, Bulletin of the Geological Society of Denmark, 46(1), 101-114, 1999.
- Hourdin F., Musat I., Bony S., Braconnot P., Codron F., Dufresne J.-L., Fairhead L., Filiberti M.-A., Friedlingstein P., Grandpeix J.-Y. and Others: The LMDZ4 general circulation model: climate performance and sensitivity to parametrized physics with emphasis on tropical convection, Climate Dynamics, 27(7-8), 787-813, 2006.
- Hourdin F., Foujols, M.-A., Codron F., Guemas V., Dufresne J.-L., Bony S., Denvil S., Guez L., Lott F.,
   Ghattas J., Braconnot P., Marti O., Meurdesoif Y., Bopp L.: Impact of the LMDZ atmospheric
   grid configuration on the climate and sensitivity of the IPSL-CM5A coupled model, Climate
   Dynamics, 40, 2167-2192, doi:10.1007/s00382-012-1411-3, 2013.
- Hutter K.: Theoretical Glaciology. Dordrecht, Holland, D. Reidel, 1983.
- Kageyama M., Braconnot P., Bopp L., Caubel A., Foujols M.-A., Guilyardi E., Khodri M., Lloyd J., Lombard F., Mariotti V., Marti O., Roy T. and Woillez M.-N. e.: Mid-Holocene and Last Glacial Maximum climate simulations with the IPSL model-part I: comparing IPSL-CM5A to IPSL-CM4, Climate Dynamics, 40(9-10), 2447-2468, 2013.
- Kageyama M. and Valdes P.: Impact of the North American ice-sheet orography on the Last Glacial Maximum eddies and snowfall, Geophysical Research Letters, 27(10), 1515-1518, 2000.
- Kasahara A.: Computational aspects of numerical models for weather prediction and climate simulation,
  General circulation models of the atmosphere.(A 78-10662 01-47) New York, Academic Press,
  Inc., 1-66, 1977.
- Kleman J., Fastook J., Ebert K., Nilsson J. and Caballero R.: Pre-LGM Northern Hemisphere ice sheet topography, Climate of the Past, 9(5), 2365-2378, 2013.
- Laîné A., Kageyama M., Salas-Mélia D., Voldoire A., Rivière G., Ramstein G., Planton S., Tyteca S.
   and Peterschmitt J. Y.: Northern hemisphere storm tracks during the last glacial maximum in the
   PMIP2 ocean-atmosphere coupled models: energetic study, seasonal cycle, precipitation, Climate
   Dynamics, 32, 593–614, doi: 10.1007/s00382-008-0391-9, 2009.
- Lambeck K.: Sea Level Change Through the Last Glacial Cycle, Science, 292(5517), 679-686, 2001.
- Lambeck K., Purcell A., Funder S., Kjaer K., Larsen E. and M\"Oller P.: Constraints on the Late Saalian to early Middle Weichselian ice sheet of Eurasia from field data and rebound modelling, Boreas, 35(3), 539-575, 2006.
- Landvik J. Y., Bondevik S., Elverhoi A., Fjeldskaar W., Mangerud J., Salvigsen O., Siegert M. J.,
   Svendsen J. I. and Vorren T. O.: The last Glacial Maximum of Svalbard and the Barents Sea area:
   ice sheet extent and configuration, Quaternary Science Reviews, 17, 43-75, 1998.
- Treut H., Forichon M., Boucher O. and Li Z.-X.: Sulfate Aerosol Indirect Effect and CO2 Greenhouse Forcing: EquilibriumResponse of the LMD GCM and Associated Cloud Feedbacks, Journal of climate, 11(7), 1673-1684, 1998.
- Le Treut H., Li Z. X. and Forichon M.: Sensitivity of the LMD general circulation model to greenhouse forcing associated with two different cloud water parameterizations, Journal of Climate, 7(12), 1827-1841, 1994.
- Liakka J., Nilsson J. and L\"Ofverstr\"Om M.: Interactions between stationary waves and ice sheets: linear versus nonlinear atmospheric response, Climate Dynamics, 38(5-6), 1249-1262, 2011.

- Löfverström M., Caballero R., Nilsson J. and Kleman J.: Evolution of the large-scale atmospheric circulation in response to changing ice sheets over the last glacial cycle, Climate of the Past, 10(4), 1453-1471, 2014.
- Loulergue L., Schilt A., Spahni R., Masson-Delmotte V., Blunier T., Lemieux B., Barnola J.-M.,
   Raynaud D., Stocker T. F. and Chappellaz J.: Orbital and millennial-scale features of atmospheric
   CH<sub>4</sub> over the past 800,000 years, Nature, 453, doi: 10.1038/nature06950, 2008.
- Lundqvist J.: Glacial stratigraphy in Sweden, Geological Survey of Finland, Special Paper, 15, 43-59,
   1992.
- Lüthi D., Le Floch M., Bereiter B., Blunier T., Barnola J.-M., Siegenthaler U., Raynaud D., Jouzel J., Fischer H., Kawamura K. and Stocker T. F.: High-resolution carbon dioxide concentration record 650,000-800,000 years before present, Nature, 453(7193), 379-382, 2008.
- Macayeal D. R.: Large-scale ice flow over a viscous basal sediment: Theory and application to ice stream B, Antarctica, Journal of Geophysical Research: Solid Earth (1978-2012), 94(B4), 4071-4087, 1989.
- Manabe S. and Broccoli A. J.: The influence of continental ice sheets on the climate of an ice age, Journal of Geophysical Research, 90(D1), 2167-2190, 1985.
- Mangerud J., Dokken T., Hebbeln D., Heggen B., Ingolfsson O., Landvik J. Y., Mejdahl V., Svendsen J.
   I. and Vorren T. O.: Fluctuations of the Svalbard-Barents Sea Ice Sheet during the last 150 000 years, Quaternary Science Reviews, 17(1-3), 11-42, 1998.
- Morland L., Smith G. and Boulton G.: Basal sliding relations deduced from ice sheet data, J. Glaciol, 30(105), 131-139, 1984.
- Pausata F. S. R., Li C., Wettstein J. J., Kageyama M. and Nisancioglu K. H.: The key role of topography
   in altering North Atlantic atmospheric circulation duringthe last glacial period, Climate of teh
   Past, 7, 1089-1101, doi: 10.5194/cp-7-1089-2011, 2011.
- Peltier W. R.: Global glacial isostasy and the surface of the ice-age Earth: the ICE-5G (VM2) model and GRACE, Annual Reviews of Earth and Planetary Science, 32, 111-149, 2004.
- Peyaud V., Ritz C. and Krinner G.: Modelling the early Wechselian Eurasian ice sheets: role of iceshelves and influence of ice-dammed lakes, Climate of the Past, 3, 375-386, 2007.
- Reeh N.: Parameterization of melt rate and surface temperature on the Greenland ice sheet, Polarforschung, 59, 113-128, 1991.
- Ritz C., Rommelaere V. and Dumas C.: Modeling the Antarctic ice sheet evolution of the last 420 000 years: implication for altitude changes in the Vostok region, Journal of Geophysical Research, 106(D23), 31,943-931,964, 2001.
- Rivière G., Laîné A., Lapeyre G., Salas Y Mélia D. and Kageyama M.: Links between Rossby wave breaking and the North Atlantic Oscillation-Arctic Oscillation in present-day and Last Glacial Maximum simulations, Journal of Climate, 23, 2987-3008, doi: 10.1175/2010JCLI3372.1, 2010.
- Roberts, W. H. G., Valdes, P. J., Payne, A. J.: Topography's crucial role in Heinrich events, Proceedings of the National Academy of Sciences, doi: 10.1073/pnas.1414882111, 2014.
- Roe G. H. and Lindzen R. S.: The Mutual Interaction between Continental-Scale Ice Sheets and Atmospheric Stationary Waves, Journal of Climate, 14(7), 1450-1465, 2001a.
- Roe G. H. and Lindzen R. S.: A one-dimensional model for the interaction between continental-scale ice sheets and atmospheric stationary waves, Climate Dynamics, 17, 479-487, 2001d.
- Sadourny R.: Compressible model flows on the sphere, Journal of the Atmospheric Sciences, 32(11),
   2103-2110, 1975a.
- Sadourny R.: The dynamics of finite-difference models of the shallow-water equations, Journal of the Atmospheric Sciences, 32(4), 680-689, 1975b.
- Sadourny R. and Laval K.: January and July performances of the LMD general circulation model. New persepectives in climate modelling, Dev. Atmos. Sci. A. Berger and C. Nicolis: 173-198, 1984.

- Sanberg J. and Oerlemans J.: Modelling of pleistocene European ice sheets: the effect of upslope precipitation, Geologie en Mijnbouw, 62, 267-273, 1983.
- Spahni R., Chappellaz J., Stocker T. F., Loulergue L., Hausammann G., Kawamura K., Flückiger J.,
   Schwander J., Raynaud D., Masson-Delmotte V. and Jouzel J.: Atmospheric methane and nitrous
   oxide of the late Pleistocene from Antarctic ice cores, Science, 310, 1317-1320, 2005.
- 798 Storch von H. and Zwiers, F. W., Statistical Analysis in Climate Research, Cambridge University Press, 2003.
- Svendsen J. I., Alexanderson H., Astakhov V. I., Demidov I., Dowdeswell J., Funder S., Gataullin V.,
  Henriksen M., Hjort C., Houmark-Nielsen M., Hubberten H. W., Ingolfsson O., Jakobsson M.,
  Kjaer K. H., Larsen E., Lokrantz H., Lunkka J. P., Lysa A., Mangerud J., Matiouchkhov A.,
  Murray A., Möller P., Niessen F., Nikolskaya O., Polyak L., Saarnisto M., Siegert C., Siegert M.
  J., Spielhagen R. F. and Stein R.: Late Quaternary ice sheet history of Northern Eurasia,
  Quaternary Science Reviews, 23, 1229-1271, 2004.
- Tarasov L., Dyke A. S., Neal R. M. and Peltier W. R.: A data-calibrated distribution of deglacial chronologies for the North American ice complex from glaciological modeling, Earth and Planetary Science Letters, 315-316, 30–40, doi: 10.1016/j.epsl.2011.09.010, 2012.
- Toscano M. A., Peltier W. R. and Drummond R.: ICE-5G and ICE-6G models of postglacial relative sea-level history applied to the Holocene coral reef record of northeastern St Croix, U.S.V.I.: investigating the influence of rotational feedback on GIA processes at tropical latitudes, Quaternary Science Reviews, 30, 3032-3042, doi: 10.1016/j.quascirev.2011.07.018, 2011.
- Ullman D. J., Legrande A. N., Carlson A. E., Anslow F. S. and Licciardi J. M.: Assessing the impact of Laurentide Ice Sheet topography on glacial climate, Climate of the Past, 10, 487-507, doi: 10.5194/cp-10-487-2014, 2014.
- 816 Wunsch C., Abrupt climate change: An alternative view, Quaternary research, 65, 191-203, 2006.

819


Zhang X., Lohmann G., Knorr G. and Purcell C.: Abrupt glacial climate shifts controlled by ice sheet changes, Nature, 512, 290-297, doi: 10.1038/nature13592, 2014.

#### 820 Figure captions

- Figure 1: Ice-sheet and oceanic boundary conditions used for the LMDZ experiments.
- a) Ice-sheet height (light blue scale) and land height (brown scale) given as boundary conditions
- for the five selected experiments out of twelves. Dark blue-brown limits represent the sea-land
- mask, and brown-light blue limits represent the land-ice mask.
- b) Averaged annual sea-surface temperature (shading) and sea-ice extent for boreal summer
- (red contours) and winter (blue contours) from the PMIP3 IPSL-CM5A-LR simulations. The
- averages are computed over 50 years which have been used to create the boundary conditions
- for the LMDZ experiments.
- 829 Figure 2: Laurentide ice sheet height for each LMDZ experiment (with name of the
- 830 corresponding experiment). The height of the LIS is defined as a relative difference between
- PMIP3 Laurentide ice-sheet height and the present-day topography (Δh LIS PMIP3).
- Figure 3: 2m summer (left) and winter (right) temperature without the LIS (noIS) and anomaly
- of the 2-meter temperature between the xxdhL experiments and the noIS experiment: we can
- therefore quantify the impact of the LIS presence, depending on its topography. The visible
- 835 surface albedo difference between xxdhL and noIS experiments are represented by orange
- contours (albedo difference = 0.4) and yellow contours (albedo difference = 0.2). The statistical
- 837 significance of the patterns of temperature anomalies has been checked using a two-sided
- Student t-test with a p-value =0.05. The white shading indicates areas where the anomaly is not
- 839 significant.
- Figure 4: 500 hPa summer geopotential height anomaly from zonal mean (left panels) and 500
- hPa horizontal wind (arrows) and meridional wind (shaded, right panels).
- Figure 5: Same as Fig. 4 for the winter season
- 843 Figure 6: Same as Fig. 3 for annual precipitation (left column) and annual snowfall (right
- 844 column).
- Figure 7: Large-scale (left pannels) and convective (right pannels) annual precipitation without
- 846 LIS (noIS experiment, top panels), and anomalies of large-scale and convective precipitation
- between the xxdhL and noLIS experiments. The statistical significance of the patterns of
- temperature anomalies has been checked using a two-sided Student t-test with a p-value = 0.05.
- The white shading indicates areas where the anomaly is not significant.
- 850 **Figure 8:** (a) Scandinavian and (b) Barents-Kara precipitation anomalies (xxdhL noIS)
- during winter (circles) and summer (triangles) versus annual precipitation anomaly. The lower
- the Laurentide ice sheet altitude, the lighter grey the symbols are.
- Figure 9: Left column: same as Fig. 3 for winter precipitation (shaded) and position of the
- North Atlantic jet stream without LIS (dashed) and for the xxdhL experimenst (solid line).

- 855 Right column: summer precipitation (shaded) and snowfall (contour every 0.1 mm/day, blue
- 856 contour for positive values).
- Figure 10: 850 hPa winter jet-stream shift (in latitude) versus winter precipitation anomaly
- over the Scandinavian region. The range of results has been obtained by bootstrapping. Up and
- down triangles represent the 5<sup>th</sup> precipitation percentile versus the 95<sup>th</sup> jet position percentile,
- and the 95th precipitation percentile versus the 5th jet position percentile respectively. Circles
- are the average values and crosses indicate the 5th and 95th percentiles.
- Figure 11: Meridional cross sections of the 500hPa geopotential height anomalie from zonal
- means over the East Laurentide (90°W-60°W, top panel) and the Atlantic sectors (50°W-10°W,
- middle) in meters, and of the 500hPa zonal wind (in m/s) for the Atlantic sector.
- Figure 12: (a) Snow accumulation computed by the GRISLI ice-sheet model for the noIS
- experiment (top panel), and differences of snow accumulation between the xxdhL and the noIS
- experiments (in m/yr). (b) same as (a) for the ablation (in m/yr).
- Figure 13: (a) Surface mass balance at the beginning of the simulation (in m/yr) and (b) surface
- 869 elevation at the end of the simulation computed by the ice-sheet model for the selected
- experiments (in meters).
- Figure 14: Summer 2m temperatures for the a) 00dhL (flat LIS, no FIS) experiment, b)
- 00dhLFIS (flat LIS, high FIS), c) 100dhL (high LIS, no FIS) and d)100dhLFIS (high LIS, high
- 873 FIS). The extents of the ice sheets (blue contours) are superimposed on right panels for
- 874 00dhLFIS and 100dhLFIS runs.
- Figure 15: Same as Fig. 14 for the vertically integrated meridional (shaded) and the horizontal
- heat transport (arrows) in summer. Numerical values are given in 10<sup>10</sup> J m<sup>-1</sup> s<sup>-1</sup>.
- Figure 16: Same as Fig. 14 for the 850hPa meridional wind (shaded) and the 850hPa horizontal
- winds (arrows) for the summer season (in m/s).
- Figure 17: Differences in snow accumulation (in m/yr) computed by the GRISLI ice-sheet
- model between a) 00dhLFIS and 00dhL experiments b) 00dhLFIS and 100dhL experiments, c)
- 100dhLFIS and 100dhL experiments and d) 00dhLFIS and 100dhLFIS experiments.
- Figure 18: Same as Fig 14 for the precipitation (mm/day) and 850 hPa horizontal winds in
- winter.
- Figure 19: Difference of simulated ice thickness of the Eurasian ice sheet between 00dhLFIS
- and 100dhLFIS after 100 kyr (in meters).

# a) Ice-sheet boundary conditions



# b) Annual SSTs and sea-ice extent averaged over 50 years





Figure 1








Figure 3








Figure 6




Figure 7





Figure 8






Figure 10






Figure 11








Figure 14




Figure 15




Figure 16




Figure 17




Figure 18

## 00dhLFIS - 100dhLFIS




Figure 19