

Taxonomic Notes

The supra-generic classification applied here is adapted from Riedel (1967a,b) and Petrushevskaya (1971a,b) and is preferred to more recent alternatives (e.g. De Wever et al., 2001; Suzuki and Aita, 2011) because it is more practical when endeavouring to assign specimens of variable preservation to counting group categories for statistical analysis. In addition to the taxa listed below, the counting group categories include undifferentiated families and the following selected genera: *Ceratocyrtis*, *Cycladophora*, *Eucyrtidium*, *Larcopyle*, *Lithomellissa*, *Lophocyrtis* and *Pseudodictyophimus*. These genus-level categories are used for specimens that cannot be confidently ascribed to one of the species listed below due to poor or incomplete preservation. As all the species ascribed to these genera have high latitude affinities, the genus-level counts are included in the percentage of high latitude taxa individuals present in each sample.

Subclass RADIOLARIA

Order SPUMELLARIA

Family ACTINOMMIDAE Haeckel

Actinommidae n.sp. A sensu Hollis [Plate 1, fig. 1]

Actinommidae n.sp. A, Hollis et al. 1997, p. 45, pl. 1, fig. 18-19.

Medium-sized actinommid with slightly irregular spheroidal cortical shell. Pores numerous, closely spaced, circular in regular hexagonal frames. 8 evenly spaced, slender, bladed radial spines. 1-2 medullary shells.

Genus *AMPHISPHAERA* Haeckel, emend. Petrushevskaya 1975

The genus name is applied in the sense of Petrushevskaya (1975). See Hollis (1997) for further discussion.

***Amphisphaera coronata* group (Ehrenberg), New Group** [Plate 1, fig. 2]

Stylosphaera coronata Ehrenberg 1874, p. 258; 1876, pl. 25, fig. 4; Ogane et al. 2009, pl. 12, fig. 1a-d; Suzuki et al. 2009, p. 244, pl. 1, fig. 13a-b.

Stylosphaera radiosa Ehrenberg 1876, pl. 24, fig. 5.

Ellipsostylus anisoxiphos Clark and Campbell 1942, p. 32, pl. 5, fig. 7, 11.

Lithatractus hederæ Clark and Campbell 1942, p. 33, pl. 5, fig. 3.

Lithatractus pterosphaerus Clark and Campbell 1942, p. 33, pl. 5, fig. 33.

Stylosphaera coronata coronata (Ehrenberg), Sanfilippo and Riedel 1973, p. 520, pl. 1, fig. 13-17, pl. 25, fig. 4; Nishimura 1992, p. 329, pl. 1, fig. 2, pl. 11, fig. 9.

Stylosphaera coronata laevis (Ehrenberg), Sanfilippo and Riedel 1973, p. 520, pl. 1, fig. 19,

pl. 25, fig. 5-6.
Stylosphaera coronata sabaca (Ehrenberg), Sanfilippo and Riedel 1973, p. 521, pl. 1, fig. 18,
 pl. 25, fig. 57-8.
Amphisphaera spp. gr. D. Hollis 1991, p. 71, pl. 3, fig. 20; 1993, p. 319.
Amphisphaera coronata Ehrenberg s.l., Hollis et al. 1997, p. 42.
Stylosphaera coronata Ehrenberg, Suzuki et al. 2009, p. 244, pl. 1, fig. 13.

This species group includes all members of the genus with a subspherical to ellipsoidal outer shell and opposing polar spines of unequal length. Sanfilippo and Riedel (1973) distinguished *A. coronata* s.s. by the shorter polar spine having an arrow-shaped termination. Reillustration of the type specimen (Ogane et al., 2009) indicates that the species may be even more narrowly defined: the shorter polar spine has distinct ridges and increases in width prior to the arrow-like termination. The morphotype Sanfilippo and Riedel (1973) consider typical *S. coronata* compares well with *Lithatractus hederæ* Clark and Campbell. Many of the other bipolar spumellarians described by Clark and Campbell (1942) are included in *A. coronata* group here. However, these narrow definitions do not reflect how the name has been applied in the past (e.g. Nishimura 1992; Suzuki et al., 2009) and so we use a broad definition of the species group that accommodates a wide range of variation. *A. macrosphaera* is distinguished by a spherical outer shell and the shorter spine being no longer than the width of the outer shell. *A. goruna* is distinguished by having at least one additional spine outside the polar plane.
 Cosmopolitan.

***Amphisphaera radiosa* (Ehrenberg) [Plate 1, fig. 3]**

Stylosphaera radiosa Ehrenberg, 1854a, p. 246; 1876, pl. 24, fig. 5; Ogane et al. 2009, pl. 12,
 fig. 3a-4e (non Abelman 1990).
Ellipsoxiphus? sp., Hollis et al. 1997, pl. 1, fig. 3-4.

This distinctive species is characterised by a strongly ellipsoidal outline, thickened outer shell and two polar spines of unequal length, the length of the shorter spine being equivalent to the thickness of the outer shell.

Amphisphaera* aff. *radiosa* (Ehrenberg) [Plate 1, fig. 4a, b] = *Stylosphaera radiosa sensu Abelman

Amphisphaera radiosa (Ehrenberg), Petrushevskaya 1975, p. 570, pl. 2, fig. 18-20; Hollis et al., 1997, p. 43, pl. 1, fig. 5-6 (non Crouch and Hollis 1996, p. 26).
Stylosphaera radiosa Ehrenberg, Abelman 1990, p. 692, pl. 2, fig. 4a, b, c, 7; (?) Caulet 1991, p. 539 (non Ehrenberg 1854).

The species ascribed to *S. radiosa* by Abelman (1990) bears little resemblance to the type specimen reillustrated by Ogane et al. (2009). Abelman (1990) suggests that *A. radiosa* has a more spherical shell than *A. coronata* but this is clearly not the case. *A. radiosa* s.s. has a more ellipsoidal shell than *A. coronata* and is further distinguished by a thicker outer shell and more smoothly tapering polar spines, the shorter one of which is no longer than the thickness of the outer shell. The species referred to *A. aff. radiosa* here has a relatively small but thickened outer shell, which is subspheroidal, and one very short polar spine. Additional short spines or thorns may be present. This species compares well with the species reported by Petrushevskaya (1975) but is only tentatively synonymised with the species described by Abelman (1990) in which the

shorter spine varies in length. The Late Oligocene FAD of this species at ODP Sites 689 and 690 is used to define the base of Abelmann's (1990) *Stylophaera radiosa* zone. The wider utility of this zone is questionable as *A. radiosa* first appears in the late Eocene at DSDP Site 277.

***Amphisphaera spinulosa* (Ehrenberg) = *A. goruna* (Sanfilippo and Riedel)** [Plate 1, fig. 5]
Stylosphaera spinulosa Ehrenberg 1874, p. 259; 1876, pl. 25, fig. 8; Ogane et al. 2009, pl. 12, 6a-f (erroneously recorded as *Stylosphaera liostylus*).

Druppattractus cf. *coronatus* (Squinabol), Dumitrica 1973, p. 787, pl. 6, fig. 4,6; pl. 12, fig. 11.

Stylosphaera goruna Sanfilippo and Riedel 1973, p. 521, pl. 1, fig. 20-22; pl. 25, fig. 9, 10 (? fig. 11); Nishimura 1987, pl. 1, fig. 1; 1992, p. 729, pl. 1, fig. 1; p. 11, fig. 8.

Amphisphaera spinulosa (Ehrenberg), Petrushevskaya 1975, p. 570, pl. 15, fig. 8; Hollis 1991, p. 69, pl. 3, fig. 11-15; 1993, p. 318, pl. 1, fig. 5.

Amphisphaera goruna (Sanfilippo and Riedel). Strong et al. 1995, p. 208, fig. 8G, 9A; Hollis 1997, p. 34, pl. 2, fig. 10, 11; Hollis et al. 1997, p. 42.

With the reillustration of Ehrenberg's type specimen of this species (Ogane et al. 2009) there can be little doubt the *A. goruna* is a junior synonym of *A. spinulosa*. The discontinuous range observed at Mead Stream (Strong et al. 1995; Hollis 1997) and possibly other regions may be a response to climatic changes, with the species appearing to show a preference for the cooler conditions of the Paleocene and the late Middle Eocene to Oligocene. Cosmopolitan.

***Amphisphaera? megapora* (Ehrenberg)** [Plate 1, fig. 6]

Cenosphaera megapora Ehrenberg 1874, p. 218, 1876, pl. 3, fig. 1; Ogane et al. 2009, pl. 3, fig. 1a-c.

Amphisphaera aff. *spinulosa* (Ehrenberg), Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 43, pl. 1, fig. 1,2.

? *Amphisphaera spinulosa* (Ehrenberg) group. Petrushevskaya 1975 (in part), p. 570.

Distinguished from *A. spinulosa* by a spherical cortical shell and the absence of distinct polar spines. All radial spines are of similar length. As *A. spinulosa* was not encountered in Site 281 material (Crouch and Hollis 1996) it seems likely that the form recorded from the Eocene of DSDP Site 280 by Petrushevskaya (1975) is *A. megapora*.

Genus *ANOMALACANTHA* Loeblich & Tappan 1961

***Anomalocantha dentata* (Mast)**

Heteracantha dentata Mast 1910 p. 159, pl. 37, fig. 47,

Anomalocantha dentata (Mast), Benson 1966, p. 170, pl. 5, fig. 10-11.

Anomalocantha dentata (Mast), Petrushevskaya and Kozlova 1979, fig. 219, 427.

Cladococcus eocenica (Petrushevskaya and Kozlova 1979), Suzuki et al. 2009, p. 244, pl. 2, fig. 9.

Genus *STYLOSPHAERA* Ehrenberg, emend. Petrushevskaya 1975

Stylosphaera minor Clark and Campbell group [Plate 1, fig. 7]

Stylosphaera minor Clark and Campbell 1942, p. 27, pl. 5, fig. 1, 2, 2a, 12; Petrushevskaya 1975, p. 569; Caulet 1986, p. 853; 1991, p. 539; Strong *et al.* 1995, p. 209.

Amphisphaera minor (Clark and Campbell), Sanfilippo and Riedel 1973, p. 486, pl. 1, fig. 1-5; pl. 22, fig. 4; Chen 1975, p. 452, pl. 3, fig. 1.

Axoprunum minor (Clark and Campbell), Suzuki *et al.* 2009, p. 241, pl. 4, fig. 5a-6c.

Included in this complex are forms with a moderately large spherical, outer shell with numerous small circular pores, slender conical (rarely bladed) polar spines. Suzuki *et al.* (2009) observe the presence of equatorial beams and on this basis transfer the species to *Axoprunum*. These structures are not evident in the specimens we refer to *S. minor* here. Similar specimens that have equatorial beams characteristic of *Axoprunum* are placed in *A. pierinae* gr. here. Cosmopolitan.

Genus *ZEALITHAPIUM* O'Connor 1999a

Zealithapium mitra (Ehrenberg) [Plate 1, fig. 8]

Cornutella mitra Ehrenberg, 1874, p. 221; 1876, pl. 2, fig. 8.

Lithapium mitra (Ehrenberg). Riedel and Sanfilippo 1970, p. 520, pl. 4, fig. 6, 7; Sanfilippo and Riedel 1973, p. 516; Takemura 1992 (cf.), p. 742, pl. 7, fig. 2.; Hollis *et al.* 1997, p. 45, pl. 1, fig. 20.

Distinguished by its single cortical shell in the shape of an inflated cone, narrowing slightly at the open end. Meshwork is coarse with thorns arising from junctions of pore frames. Cosmopolitan.

Subfamily SATURNALINAE Deflandre

Genus *AXOPRUNUM* Haeckel

Axoprunum bispiculum (Popofsky) [Plate 1, fig. 9]

Stylocontarium bispiculum Popofsky 1912, p. 91, pl. 2, fig. 2; Chen 1975, p. 454, pl. 21, fig. 1-2.

Axoprunum bispiculum (Popofsky), Takemura 1992, p. 741, pl. 1, fig. 1-2; Hollis *et al.* 1997, p. 43, pl. 1, fig. 14; Takemura and Ling 1997, p. 111 (*non* Suzuki *et al.* 2009).

Distinguished from *A. pierinae* by a larger, almost spherical cortical shell and shorter polar spines, which often diverge slightly from main axis. Intermediate forms occur in the early Oligocene. High-latitude.

Axoprunum pierinae (Clark and Campbell) group, New Group [Plate 1, fig. 10, 11]

Lithatractus pierinae Clark and Campbell 1942, p. 34, pl. 5, fig. 25.

Sylatractus pictus Mamedov 1969, p. 99, pl. 2, fig. 4-4a.

Axoprunum pierinae (Clark and Campbell), Sanfilippo and Riedel 1973, p. 488, pl. 1, fig. 6-

12, pl. 23, fig. 3; Petrushevskaya 1975, p. 571; Caulet 1991, p. 537; Takemura. 1992, p. 742, pl. 6, fig. 3-6; Strong *et al.* 1995, p. 208, fig. 10c; Crouch and Hollis 1996, p. 26; Hollis *et al.* 1997, p. 44, pl. 1, fig. 7-13; Takemura and Ling, 1997, pl. 1, fig. 1.
Axoprunum liostylum (Ehrenberg). Petrushevskaya and Kozlova 1972, p. 521, pl. 10, fig. 3; Petrushevskaya 1975, p. 571, pl. 2, fig. 22; Crouch and Hollis 1996, p. 26.
Axoprunum bispiculum (Popofsky), Suzuki *et al.* 2009, p. 241, pl. 1, fig. 3a-b.
 ? *Axoprunum euterpe* (Haeckel), Caulet 1986, p. 852.
non Stylosphaera liostylus Ehrenberg 1875, pl. 25, fig. 2; Ogane *et al.* 2009, pl. 12, fig. 6a-f.

Ellipsoidal cortical shell with 2 conical polar spines, usually of unequal length; typically slender but may be thickened throughout or medially. 1-2 medullary shells supported by 2 strong polar beams and 4 evenly spaced equatorial beams. It is uncertain if this species group comprises a single polymorphic species or several distinct species. Variations in cortical shell thickness and size, polar spine thickness and shape and shape of outer medullary shell may have some ecological significance. Morphotypes with thicker polar spines appear to be restricted to southern high latitude locations (e.g. Takemura 1992, pl. 6, fig. 3-4; Hollis *et al.*, pl. 1, fig. 7; Takemura and Ling 1997, pl. 1, fig. 1; Suzuki *et al.*, 2009, pl. 1, fig. 1), whereas morphotypes with medially inflated polar spines appear to have had a bipolar distribution (e.g. Mamedov 1969, pl. 2, fig. 4-4a; Kozlova 1999, pl. 33, fig. 10, pl. 38, fig. 2; Suzuki *et al.* 2009, p. 1, fig. 2). At Site 277, however, there appears to be a gradation between these specimens with thickened spines, specimens that conform with the type species and specimens with near-spherical cortical shells (Hollis *et al.* 1997, pl. 1, fig. 9). For this reason, we combine this range of morphotypes into a single species group here. This group accommodates the forms ascribed to *A. liostylum* by Petrushevskaya (1975) and probably several of the species ascribed to *Axoprunum* by Petrushevskaya and Kozlova (1972). Ehrenberg's species bears little resemblance to the genus and instead may well be a synonym for *Amphisphaera spinulosa*. No figure is available for *A. euterpe* but Haeckel's (1887, p. 135) description conforms with *A. pierinae*. We include following species from Suzuki *et al.* (2009) in our concept of *A. pierinae* (*A. venustum*, *A. aff. venustum*). Cosmopolitan.

***Axoprunum? irregularis* Takemura** [Plate 1, fig. 12]

Actinomma medusa (Ehrenberg) gr., Petrushevskaya 1975 (in part), p. 568, pl. 2, fig. 6-8.

? *Amphisphaera* sp. Chen 1975, pl. 6, fig. 1.

Axoprunum? irregularis Takemura 1992, p. 742, pl. 3, fig. 8-11; Hollis *et al.* 1997, p. 44, pl. 1, fig. 15; Takemura and Ling, 1997, pl. 1, fig. 2.

Irregular-shaped, almost polygonal cortical shell with two slender, conical polar spines, usually set askew to main axis. Two medullary shells are supported by more than 6 radial bars. Lacks diagnostic features of the genus (ellipsoidal cortical shell, 6 symmetrically disposed internal beams) but provisionally retained in *Axoprunum* because of a possible phylogenetic link with *A. bispiculum* (see Takemura 1992). High-latitude.

Genus SATURNALIS Haeckel 1881, emend. Nigrini 1967

***Saturnalis circularis* Haeckel**

Saturnalis circularis Haeckel 1887, p. 131; Nigrini 1967, p. 24, pl. 1, fig. 9; Chen 1975, p.

454, pl. 24, fig. 2.

Genus *SPONGATRACTUS* (Ehrenberg)

Spongatractus pachystylus (Ehrenberg)

Spongosphaera pachystyla Ehrenberg, 1874, p. 256; 1876, pl. 26, fig. 3.

Spongatractus pachystylus (Ehrenberg), Sanfilippo and Riedel, 1973, p. 519, pl. 2, figs. 4-6; pl. 25, fig. 3.

Thick-walled ellipsoidal shell covered with spongy meshwork, with two robust, bladed polar spines. Our single specimen was in a picked sample (DSDP 277-25, CC).

Family PHACODISCIDAE Haeckel

Genus *HELIODISCUS* Haeckel emend. O'Connor 1999b

= *Astrophacus* Haeckel (see O'Connor, 1999b)

Heliodiscus inca Clark and Campbell

Heliodiscus inca Clark and Campbell 1942, p. 38, pl. 3, fig. 17; Hollis et al. 1997, p. 45, pl. 1, fig. 25, 26.

Heliodiscus hexasteriscus Clark and Campbell 1942 (in part), pl. 3, fig. 15.

Astrophacus inca (Clark and Campbell). Caulet 1986, p. 851; 1991, p. 537.

Lenticular cortical shell with 6-10 moderately long, bladed marginal spines. Surface of cortical shell roughened by raised sepaloid points or thorns at junctions between well-developed hexagonal pore frames. Distinguished from *H. linckiaformis* and *H. hexasteriscus* s.s. by much longer spines.

Genus *PERIPHAENA* Ehrenberg

Periphaena decora Ehrenberg

Periphaena decora Ehrenberg, 1874, p. 246; 1876, pl. 28, fig. 6; Sanfilippo and Riedel 1973, p. 523, pl. 8, fig. 8-10; pl. 27, fig. 2-5; Caulet 1986, p. 853; 1991, p. 538; Takemura 1992, p. 743, pl. 6, fig. 8; Strong et al. 1995, p. 209, fig. 10j; Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 46, pl. 1, fig. 21-24.

Lenticular cortical shell with a marginal girdle of varying width, with or without marginal spines. Surface of cortical shell smooth, with numerous small pores. We include *Heliodiscus pertsus* of Suzuki et al. (2009) in our concept of *P. decora*. Cosmopolitan.

Periphaena heliastericus (Clark and Campbell)

Heliodiscus heliastericus Clark and Campbell, p. 39, pl. 3, fig. 10, 11.

Periphaena heliasteriscus [sic] (Clark and Campbell). Sanfilippo and Riedel 1973, p. 523, pl.

9, fig. 1-6; pl. 27, fig. 8, 9. Caulet 1986, p. 853; 1991, p. 538. Takemura 1992, p. 743, pl. 4, fig. 13. Strong *et al.* 1995, p. 209, fig. 10h. Crouch and Hollis 1996, p. 26.
Periphaena heliastericus (Clark and Campbell), Hollis *et al.* 1997, p. 46
11.

Distinguished from *P. decora* by absence of marginal girdle. Usually has marginal spines. We include *Heliodiscus perplexus* of Suzuki *et al.* (2009) in our concept of *P. heliastericus*. Cosmopolitan.

Family SPONGODISCIDAE Haeckel

Genus *PLECTODISCUS* Kozlova

Plectodiscus circularis (Clark and Campbell)

Porodiscus circularis Clark and Campbell 1945, p. 42, pl. 11, fig. 2, 6, 10.

Plectodiscus circularis (Clark and Campbell), Petrushevskaya and Kozlova 1972, p. 526, pl. 19, fig. 9-12; Hollis 2002, p. 288, pl. 2, fig. 20-22.

Xiphospira circularis (Clark and Campbell), Sanfilippo and Riedel 1973, p. 526, pl. 14, fig. 5-12; pl. 31, fig. 4-7.

Circodiscus circularis (Clark and Campbell), Suzuki *et al.* 2009, p. 254, pl. 16, fig. 12-13b.

Circodiscus sp. A and C, Suzuki *et al.* 2009, p. 254, pl. 17, fig. 1a-9.

Our broad definition of this species follows Sanfilippo and Riedel (1973). We include *Circodiscus* sp. A and C of Suzuki *et al.* in our concept of *P. circularis*. Cosmopolitan.

Genus *PERICHLAMYDIUM* Ehrenberg

Perichlamyidium limbatum Ehrenberg

Perichlamyidium limbatum Ehrenberg 1847, p.43; 1854b, pl 22, fig 20; Haeckel, 1887, p. 514; Petrushevskaya, 1975, p. 575, pl. 6, fig. 11; pl. 39, figs. 1-4; Kruglikova 1977, pl 110, fig. 15.

Genus *SPONGODISCUS* Ehrenberg

Spongodiscus craticulatus (Stöhr)

Spongodiscus craticulatus (Stöhr), Petrushevskaya 1975, pl. 5, fig. 9-10.

Spongodiscus cruciferus (Clark and Campbell)

Spongasteriscus (*Spongasteriscinus*) *cruciferus* Clark and Campbell 1942, p. 50, pl. 1, fig. 1-6, 8, 10, 11, 16-18; Sanfilippo and Riedel 1973, p. 524, pl. 11, fig. 14-17; pl 28, fig. 10, 11; Crouch and Hollis 1996, p. 26; Hollis *et al.*, 1997, p. 50; Suzuki *et al.* 2009, p. 252, pl. 14, fig. 1a-5b.

Cosmopolitan.

***Spongodiscus festivus* (Clark and Campbell)**

Stylotrochus festivus Clark and Campbell 1942, p. 48, pl. 2, fig. 5-8; Suzuki et al. 2009, p. 251, pl. 13, fig. 4-7.

Stylotrochus nitidus Sanfilippo and Riedel 1973, p. 525, pl. 13, fig. 9-14, pl. 30, fig. 7-10.

This species is distinguished by having numerous cylindrical spines radiating from the margin. We include forms with an internal meshwork that ranges from densely spongy (*S. festivus* s.s.) to being arranged in closely-spaced concentric rings (*S. nitidus* s.s.).

Genus *SPONGOPYLE* Dreyer

***Spongopyle osculosa* Dreyer** [Plate 1, fig. 13]

Spongopyle osculosa Dreyer, 1889, p. 42, pl. 6, fig. 99, 100. Riedel 1958, p. 226, pl. 1, fig. 12.

Spongodiscus osculosus (Dreyer), Chen 1975, p. 455, pl. 24, fig. 4; Hollis et al. 1997, p. 50; Suzuki et al. 2009, p. 251, pl. 12, fig. 12-14.

Spongodiscus resurgens Ehrenberg *osculosa* (Dreyer). Petrushevskaya 1975, p. 574, pl. 5, fig. 11; pl. 36, fig. 1-4.

Spongodiscus sp. D Suzuki et al. 2009, p. 252, pl. 12, fig. 10-11.

We include all forms with a circular to oval outline, covered with a spongy meshwork, no internal structure visible and a pylom or notch on the rim. We include *Spongodiscus* sp. D of Suzuki et al. (2009) into our concept of *S. osculosa*. High-latitude.

Family SPONGURIDAE Haeckel, sensu Petrushevskaya, 1975

Genus *AMPHICRASPEDUM* Haeckel, sensu Sanfilippo and Riedel 1973

***Amphicraspedum murrayanum* Haeckel** [Plate 1, fig. 14]

Amphicraspedum murrayanum Haeckel 1887, p. 523, pl. 44, fig. 10; Sanfilippo and Riedel 1973, p. 524, pl. 10, fig. 3-6, pl. 28, fig. 1.

Low-latitude.

***Amphicraspedum prolixum* Sanfilippo and Riedel group** [Plate 1, fig. 15-17]

Amphibrachium gracilis Lipman 1972, p. 55, pl. 9, fig. 11-13.

Amphicarydiscus gracilis (Lipman), Kozlova 1999 (in part), p. 98, pl. 15, fig. 5; Pl. 20, fig. 5 (not fig. 3), pl. 22, fig. 15, pl. 42, fig. 1, 3.

Amphicraspedum prolixum Sanfilippo and Riedel gr. 1973, p. 524, pl. 11, fig. 1-5, 7-11, pl. 28, fig. 3-5. Strong *et al.* 1995, p. 208.

Amphicraspedum? *prolixum* Sanfilippo and Riedel, Kozlova 1999, p. 100, pl. 20, fig. 6, pl.

22, fig. 16, pl. 42, fig. 2 (also fig. 5).

Spongurids with elongated cylindrical arms, thickened at the ends, and usually with longitudinally aligned meshwork; may or may not be inflated medially. This range in variation appears to conform to the description of *A. gracilis*, which may prove to be the senior synonym. Low-latitude.

Genus *AMPHYMENIUM* Haeckel

***Amphymenium splendiaratum* Clark and Campbell** [Plate 1, fig. 18, 19]

Amphymenium splendiaratum Clark and Campbell 1942, p. 46, pl. 1, fig. 12, 14; Sanfilippo and Riedel 1973, p. 524, pl. 11, fig. 6-8; pl. 28, fig. 6-8; Petrushevskaya 1975, p. 577, pl. 7, fig. 1; pl. 37, fig. 1-3; Caulet 1991, p. 537; O'Connor 1993, p. 40, pl. 2, fig. 16, 17; Suzuki et al. 2009, p. 253, pl. 6, fig. 9-12.

Amphymenium cf. *splendiaratum* Clark and Campbell Hollis et al. 1997, p. 49, pl. 3, fig. 9.
? *Tessarastrum* sp. Caulet 1986, pl. 2, fig. 5.

Cosmopolitan

Genus *SPONGURUS* Haeckel

***Spongurus bilobatus* Clark and Campbell**

Spongurus bilobatus Clark and Campbell 1942, p. 36, fig. 7-9. Blueford 1988, p. 252, pl. 6, fig. 9; pl. 7, fig. 1; Hollis et al. 1997, p. 50, pl. 3, fig. 5-6.; Suzuki et al. (2009), p. 247, pl. 6, fig. 13-19.

Family LITHELIIDAE Haeckel

We include in this family all spumellarians with spherical to ellipsoidal cortical shells, an inner shell structure of closely spaced spiral or concentric layers, and with or without a pylome.

Genus *LARCOPYLE* Dreyer, emend. Lazarus et al. 2005

Larcopyle Dreyer 1889 – preferred name for prune-shaped spumellarians (“prunoids”)

Prunopyle Dreyer 1889 – type species is probably a stylosphaerin

Lithocarpium Stöhr, *sensu* Petrushevskaya 1975, p. 572. This is a nassellarian genus!

Spumellarians with an ellipsoidal or flattened ellipsoidal cortical shell (prune-shaped), usually bearing a pylome at one pole. A spongy, spiral, tightly concentric or pylonid inner shell structure is connected to the out shell either directly (last whorl in spiral) or by randomly distributed radial beams that may form radial spines. High-latitude.

***Larcopyle frakesi* (Chen)** [Plate 1, fig. 20]

Prunopyle frakesi Chen 1975, p.454, pl.10, fig.1-3.

Lithocarpium monikae Petrushevskaya 1975, p. 572, pl.4, fig.6-10, pl.30, fig.1-7.

Cortical shell prune-shaped with small pylome, internal shell spongy. High-latitude.

***Larcopyle hayesi* (Chen)** [Plate 1, fig. 21]

Prunopyle hayesi Chen 1975, p. 454, pl. 9, fig. 3-5; Caulet 1986, p. 853; 1991, p. 539;

Abelmann 1990, p. 693, pl. 3, fig. 14; Takemura 1992, p. 742, pl. 1, fig. 13, 14; Hollis et al. 1997, p. 48, pl. 2, fig. 12-14; Takemura and Ling, 1997, p. 114.

Ommatodiscus haeckeli Stöhr. Petrushevskaya 1975 (in part), p. 572, pl. 3, fig. 9, 12, 13; pl. 32, fig. 1-8.

Prunopyle cf. *decipiens* (Stöhr), Hollis et al. 1997, p. 47, pl. 2, fig. 18-24.

Larcopyle hayesi (Chen), Lazarus et al. 2005, p. 119, pl. 11, fig. 1-21.

Larcopyle occidentalis (Clark and Campbell 1942), Suzuki et al. (2009), p. 249, pl. 8, fig. 1-3, pl. 9, fig. 1-13.

Inflated ellipsoidal outer shell, with a densely spiral or concentric inner shell structure. Broad pylome is surrounded by short spines. High-latitude.

***Larcopyle labyrinthusa* Lazarus** [Plate 1, fig. 22]

Larcopyle labyrinthusa Lazarus et al. 2005, p.111, pl.5, figs. 14-24.

Subspherical whorl-shaped shell, beams connecting the spirals, pylome present with rather long teeth. Described from middle Miocene of Weddell Sea (Lazarus et al., 2005).

***Larcopyle polyacantha* (Campbell and Clark) group** [Plate 1, fig. 23, 24]

Ommatodiscus fragilis Stöhr, 1880, p. 116, pl. 6, fig. 10.

Larnacantha polyacantha Campbell and Clark 1944 (in part), p. 30, pl. 5, fig. 4-6 (not fig. 7).

Prunopyle polyacantha (Campbell and Clark) Caulet 1991, p. 539; Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 48, pl. 2, fig. 25-27 (also fig. 11).

Prunopyle titan Campbell and Clark 1944, p. 20, pl. 3, fig. 1-3; Hays 1965, p. 173, pl. 2, fig. 4; Caulet 1986, p. 853. Abelmann 1990, p. 693, pl. 3, fig. 16.

Lithocarpium polyacantha (Campbell and Clark). Petrushevskaya 1975, p. 572, pl. 3, fig. 6-8 (also fig. 9), pl. 29, fig. 6; Abelmann 1990, p. 694, pl. 4, fig. 2; O'Connor 1993, p. 37, pl. 2, fig. 12, 13.

Lithocarpium fragilis (Stöhr). Petrushevskaya 1975, p. 572, pl. 4, fig. 2-4.

Prunopyle sp. B gr. Abelmann 1990, p. 693, pl. 4, fig. 3a, b.

Prunopyle fragilis (Stöhr). Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 47, pl. 2, fig. 28-29.

Prunopyle cf. *titan* Campbell and Clark. Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 49, pl. 2, fig. 31-32.

Larcopyle polyacantha (Campbell and Clark) group, Lazarus et al. 2005, p. 106, pl. 2, fig. 1-17; pl. 3, fig. 1; pl. 4, fig. 1-10.

Subcylindrical to ellipsoidal cortical shell, usually with clusters of spines at one or both poles. Inner shell structure composed of weakly increasing single or double spirals; spongy meshwork may obscure spiral structure. Caps may develop at poles. A pylome is usually present.

Larcopyle cf. pylomaticus (Riedel) [Plate 1, fig. 25a, b]

Larcopyle pylomaticus (Riedel), Lazarus et al. 2005, p.115, pl.9, fig.1-12.

The species differ from true *L. pylomaticus* by not having a double spiral with constantly spaced whorls. Our specimens have a thick shell with small pores, a diffuse, spongy inner structure with sometimes and a strong pylome with teeth. Our specimen differ from Lazarus et al. 2005, pl. 9, fig. 6, by having a more elliptical shape.

Genus *LITHELIUS* Haeckel

Lithelius foremanae Sanfilippo and Riedel

Lithelius foremanae Sanfilippo and Riedel 1973, p. 522, pl. 7, fig. 1-6, pl. 26, fig. 4-5.

? *Lithelius* aff. *foremanae* Sanfilippo and Riedel, Hollis 1997, p. 44, pl. 5, fig. 7-9.

***Lithelius minor* Jörgensen group** [Plate 1, fig. 26-28]

Litheliid gen. gr. A. Petrushevskaya and Kozlova 1972, p. 529, pl. 5, fig. 14, 15.

Lithelius minor Jörgensen. Nigrini and Moore 1979, p. S135, pl. 17, fig. 3, 4a, b. O'Connor 1993, p. 37, pl. 10, fig. 3, 4.

Lithelius minor Jörgensen gr. Hollis, 1997, p. 45. pl. 6, fig. 6-16; Hollis et al., 1997, p. 47.

Ommatodiscus aff. *haeckeli* Stöhr, Hollis et al. 1997, p. 47, pl. 2, fig. 16, 17.

We include in this group following species by Suzuki et al. 2009, as we consider these types to be *L. minor* gr.: *Larcopyle compositus*, *Larcopyle hayesi*, *Lithelius* spp., *Lithelius* sp. A and *Lithelius* sp.

High-latitude.

Genus *SPHAEROPYLE* Dreyer, 1888

Spheroidal test with concentric inner shells and distinct pylome.

Sphaeropyle tetrapila* (Hays) = *Larcopyle tetrapila* / *Prunopyle tetrapila [Plate 1, fig. 29]

Prunopyle tetrapila Hays 1965, p. 172, pl. 2, fig. 5; Abelman 1990, p. 693, pl. 3, fig. 16;

Caulet 1991, p. 539; Takemura 1992, p. 742, pl. 2, fig. 1-2; Hollis et al. 1997, p. 48, pl. 2, fig. 30; Takemura and Ling 1997, p. 114.

Transferred to *Sphaeropyle* based on spherical outline of test. High-latitude.

Order NASSELLARIA

Family TRISSOCYCLIDAE Haeckel

Members of this family have not been differentiated, except the taxon below.

Genus *ZYGOCIRCUS* Bütschli

Zygocircus bütschli Haeckel

Zygocircus bütschli Haeckel, Petrushevskaya and Kozlova 1972, p. 534, pl. 41, fig. 8-11;
Takemura 1992, p. 743, pl. 5, fig. 4; Hollis et al., 1997, pl. 3, fig. 11; Takemura and Ling,
1997, pl. 1, fig. 4.

Family PLAGIACANTHIDAE Haeckel, emend. Petrushevskaya, 1971a

Genus *AMPHICENTRIA* Ehrenberg 1861

Amphicentrica sp. 1 sensu Suzuki [Plate 2, fig. 1]

Amphicentria sp. 1, Suzuki et al. 2009, p. 257, pl. 22, fig. 4.

This species is characterized by a large cephalis with a large open area on its top.

Genus *ANTARCTISSA* Petrushevskaya 1967

Antarctissa cylindrica Petrushevskaya

Antarctissa cylindrica Petrushevskaya 1975, p. 591. pl. 11, fig. 19-20.

High-latitude.

Antarctissa robusta Petrushevskaya

Antarctissa robusta Petrushevskaya 1975, p. 591. pl. 11, fig. 21-22.

High-latitude.

Genus *ARCHIPILIUM* Haeckel 1881

Archipilium macropus (Haeckel)

Sethopilium macropus Haeckel 1887 p.1203; pl.97, fig.9.

Archipilium spp. aff. *A. macropus* (Haeckel), Petrushevskaya and Kozlova 1972, pl. 29, fig.

14.

Archipilium macropus (Haeckel), Petrushevskaya 1981, p. 249

Genus *CALLIMITRA* Haeckel, emend. Goll 1979

Callimitra aff. *atavia* Goll [Plate 2, fig. 2]

? *Callimitra atavia* Goll 1979, p. 338, pl. 5, fig. 1, 5-9, 11. O'Connor 1993, p. 60, pl. 5, fig. 26, 27; Hollis et al. 1997, pl. 3, fig. 30.

Differs from *C. atavia* by having a latticed cephalic structure connecting four branched spines, rather than a solid perforate cephalis.

Genus *CERATOCYRTIS* Bütschli, sensu Petrushevskaya 1971b

Ceratocyrtis spp. [Plate 2, fig. 3-5]

We include all Ceratocyrtids, a small spherical cephalis, sometimes with short spines, a robust shell often spinose in the upper part, irregular pores. All ceratocyrtids have a high-latitude affinity.

Genus *CLADOSCENIUM*

Cladoscenium ancoratum Haeckel

Cladoscenium ancoratum Haeckel 1887, pl. 53, fig. 13.

Cladoscenium ancoratum Haeckel, Abelman 1992, pl. 4, fig. 9.

Genus *CORYTHOMELISSA* Campbell

Corythomelissa adunca (Sanfilippo and Riedel)

Spongomelissa adunca Sanfilippo and Riedel 1973, pl. 34, fig. 1.

Corythomelissa horrida Petrushevskaya. Takemura 1992, p. 744, pl. 3, fig. 14. Takemura and Ling 1997, p. 111.

Corythomelissa adunca (Sanfilippo and Riedel), Strong et al. 1995, p. 208, fig. 9b; Hollis et al., 1997, p. 51.

This species is distinguished from *C. cucumella* Sanfilippo and Riedel and *C. horrida* Petrushevskaya by the thorax being open at the base, although it may terminate in a thickened peristome with an interned rim. In these other two species, the thorax contracts distally and terminates in a constricted peristome, which may be closed with a latticed plate. In the specimen illustrated by Takemura (1992), the thorax is not constructed distally, terminating with a ragged margin. We assume the same species was also recorded by Takemura and Ling (1997). *C. adunca* bears some resemblance to *Pseudictyophimus gracilipes* but is distinguished by its large hemispherical cephalis, comprising 2-3 lobes, and the absence of a well-developed vertical spine, extending outside the cephalis. If a vertical spine does protrude from the cephalis in *C. adunca*, it is surrounded by a short latticed tube.

Genus *LITHOMELISSA* Ehrenberg

All *Lithomelissa* spp. we identified here are considered to have a high-latitude affinity.

Therefore all species identified in Suzuki et al. (2009) are included here (*Lithomelissa* sp. 6, *L.* sp. 7, *L.* sp. H, *L.* sp. G, *L.* sp. 2, *L.* sp. 3, *L.* sp. I).

Lithomelissa challengeræ Chen [Plate 2, fig. 6-8]

Lithomelissa challengeræ Chen 1975, p. 457, pl. 8, fig. 3; Takemura 1992, p. 744, pl. 4, fig. 11-12; Takemura and Ling 1997, p. 113.

Lithomelissa cf. *challengeræ* Chen [Plate 2, fig. 9]

Lithomelissa challengeræ Chen, Hollis et al. 1997, p. 51, pl. 3, fig. 21

? *Lithomelissa challengeræ* Chen 1975, p. 457, pl. 8, fig. 3; Takemura 1992, p. 744, pl. 4, fig. 11, 12; Takemura and Ling 1997, p. 113.

This species was poorly illustrated by Chen (1975) but, assuming the same species was illustrated by Takemura (1992), the species encountered at DSDP Site 277 differs in having perforated lateral and dorsal wings and a broadly bladed apical horn (cf. *Lithomelissa spongiosa* Bütschli 1882, p. 33, fig. 25).

Lithomelissa ehrenbergi Bütschli [Plate 2, fig. 10, 11]

Lithomelissa macroptera Ehrenberg 1876 (in part), pl. 3, fig. 8 (not fig. 9-10).

Lithomelissa ehrenbergi Bütschli 1882, p. 519, pl. 33, fig. 21; Caulet 1991, p. 538, pl. 2, fig. 3; Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 51, pl. 3, fig. 17-20; Suzuki et al., 2009, p. 256, pl. 20, fig. 1a-b.

Lithomelissa sp. B aff. *L. mitra* Bütschli, Chen 1975, p. 458, pl. 8, fig. 4-5.

Characterised by a bulbous, perforate cephalis with well-developed apical, dorsal and lateral spines (see comments in Caulet 1991, p. 533, and figure in Ehrenberg 1875, pl. 3, fig. 8). Distinguished from *L. haeckeli* by having straight or slightly curved dorsal and lateral spines; spines in *L. haeckeli* are strongly convex. We include *L. haeckeli* of Suzuki et al. (2009) in our concept of *L. ehrenbergi*.

Lithomelissa gelasinus O'Connor [Plate 2, fig. 12, 13]

Lithomelissa gelasinus O'Connor, 1997, pl. 2, fig. 3-6, pl. 6, fig. 6-9; Hollis et al. 1997, p. 52, pl. 3, fig. 15-16.

Poreless, but dimpled cephalis, irregular pore size on abdomen. Very similar to *L. sphaerocephalis* from which it is distinguished by a thorax which is cylindrical rather than conical and of about as wide as the cephalis.

Lithomelissa cf. *haeckeli* Bütschli [Plate 2, fig. 14]

Lithomelissa cf. *haeckeli* Bütschli, Hollis et al. 1997, p. 52, pl. 3, fig. 22.

cf. *Lithomelissa haeckeli* Bütschli, 1882, p. 519, pl. 33, fig. 23.

Uncertainly ascribed to *L. haeckeli* based on the strongly curved lateral spines.

***Lithomelissa macroptera* Ehrenberg** [Plate 2, fig. 15a, b]

Lithomelissa macroptera Ehrenberg 1874, p. 241; 1876, pl. 3, fig. 9-10 (not fig. 8).
non *Dictyophimus macropterus* (Ehrenberg), Takahashi p.116; pl.39, fig.8-11.

Haeckel 1887 does not illustrate this species and modern specimens (e.g. Takahashi) do not appear to be in the concept of Ehrenberg's illustration.

***Lithomelissa robusta* Chen** [Plate 2, fig. 16]

Lithomelissa robusta Chen 1975, p. 457, pl. 8, fig. 1, 2; Caulet 1991, p. 538;
non *Lithomelissa robusta* Chen 1975, Hollis et al. 1997, p. 52, pl. 3. Fig. 23.

Cephalis is markedly smaller than thorax. The specimen of Hollis et al. 1997 is *Lithomelissa ehrenbergi*.

***Lithomelissa sphaerocephalis* Chen** [Plate 2, fig. 17]

Lithomelissa sphaerocephalis Chen 1975, p. 457, pl. 8, fig. 1, 2; Takemura 1992, p. 744, pl. 4, fig. 8, 9; Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 52, pl. 3. Fig. 12-13;
Takemura and Ling 1997, p. 113.

Spheroidal cephalis, dimpled, poreless, bears a short apical spine. Short downward directed dorsal and lateral spines. Thorax conical, with subcircular to irregular pores, often covered with spongy meshwork.

***Lithomelissa tricornis* Chen** [Plate 2, fig. 18]

Lithomelissa tricornis Chen 1975, p. 458, pl. 8, fig. 6-7; Takemura 1992, p. 744, pl. 2, fig. 11-12; Hollis et al. 1997, p. 53; Takemura and Ling, 1997, p. 113.

***Lithomelissa? sakai* O'Connor** [Plate 2, fig. 19]

Lithomelissa sakai O'Connor 2000, p. 200, pl.2, fig. 9a-15

Robust shell, latticed feet and apical horn, irregular pore size.

Genus *LOPHOPHAENA* Ehrenberg, emend. Petrushevskaya 1971b

***Lophophaena capito* Ehrenberg**

Lophophaena capito Ehrenberg 1874, p. 242; 1876, pl. 8, fig. 6; Petrushevskaya and Kozlova 1972, p. 535, pl. 33, fig. 20-23; Petrushevskaya 1975, pl. 9, fig. 21 (not from Leg 29);
Crouch and Hollis 1996, p. 26.

Cosmopolitan.

***Lophophaena simplex* Funakawa**

Lophophaena simplex Funakawa 1994, p.465-466, pl.8, fig.1a-2c

Genus *PSEUDODICTYOPHIMUS* Petrushevskaya

All specimens assigned to the genus *Pseudodictyophimus* are considered to be high-latitude.
[Plate 2, fig. 24-27]

Pseudodictyophimus galeatus Caulet [Plate 2, fig. 20]

Pseudodictyophimus galeatus Caulet 1991, p. 534, pl. 2, fig. 9-10.

This species differs from *Pseudodictyophimus gracilipes* gr. by having a thick-walled, spongy shell.

Pseudodictyophimus gracilipes (Bailey) group, New group [Plate 2, fig. 21-23]

Dictyophimus gracilipes Bailey, 1856, p. 4, pl. 1, fig. 8.

Pseudodictyophimus gracilipes (Bailey). Petrushevskaya 1971b, p. 93, fig. 47-49; 1975, p. 592, pl. 11, fig. 17. Caulet 1986, p. 853.

Lithomelissa dupliphysa Caulet, Crouch and Hollis 1996, p. 26.

Petrushevskaya (1971b) described two subspecies of *P. gracilipes*: *P. gracilipes gracilipes* and *P. gracilipes tetracanthus*. The latter is said to be confined to the tropics whereas the former is restricted to high latitudes (bipolar). However, the type specimen appears to bear closest resemblance to *P. g. tetracantha* despite being described from the boreal Kamchatka Sea (Bailey 1856). Nevertheless, the specimens encountered in our study are consistent with Petrushevskaya's inference as they most closely resemble the high latitude morphotype: *P. g. gracilipes*. This subspecies is distinguished by a lobate outline, a cephalis that is subcircular in outline, and large irregularly distributed pores. *P. g. tetracantha* is distinguished by a subtriangular outline, thin tapering feet, and smaller regularly distributed pores.

Genus *TRIPODISCINUS* Haeckel, *sensu* Petrushevskaya and Kozlova 1979

Tripodiscinus clavipes (Clark and Campbell)

Tripilidium clavipes Clark and Campbell 1942, p. 64, pl. 9, fig. 29.

Tripodiscinus clavipes (Campbell). Petrushevskaya and Kozlova 1979, p. 115, fig. 302. Caulet 1986, p. 854.

? *Clathrocanium* sp. O'Connor 1993, p. 60, pl. 10, fig. 16.

Cosmopolitan.

Family *CANNOBOTRYIDAE* Haeckel

Genus *ARTOBOTRYS* Petrushevskaya

Placed in the Cannobotryidae by Petrushevskaya (1971).

***Artobotrys auriculaleporis* (Clark and Campbell)**

Lophophaena auriculaleporis Clark and Campbell 1942, p. 76, pl. 8, fig. 20, 27-29; Blueford 1988, p. 246, pl. 3, fig. 1-3.

Lophocyrtis biaurita (Ehrenberg), Foreman 1973 (in part), p. 442, pl. 8, fig. 23; Chen 1975, p. 461, pl. 3, fig. 2; (?) Caulet 1986, p. 853; Takemura 1992, p. 747, pl. 7, fig. 8; Takemura and Ling 1997, p. 113.

Artobotrys auriculaleporis (Clark and Campbell), Caulet 1991, p. 537.

Lophocyrtis? auriculaleporis (Clark and Campbell), Strong *et al.* 1995, p. 208, fig. 10s, t.

Cycladophora auriculaleporis (Clark and Campbell), Hollis *et al.*, 1997, p. 59 pl. 3, fig. 31.

Distinguished from *Artobotrys biaurita* by a stout thorax with regularly arranged pores, at least in the central area, and a short peristome that is not flared. Suzuki *et al.* (2009) specimen of *A. norvegiensis* are included here. Cosmopolitan.

***Artobotrys titanothericeraos* (Clark and Campbell)**

Lophoconus titanothericeraos Clark and Campbell 1942, p. 89, pl. 8, fig. 24-26, 28, 30-37.

Artobotrys titanothericeraos (Clark and Campbell), Suzuki *et al.* (2009), p. 259, pl. 21, fig. 4a-5b.

Distinguished from *Artobotrys auriculaleporis* by having three segments and a well-defined lumbar stricture. Possibly restricted to cool-water mid-latitude affinities, as so far only known from California and New Zealand.

Genus *BOTRYOCELLA* Haeckel 1881

***Botryocella?* sp. A, sensu Apel 2002 [Plate 3, fig. 1-4]**

Botryocella? sp. A, Apel, p. 44, pl. 6, fig. 12.

Genus *GLYCOBOTRYS* Campbell

This genus includes many species that have previously been placed in *Botryocella*. Campbell (1951) notes that the type species of the latter genus has a closed thorax and lacks cephalic tubes. Campbell described *Glycobotrys* as having cephalic tubules and a fenestrated (open) thorax. We apply the name to all cannobotryiids with simple tubes extending from the antecephalic or postcephalic lobe, with or without a closed thorax. This definition encompasses the range of variation exhibited by the type species, *Lithobotrys geminata* Ehrenberg (1876, pl. 3, fig. 19) and related species, *Lithobotrys nasuta* Ehrenberg (1876, pl. 3, fig. 21).

***Glycobotrys nasuta* (Ehrenberg) group, New Group [Plate 3, fig. 5-7]**

Lithobotrys nasuta Ehrenberg, 1874, p. 237; 1876, pl. 3, fig. 21.

Botryopyle dictyocephalus (Haeckel) gr. Riedel and Sanfilippo 1971 (in part), p. 1602, pl. 2J, fig. 18; pl. 3F, fig. 12. Takemura 1992, p. 743, pl. 3, fig. 7. O'Connor 1993, p. 56, pl. 10, fig. 15. Takemura and Ling, 1997, p. 111.

Botryocella aff. *cribrosa* (Ehrenberg) gr. Petrushevskaya and Kozlova 1972, p. 554, pl. 39, fig. 4-6.

Botryocella? appenninica Vinassa de Regny gr. Petrushevskaya 1975, p. 589, pl. 13, fig. 16.
Botryocella cribrosa (Ehrenberg) gr. Caulet 1986, p. 852.
Botryopyle sp. Crouch and Hollis 1996, p. 26.
Botryocella pauciperforata O'Connor 1999a, p. 10, pl. 1, fig. 21a-24; pl. 5, fig. 20a-24.
Glycobotrys geminata (Ehrenberg), Suzuki et al. 2009, p. 258, pl. 18, fig. 1a-b.
non Lithobotrys cribrosa Ehrenberg 1874, p. 237; 1876, pl. 3, fig. 20 (no cephalic tubes).

We include in this group all morphotypes with a large antecephalic lobe and a postcephalic tube. Thorax may be open or closed. Distinguished from *Botryopyle dictyocephalus* by a postcephalic tube at base of cephalis. Cosmopolitan.

Family ARTOSTROBIIDAE Riedel

Genus *BURYELLA* Foreman

Buryella granulata (Petrushevskaya)

Lithocampe sp. A, Dumitrica 1973, p. 789, pl. 10, fig. 3; pl. 11, fig. 3.
Lithocampe? granulata Petrushevskaya 1977, p. 18, pl. 3, fig. a, b, v.
Stichomitra granulata (Petrushevskaya). Hollis 1991, p. 134, pl. 20, fig. 7-12; 1993, p. 321, pl. 1, fig. 10, 11. Strong et al. 1995, p. 209, fig. 8j.
Buryella granulata (Petrushevskaya), Hollis 1997, p. 80, pl. 21, fig. 1-5; Hollis et al. 1997, p. 57, pl. 5, fig. 6.

Genus *SIPHOCAMPE* Haeckel, emend. Nigrini 1977

= *Lithomitra* Bütschli, 1882; *Lithomitrella* Haeckel, 1887.

Siphocampe lineata (Ehrenberg)

Lithocampe lineata Ehrenberg, 1838, p. 130 (in part); 1854b, pl. 22, fig. 26 (not pl. 36, fig. 16).
Lithomitra lineata (Ehrenberg), Haeckel 1887, p. 1484 (in part); Riedel and Sanfillippo 1971 (in part), p. 1600, pl. 11, fig. 2, 8, 11 (not fig. 1, 3-7, 9-10); pl. 21, fig. 15 (not fig. 14, 16).
Siphocampe lineata (Ehrenberg), Nigrini 1977, pl. 3, fig. 9-10.

Distinguished by its slender abdomen, which has a smooth outline and pores arranged in widely spaced transverse rows.

Siphocampe nodosaria (Haeckel)

Lithomitra nodosaria Haeckel 1887, p. 1484, pl. 79, fig. 1; Petrushevskaya 1967, pl. 83, figs. 8-9; Kruglikova 1969, pl. 4, fig. 3; Petrushevskaya and Kozlova 1972, pl. 24, figs. 29-30.
? *Lithomitra embrionalis* Vinassa de Regny 1900, pl. 3, fig. 21.
Lithomitra eruca Haeckel. 1887, p. 1485, pl. 79, fig. 3; Petrushevskaya 1971b, pl. 122, fig. 6; Petrushevskaya and Kozlova 1972, p. 539, pl. 24, figs. 32, 33.
? *Siphocampe annulosa* Haeckel. 1887, p. 1500, pl. 79, fig. 10; cf. Nakaseko, 1963, p. 195, pl. 4, fig. 8.

Lithomitra lineata (Ehrenberg) Riedel and Sanfillippo 1971 (in part), p. 1600, pl. 11, fig. 1, 3-7, 9-10 (not 2, 8, 11); pl. 21, fig. 14, 16 (not fig. 15); Foreman 1973, p. 431, pl. 8, fig. 18-19.

Siphocampe nodosaria (Haeckel). Nigrini 1977, p. 256, pl. 3, fig. 11; Takemura 1992, p. 743, pl. 3, fig. 15; O'Connor 1993, p. 52, pl. 4, fig. 23-24; Takemura and Ling, 1997, p. 114.

Siphocampe arachnaea (Ehrenberg). Abelmann 1990, p. 698, pl. 8, fig. 4a, b. Strong *et al.* 1995, p. 209.

Siphocampe imbricata (Ehrenberg), Caulet 1991, p. 539, pl. 3, fig. 13; Suzuki *et al.* 2009, p. 264, pl. 18, fig. 12a-b.

Siphocampe elegans (Ehrenberg), Suzuki *et al.* 2009, p. 263, pl. 18, fig. 10.

Distinguished by abdominal indentations and longitudinal ridges between pores. Cosmopolitan.

***Siphocampe quadrata* (Petrushevskaya and Kozlova)**

Lithamphora sacculifera quadrata Petrushevskaya and Kozlova 1972, p. 539, pl. 30, fig. 4-6 (also fig. 1, 2 and Pl. 24, fig. 7).

Lithomitra docilis Foreman 1973, p. 431, pl. 8, fig. 20-22; pl. 9, fig. 3-5.

Siphocampe? quadrata (Petrushevskaya and Kozlova), Nigrini 1977, p. 257, pl. 3, fig. 12; Caulet 1991, p. 539; Takemura 1992, p. 743, pl. 7, fig. 7; Takemura and Ling, 1997, p. 114; Suzuki *et al.*, 2009, p. 264, pl. 18, fig. 9.

Siphocampe? sp. Caulet 1986, pl. 2, fig. 11.

Siphocampe sacculifera (Clark and Campbell), Suzuki *et al.*, 2009, p. 264, pl. 18, fig. 11a-b.

Distinguished by smooth outline and abdominal pores arranged in closely spaced transverse row, with no longitudinal ridges. Abdomen is widest medially. Cosmopolitan.

***Siphocampe* (?) *acephala* (Ehrenberg) group** [Plate 3, fig. 8-10]

? *Eucyrtidium elegans* Ehrenberg, 1854b, pl. 36, fig. 17; 1876, pl. 11, fig. 12.

? *Eucyrtidium acephalum* Ehrenberg, 1876, p. 70, pl. 11, fig. 5.

Lithomitra laevigata Principi 1909, p. 17, pl. 1, fig. 55.

Lithomitra urnula Clark and Campbell 1942, p. 91, pl. 9, fig. 19.

Lithomitra elizabethae Clark and Campbell 1942, p. 92, pl. 9, fig. 18.

Lithocampe minuta Clark and Campbell 1942, p. 93, pl. 9, fig. 17.

Theocampe spp. Johnson 1974, pl. 3, fig. 13-15 (also fig. 11).

Theocampe amphora Haeckel. Chen 1975 (in part), p. 456, pl. 2, fig. 2.

Theocampe minuta (Clark and Campbell), Petrushevskaya 1975, p. 578, pl. 26, figs. 5, 6.

Siphocampe acephala (Ehrenberg). Nigrini 1977, p. 254, pl. 3, fig. 5. Takemura 1992, p. 743, pl. 6, fig. 9. Takemura and Ling, p. 114.

Lithomitrella minuta (Clark and Campbell) Petrushevskaya and Kozlova 1979, p. 154, fig. 412, 413, 478; Shilov 1995, p. 127, pl. 2, fig. 5.

Lithomitrella elizabethae (Clark and Campbell), Petrushevskaya and Kozlova 1979, p. 153, fig. 418, 419, 483-486, 540-544; Shilov 1995, p. 127.

Artostrobis elegans (Ehrenberg). Caulet 1986, p. 851.

Lithomitrella acephala (Ehrenberg). Caulet 1986, p. 853.

Lithomitrella minuta (Clark and Campbell). Caulet 1986, p. 853.

Siphocampe pachyderma (Ehrenberg). Caulet 1991, p. 539, pl. 3, fig. 12.

Siphocampe missilis O'Connor 1994, p. 340, pl. 1, fig. 7, 9-12. pl. 3, fig. 8-12.

Siphocampe acephala (Ehrenberg) group, Hollis et al. 1997, p. 54, pl. 4, fig. 8-20.
Siphocampe minuta (Clark and Campbell), Suzuki et al. 2009, p. 264, pl. 18, fig. 5a-6b.
non *Siphocampe elizabethae* (Clark and Campbell), Nigrini 1977, p. 256, pl. 3, fig. 6 (= *S.?*
amygdala).

Bullet-shaped test of 3 segments. Cephalis partially submerged in thorax, becoming more prominent in later part of range, lacks vertical tube, rarely bears a small apical horn. Thorax hemispherical, with 3-5 rows of circular pores. Lumbar stricture absent to distinct. Abdomen cylindrical, may narrow distally, with 4-8 rows of small subcircular pores, becoming large and subquadrangular in later part of range.

Specimens referred to this species by Nigrini (1977) and subsequent workers appear closer to Principi's species than *E. acephalum* s.s. Under Nigrini's usage the species ranges from Early to Late Eocene (*Phormocyrtis striata striata* to *Theocyrtis bromia* Zones). Takemura reported the species in the Middle Eocene, with an isolated occurrence in Early Oligocene. For practical considerations co-occurring variants have not been differentiated, i.e. *S. elegans* (Ehrenberg), which has a well-developed apical horn, *S. missilis* O'Connor, which has a perforate, centred, rather than asymmetrically-placed, cephalis and a longer abdomen with 8-17 rows of pores.

This species group is tentatively placed in *Siphocampe* because its artostrobiid cephalic structure has not been confirmed. It appears to be closely related to the multisegmented *S.?*
amygdala.

***Siphocampe? amygdala* (Shilov) [Plate 3, fig. 11, 12]**

Theoperid gen. et sp. indet. Johnson, 1974, pl. 3, fig. 12.

Siphocampe elizabethae (Clark and Campbell). Nigrini 1977, p. 256, pl. 3, fig. 6.

Lithomitrella elizabethae (Clark and Campbell). Caulet 1986, p. 853.

Archaeodictyomitra? sp. Takemura 1992, p. 744, pl. 3, fig. 1-2.

Mita sp. Strong et al. 1995, p. 209, fig. 10x.

Dictyomitra amygdala Shilov 1995, p. 126, pl. 1, fig. 4-6b.

Siphocampe? “*elizabethae*” sensu Nigrini, Hollis et al. 1997, p. 55, pl. 4, fig. 21-26.

Dictyoprora? amygdala (Shilov), Suzuki 2009, p. 263, pl. 18, fig. 3.

non *Lithomitra elizabethae* Clark and Campbell 1942, p. 92, pl. 9, fig. 18.

Fusiform test of 5-6 segments with indistinct strictures. Thorax hemi-spherical with pores evenly distributed and smaller than in subsequent segments. Pores of post-thoracic segments longitudinally aligned and arranged in 3 rows on each segment. Outline smooth to slightly lobate, narrows distally, basal margin ragged. Differs from *S. acephala* gr. in that the post-thoracic part is not a single abdomen but 2-5 segments, each having 2-3 rows of subcircular pores. High-latitude.

Genus *SPIROCYRTIS* Haeckel, emend. Nigrini

***Spirocyrtis greeni* O'Connor**

Spirocyrtis greeni O'Connor 1999a, p. 8, pl. 1, fig. 15-20b, pl. 5, fig. 16a-19.

***Spirocyrtis joides* (Petrushevskaya)**

Botryostrobus sp. P, Petrushevskaya and Kozlova 1972, p. 539, pl. 24, fig. 8-11.

Botryostrobus joides Petrushevskaya 1975, p. 585, pl. 1-, fig. 37; Hull 1996, p. 137, pl. 4, fig. 5, 6, 20; Sanfilippo and Fourtanier 2003, p. 11, pl. P1, fig. 3, 12.
Spirocyrtilis sp. A. Hollis et al. 1997, p. 56, pl. 4, fig. 33-35.

Narrowly conical test of 6-10 segments. Thorax truncate-conical with 3 rows of pores. Cephalis hemispherical, internal features obscured by a "cockscomb" of webbed spines linking apical horn and vertical tube. Post-thoracic segments are rounded with 4 rows of closely-spaced subcircular pores. Final segment narrows, may terminate with a narrow peristome. Cosmopolitan.

Genus **THEOCAMPE** Haeckel (= **DICTYOPRORA** Haeckel)

***Theocampe amphora* (Haeckel)**

Dictyocephalus amphora Haeckel, 1887, p. 1305, pl. 62, fig. 4.
Theocampe amphora (Haeckel), Foreman 1973, p. 431, pl. 8, fig. 7, 9-13; pl. 9, fig. 8, 9; Chen 1975 (in part), p. 456, pl. 2, fig. 2; Hollis et al. 1997, p. 56, pl. 4, fig. 38-39.
Dictyoprora amphora (Haeckel). Nigrini 1977, p. 250, pl. 4, fig. 1, 2; Caulet 1991, p. 538; O'Connor 1993, p. 48, pl. 4, fig. 14, pl. 10, fig. 10.

Cosmopolitan.

***Theocampe urceolus* (Haeckel)**

Dictyocephalus urceolus Haeckel, 1887, p. 1305.
Theocampe urceolus (Haeckel), Foreman 1973, p. 432, pl. 8, fig. 14-17; pl. 9, fig. 6, 7; Chen 1975, p. 456, pl. 3, fig. 7; Hollis et al. 1997, p. 56, pl. 4, fig. 36-37.
Dictyoprora urceolus (Haeckel), Nigrini 1977, p. 251, pl. 4, fig. 9, 10; O'Connor 1993, p. 50, pl. 10, fig. 11.
? *Dictyoprora physothorax* Caulet 1991, p. 535, pl. 3, fig. 11.
Dictyoprora pirum (Ehrenberg), Takemura 1992, p. 743, pl. 5, fig. 11; Takemura and Ling, 1997, p. 111.

Specimens with a thorax as wide as the abdomen (= *T. physothorax*) are included in *T. urceolus* here. Takemura's figured specimen lacks the flattened abdomen characteristic of *T. pirum*. Cosmopolitan.

Family **ACROPYRAMIDIDAE** Haeckel

Genus **ARTOSTROBUS** Haeckel

***Artostrobos annulatus* (Bailey)**

Cornutella annulatus Bailey 1856, p. 3, fig. 5.
Artostrobos annulatus (Bailey), Petrushevskaya 1975, p. 579, pl. 10, fig. 4, 5; Hollis et al. 1997, p. 57, pl. 5, fig. 1-2.

High-latitude.

***Artostrobus* cf. *pretabulatus* Petrushevskaya** [Plate 3, fig. 13]

Artostrobus cf. *pretabulatus* Petrushevskaya. Takemura 1992, p. 745, pl. 5, fig. 12. Crouch and Hollis 1996, p. 26; Hollis et al., 1997, p. 57, pl. 5, fig. 3; Takemura and Ling, 1997, p. 111.

cf. *Artostrobus*? *pretabulatus* Petrushevskaya 1975, p. 580, pl. 10, fig. 2-3.

Distinguished from *A. pretabulatus* by a broader more conical thorax. High-latitude.

Genus *CINCLOPYRAMIS* Haeckel

Cinclopyramis Haeckel 1879; *Bathropyramis* Haeckel 1882; *Peripyramis* Haeckel 1882; *Sethopyramis* Haeckel 1882; *Enneapleuris* Haeckel 1887

As noted by Suzuki et al. (2009), the synonymised generic names have been applied inconsistently in the past and serve only to confuse species level discrimination within what is here considered to be a single long-ranging genus. *Cinclopyramis* is the senior synonym.

***Cinclopyramis circumtexta* (Haeckel)**

Peripyramis circumtexta Haeckel 1887. p. 1162, pl. 54, fig. 5; Riedel 1958, p. 231, pl. 2, fig. 8; Petrushevskaya 1975, pl. 13, fig. 29, pl. 44, fig. 5-6.

Sethopyramis magnifica Clark and Campbell 1942, p. 72, pl. 8, fig. 1, 5, 9.

Bathropyramis magnifica (Clark and Campbell), Hollis et al. 1997, p. 57, pl. 6, fig. 22.

Cinclopyramis quadrata (Haeckel), Suzuki et al., 2009, p. 262, pl. 22, fig. 13.

non *Bathropyramis quadrata* Haeckel 1887, p. 1159, pl. 54, fig. 1.

The specimens of *C. circumtexta* illustrated by Riedel (1958) indicate that the species is the senior synonym for *S. magnifica* Clark and Campbell. Both species are distinguished from *C. quadrata* by having a small subcircular cephalis with a distinct collar stricture, transverse pore frames being slightly offset between longitudinal beams, and a change in contour in the proximal thorax where pores change from being circular to rectangular. In *C. quadrata*, transverse beams are closely aligned along the length of the shell. According to Petrushevskaya, the distinctive outer layer of coarsely spongy meshwork is only seen from the middle Miocene onwards. However, Clark and Campbell (1942) mention this secondary mesh also occurring in some Eocene specimens. Cosmopolitan.

Genus *CORNUTELLA* Ehrenberg

***Cornutella profunda* Ehrenberg 1854**

Cornutella clathrata profunda Ehrenberg 1854a, p. 241; Ehrenberg 1854b, pl. 35B, fig. 21

Cornutella profunda Ehrenberg, Riedel 1958, p. 232, pl. 3, figs. 1-2; Petrushevskaya 1975, p. 587, pl. 13, fig. 29, pl. 44, fig. 5-6; Suzuki et al. 2009, p. 263, pl. 22, fig. 12a-b.

Cosmopolitan.

Family AMPHIPYNDACIDAE Riedel

We included in this family multisegmented nassellarians with the cephalis divided into two superposed chambers by a transverse internal ledge.

Family EUCYRTIDIIDAE Ehrenberg

Genus *ASPIS* Nishimura 1992

Aspis sp. A, sensu Hollis 2002 [Plate 3, fig. 14-16]

Ceratocyrtis sp., Dumitrica 1973, p. 788, pl. 3, fig. 2-4.

Aspis sp. A, Hollis 2002, p. 303, pl. 6, fig. 1-2.

Genus *CALOCYCLOMA* Haeckel

Calocycloma ampulla (Ehrenberg)

Calocycloma ampulla (Ehrenberg), Foreman 1973, p. 434, pl. 1, fig. 1-5; pl. 9, fig. 20; Strong *et al.* 1995, p. 208, fig. 9v-x; Hollis *et al.* 1997, p. 58.

Clathrocyclas universa Clark and Campbell

Clathrocyclas universa Clark and Campbell 1942, p. 86, pl. 7, fig. 8-12, 14-21, 25; Chen 1975, p. 459, pl. 1, fig. 2, 3; Petrushevskaya 1975, pl. 15, fig. 15-16; Blueford 1988, p. 244, pl. 2, fig. 1-3; Takemura 1992, p. 745, pl. 7, fig. 8; Strong *et al.* 1995 (in part), p. 208; Hollis *et al.* 1997, p. 59, pl. 5, fig. 17-20; Takemura and Ling 1997, p. 111.

We included *Clathrocyclas* ex. gr. aff. *extensa* and *C. ex. gr. extensa* from Suzuki *et al.* (2009). Cosmopolitan.

Genus *CYCLADOPHORA* Ehrenberg, emend. Lombari and Lazarus 1988

Distinguished from *Clathrocyclas* by a smaller, subspherical, cephalis with pores smaller and/or less numerous than those of the thorax, and flared thorax divided into upper and lower parts.

Cycladophora cosma cosma Lombari and Lazarus [Plate 3, fig. 17]

Cycladophora cosma cosma Lombari and Lazarus 1988, p. 104, pl. 1, figs. 1-6.

High-latitude.

Cycladophora humerus (Petrushevskaya) [Plate 3, fig. 18]

Clathrocyclas humerus Petrushevskaya 1975, p. 586, pl. 15, fig. 17, 12, 22, 23; pl. 43, fig. 1-2.

Cycladophora humerus (Petrushevskaya), Lombari and Lazarus 1988, p. 123, pl. 9, fig. 1-6.

Cycladophora cf. *humerus* (Petrushevskaya), Hollis et al. 1997, p. 59, pl. 5, fig. 21.

The form recorded by Hollis et al. (1997) is distinguished from *C. humerus* s.s. by a higher thorax with a subcylindrical lower part. In this respect this species is similar to the Paleocene species illustrated by Petrushevskaya (1975, pl. 15, fig. 18-20) but the latter has fewer pores (6-7 in a transverse row).

High-latitude.

Genus CYMAETRON Caulet 1991

***Cymaetron sinolampas* Caulet**

Cymaetron sinolampas Caulet 1991, p. 536, pl. 4, fig. 10-12.

Genus CYRTOLAGENA Haeckel 1887

***Cyrtolagena laguncula* Haeckel**

Cyrtolagena laguncula Haeckel 1887, p. 1451; pl. 75, fig. 10.

Cosmopolitan.

Genus EUCYRTIDIUM Ehrenberg

***Eucyrtidium antiquum* Caulet [Plate 3, fig. 19]**

Eucyrtidium sp., Chen 1975, p. 461, pl. 7, fig. 6-8; O'Connor 1993 (in part), p. 73, pl. 8, fig. 6, 7.

Eucyrtidium antiquum Caulet 1991, pl. 4, fig. 1, 2 (also fig. 5); Hollis et al. 1997, p. 60, pl. 5, fig. 30-31; Takemura and Ling 1997, p. 113, pl. 1, fig. 9.; Suzuki et al. (2009), p.259, pl. 21, fig. 14.

Eucyrtidium cheni Takemura 1992, p. 746, pl. 4, fig. 1-4.

E. cheni was synonymised with *E. antiquum* by Takemura and Ling (1997).

High-latitude.

***Eucyrtidium mariae* Caulet**

Eucyrtidium mariae Caulet 1991, p. 536, pl. 4, fig. 3, 4. Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 61 (as *E. ? mariae*).

Distinguished from *E. spinosum* by a more slender and thin-walled test, with small circular pores lacking well-developed frames. Internal segmentation of post-thoracic chambers frequently asymmetric. High-latitude.

***Eucyrtidium microporum* Ehrenberg**

Eucyrtidium microporum Ehrenberg, 1874, p. 230; 1876, pl. 11, fig. 20.

Stichopodium? microporum Ehrenberg, Funakawa et al., 2006, pl. 13, fig. 3-4.

***Eucyrtidium nishimurae* Takemura and Ling** [Plate 3, fig. 20a, b]

Eucyrtidium nishimurae Takemura and Ling 1997, p. 113, pl. 2, fig. 1-6; Suzuki et al. (2009), p. 259, pl. 21, fig. 15.

Spongy fourth segment. High-latitude.

***Eucyrtidium spinosum* Takemura** [Plate 3, fig. 21]

Eucyrtidium sp. A, Petrushevskaya 1975, p. 581, pl. 14, fig. 21-22.

Eucyrtidium? aff. *montiparium* Ehrenberg, Caulet 1991, p. 538, pl. 4, fig. 5-7.

Eucyrtidium spinosum Takemura 1992, p. 746, pl. 5, fig. 5-8; Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 61, pl. 5, fig. 38-31; Takemura and Ling 1997, p. 113, pl. 1, fig. 10.

High-latitude.

***Eucyrtidium ventriosum* O'Connor** [Plate 3, fig. 22]

Eucyrtidium montiparum Ehrenberg, Hollis et al. 1997, p. 61, pl. 5, fig. 25-27.

Eucyrtidium cf. *cheni* Takemura, Strong et al. 1995, p. 208, fig. 11x, y.

? *Eucyrtidium* aff. *montiparum* Ehrenberg, Petrushevskaya and Kozlova 1972, p. 548, pl. 26, fig. 2-4.

Eucyrtidium ventriosum O'Connor 1999a, p. 21, pl. 3, fig. 17-21b, pl. 6, fig. 28a-31.

Differs from *E. nishimurae* by lacking an elongate, subcylindrical, spongy fourth segment. High-latitude.

Genus EURYSTOMOSKEVOS Caulet

***Eurystomoskevos cauleti* O'Connor** [Plate 3, fig. 23a, b]

Eurystomoskevos "cauleti" O'Connor, Hollis et al. 1997, p. 61, pl. 5, fig. 4-5 (nom. Nud.).

Eurystomoskevos cauleti O'Connor 1999a, pl. 3, fig. 22-26; pl. 7, fig. 1a-3.

Distinguished from *E. petrushevskae* by a smaller apical horn, a more distinct cephalis and a thorax that usually is not flared. High-latitude.

***Eurystomoskevos petrushevskae* Caulet** [Plate 3, fig. 24]

Diplocyclas sp. Petrushevskaya and Kozlova 1972, p. 541, pl. 33, fig. 14-16; Petrushevskaya 1975, p. 587, pl. 24, fig. 4; Chen 1975, p. 460, pl. 7, fig. 4, 5; Takemura 1992, p. 746, pl. 3, fig. 16.

Eurystomoskevos petrushevskae Caulet 1991, p. 536, pl. 3, fig. 14, 15; O'Connor 1993, p. 74, pl. 8, fig. 12-14; Crouch and Hollis 1996, p. 26; Hollis et al. 1997, p. 62, pl. 5, fig. 22-23; Suzuki et al. (2009), p. 265, pl. 22, fig. 5a-6.

High-latitude.

Genus *EUSYRINGIUM* Haeckel

Eusyringium fistuligerum (Ehrenberg) [Plate 3, fig. 25]

Eucyrtidium fistuligerum Ehrenberg 1874, p. 229; 1876, pl. 9, fig. 3

Eusyringium fistuligerum (Ehrenberg), Foreman 1973, p. 435, pl. 11, fig. 6; Chen 1975, p. 461, pl. 3, fig. 3; Sanfilippo *et al.* 1985, p. 672, fig. 17.2a-c; Strong *et al.* 1995, p. 208, fig. 11C; Suzuki *et al.* (2009), pl. 22, fig. 14.

Cosmopolitan.

Eusyringium lagena (Ehrenberg)

Eusyringium lagena (Ehrenberg), Foreman 1973, p. 435, pl. 11, fig. 4-5; Chen 1975, p. 461; Sanfilippo *et al.* 1985, p. 672, fig. 17.2a-c; Strong *et al.* 1995, p. 208, fig. 11c; Hollis *et al.* 1997, p. 62, pl. 5, fig. 15-16.

Cosmopolitan.

Genus *PHORMOCYRTIS* Haeckel 1887

Phormocyrtis striata striata Brandt

Phormocyrtis striata striata Brandt, Foreman 1973, p. 438, pl. 7, fig. 5, 6, 9; Chen 1975, p. 456, pl. 3, fig. 8; Sanfilippo *et al.* 1985, p. 679, fig. 20.1a, b; O'Connor 1993, p. 78, pl. 9, fig. 8; Strong *et al.* 1995, p. 209, fig. 9o, p; Hollis *et al.* 1997, p. 65.

Cosmopolitan.

Genus *SETHOCYRTIS* Haeckel

Sethocyrtis chrysallis Sanfilippo and Blome [Plate 3, fig. 26a, b]

Sethocyrtis sp. Chen 1975, p. 459, pl. 1, fig. 4, 5; Takemura 1992, p. 747, pl. 7, fig. 14, 15; Strong *et al.* 1995 (in part), p. 209, fig. 11w; Takemura and Ling 1997, p. 114, pl. 1, fig. 11.

Sethocyrtis sp. A Hollis *et al.* 1997, p. 65, pl. 6, fig. 7.

Sethocyrtis chrysallis Sanfilippo and Blome 2001, p. 206, fig. 6j-n.

Cosmopolitan.

Genus *THYRSOCYRTIS* Ehrenberg

Thyrsoyrtis pinguisicoides O'Connor [Plate 3, fig. 27]

Thyrsoyrtis sp. Dinkelman 1973, p. 788, pl. 3, fig. 7, 8.

Theocotyle "pinguisicoides" O'Connor, Hollis *et al.* 1997, p. 65, pl. 6, fig. 10-12 (*nom. nud.*).

Thyrsoyrtis pinguisicoides O'Connor 1999a, p. 29, pl. 4, fig. 28-32; pl. 7, fig. 28a-31.

Distinguished from other members of the genus by the absence of terminal feet. High-latitude.

Family LYCHNOCANIIDAE Haeckel

Genus *DICTYOPHIMUS* Ehrenberg, emend. Nigrini 1967

Three-segmented, with 3 thoracic ribs that form divergent feet. Distinguished from *Pterocyrtidium* by indistinct separation between thorax and abdomen, triangular outline and large pores. *Pterocyrtidium* has a bullet-shaped outline. *Rhopalocanium* is similar to *Pterocyrtidium* but the feet extend from the abdomen and are often connected to it by transverse bars.

Dictyophimus infabricatus Nigrini

Dictyophimus infabricatus Nigrini 1968, p. 56, pl. 1, fig. 6; Takahashi 1991, pl. 37, fig. 3-5.

Dictyophimus? archipilium Petrushevskaya [Plate 4, fig. 1a, b, 2]

Dictyophimus? archipilium Petrushevskaya 1975, p. 583, pl. 25, fig. 1, 2.

Dictyophimus cf. *archipilium* Petrushevskaya, Crouch and Hollis 1996, p. 26 (in part); Hollis et al. 1997, p. 60 (in part).

Dictyophimus sp. 1 Suzuki et al. 2009, pl.22, fig.7.

This species is distinguished by its small size, weak apical horn and terminal feet that extend out from the lower thorax, and typically are short protuberences. Hollis et al (1997) failed to identify the distinguishing features of this species and erroneously included some specimens in which the feet were attached to the abdomen. Such specimens are referred to *D.? aff. archipilium* here (sensu Hollis 1997). This species may be better referred to *Pterocyrtidium*. High-latitude.

Dictyophimus? aff. archipilium Petrushevskaya [Plate 4, fig. 3a,b-8]

Dictyophimus? spp. Dumitrica 1973, p. 788, pl. 7, fig. 3-9; pl. 8, fig. 4; pl. 9, fig. 8; pl. 12, fig. 6, 7; pl 13, fig. 14. Hollis 1993, p. 322.

Dictyophimus sp. A, Strong et al. 1995, p. 208, fig. 8o; Hollis et al. 1997, p. 60.

Dictyophimus cf. *archipilium* Petrushevskaya, Crouch and Hollis 1996, p. 26 (in part); Hollis et al. 1997, p. 60 (in part), pl. 5, fig. 38-19.

Dictyophimus aff. *archipilium* Petrushevskaya, Hollis 1997, p. 82, pl. 22, fig. 5-7.

aff. *Dictyophimus? archipilium* Petrushevskaya 1975, p. 583, pl. 25, fig. 1, 2.

This Paleocene-Eocene species is distinguished from *D.? archipilium* by its distinctive feet, which extend from the base of the thorax and are connected to the abdomen by transverse bars. These bars are evident in the specimens referred to *D. cf. archipilium* by Hollis et al. (1997), indicating that the species persists into the Early Oligocene. This species may be better referred to *Rhopalocanium*. High-latitude.

Dictyophimus? aff. constrictus Nishimura [Plate 4, fig. 9, 10]

aff. *Dictyophimus? constrictus* Nishimura 1992, O'Connor 1999a, pl. 9, fig. 13.

Genus *LYCHNOCANIUM* Ehrenberg, sensu O'Connor 1999

Lychnocanium amphitrite (Foreman) [Plate 4, fig. 11a, b, c, 12]

Lychnocanoma amphitrite Foreman 1973, p. 437, pl. 11, fig. 10; Chen 1975, p. 462, pl. 2, fig. 7; Caulet 1986, p. 853; Takemura 1992, p. 747, pl. 7, fig. 9-10; O'Connor 1993, p. 76, pl. 8, fig. 29; Strong *et al.* 1995, p. 208, fig. 11k-l; Takemura and Ling 1997, p. 114, pl. 1, fig. 21; Suzuki *et al.* 2009, p. 261, pl. 19, fig. 5a-6b.

Lychnocanoma bellum (Clark and Campbell), Foreman 1973 (in part), pl. 1, fig. 17.

Lychnocanium amphitrite (Foreman), Hollis *et al.* 1997, p. 63, pl. 6, fig. 1-4.

Specimens with any vestige of a third segment, such as that figured by Foreman (1973, pl. 1, fig. 17), are here included in *L. amphitrite*. Cosmopolitan.

Lychnocanium babylonis (Clark and Campbell) [Plate 4, fig. 13a, b, 14]

Dictyophimus babylonis Clark and Campbell 1942, p. 67, pl. 9, fig. 32, 36.

Lychnocanoma babylonis (Clark and Campbell), Foreman 1973, p. 437, pl. 2, fig. 1; Chen 1975, p. 462, pl. 2, fig. 8; Caulet 1986, p. 853; Strong *et al.* 1995, p. 208, fig. 11a, b.

Lychnocanoma cf. *babylonis* Clark and Campbell, Takemura 1992, p. 747, pl. 7, fig. 13; Takemura and Ling, 1997, p. 114.

Sethochytris babylonis (Clark and Campbell). Hollis *et al.*, 1997, p. 65, pl. 5, fig. 32; Hollis, 2002.

Cephalis may be partially encased in thoracic wall. Distinguished from *Lychnocanium auxillum* and *Rhopalocanium ornatum* by the absence of an abdomen. Cosmopolitan.

Lychnocanium bellum Clark and Campbell [Plate 4, fig. 15, 16]

Lychnocanium bellum Clark and Campbell 1942, p. 72, pl. 9, fig. 35, 39; Hollis *et al.* 1997, p. 63, pl. 6, fig. 5-6.

Lychnocanoma bellum (Clark and Campbell); Foreman 1973 (in part), p. 437, pl. 11, fig. 9; Caulet 1986, p. 853; Strong *et al.* 1995, p. 208, fig. 11i, j; Suzuki *et al.* 2009, p. 261, pl. 19, fig. 1a-b.

Lychnocanium aff. *carinatum* Ehrenberg [Plate 4, fig. 17]

? *Lychnocanium carinatum* Ehrenberg 1876, p. 78, pl. 8, fig. 5; Haeckel 1887, p. 1226.

Theopodium sp. aff. *Lychnocanium carinatum* Ehrenberg, Nishimura, 1992, p. 7, fig. 11.

This Paleocene species was ascribed to *Theopodium* by Nishimura (1992) because it consists of three segments and has three solid feet. As the third segment is not fully developed and in other respects this species closely resembles *L. auxilla*, we transfer the species to *Lychnocanium*. This species is distinguished from *L. sphinx* by the presence of a third segment. It is distinguished from *L. auxilla* by having longitudinal costae on the thorax. It is distinguished from both species by having feet that curve inwards.

Lychnocanium conicum Clark and Campbell

Lychnocanium conicum Clark and Campbell 1942, p. 71, pl. 9, fig. 38.

Lychnocanoma? *conica* (Clark and Campbell), Suzuki *et al.* 2009, p. 261, pl. 19, fig. 1a-b.

non *Lychnocanoma conica* (Clark and Campbell). Abelman 1990, p. 697, pl. 6, fig. 8; pl. 7,

fig. 1a, b; Takemura 1992, p. 747, pl. 2, fig. 13-14; Takemura and Ling 1997, p. 114, pl. 1, fig. 20.

L. conicum s.s. differs from the morphotypes ascribed to the species by Abelmann (1990) and subsequent researchers, by having a pear-shaped thorax with and small aperture. It differs from *L. babylonis* by having a more inflated thorax and more slender feet. Cosmopolitan.

***Lychnocanium continuum* Ehrenberg**

Lychnocanium continuum Ehrenberg, 1874, p. 243; 1876, pl. 7, fig. 11; Funakawa et al. (2006), pl. 12, fig. 7.

***Lychnocanium tetrapodium* Ehrenberg** [Plate 4, fig. 18a, b]

Lychnocanium tetrapodium Ehrenberg 1874, p. 244; Ogane et al. 2009, pl. 7, fig. 9.

Lychnocanoma tetrapodium (Ehrenberg), Kamikuri et al. 2012, p. 103, pl. 2, fig. 9, 10.

Distinguished by having four feet. Low-latitude.

***Lychnocanium waiareka* O'Connor**

Lychnocanium waiareka O'Connor 1999a, p. 25, pl. 4, fig. 6-11, pl. 7, fig. 12a-15.

Lychnocanium tripodium Ehrenberg, Hollis et al. 1997, p. 64, pl. 5, fig. 34-37.

? *Lychnocanium hamosum* Ehrenberg 1876, pl. 7, fig. 9; Ogane et al. 2009, pl. 7, fig. 8a-b.

? *Lychnocanium tripodium* Ehrenberg 1876, pl. 7, fig. 2.

? *Lychnocanium neptunei* O'Connor 1997, pl. 3, fig. 7-10, pl. 9, fig. 7-12.

This species is distinguished by a campanulate thorax, which is almost as wide as it is long, and terminal feet that are only slightly flared. A delicate abdomen is preserved on well-preserved specimens. In less well-preserved specimens, all that remains of the abdomen are short thorns on the inner sides of the feet (e.g. Hollis et al. 1997, pl. 5, fig. 35-37). Such forms closely resemble *L. hamosum* Ehrenberg. We provisionally include specimens with smooth feet (e.g. Hollis et al. 1997, pl. 5, fig. 34), which may be *L. tripodium* Ehrenberg or *L. neptunei* O'Connor.

Genus *PTEROCODON* Ehrenberg 1847

***Pterocodon apis* Ehrenberg** [Plate 4, fig. 19, 20a, b]

Pterocodon apis Ehrenberg 1874, p. 255, Ehrenberg 1876, pl. 19, fig. 3; Ogane et al. 2009, pl. 5, fig. 8a-d.

? *Pteropilium* sp. B, O'Connor 1999b, p. 507, pl. 6, fig. C.

Three segmented, strong apical horn, two lateral wings from thorax, described from Barbados, Tertiary deposits. O'Connor's species is similar, but seems to have fewer pores on thorax and abdomen. Our specimens have larger, hexagonal pores on the abdomen and smaller, rounded pores on the thorax.

Genus *PTEROPILIUM* Haeckel 1882

The genus *Pteropilium* differs from the closely allied *Dictyoceras* in the development of three

free, vertical, latticed cephalic wings, expanded between the apical horn of the cephalis and the three wings of the torax; they may be direct prolongations of the latter. *Pteropilium* may have evolved from *Callimitra* or *Clathrocorys* by development of an abdomen.

***Pteropilium* aff. *contiguum* (Ehrenberg) [Plate 4, fig. 21]**

Pteropilium aff. *Pterocanium contiguum* Caulet 1991, p. 539, pl. 2, fig. 11..

aff. *Pterocanium contiguum* Ehrenberg 1874, p. 255, 1876, pl. 17, fig. 7.

Family LOPHOCYRTIIDAE sensu Sanfilippo and Caulet

Genus *APHETOCYRTIS* Sanfilippo and Caulet

The genus comprises the lineage: *Aphetocyrtis gnomabax* – *A. rossi* – *A. catalexis*. All three species have a porous subspherical cephalis with a small apical horn; a hemispherical to campanulate (bell-shaped) thorax with a rough thorny surface; and a subcylindrical to inverted truncate conical abdomen that may be closed and lacks a peristome or terminal feet.

Sanfilippo and Caulet (1998) differentiate species in this genus by skeletal features within the cephalis that are difficult to apply consistently to variably preserved material and to published illustrations. We focus on more general features of species that facilitate discrimination in our census studies.

This genus appears to arise in low latitudes in the Middle Eocene (*A. gnomabax*) and migrate to high latitudes in the late Middle to Late Eocene, with *A. rossi* (Late Eocene – Late Oligocene) largely and *A. catalexis* (Late Oligocene – Miocene) completely restricted to high latitudes.

***Aphetocyrtis bianulus* (O'Connor) [Plate 5, fig. 1]**

Theocorys bianulus O'Connor 1997, p. 84, pl. 4, fig. 1-4; pl. 10, fig. 1-4; pl. 11, fig. 5;

Sanfilippo and Fourtanier 2003, p. 12, pl. P1, fig. 9-13.

Aphetocyrtis bianulus (O'Connor), Suzuki et al. 2009, pl. 18, fig. 16a-b.

Closely resembles *A. rossi*, differing by having two distinctive bulges in the abdomen. High-latitude.

***Aphetocyrtis gnomabax* Sanfilippo and Caulet [Plate 5, fig. 2-7]**

Calocyclus semipolita Clark and Campbell, Caulet 1991, p. 537, pl. 4, fig. 9.

Aphetocyrtis gnomabax Sanfilippo and Caulet 1998, p. 16, pl. 2, fig. 6, 7, 10, 11, 14-17; pl. 7, fig. 10-13.

? *Cyrtocapsella* aff. *japonica* (Nakaseko), Takemura 1992, p. 746, pl. 1, fig. 11-12.

non *Calocyclus semipolita* Clark and Campbell, Strong et al. 1995, p. p. 208, fig. 10V [*L. semipolita*?].

non *Calocyclus* cf. *semipolita* Clark and Campbell, Abelman 1990, p. 697, pl. 7, fig. 4 [*A. catalexis*].

Thorax is bell-shaped to narrowly hemispherical. Pores of thorax and abdomen are of similar size. Abdomen is subcylindrical to inverted conical with a ragged basal margin.

***Aphetocyrtis rossi* Sanfilippo and Caulet** [Plate 5, fig. 8-11]

Calocyclus semipolita? Clark and Campbell, Chen 1975, p. 459, pl. 6, fig. 3-6.

Calocyclus asperum (Ehrenberg), Caulet 1992, p. 537, pl. 4, fig. 8, fig. 4, no. 4 (not no. 3)

Calocyclus cf. *semipolita* Clark and Campbell, Takemura 1992, p. 745, pl. 4, fig. 5-6;

Takemura and Ling 1997, p. 111, pl. 1, fig. 16.

Calocyclus semipolita Clark and Campbell group, Crouch and Hollis 1996, p. 26; Hollis et al.

1997, p. 58, pl. 5, fig. 11-12.

Cyrtocapsella robusta Abelman, Hollis et al., 1997, p. 60, pl. 5, fig. 13-14.

Amphetocyrtis rossi Sanfilippo and Caulet 1998, p. 18, pl. 2, fig. 8-9, 12-13; pl. 7, fig. 1-9.

Aphetocyrtis cf. *rossi* Sanfilippo and Caulet, Suzuki et al. (2009), p. 260, pl. 18, fig. 17.

Hemispherical thorax. Subcylindrical abdomen with pores slightly larger than on thorax, narrows distally, may be closed with a lattice plate. Distinguished from *A. gnomabax* and *C. robusta* by its hemispherical thorax. Distinguished from *Lophocyrtis* (*Apoplanius*) *aspera* by thorax and abdomen being of similar thickness. The two morphotypes illustrated by Hollis et al. (1997) appear to represent end-member variation within the species from slender forms with the cephalis free of the thoracic wall and a long cylindrical abdomen, open distally ("*C. semipolita*"), to more squat forms with the cephalis partly enclosed in the thoracic wall and a short abdomen, which is closed or has a basal lattice plate ("*C. robusta*"). High-latitude.

Genus *CLINORHABDUS* Sanfilippo and Caulet

Comprises the lineage *Clinorhabdus anantomus* -- *C. ocymora* -- *C. robusta* -- *C. longithorax*. Distinguished from other lophocyrtids by a short apical horn (when present), a bell-shaped or truncate conical thorax, and an abdomen that is subcylindrical and commonly closed. This genus is endemic to the Antarctic and ranges from the Middle Eocene to the Early Miocene.

***Clinorhabdus anantomus* Sanfilippo and Caulet** [Plate 5, fig. 12, 13]

Clinorhabdus anantomus Sanfilippo and Caulet 1998, p. 20, pl. 1, fig. 14-25; pl. 8, fig. 1a-b, 3a-b.

Apical spine, vertical spine and apophyses from mitral arches protrude from cephalis as, respectively, a moderately robust apical horn, a short spines and several short thorns. Thorax has a distinct neck and primary dorsal and lateral spines may protrude as short spines. Abdomen thin-walled with a ragged distal margin. High-latitude.

Genus *LOPHOCYRTIS* Haeckel 1887 sensu Sanfilippo 1990

The genus includes *Lophocyrtis jacchia* and all of its descendants. Most species have three segments, pores of thorax quincuncially arranged; many have an apical horn, three feet and a closed abdomen.

Subgenus *APOPLANIUS* Sanfilippo and Caulet

The subgenus comprises the lineage *L. (A.) klydus* -- *L. (A.) keraspera* -- *L. (A.) aspera* -- *L. (A.)*

nomas. The transition from *L. (A.) klydus* to *L. (A.) aspera* includes a shortening and reduction in number of spines protruding from the cephalis; an expansion of the proximal thorax with overall shape changing from narrowly campanulate to hemispherical, and reduction in the abdomen, in length, width and test thickness.

The first three members of the lineage span the early Eocene to Oligocene in low latitudes and co-occur in the Middle Eocene in high latitudes.

***Lophocyrtis (Apoplanius) aspera* (Ehrenberg), emend. Sanfilippo and Caulet** [Plate 5, fig. 14a, b-16]

Eucyrtidium asperum Ehrenberg 1874, p. 226; 1876, pl. 8, fig. 15.

Calocyclus asperum (Ehrenberg), Petrushevskaya and Kozlova 1972, p. 548, pl. 28, fig. 16-18; Strong et al. 1995, p. 208, fig. 9E, Crouch and Hollis 1996, p. 26.

Theocyrtis diabloensis (Clark and Campbell), Chen 1975, p. 459, pl. 5, figs. 4-6 (not fig. 7).

Calocyclus sp. B, Takemura 1992, p. 745, pl. 5, fig. 13; Takemura and Ling 1997, fig. 11, pl. 1, fig. 15; Hollis et al. 1997, p. 58, pl. 5, fig. 7-10.

Lophocyrtis (Apoplanius) aspera (Ehrenberg), Sanfilippo and Caulet 1998, p. 14, pl. 3A, fig. 5-10, pl. 3B, fig. 1, 2, 5-9, pl. 6, fig. 6-8. C.

Hemispherical thorax. Inverted truncate conical abdomen is thin walled and has a row of larger pores below the lumbar stricture. High-latitude.

***Lophocyrtis (Apoplanius) keraspera* Sanfilippo and Caulet** [Plate 5, fig. 17-19]

Lophocyrtis (Apoplanius) keraspera Sanfilippo and Caulet 1998, p. 14, pl. 3a, fig. 13-15, pl. 3b, fig. 12-14, pl. 6, fig. 9-12.

This species is distinguished by a strong apical horn, short vertical horn, hemispherical thorax with short spines projecting from lower part, and subcylindrical to truncate conical abdomen.

Subgenus *LOPHOCYRTIS* Haeckel emend. Sanfilippo and Caulet 1998

Species have three segments. Subspherical cephalis is perforate cephalis and bears a well-developed apical horn, which may bear thorns distally. Inflated hemispherical thorax has a rough, sometimes thorny surface. Subcylindrical to inverted truncate conical abdomen has a proximal row of larger pores, no distinct peristome. Some species have three terminal or subterminal feet. Primarily a tropical subgenus with incursions in high latitudes in the Middle Eocene.

***Lophocyrtis (Lophocyrtis) jacchia hapsis* Sanfilippo and Caulet** [Plate 5, fig. 20-22]

Calocyclus sp. C, Takemura 1992, p. 745, pl. 7, figs. 3-4.

Lophocyrtis (Lophocyrtis) jacchia hapsis Sanfilippo and Caulet, 1998, p. 10, pl. 5, fig 6, 7a-b, 9.

Lophocyrtis haywardi O'Connor 1999a, pl. 3, fig. 27-31.

High-latitude variety of *L. jacchia jacchia*.

Subgenus *PARALAMPTERIUM* Sanfilippo

Lophocyrtis (Paralamptერიum) dimitricai Sanfilippo

Lophocyrtis (Paralamptერიum) dimitricai Sanfilippo 1990, p. 308, pl. 3, fig. 7-13; O'Connor 1993, p. 74, pl. 8, fig. 18 19; Strong *et al.* 1995, p. 208, fig. 11n; Crouch and Hollis 1996, p. 26; Hollis *et al.* 1997, p. 62, pl. 6, fig. 19; Suzuki *et al.* 2009, p. 260, pl. 19, fig. 8a-b.

Cosmopolitan.

Lophocyrtis (Paralamptერიum) longiventer (Chen) [Plate 5, fig. 23, 24]

Cyclamptერიum? longiventer Chen 1975, p. 459, pl. 10, fig. 7.

Lophocyrtis (Paralamptერიum) longiventer (Chen), Sanfilippo 1990, p. 309, pl. 3, fig. 1-5; Strong *et al.* 1995, p. 208, fig. 11m; Crouch and Hollis 1996, p. 26; Hollis *et al.* 1997, p. 63, pl. 6, fig. 13-17; Takemura and Ling, 1997, pl. 1, fig. 12 (and fig. 13); Suzuki *et al.* 2009, p. 260, pl. 19, fig. 7a-b.

Cyclamptერიum milowi Riedel and Sanfilippo, Takemura 1992, p. 745, pl. 5, fig. 1-3.

Family PTEROCORYTHIDAE Haeckel, emend. Moore 1972

Genus *CRYPTOCARPIUM* Sanfilippo and Riedel

Cryptocarpium bussonii (Carnevale) group, New Group [Plate 5, fig. 25a, b, 26a, b]

Sethocorys bussonii Carnevale 1908, p. 31, pl. 4, fig. 17.

Sethocorys cristata Carnevale 1908, p. 31, pl. 4, fig. 18, 19.

Dictyocephalus bergontianus Carnevale 1908, p. 32, pl. 4, fig. 20.

Dictyocephalus crassus Carnevale 1908, p. 32, pl. 4, fig. 21.

Carpocanistrum spp. Riedel and Sanfilippo 1971 (in part), p. 1596, pl. 3D, fig. 4, 8, 9; Abelman 1990, p. 695, pl. 5, fig. 13; Takemura 1992, p. 744, pl. 3, fig. 5, 6; O'Connor 1993 (in part), p. 57, pl. 5, fig. 14; Crouch and Hollis 1996, p. 26; Hollis *et al.*, 1997, p. 66, pl. 6, fig. 26-27; Takemura and Ling 1997, p. 111.

Tricolocapsa bergontiana (Carnevale). Caulet 1986, p. 854.

Subconical to ovate test of 2 segments. Cephalis weakly lobed, ovate, imperforate, no apical spine. Thorax egg-shaped to subconical, basal aperture surrounded by a thick peristome. Thoracic pores quincuncially arranged in hexagonal frames. Cosmopolitan.

Cryptocarpium ornatum (Ehrenberg)

Cryptoprora ornata Ehrenberg, Sanfilippo *et al.* 1985, p. 693, fig. 27.2a, b.

Phormocyrtis proxima (?) Clark and Campbell, Chen 1975, p. 456, pl. 2, fig. 6.

Cryptocarpium ornatum (Ehrenberg), Hollis *et al.*, 1997, p. 66, pl. 6, fig. 23-25.

Phormocyrtis proxima Clark and Campbell, Suzuki *et al.* (2009), p. 264, pl. 22, fig. 10-11

Cosmopolitan.

Genus *LAMPROCYCLAS* Haeckel

Lamprocyclas particollis O'Connor [Plate 5, fig. 27]

? *Calocycletta parva* (Moore). Caulet 1986, p. 852.

Lamprocyclas “*particollis*” O'Connor, Hollis et al. 1997, p. 66, pl. 7, fig. 1-7 (*nom. nud.*).

Lamprocyclas particollis O'Connor, 1999a, pl. 3, fig. 5-11; pl. 6, fig. 19a-23.

Distinguished from *L. matakoe* by pores of the thorax and abdomen being of equivalent size, and a weakly differentiated peristome; from *C. parva* by an elongate, rather than ovate, cephalis with a eucephalic lobe that extends into a markedly bladed apical horn; from *Anthocyrtidium odontatum* O'Connor (1994, p. 342, pl. 2, fig. 5-8; pl. 4, fig. 1-5) by a distinct lumbar stricture and a thorax which tends to be campanulate rather than hemispherical.

Genus *STICHOPILIUM* Haeckel

Stichopilium cf. bicornе (Haeckel) [Plate 5, fig. 28a, b-29a, b]

Stichopilium bicornе Haeckel 1887, p. 1437, pl. 77, fig. 9; Takahashi, 1981, p. 254, pl. 39, figs. 13-19.

Stichopilium bicornе? Lazarus 1992, pl. 9, fig. 9-10, 12-17.

Unlike to the original description, the species encountered in this study consist of a thorax and abdomen that are fairly narrow and short. Our species are similar to Lazarus' species.

Genus *THEOCYRTIS* Haeckel

Theocyrtis tuberosa Riedel [Plate 5, fig. 30]

Theocyrtis tuberosa Riedel, 1959, p.298, pl.2, figs.10-11; Sanfilippo *et al.* 1985 (in part), p. 701, fig. 32.1a-b; Takemura 1992, p. 744, pl. 6, fig. 1, 2; O'Connor 1993 (in part), p. 70, pl. 7, fig. 22; Takemura and Ling 1997, p. 114.

Theocyrtis aff. *tuberosa* Riedel, Strong *et al.* 1995, p. 209, fig. 10y; Hollis et al. 1997, p. 67.

The early form of *T. tuberosa* described by Strong et al. (1995) has well-developed plicae but lacks tubercles on the thorax.

Cosmopolitan.

Family **SETHOPHORMIDIDAE**

All taxa assigned to this family are having a wide conical to flat dicyrtid shape with a large cephalis, which is not always preserved.

REFERENCES

- Abelmann, A.: Oligocene to Middle Miocene radiolarian stratigraphy of southern high latitudes from Leg 113, sites 689, and 690, Maud Rise, in: Baker, P.F., Kennett, J.P., et. al. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*. College Station, TX (Ocean Drilling Program), pp. 675-708, 1990.
- Apel, M., Kiessling, W., Böhm, F., & Lazarus, D.: Radiolarian faunal characteristics in Oligocene sediments of the Kerguelen Plateau, Leg 183, Site 1138. *Proc. Ocean Drill. Prog. Sci. Results*, 2002.
- Bailey, J. W.: Notice of microscopic forms found in the soundings of the Sea of Kamtschatka - with a plate, *American Journal of Science and Arts*, Series 2, 22, 1-6, 1856.
- Benson, R. N.: Recent Radiolaria from the Gulf of California, Ph.D. Thesis, Minnesota University, University, 1-577 pp., 1966.
- Bütschli, O.: Beiträge zur Kenntnis der Radiolarienskelette, insbesondere der Cyrtida, *Zeitschrift für Wissenschaftliche Zoologie*, 36, 485-540, 1882.
- Bjørklund, K. R.: Radiolaria from the Norwegian Sea, Leg 38 of the Deep Sea Drilling Project. In: Talwani, M. et al. (editors). *Initial Reports of the Deep Sea Drilling Project*. Washington (U.S. Government Printing Office), 38: 1101-1168, 1976.
- Blueford, J.: Radiolarian biostratigraphy of siliceous Eocene deposits in central California, *Micropaleontology*, 34, 236-258, 1988.
- Campbell, A. S.: New genera and subgenera of Radiolaria, *Journal of Paleontology*, 25, 527-530, 1951.
- Campbell, A. S. and Clark, B.L.: Miocene radiolarian faunas from southern California. *Geological Society of America*, special Papers 51, 1-76, 1944.
- Carnevale, P.: Radiolarie e silicoflagellati di Bergonzano (Reggio Emilia), *Memorie del Reale Istituto Veneto di Scienze, Lettere ed Arti*, 28, 1-46, 1908.
- Caulet, J. P.: Radiolarians from the Southwest Pacific, in: Kennett, J.P., Von der Borch, et. al, *Initial Reports of the Deep Sea Drilling Project*, Vol. 90, U.S. Government Printing Office, Washington, D.C., 835-861, 1986.
- Chen, P. H.: Antarctic Radiolaria, in: Hayes, D.E., Frakes, L.A., et al., *Initial Reports of the Deep Sea Drilling Project*, Vol. 28, U.S. Government Printing Office, Washington, D.C., 437-513, 1975.
- Clark, B. L., and Campbell, A. S.: Radiolaria from the Kreyenhagen Formation near Los Baños, California, *Geological Society of America*, Memoir, 101, i-vii, 1-66, 1945.
- Crouch, E. M., and Hollis, C. J.: Paleogene palynomorph and radiolarian biostratigraphy of DSDP Leg 29, sites 280 and 281 South Tasman Rise, *Institute of Geological and Nuclear Sciences science report 96/19*, 46p., 1996.
- De Wever, P., Dumitrica, P., Caulet, J. P., and Caridroit, M.: Radiolarians in the sedimentary record, *Gordon and Breach Science Publishers*, Amsterdam, 533 pp., 2001.
- Dinkelman, M. G.: Radiolarian stratigraphy: Leg 16, Deep Sea Drilling Project, in: Van Andel, T. H., Heath, G. R., and et al., *Initial Reports of the Deep Sea Drilling Project*, U.S. Government Printing Office, Washington, D.C., 747-813, 1973.
- Dreyer, F.: Die Pylombildungen in vergleichend-anatomischer und entwicklungsgeschichtlicher Beziehung bei Radiolarien und bei Protisten überhaupt, *Jenaische Zeitschrift für Naturwissenschaft*, 23, 1-138, 1889.
- Dumitrica, P.: Paleocene Radiolaria, DSDP Leg 21, in: Burns, R.E., Andrews, J.E., et al., *Initial Rep Deep Sea*. U.S. Government Printing Office, Washington, D.C., pp. 787-817, 1973.

- Ehrenberg, C. G. (1838): Über die Bildung der Kreidefelsen und des Kreidemergels durch unsichtbare Organismen, Abhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 59-147, 1839.
- Ehrenberg, C. G.: Über die mikroskopischen kieselschaligen Polycystinen als mächtige Gebirgsmasse von Barbados und über das Verhältniss der aus mehr als 300 neuen Arten bestehenden ganz eigenthümlichen Formengruppe jener Felsmasse zu den jetzt lebenden Thieren und zur Kreidebildung. Eine neue Anregung zur Erforschung des Erdlebens. Ber. Königl. Preuß. Akad. Wiss. Berlin, 40-60, 1847.
- Ehrenberg, C. G.: Die systematische Charakteristik der neuen mikroskopischen Organismen des tiefen atlantischen Oceans, Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 236-250, 1854a.
- Ehrenberg, C. G.: Mikrogeologie. Das Erden und Felsen schaffende Wirken des unsichtbar kleinen selbstständigen Lebens auf der Erde. Leopold Voss, Leipzig, Germany, xxviii + 374pp, 1854b.
- Ehrenberg, C. G. (1861): Über die Tiefgrund-Verhältnisse des Oceans am Eingange der Davisstrasse und bei Island, Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 275-315, 1862.
- Ehrenberg, C. G. (1873): Grossere Felsproben des Polycystinen-Mergels von Barbados mit weiteren Erläuterungen. Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 213-263, 1874.
- Ehrenberg, C. G. (1875): Fortsetzung der mikrogeologischen Studien als Gesamt-Uebersicht der mikroskopischen Paläontologie gleichartig analysirter Gebirgsarten der Erde, mit specieller Rücksicht auf den Polycystinen-Mergel von Barbados. Abhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 1-225, 1876.
- Foreman, H. P.: Radiolaria of Leg 10 with systematics and ranges for the families Amphipyndacidae, Artostrobiidae and Theoperidae, in: Worzel, J.L., Bryant, W., et, al., Initial Rep Deep Sea. U.S. Government Printing Office, Washington, D.C., pp. 407-474, 1973.
- Funakawa, S.: Plagiacanthidae (Radiolaria) from the Upper Miocene of eastern Hokkaido, Japan, Transactions and Proceedings of the Palaeontological Society of Japan, new Series, 174, 458-483, 1994.
- Funakawa, S., Nishi, H., Moore, T. C., & Nigrini, C. A.: Data report: Late Eocene–early Oligocene radiolarians, ODP Leg 199 Holes 1218A, 1219A, and 1220A, central Pacific, in: P. A. Wilson, M. Lyle & J. V. Firth, Proceedings of the Ocean Drilling Program, Scientific Results (Vol. 199, pp. 1-74), 2006.
- Goll, R. M.: The Neogene evolution of Zygocircus, Neosemantis and Callimitra: their bearing on nassellarian classification. A revision of the Plagiacanthoidea. Micropaleontology, 25, 365-396, 1979.
- Haeckel, E.: Naturliche Schöpfungsgeschichte, 7th ed., Reimer, Berlin, Germany, 718 pp., 1879.
- Haeckel, E.: Entwurf eines Radiolarien-Systems auf Grund von Studien der Challenger-Radiolarien, Jenaische Zeitschrift für Naturwissenschaft, 15, 418-472, 1882.
- Haeckel, E.: Report on the Radiolaria collected by H.M.S. Challenger during the years 1873-1876. Report on the Scientific Results of the Voyage of the H.M.S. Challenger, Zoology 18, clxxxviii + 1803, 1887.
- Hays, J. D.: Radiolaria and late Tertiary and Quaternary history of Antarctic Seas, in: Biology of Antarctic Seas II, American Geophysical Union, Antarctic Research Series 5, Llano, G. A., 125-184, 1965.

- Hollis, C. J.: Latest Cretaceous to Late Paleocene Radiolaria from Marlborough (New Zealand) and DSDP Site 208, The University of Auckland, ResearchSpace@Auckland, 1991.
- Hollis, C. J.: Latest Cretaceous to Late Paleocene radiolarian biostratigraphy: A new zonation from the New Zealand region. *Mar Micropaleontol* 21, 295-327, 1993.
- Hollis, C. J.: Cretaceous-Paleocene Radiolaria from eastern Marlborough, New Zealand. Institute of Geological and nuclear Sciences, Monograph 17, 1-152, 1997.
- Hollis, C. J.: Biostratigraphy and paleoceanographic significance of Paleocene radiolarians from offshore eastern New Zealand. *Mar Micropaleontol* 46, 265-316, 2002.
- Hollis, C. J., Waghorn, D. B., Strong, C. P., and Crouch, E. M.: Integrated Paleogene Biostratigraphy of DSDP Site 277 (Leg 29): Foraminifera, Calcareous Nannofossils, Radiolaria, and Palynomorphs, Institute of Geological & Nuclear Sciences Limited, 1997.
- Johnson, D. A.: Radiolaria from the eastern Indian Ocean, DSDP Leg 22, in: Von der Borch, C. C., Sclater, J. G., and et al., Initial Reports of the Deep Sea Drilling Project, U.S. Government Printing Office, Washington, D.C., 521-575, 1974.
- Kamikuri, S.-i., Moore, T. C., Lyle, M., Ogane, K., and Suzuki, N.: Early and Middle Eocene radiolarian assemblages in the eastern equatorial Pacific Ocean (IODP Leg 320 Site U1331): Faunal changes and implications for paleoceanography, *Mar. Micropaleontol.*, 77-108, 2012.
- Kozlova, G.E.: Paleogene Boreal Radiolarians from the Russia. Ministry of Natural resources of Russian Federation, All-Russian Petroleum research Exploration Institute (VNIGRI), 1999.
- Kruglikova, S. B.: Radiolarians in the core of station 4066 (northern part of the Pacific Ocean), in: Basic problems of Micropaleontology and the accumulation of organogenic sediments in oceans and seas, edited by: Jouse, A. P., Izdatelstvo Nauka, Akademiya Nauk SSSR, Okeanograficheskaya Komissiya, Moscow, USSR, 115-126, 1969.
- Kruglikova, S. B., 1977. Radiolaria. In: A.P. Jouse [ed.], Atlas of Microorganisms in Bottom Sediments of the Oceans: diatoms, radiolaria, silicoflagellates and coccoliths. Academy of sciences USSR, Publishing house "Nauka": p. 7, 13-17 (pl. 86-145).
- Lazarus, D. 1992. Antarctic Neogene radiolarians from the Kerguelen Plateau, Legs 119 and 120, in: Wise, S. W., Jr., Schlich, R., et al., 1992, Proceedings of the Ocean Drilling Program, Scientific Results, Vol. 120, pp. 785-809.
- Lazarus, D. B., Faust, K. and Popova-Goll, I.: New species of prunoid radiolarians from the Antarctic Neogene. *Journal of Micropaleontology*, vol. 24(2): 97-121, 2005.
- Lipman, R. K.: New Eocene Radiolaria from the Turgai trough and northern Aral region, in: New species of ancient plants and invertebrates in the USSR, Nauka, Moscow, USSR, 42-56, 1972.
- Loeblich, A. R., and Tappan, H.: Remarks on the systematics of the Sarkodina (Protozoa), renamed homonyms and new validated genera, *Proceedings of the Biological Society of Washington*, 74, 213-234, 1961.
- Lombardi, G., and Lazarus, D. B.: Neogene cycladophorid radiolarians from the North Atlantic, Antarctic, and North Pacific deep-sea sediments, *Micropaleontology*, 34, 97-135, 1988.
- Mamedov, N. A.: New representatives of Eocene radiolarians in Azerbaidzhan, in: Fossil and Recent Radiolarians: Materials of the Second All Union Seminar on Radiolaria, edited by: Vyalov, O. S., Izdatelstvo Lvovskogo Universiteta (Lvov University), Lvov, USSR, 94-101, 1969.
- Mast, H.: Die Astrosphaeriden. *Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition auf dem Dampfer "Valdivia" 1898-1899*, vol. 19 (4): 123-190, 1910.
- Moore, T. C.: Mid-Tertiary evolution of the radiolarian genus *Calocycletta*, *Micropaleontology*, 18, 144-152, 1972.

- Nakaseko, K.: Neogene Cyrtioidea (Radiolaria) from the Isozaki Formation in Ibaraki Prefecture, Japan, *Science Reports, College of General Education Osaka University*, 12, 165-198, 1963.
- Nigrini, C.: Radiolaria in pelagic sediments from the Indian and Atlantic Oceans, *Bulletin of the Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California*, 11, 1-125, 1967.
- Nigrini, C.: Radiolaria from eastern tropical Pacific sediments, *Micropaleontology*, 14, 51-63, 1968.
- Nigrini, C.: Tropical Cenozoic Artostrobiidae (Radiolaria). *Micropaleontology* 23, 241-269, 1977.
- Nigrini, C., and Moore, T. C.: A guide to modern Radiolaria, Cushman Foundation for foraminiferal Research, special Publication, 257 pp., 1979.
- Nishimura, A.: Cenozoic Radiolaria in the western North Atlantic, Site 603, Leg 93 of the Deep Sea Drilling Project, in: *Initial Reports of the Deep Sea Drilling Project*, edited by: Van Hinte, J. E., Wise, S. W. J., and et, a., U.S. Government Printing Office, Washington, D.C., 713-737, 1987.
- Nishimura, A.: Paleocene radiolarian biostratigraphy in the northwest Atlantic at Site 384, Leg 43, of the Deep Sea Drilling Project. *Micropaleontology* 38, 317-362, 1992.
- O'Connor, B. M.: Radiolarian from the Mahurangi Limestone, Northland, New Zealand, University of Auckland, Auckland, 136 pp., 1993.
- O'Connor, B. M.: Seven new radiolarian species from the Oligocene of New-Zealand. *Micropaleontology* 40, 337-350, 1994.
- O'Connor, B. M.: New Radiolaria from the Oligocene and Early Miocene of Northland, New Zealand. *Micropaleontology* 43, 63-100, 1997.
- O'Connor, B. M.: Radiolaria from the Late Eocene Oamaru Diatomite, South Island, New Zealand. *Micropaleontology* 45, 1-55, 1999a.
- O'Connor, B. M.: Distribution and biostratigraphy of latest Eocene to latest Oligocene Radiolaria from the Mahurangi Limestone, Northland, New Zealand. *New Zealand Journal of Geology and Geophysics* 42, 489-511, 1999b.
- O'Connor, B. M.: Stratigraphic and geographic distribution of Eocene Miocene Radiolaria from the southwest Pacific. *Micropaleontology* 46, 189-228, 2000.
- Ogane, K., Suzuki, N., Aita, Y., Sakai, T., Lazarus, D.: Ehrenberg's Radiolarian Collections from Barbados, in: Tanimura, Y., Aita, Y. (Eds.), *JOINT HAECKEL and EHRENBURG PROJECT: Reexamination of the Haeckel and Ehrenberg Microfossil Collections as a Historical and Scientific Legacy*. National Museum of Nature and Science Monographs, pp. 97-106, 2009.
- Petrushevskaya, M. G.: Antarctic Spumelline and Nasselline radiolarians, in: *Issledovaniya Fauny Morei, Resultaty Biologicheskikh Issledovaniy Sovetskoi Antarkticheskoi Ekspeditsii 1955-1958*, Zoologicheskii Institut Akademii Nauk SSSR, 5-186, 1967.
- Petrushevskaya, M. G.: On the natural system of polycystine Radiolaria (Class Sarcodina), in: *Proceedings of the II Planktonic Conference, Roma 1970*, edited by: Farinacci, A., Edizioni Tecnoscienza, Roma, Italy, 981-992, 1971a.
- Petrushevskaya, M. G.: Nassellarian radiolarians in the plankton of the World Ocean, *Akademiya nauk SSSR, Zoologicheskii Institut, Issledovaniya Fauny Morei*, 9, 294, 1971b.
- Petrushevskaya, M. G.: Nassellarian radiolarians from the world oceans, in: *Publications of the Zoological Institute, Academy of Sciences of the USSR, Description of the Fauna of the USSR*, Nauka, Leningradskoe Otdelenie, Leningrad, USSR, 405, 1981.
- Petrushevskaya, M. G.: Cenozoic radiolarians of the Antarctic, Leg 29, DSDP, in: Kennet, J.P.,

- Houtz, R.E., et. al. (Eds.), Initial Rep Deep Sea. U.S. Government Printing Office, Washington, D.C., pp. 541-675, 1975.
- Petrushevskaya, M. G.: New species of the radiolarians of the order Nassellaria, in: Exploration of the Fauna of the Seas - New Species and Genera of Marine Invertebrates, *Zoologicheskii Zhurnal*, Akademia Nauk SSSR, Leningrad, USSR, 10-19, 1977.
- Petrushevskaya, M. G. and Kozlova, G.E.: Radiolaria: Leg 14, Deep Sea Drilling Project, in: Hayes, D.E., Pimm, A.C., et. al. (Eds.), Initial Rep Deep Sea. U.S. Government Printing Office, Washington, D.C., pp. 495-648, 1972.
- Petrushevskaya, M. G., and Kozlova, G. E.: Description of the radiolarian genera and species, in: The history of the microplankton of the Norwegian Sea (on the Deep Sea Drilling materials), edited by: Sterlkov, A. A., and M.G., P., Exploration of the Fauna of the Seas, Nauka Academy of Sciences of the USSR, Zoological Institute, Leningrad, USSR, 86-157, 1979.
- Popofsky, A.: Die Sphaerellarien des Warmwassergebietes, in: Deutsche Sudpolar-Expedition, 1901-1903, edited by: Drygalski, E., Georg Reimer, Berlin, Germany, 73-159, 1912.
- Principi, P.: Contributo allo studio dei Radiolari Miocenici Italiani, *Bollettino della Societa Geologica Italiana*, 28, 1-22, 1909.
- Riedel, W. R.: Radiolaria in Antarctic sediments, Reports of the B.A.N.Z. Antarctic Research Expedition, series B, 6, 217-255, 1958.
- Riedel, W. R.: Oligocene and Lower Miocene Radiolaria in tropical Pacific sediments, *Micropaleontology*, 5, 285-302, 1959.
- Riedel, W. R.: Subclass Radiolaria, in: The Fossil Record. A symposium with documentation, edited by: Harland, W. B., Holland, C. H., House, M. R., Hughes, N. F., Reynolds, A. B., Rudwick, M. J. S., Satterthwaite, G. E., Tarlo, L. B. H., and Willey, E. C., Geological Society of London, London, 291-298, 1967a.
- Riedel, W. R.: Some new families of Radiolaria, *Proceedings of the Geological Society of London*, 1640, 148-149, 1967b.
- Riedel, W. R. and Sanfilippo, A.: Radiolaria, Leg 4, Deep Sea Drilling Project, in: Bader, R.G., et. al. (Eds.), Initial Rep Deep Sea. U.S. Government Printing Office, Washington, D.C., pp. 503-575, 1970.
- Riedel, W. R. and Sanfilippo, A.: Cenozoic Radiolaria from the western tropical Pacific, Leg 7, in: Winterer, E.L., Riedel, W.R., et. al. (Eds.), Initial Rep Deep Sea. U.S. Government Printing Office, Washington, D.C., pp. 1529-1672, 1971.
- Sanfilippo, A.: Origin of the subgenera *Cyclampterium*, *Paralampterium* and *Sciadiopeplus* from *Lophocyrtis* (*Lophocyrtis*) (*Radiolaria*, *Theoperidae*). *Mar Micropaleontol* 15, 287-312, 1990.
- Sanfilippo, A., and Blome, C. D.: Biostratigraphic implications of mid-latitude Palaeocene-Eocene radiolarian faunas from Hole 1051A, ODP Leg 171B, Blake Nose, western North Atlantic, Geological Society, London, Special Publications, 183, 185-224, 2001.
- Sanfilippo, A., and Caulet, J. P.: Taxonomy and evolution of Paleogene Antarctic and Tropical *Lophocyrtid* radiolarians, *Micropaleontology*, 44, 1-43, 1998.
- Sanfilippo, A. and Fourtanier, E.: Oligocene radiolarians, diatoms, and ebridians from the Great Australian Bight (ODP Leg 182, Site 1128), in: Hine, A.C., Feary, D.A., Malone, M.J. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*, pp. 1-24, 2003.
- Sanfilippo, A. and Riedel, W.R.: Cenozoic Radiolaria (exclusive of theoperids, artostrobiids and amphipyndacids) from the Gulf of Mexico, DSDP Leg 10, in: Worzel, J.L., Bryant, W., et. al. (Eds.), Initial Rep Deep Sea. U.S. Government Printing Office, Washington, D.C., pp. 475-611, 1973.

- Sanfilippo, A., Westberg-Smith, M.J., Riedel, W.R.: Cenozoic Radiolaria, in: Bolli, H.M., Saunders, J.B., Perch-Nielsen, K. (Eds.), *Plankton Stratigraphy*. Cambridge University Press, Cambridge/ New York/New Rochelle/Melbourne/Sidney, pp. 631-712, 1985.
- Shilov, V. V.: Eocene-Oligocene Radiolarians from Leg 145, North Pacific, in: *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by: Rea, D. K., Basov, I. A., Scholl, D. W., and Allan, J. F., College Station, TX (Ocean Drilling Program), 117-132, 1995.
- Stöhr, E.: Die Radiolarienfauna der Tripoli von Grotte, Provinz Girgenti in Sicilien, *Palaeontographica*, 26, 71-124, 1880.
- Suzuki, N., and Aita, Y.: Radiolaria: achievements and unresolved issues: taxonomy and cytology, *Plankton Benthos Researchs*, 6, 69-91, 2011.
- Suzuki, N., Ogane, K., Chiba, K.: Middle to Late Eocene polycystine radiolarians from the Site 1172, Leg 189, Southwest Pacific. *News of Osaka Micropaleontologists*, special volume 14, 239-296, 2009.
- Takahashi, K.: Radiolaria: Flux, Ecology, and Taxonomy in the Pacific and Atlantic, in: *Ocean Biocoenosis Series*, edited by: Honjo, S., Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, 1-303, 1991.
- Takemura, A.: Radiolarian Paleogene biostratigraphy in the southern Indian Ocean, Leg 120, in: Wise, S.W.J., Schlich, R., et. al. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*. College Station, TX (Ocean Drilling Program), pp. 735-756, 1992.
- Takemura, A. and Ling, H.Y.: Eocene and Oligocene radiolarian biostratigraphy from the Southern Ocean - correlation of ODP Legs 114 (Atlantic Ocean) and 120 (Indian Ocean). *Mar Micropaleontol* 30, 97-116, 1997.
- Vinassa de Regny, P. E.: Radiolari Miocenici Italiani, *Memorie della Reale Accademia delle Scienze dell'Istituto di Bologna*, 8, 227-257 (565-595), 1900.






Plate 1. Scale bars equal 100µm. Images of selected radiolarian species:

1) Actinommidae n.sp. A sensu Hollis (DSDP 277, 23-1, 33cm, slide A), **2) *Amphisphaera coronata* gr.** (DSDP277, 25-1, 60cm, slide A), **3) *Amphisphaera radiosa*** (DSDP277, 24-2, 20cm, slide B), **4a,b) *Amphisphaera* aff. *radiosa*** (DSDP277, 17-3, 33cm), **5) *Amphisphaera spinulosa*** (DSDP283, 8-2, 25cm, slide A), **6) *Amphisphaera?* *megapora*** (DSDP277, 25-1, 60cm, slide C), **7) *Stylosphaera minor*** (DSDP277, 32-3, 100cm), **8) *Zealithapium mitra*** (DSDP277, 23-1, 33cm, slide B), **9) *Axoprunum bispiculum*** (DSDP 277, 17-3, 33cm), **10, 11) *Axoprunum pierinei* gr.** (10: DSDP277, 25-1, 60cm, slide A; 11: DSDP281, 14-1, 100cm, slide A), **12) *Axoprunum?* *irregularis*** (DSDP277, 15-3, 100cm), **13) *Spongopyle osculosa*** (DSDP277, 24-2, 20cm, slide B), **14) *Amphicraspedum murrayanum*** (DSDP277, 32-3, 100cm), **15-17) *Amphicraspedum prolixum* gr.** (15: DSDP277, 22-1, 82cm, slide A; 16: DSDP280A, 6-2, 60cm, slide A; 17: DSDP277, 33-2, 100cm), **18, 19) *Amphymenium splendiararmatum*** (DSDP277, 32-3, 100cm), **20) *Larcopyle frakesi*** (DSDP280A, 5-1, 104cm, slide C), **21) *Larcopyle hayesi*** (DSDP277, 24-2, 20cm, slide B), **22) *Larcopyle labyrinthusa*** (DSDP280A, 5-1, 104cm, slide C), **23, 24) *Larcopyle polyacantha* gr.** (23: DSDP280A, 5-1, 104cm, slide C; 24: DSDP277, 24-2, 20cm, slide B), **25a,b) *Larcopyle* cf. *pylomaticus*** (DSDP280A, 5-1, 104cm, slide C), **26-28) *Lithelius minor* gr.** (26: DSDP277, 32-3, 100cm; 27-28: DSDP277, 24-2, 20cm, slide B), **29) *Sphaeropyle tetrapila*** (DSDP277, 18-3, 20cm).




1




2




3




4




5




6




7




8




9




10



11



12




13




14




15a



16



17




18




22



15b




19




20



21



23




24



25



26



27

Plate 2. Scale bars equal 100µm. Images of selected radiolarian species:

1) *Amphicentria* sp. 1 sensu Suzuki (DSDP277, 22-1, 82cm, slide B), **2) *Callimitra* aff. *atavia*** (DSDP277, 22-1, 82cm), **3-5) *Ceratocyrtis* spp.** (3-4: DSDP280A, 5-1, 104cm, slide C, B; 5: DSDP283, 7-1, 20cm, slide A), **6-8) *Lithomelissa challengeræ*** (DSDP280A, 6-2, 60cm, slide A), **9) *Lithomelissa* cf. *challengeræ*** (DSDP277, 23-1, 33cm, slide B), **10-11) *Lithomelissa ehrenbergi*** (10: DSDP280A, 5-1, 104cm, slide B; 11: DSDP283, 6-2, 40cm, slide A), **12, 13) *Lithomelissa gelasinus*** (DSDP280A, 5-1, 104cm, slide A), **14) *Lithomelissa* cf. *haeckeli*** (DSDP277, 22-1, 82cm, slide B), **15a,b) *Lithomelissa macroptera*** (DSDP283, 5-2, 25cm, slide A), **16) *Lithomelissa robusta*** (DSDP280A, 7-1, 90cm, slide A), **17) *Lithomelissa sphaerocephalis*** (DSDP280A, 7-3, 40cm, slide A), **18) *Lithomelissa tricornis*** (DSDP280A, 7-1, 90cm, slide A), **19) *Lithomelissa?* *sakai*** (DSDP280A, 5-1, 104cm, slide B), **20) *Pseudodictyophimus galeatus*** (DSDP280A, 7-3, 40cm, slide A), **21-23) *Pseudodictyophimus gracilipes* gr.** (21-22: DSDP280A, 7-1, 90cm, slide A; 23: DSDP283, 5-2, 25cm, slide B), **24-27) *Pseudodictyophimus* spp.** (24: DSDP280A, 7-3, 40cm, slide A; 25-26: DSDP280A, 7-1, 90cm, slide A; 27: DSDP280A, 5-1, 104, slide A).




Plate 3. Scale bars equal 100µm. Images of selected radiolarian species:

1-4) *Botryocella* sp. A sensu Apel (DSDP280A, 5-1, 104cm, slide A), **5-7) *Glycobotrys nasuta* gr.** (5: DSDP280A, 5-1, 104cm, slide A; 6: DSDP277, 25-1, 60cm, slide A; 7: DSDP277, 24-2, 20cm, slide A), **8-10) *Siphocampe* (?) *acephala* gr.** (DSDP277, 25-1, 60cm, slide A), **11, 12) *Siphocampe?* *amygdala*** (11: DSDP277, 24-2, 20cm, slide B; 12: DSDP277, 23-1, 33cm, slide A), **13) *Artostrobos* cf. *pretabulus*** (DSDP277, 23-1, 33cm, slide A), **14-16) *Aspis* sp. A** (14: DSDP 277, 22-1, 82; 15: DSDP 283, 7-5, 94cm, slide A; 16: DSDP 280A, 6-2, 60cm, slide A), **17) *Cycladophora cosma cosma*** (DSDP280A, 7-3, 40cm, slide A), **18) *Cycladophora humerus*** (DSDP280A, 5-1, 104cm, slide A), **19) *Eucyrtidium antiquum*** (DSDP280A, 7-1, 90cm, slide A), **20a,b) *Eucyrtidium nishimurae*** (DSDP283, 8-2, 25cm, slide A), **21) *Eucyrtidium spinosum*** (DSDP283, 7-5, 94cm, slide A), **22) *Eucyrtidium ventriosum*** (DSDP277, 23-1, 33cm, slide A), **23a,b) *Eurystomoskevos cauleti*** (DSDP280A, 7-3, 40cm, slide A), **24) *Eurystomoskevos petrushevskaae*** (DSDP277, 23-1, 33cm, slide A), **25) *Eusyringium fistuligerum*** (DSDP277, 33-2, 20cm, slide A), **26a,b) *Sethocyrtis chrysallis*** (DSDP277, 32-3. 100cm), **27) *Thyrsocyrtis pinguisicoides*** (DSDP277, 23-1, 33cm, slide A).




Plate 4. Scale bars equal 100µm. Images of selected radiolarian species:

1a,b, 2) *Dictyophimus? archipilium* (DSDP283, 7-1, 20cm, slide A), **3a,b-8) *Dictyophimus? aff archipilium*** (3: DSDP277, 17-3, 33cm; 4-6: DSDP283, 8-1, 40cm, slide B; 7-8: DSDP277, 23-1, 22cm, slide A), **9, 10) *Dictyophimus aff. constrictus*** (9: DSDP280A, 9-1, 90cm, slide A, 10: DSDP280A, 6-2, 60cm), **11a-c, 12) *Lychnocanium amphitrite*** (11: DSDP283, 5-2, 25cm, slide B; 12: DSDP277, 33-2, 20cm slide A), **13a,b, 14) *Lychnocanium babylonis*** (13: DSDP283, 8-2, 25cm, slide A; 14: DSDP277, 23-1, 33cm, slide A), **15, 16) *Lychnocanium bellum*** (DSDP277, 25-1, 60cm, slide A), **17) *Lychnocanium aff. carinatum*** (DSDP277, 26-1, 60cm, slide A), **18a,b) *Lychnocanium tetrapodium*** (DSDP277, 23-1, 33cm, slide A), **19, 20a,b) *Pterocodon apis*** (19: DSDP283, 7-1, 20cm, slide B; 20: DSDP283, 6-2, 40cm, slide A), **21) *Pteropilium aff. contiguum*** (DSDP277, 23-1, 33cm, slide A).




Plate 5. Scale bars equal 100µm. Images of selected radiolarian species:

1) *Aphetocyrtis bianulus* (DSDP277, 22-1, 82cm, slide A), **2-7)** *Aphetocyrtis gnomabax* (2: DSDP280A, 6-2, 60cm, slide A; 3-6: DSDP280A, 5-1, 104cm, slide A; 7: DSDP277, 24-2, 20cm, slide A), **8-11)** *Aphetocyrtis rossi* (8: DSDP283, 6-2, 40cm, slide A; 9: DSDP280A, 5-1, 104cm, slide A; 10: DSDP280A, 7-1, 90cm, slide A; 11: DSDP277, 23-1, 33cm, slide A), **12, 13)** *Clinorhabdus anantomus* (DSDP283, 7-1, 20cm, slide A), **14a,b-16)** *Lophocyrtis aspera* (14-15: DSDP283, 7-1, 20cm, slide A; 16: DSDP277, 24-2, 20cm, slide A), **17-19)** *Lophocyrtis keraspera* (DSDP277, 23-1, 33cm, slide A), **20-22)** *Lophocyrtis jacchia hapsis* (20: DSDP283, 8-1, 40cm, slide A; 21-22: DSDP277, 32-3, 100cm), **23, 24)** *Lophocyrtis longiventer* (23: DSDP280A, 5-1, 104cm, slide A; 24: DSDP283, 5-2, 25cm, slide B), **25a,b, 26a,b)** *Cryocarpium bussonii* gr. (DSDP277, 18-3, 20cm), **27)** *Lamprocyclas particollis* (DSDP277, 25-1, 60cm, slide A), **28a,b, 29a,b)** *Stichopilium cf. bicornis* (28: DSDP283, 7-1, 20cm, slide A; 29: DSDP283, 6-2, 40cm, slide A), **30)** *Theocyrtis tuberosa* (DSDP283, 5-2, 25cm, slide B).