

Revisiting carbonate chemistry controls on planktic foraminifera Mg/Ca: implications for sea surface temperature and hydrology shifts over the Paleocene–Eocene Thermal Maximum and Eocene–Oligocene Transition

D. Evans^{1,2}, B. S. Wade³, M. Henehan^{2,4}, J. Erez⁵, and W. Müller¹

¹Department of Earth Sciences, Royal Holloway University of London, Egham, TW20 0EX, UK

²Department of Geology and Geophysics, Yale University, New Haven, CT 06511, USA

³Department of Earth Sciences, University College London, London, UK

⁴Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton Waterfront Campus, European Way, Southampton SO14 3ZH, UK

⁵Earth Science Institute, The Hebrew University of Jerusalem, Israel

Correspondence to: D. Evans (d.evans@yale.edu)

Abstract. Much of our knowledge of past ocean temperatures comes from the foraminifera Mg/Ca palaeothermometer. Several non-thermal controls on foraminifera Mg incorporation have been identified, of which vital-effects, salinity and secular variation in seawater Mg/Ca are the most commonly considered. Ocean carbonate chemistry is also known to influence Mg/Ca, yet this is rarely examined as a source of uncertainty either because (1) precise pH and $[\text{CO}_3^{2-}]$ reconstructions are sparse, or (2) it is not clear from existing culture studies how a correction should be applied. We present new culture data of the relationship between carbonate chemistry for the surface-dwelling planktic species *Globigerinoides ruber*, and compare our results to data compiled from existing studies. We find a coherent relationship between Mg/Ca and the carbonate system and argue that pH rather than $[\text{CO}_3^{2-}]$ is likely to be the dominant control. Applying these new calibrations to datasets for the Paleocene–Eocene Thermal Maximum (PETM) and Eocene–Oligocene Transition (EOT) enable us to produce a more accurate picture of surface hydrology change for the former, and a reassessment of the amount of subtropical precursor cooling for the latter. We show that pH-adjusted Mg/Ca and $\delta^{18}\text{O}$ datasets for the PETM are within error of no salinity change, and that the amount of precursor cooling over the EOT has been previously underestimated by $\sim 2^\circ\text{C}$ based on Mg/Ca. Finally, we present new laser-ablation data of EOT-age *Turborotalia ampliapertura* from St. Stephens Quarry (Alabama), for which a solution ICPMS Mg/Ca record is available (Wade et al., 2012). We show that the two datasets are in excellent agreement, demonstrating that fossil solution and laser-ablation data may be directly comparable. Together with an advancing understanding of

20 the effect of $\text{Mg}/\text{Ca}_{\text{sw}}$, the coherent picture of the relationship between Mg/Ca and pH that we outline here represents a step towards producing accurate and quantitative palaeotemperatures using this proxy.

1 Introduction

The relationship between foraminifera Mg/Ca and temperature has been widely applied as a method
25 of reconstructing the thermal evolution of the oceans across a range of timescales (e.g. Mashiotta et al., 1999; Lear et al., 2000; Sosdian and Rosenthal, 2009; Wade et al., 2012). Whilst a substantial portion of our understanding of the response of ocean temperature, hydrology and circulation to external forcing is derived from this proxy, several non-thermal controls have been identified (aside from the well-known vital effects), which complicate the translation of measured Mg/Ca ratios to
30 temperature. Principal amongst these are dissolution (Regenberg et al., 2014), salinity (Kisakürek et al., 2008; Hönisch et al., 2013), and deeper in geological time, bias derived from secular variation in seawater Mg/Ca (Evans and Müller, 2012; Evans et al., 2015). Finally, the effect of seawater carbonate chemistry on Mg incorporation has been investigated by several culture studies (Lea et al., 1999; Russell et al., 2004; Kisakürek et al., 2008). However, the potential effect of differ-
35 ential palaeo-carbonate chemistry on Mg/Ca -derived reconstructions is rarely noted for planktic foraminifera, although a carbonate ion correction is routinely applied to some benthic foraminifera species (Sosdian and Rosenthal, 2009; Yu and Broecker, 2010). All of these studies indicate that the relationship between pH or $[\text{CO}_3^{2-}]$ and Mg/Ca is large, on the order of a 30–60 % increase in Mg/Ca for a 0.4 unit pH decrease from 8.0 (total scale). This implies that absolute Mg/Ca -derived
40 temperatures may be systematically biased when ocean pH or $[\text{CO}_3^{2-}]$ was significantly different to modern, and relative Mg/Ca -derived temperature shifts over climatic events associated with a large perturbation of the carbonate system may be inaccurate.

Here we focus on the implications of variable carbonate chemistry for Mg/Ca -derived sea surface temperature (SST) shifts across two key intervals of geologically rapid climatic change, the
45 Paleocene–Eocene Thermal Maximum (PETM) and Eocene–Oligocene Transition (EOT). Both of these events are associated with a significant (0.1–0.3 unit) pH shift (Penman et al., 2014; Pearson et al., 2009), therefore a re-examination of previously published records is warranted.

The PETM is characterised by a massive release of ^{13}C -depleted carbon over a geologically brief period of time (Kennett and Stott, 1991). The event is associated with a surface warming of several
50 °C in most locations that this has been reconstructed, using TEX_{86} or planktic foraminifera Mg/Ca and $\delta^{18}\text{O}$ (e.g. Sluijs et al., 2007; Zachos et al., 2003). Because there is evidence that the PETM was also associated with a shift in the terrestrial and oceanic hydrological cycle (Schmitz and Pujalte, 2007; John et al., 2008; D’haenens et al., 2012), $\delta^{18}\text{O}$ is frequently coupled with an independent temperature proxy in order to reconstruct surface ocean salinity change (Zachos et al., 2003; Tri-

55 pati and Elderfield, 2004). Both these, and the results of coupled climate models suggest that the hydrological response was complex and spatially variable (Sluijs et al., 2011; Winguth et al., 2010), indicating more saline surface waters in the (sub)tropical Pacific, whilst the subtropical Atlantic and coastal-proximal sites may have been characterised by increased freshwater flux to the surface ocean. These proxy reconstructions are sensitive to the control that pH exerts on foraminifera $\delta^{18}\text{O}$ 60 (Uchikawa and Zeebe, 2010), and also require revision in light of our analysis of the relationship between carbonate chemistry and Mg/Ca. Our data and reanalysis of existing records are timely given the recent publication of a high resolution $\delta^{11}\text{B}$ record across the PETM (Penman et al., 2014).

The EOT is marked by a stepwise transition to a climate state characterised by a continental-scale ice sheet on Antarctica, as evidenced by $\delta^{18}\text{O}$ measurements of benthic foraminifera (Zachos et al., 65 1996; Katz et al., 2008; Coxall et al., 2005). Sequence stratigraphic studies and Mg/Ca-derived deep ocean temperatures indicate that approximately 50% of this shift can be attributed to temperature (Miller et al., 2008; Lear et al., 2008). A $\delta^{11}\text{B}$ -pH record is now available (Pearson et al., 2009), which means that both planktic foraminiferal $\delta^{18}\text{O}$ and Mg/Ca may be corrected for the effects of a significant (~ 0.15 pH unit), shift in the ocean carbonate system associated with CO_2 70 drawdown. There are relatively few high-resolution SST records across the transition. In the subtropics, the St. Stephens Quarry (SSQ, Alabama) *Turborotalia ampliapertura* Mg/Ca record shows a large precursor cooling event beginning ~ 250 ka before the Eocene–Oligocene boundary (Wade et al., 2012). In the Southern Hemisphere, a similar albeit smaller event has been reconstructed from Tanzania Drilling Project samples of the same species (Lear et al., 2008). Finally, several sites 75 in the Southern Ocean show Mg/Ca cooling that occurs prior to the largest benthic foraminifera $\delta^{18}\text{O}$ shift, “Oi-1” (Bohaty et al., 2012). Whilst this indicates a spatially coherent picture of surface ocean cooling, a large portion of our current knowledge is derived from Mg/Ca analyses of planktic foraminifera, although biomarker-derived proxies have also been utilised (Liu et al., 2009). Therefore, improving the accuracy of these reconstructions is of critical importance to our understanding 80 of this major climatic transition.

As well as examining the implications of this pH change on published Mg/Ca-derived subtropical SST across the EOT, we also present new laser-ablation ICPMS data of *Turborotalia ampliapertura* from SSQ, for which a solution ICPMS Mg/Ca record is already available (Wade et al., 2012). This provides a methodological comparison for deep-time fossil samples: laser-ablation is spatially 85 resolved (inter- and intra-chamber) and allows the exclusion of data from poorly preserved areas of the test, whilst samples crushed for solution analysis may be easier to clean.

In order to form the basis of more accurate Mg/Ca-derived absolute and relative temperature shifts throughout the Cenozoic, we present a compilation of new and previously published data of the relationship between Mg/Ca, pH and $[\text{CO}_3^{2-}]$ for planktic foraminifera. Whilst we use the PETM 90 and EOT as examples of climatic events for which ignoring this effect may lead to substantial bias, as $\delta^{11}\text{B}$ -derived reconstructions of temporal changes in pH are available for both, our findings are

applicable to any time period with a non-modern and/or temporally variable surface ocean carbonate chemistry.

2 Materials and methods

95 We present new culture data of the relationship between Mg/Ca and carbonate system parameters for *Globigerinoides ruber*, and a reanalysis and compilation of literature data for this species, *Globigerina bulloides* and *Orbulina universa*. We utilise data from culture experiments of *G. ruber* grown at the Inter-University Institute for Marine Science, Eilat, between January 2010 and November 2013. The primary purpose of these experiments was to determine the pH-sensitivity of $\delta^{11}\text{B}$ in *G. ruber* for different $[\text{B}]_{\text{sw}}$ (with some of these cultures described in Henehan et al., 2013), and to examine the relationship between $(\text{Mg}/\text{Ca}_{\text{sw}})$ and shell Mg/Ca (Evans et al., 2016). Because seawater Mg/Ca ($\text{Mg}/\text{Ca}_{\text{sw}}$) was varied in the cultures of Evans et al. (2016), this enables the consistency of the relationship between carbonate chemistry and foraminifera Mg/Ca ($\text{Mg}/\text{Ca}_{\text{test}}$) to be examined at below-modern $\text{Mg}/\text{Ca}_{\text{sw}}$.

105 For both new and previously published data, Mg/Ca is normalised to the ratio that would be expected at 26 °C and a salinity of 35‰, specific to each species. In order to account for different temperatures, salinities and in one case $\text{Mg}/\text{Ca}_{\text{sw}}$ between the cultures, all data were normalised to these conditions where they differed and/or where $\text{Mg}/\text{Ca}_{\text{sw}}$ was not the same as in the modern ocean. For *G. ruber* this was achieved using the Mg/Ca-temperature calibration of Kisakürek et al. (2008), the Mg/Ca-salinity relationship of Hönisch et al. (2013) and the seawater-test Mg/Ca calibration of Evans et al. (2016). Between the three sets of *G. ruber* cultures, temperature varied between 26–27 °C and salinity varied between 35.0–37.2‰. For *G. bulloides* and *O. universa*, data were normalised to 26 °C assuming both species are characterised by Mg/Ca-temperature relationships with an exponential coefficient of 0.09 in the modern ocean (Anand et al., 2003). Mg/Ca was then normalised to that of the culture(s) with modern seawater pH (~ 8.02 total scale) within the same study. No salinity correction was necessary as this was consistent within the studies of Russell et al. (2004) and Lea et al. (1999) and would therefore cancel. Where measurements were not available, DIC and $[\text{CO}_3^{2-}]$ were calculated using co2sys (Lewis and Wallace, 2006), based on the same constants as used by Raitzsch et al. (2010).

120 2.1 Laboratory culture calibration

Whilst a more detailed description of culturing methods is provided elsewhere (Henehan et al., 2013; Evans et al., 2016), a brief summary is given here. Foraminifera were towed from the Gulf of Eilat, and cultured in individual 120 mL airtight flasks. Individuals were fed a juvenile brine shrimp every 1–2 days until gametogenesis took place or calcification ceased, after which they were rinsed in deionised water, dried and stored for analysis.

Within experiments intended for boron isotope measurement, the pH of flask solutions was measured at the beginning of each experiment, and then every 2–3 days using an electrode calibrated against NBS buffers. At the end of experiments, individual flask pH was measured and a composite sample of all flasks combined, measured and then poisoned for Dissolved Inorganic Carbon (DIC) and Total Alkalinity (TA) analysis at the UKOARP, Southampton (see Henehan et al., 2013). For experiments designed around varying Mg/C_{a,sw} (Evans et al., 2016), pH was measured at the start of culture. Based on repeat measurements of the reservoir seawater used to fill the individual culture flasks, we apply a conservative pH uncertainty of ± 0.1 to these experiments. pH electrode measurements were corrected to total scale pH via calibration against pHs derived from analysis of multiple DIC and TA water samples.

2.1.1 Laser-ablation analysis

The cultures outlined in Evans et al. (2016) were analysed using the 193 nm RESolution M50 laser-ablation system at Royal Holloway (Müller et al., 2009), as described previously (Evans et al., 2015). Briefly, foraminifera were cleaned in 10 % NaOCl, rinsed in deionised water and mounted on carbon tape. Individual chambers were identified as those precipitated in culture if they were characterised by elevated ¹³⁵Ba/¹³⁸Ba, as the seawater used in these experiments was spiked with 74 nM ¹³⁵Ba. NIST610 was used as a calibration standard for Mg, based on the NIST Mg value of Pearce et al. (1997) which results in Mg/Ca accuracy $\sim 7\%$ better than the value of Jochum et al. (2012) on our system. Data reduction followed established procedures (Longerich et al., 1996) using in-house Matlab data reduction software. ⁴³Ca was used as an internal standard. Long-term Mg/Ca accuracy and precision is better than 5 % (2SD) as assessed by standardising the MPI-DING komatiite glass GOR132 (Jochum et al., 2006) to NIST610.

2.1.2 Solution ICPMS

Prior to analysis, foraminifera were cleaned as described in Henehan et al. (2013). Foraminiferal culture samples of 1–3 mg (to permit boron isotope analysis on the same samples where possible) were lightly crushed between two clean glass slides, ultrasonicated and rinsed repeatedly with Milli-Q ultrapure water (18.2 M Ω). Samples were subject to intensified oxidative cleaning (3×20 – 30 min treatments of 250–400 μ L 1 % H₂O₂ in 0.1 M NH₄OH at 80 °C) to remove organic material, followed by a brief weak acid leach in 0.0005 M HNO₃ to remove any readsorbed contaminants. Finally, 200 μ L of Milli-Q was added to each sample (to slow subsequent dissolution and reduce the likelihood of leaching of B off any remnant contaminants) and 0.5 M HNO₃ (normally < 300 μ L) added incrementally until the sample was fully dissolved. Samples were centrifuged for > 5 min at 1400 rpm, and the supernatant removed to a clean vial, leaving the last ~ 20 μ L to be discarded.

Solution ICPMS measurements of Mg/Ca were performed on a Thermo Element ICP-MS at the University of Southampton and the Bristol Isotope Group. Analysis of common consistency

standards ensures no bias exists between these two laboratories. Tuning is performed on a 0.1 ppb multielement tune solution to optimize sensitivity while minimising the presence of oxides (the ratio of m/z 248/232 was maintained below 7%). During each analytical session, in-house consistency standards (Mg/Ca ratios of 1.277, 3.289 and 7.573 mmol mol⁻¹ respectively) were analysed at
165 a range of concentrations (typically 0.5, 1, and 2 mM Ca) to monitor accuracy and precision. Reproducibility of element ratio measurements is optimised (following Yu et al., 2005) by matrix-matching samples and their bracketing standards (the in-house gravimetric standard BSGS). This is achieved by analysing for [Ca] in a dilute aliquot of each sample (typically 20 in 200 µL 0.5 M HNO₃) prior to full elemental determination. Samples are then diluted with 0.5 M HNO₃ to match standard [Ca]
170 and analysed for the full trace metal suite. Long-term reproducibility of Mg/Ca ratio measurements is better than 3% (2SD), based on repeat measurements of in-house consistency standards.

Because foraminifera did not precipitate calcite exclusively in culture, final Mg/Ca ratios were obtained via mass-balance calculations. To determine this, initial pre-culture mass was estimated from starting diameter using the calculated size-mass relationship from Henehan et al. (2013).

175 2.2 Fossil samples: St. Stephen's Quarry

St. Stephens Quarry (Alabama) is a shallow water site characterised by a palaeodepth of 50–100 m (Miller et al., 2008) and exceptional foraminifera preservation (Wade et al., 2012). In order to compare laser-ablation-derived trace element data with the solution ICPMS Mg/Ca record of Wade et al. (2012), the planktic foraminifera *Turborotalia ampliapertura* was sampled from selected intervals
180 across the Eocene–Oligocene Transition. The laser-ablation analytical procedure differed from above only in that foraminifera were cleaned by ultrasonication for one minute in acetone, one minute in methanol and four minutes in deionised water to maximise removal of clay particles without damaging the specimens. Every chamber available for analysis was targeted. Mean element/Ca ratios for each foraminifera were calculated by taking the average of all measurements for that individual.
185 Because laser-ablation depth profiling is a highly spatially-resolved technique, poorly preserved areas of the test or those associated with clay mineral contamination could be excluded before the calculation of these means. We use the qualitative coherence of Mg/Ca and elements indicative of diagenetic overgrowths (e.g. the REE which are present at very low concentrations in primary foraminifera calcite, see e.g. Palmer, 1985). A sample-specific Al/Ca cut-off was used to exclude
190 foraminifera that presumably contained clay minerals which were not removed during ultrasonication. This was either 2 or 4 mmol mol⁻¹, based on the point at which elevated Al/Ca resulted in elevated Mg/Ca.

Table 1. New and compiled foraminifera culture data of the relationship between shell Mg/Ca and seawater carbonate chemistry.

Exp.	ref. ¹	pH (total scale)	[CO ₃ ²⁻] (μM)	DIC (μM)	T (°C)	salinity (‰)	Mg/Ca (mmol mol ⁻¹)	norm. Mg/Ca ²
<i>Globigerinoides ruber</i>								
DE3-5-26	1	7.90 ± 0.1	134	2089	26.3	37	6.37 ± 0.25	1.23
DE3-3-26	1	7.98 ± 0.1	162	2137	26.3	37	4.38 ± 0.25	1.34
C1	2	8.18 ± 0.01	297	1957	26.0	37	4.99 ± 0.15	0.99
C2	2	7.90 ± 0.01	164	1954	26.0	37	6.92 ± 0.21	1.38
C3	2	7.56 ± 0.01	79	1942	26.0	37	7.05 ± 0.21	1.40
MH13-1	–	7.62 ± 0.02	94	2175	26.0	37	7.99 ± 0.24	1.59
MH13-2	–	7.93 ± 0.01	175	2092	26.0	37	5.21 ± 0.16	1.04
MH13-3	–	8.48 ± 0.00	526	2026	26.0	37	3.56 ± 0.11	0.71
BK3-7.9	3	7.77 ± 0.06	119	1881	27	35	7.5 ± 0.3	1.46
BK3-8.1R	3	7.95 ± 0.08	175	1885	27	35	4.2 ± 0.2	0.82
BK3-8.3R	3	8.13 ± 0.07	255	1893	27	35	4.2 ± 0.2	0.82
BK3-8.4R	3	8.29 ± 0.08	353	1920	27	35	3.1 ± 0.2	0.60
<i>Globigerina bulloides</i>								
BH1	4	8.17 ± 0.02	185	1996	22	33.6	4.23 ± 0.13	1.06
BH2	4	8.2 ± 0.02	197	1992	22	33.8	3.82 ± 0.11	0.96
BH3	4	7.74 ± 0.02	71	1974	22	33.7	7.75 ± 0.23	1.95
BH4	4	8.65 ± 0.02	469	1980	22	33.7	2.69 ± 0.08	0.68
–	5	7.47	51	1987	22	33	6.33 ± 0.13	1.33
–	5	8.37	282	1591	22	33	3.67 ± 0.07	0.77
<i>Orbulina universa</i>								
–	5	7.67	78	1927	22	33	10.12 ± 0.20	1.14
–	5	8.47	325	1523	22	33	6.15 ± 0.12	0.69
BH3	4	7.74 ± 0.03	72	1523	22	33.7	9.41 ± 0.9	1.36
BH6	4	7.93 ± 0.01	110	1974	22	33.7	7.92 ± 1.49	1.15
AR1	4	8.24 ± 0.02	213	1981	22	33.9	6.82 ± 0.27	0.99
AR3	4	8.41 ± 0.02	300	1977	22	33.8	6.68 ± 0.33	0.97
AR4	4	8.56 ± 0.01	399	1988	22	33.6	6.42 ± 0.46	0.93
AR5	4	8.66 ± 0.01	482	2004	22	33.6	6.49 ± 0.34	0.94
AR6	4	8.22 ± 0.01	170	2030	15	33.8	3.53 ± 0.11	0.96
AR27	4	8.17 ± 0.03	167	2029	18	33.7	5.02 ± 0.33	1.04
AR10	4	8.21 ± 0.01	217	1962	25	34.0	9.6 ± 0.39	1.06

¹ 1: Evans et al. (2016), 2: Henehan et al. (2013), 3: Kisakürek et al. (2008), 4: Russell et al. (2004), 5: Lea et al. (1999).

² Normalised (ratioed) to Mg/Ca in mmol mol⁻¹ expected at 26 °C, 35‰, modern Mg/Ca_{sw} for a given species.

3 Results

3.1 Carbonate chemistry control on foraminifera Mg/Ca

195 New and published Mg/Ca-carbonate chemistry data are shown in Table 1, and our *G. ruber* culture data is shown in the context of a previous study examining the relationship between pH and Mg/Ca in this species (Kisakürek et al., 2008), as well as data compiled for *G. bulloides* and *O. universa* (Lea et al., 1999; Russell et al., 2004), in Fig. 1. Because it has been argued that there is a limited control on foraminifera Mg/Ca at high pH and/or [CO₃²⁻] (Russell et al., 2004), and there is no
200 relationship between pH and Mg/Ca below pH 7.8 for *G. ruber*, we fit regressions that capture the nonlinear aspects of the data. In pH space, *G. ruber* is characterised by a steeper slope between pH




Figure 1. New and compiled planktic foraminifera Mg/Ca-carbonate chemistry data. Mg/Ca is shown as a ratio to that at pH 8.02 (~ 8.15 on the NBS scale) for each species. **(a)** Species-specific Mg/Ca variation with pH. **(b)** Mg/Ca variation with $[\text{CO}_3^{2-}]$. **(c, d)** show the same datasets with regressions calculated from all data combined. Both logistic and linear relationships are shown between Mg/Ca and pH.

7.8 and 8.1 (Fig. 1a), and we use this as justification for fitting logistic functions to the data. pH and $[\text{CO}_3^{2-}]$ covaried in both our cultures and those previously published. Therefore it is not possible to identify which is the controlling factor based on these data. For this reason, normalised Mg/Ca is also shown in $[\text{CO}_3^{2-}]$ space in Fig. 1b. We fit power regressions to these data in order to capture the decreasing sensitivity of Mg/Ca to pH or $[\text{CO}_3^{2-}]$ at high values.

There are significant differences between the species-specific regressions. These may represent real differences in the varying ability of the different foraminifera to exclude Mg from the site of biomineralisation as seawater carbonate chemistry changes, or it may represent analytical and experimental noise given the different culturing methods, locations, and necessary corrections we

describe above. Whilst these calibrations should be used where these species have been utilised in the fossil record, in order to apply these relationships to reconstructions based on extinct species in the Paleogene we combine all the data in order to derive a broad picture of the likely response of foraminifera Mg/Ca to carbonate system perturbations (Fig. 1c and d). The change in slope in the relationship between Mg/Ca and pH, with two inflection points, is also manifest in the combined data and is described by the logistic function:

$$\text{Mg/Ca} = \frac{0.66}{1 + \exp(6.9 \times (\text{pH} - 8.0))} + 0.76 \quad (1)$$

$R^2 = 0.74$, $n = 27$. One outlier (the *G. bulloides* datapoint at pH 7.6) has been excluded as it has a greater than critical Cook's distance and a high residual from the regressions. Whilst a logistic function is consistent with the empirical insensitivity of Mg/Ca to pH or $[\text{CO}_3^{2-}]$ at higher values, it is not possible to statistically exclude a simpler relationship. Therefore, we also provide a linear regression through the data:

$$\text{Mg/Ca} = -0.70 \times \text{pH} + 6.7 \quad (2)$$

$R^2 = 0.71$. Finally, because it is not possible to distinguish between a $[\text{CO}_3^{2-}]$ and pH control based on these datasets, we also provide a combined power regression between Mg/Ca and $[\text{CO}_3^{2-}]$:

$$\text{Mg/Ca} = 8.1 \times [\text{CO}_3^{2-}]^{-0.068} - 4.6 \quad (3)$$

$R^2 = 0.61$. The application of these calibrations to fossil samples implicitly assumes that they describe some fundamental characteristic of the ability of the biological processes within foraminifera that exclude Mg, i.e. that the location of typical modern surface pH midway between the two inflection points of the logistic function is coincidental and does not represent an evolutionary adaptation to modern carbonate chemistry. This is important as, if a logistic function is indeed appropriate, perturbations to the carbonate system that do not shift pH outside of the range 8.2–8.6 or 7.7–7.4 are unlikely to be associated with a large pH-derived shift in $\text{Mg/Ca}_{\text{test}}$. There is no relationship between foraminifera Mg/Ca and DIC (Fig. 2), and we eliminate secular changes in DIC as a potential control on foraminifera Mg/Ca through time.

Because a proxy that relates directly to pH is available ($\delta^{11}\text{B}$), whereas $[\text{CO}_3^{2-}]$ cannot be reconstructed without another known carbonate system parameter (e.g. alkalinity, DIC), we base the implications for palaeotemperature reconstruction on the relationship between Mg/Ca and pH. Whilst we cannot eliminate $[\text{CO}_3^{2-}]$ as the primary control, we argue that pH is the most likely driver of carbonate chemistry related Mg/Ca shifts in Sect. 4.1. If this is incorrect, and changing $[\text{CO}_3^{2-}]$ is found to exert the dominant control on Mg/Ca, then our results may require minor revision.

3.2 St. Stephens Quarry laser-ablation data

Laser-ablation data of *T. ampliapertura* from St. Stephens Quarry are characteristic of excellent foraminifera preservation. Mn/Ca ratios are typically $< 0.2 \text{ mmol mol}^{-1}$, suggesting little or no




Figure 2. Foraminifera Mg/Ca variation with dissolved inorganic carbon. See Fig. 1a, b for legend.

245 bias from diagenetic overgrowths. Inter-chamber profiles (Fig. 3a) show localised areas with elevated Al/Ca which likely result from remnant clay particles, which were discounted before calculating mean shell values. Four intervals were sampled for which solution ICPMS Mg/Ca data have already been published (Wade et al., 2012), shown in Fig. 3b. Mg/Ca from both techniques are within error of each other for all intervals. Moreover, relative inter-sample differences in Mg/Ca are
 250 indistinguishable between the laser-ablation and solution Mg/Ca data. Whilst mean laser-ablation measurements (which represent the average of 10–30 individual chamber profiles) are offset to values $\sim 10\%$ higher, which would result in a 1°C temperature bias, this error is small compared to uncertainties derived from the complications we list below (see Sect. 4.4) and may be chance given the small number of samples that were analysed using both techniques. Based on these data,
 255 we infer that calibrations derived from solution analyses of planktic foraminifera are applicable to laser-ablation fossil data, and vice versa.

4 Discussion

4.1 A mechanistic understanding of Mg/Ca variation with carbonate chemistry

Foraminifera are known to elevate the pH of vacuolised seawater by up to ~ 1 unit (Bentov et al.,
 260 2009; de Nooijer et al., 2009), and seawater vacuoles have been suggested to play a fundamental role in calcification (Erez, 2003; Bentov and Erez, 2006), although the importance of their role has been questioned (Nehrke et al., 2013). This implies that the $[\text{CO}_3^{2-}]$ of the calcifying fluid is significantly higher than that of the vacuolised seawater, especially given the carbon concentrating mechanism proposed by ter Kuile and Erez (1987) and modified by Bentov et al. (2009). If Mg is pumped




Figure 3. (a) An example laser ablation profile through a *Turborotalia ampliapertura* chamber. Sub- μm variations in element/Ca ratios are identifiable, enabling poorly preserved or contaminated areas of the test to be excluded from the calculation of mean values. Here, the inner and outer surfaces of the chamber are characterised by elevated Al/Ca, implying incomplete clay particle removal. (b) A comparison of laser-ablation and solution ICPMS Mg/Ca data from the same intervals at St. Stephens Quarry. Solution data from Wade et al. (2012). Labels show the depth intervals of these samples. Solution ICPMS error bars are the long-term precision (RSD), whereas laser-ablation error bars are 2SE of all analyses from each interval.

265 out of seawater vacuoles by the vacuole-cytosol concentration gradient and membrane potential, as suggested by Bentov and Erez (2006), then the efficiency of Mg removal may be sensitive to pH. There are several reasons that this could be the case. When seawater pH is lower, then the foraminifer needs to work harder in order to elevate vacuolar pH. If this process is fundamentally linked to the ability of these organisms to remove Mg, whereby a lower pH makes it more difficult
 270 to precipitate calcite with low Mg/Ca, then a decreased pH would result in calcite with a higher Mg concentration. This may be because calcification proceeds through an intermediate mineral such as amorphous calcium carbonate (e.g. Politi et al., 2004), for which there may be a relationship between Mg and pH that is different to calcite (Burton and Walter, 1991). Alternatively, pH has been shown to significantly affect membrane potential in algae (Spanswick, 1972). If foraminifera use this gradient
 275 to pump Mg out of vacuolised seawater at some point in the calcification process, the Mg driving force would therefore also be sensitive to seawater pH.

It is more difficult to imagine a mechanistic control of seawater $[\text{CO}_3^{2-}]$ on $\text{Mg}/\text{Ca}_{\text{test}}$, because foraminifera probably source a portion of the DIC required for calcification from metabolic CO_2 (ter Kuile et al., 1989; Bentov et al., 2009). Whilst lower pH or $[\text{CO}_3^{2-}]$ is known to negatively affect
 280 calcification rates in several marine organisms (Ries et al., 2009) including *G. ruber* (Kisakürek et al., 2008; Henehan et al., 2013), in foraminifera it is more likely as a result of the reduced efficiency of the carbon concentrating mechanism when vacuole pH is lower, and not because fewer $[\text{CO}_3^{2-}]$

ions are present in the endocytosed seawater (the two are related but it is the pH of seawater that is the dominant control). However, until experiments are carried out with variable $[\text{CO}_3^{2-}]$ (DIC) at constant pH, it will be difficult to disentangle the effects of each. It may be that both pH and $[\text{CO}_3^{2-}]$ affect calcification rate and Mg/Ca in related but different ways. If pH were to be varied at constant $[\text{CO}_3^{2-}]$, the efficiency of any carbon-concentrating mechanism would be affected, whereas variable $[\text{CO}_3^{2-}]$ at constant pH would control calcite growth kinetics. Because it may be the relative rate of calcification and Mg removal from the calcifying fluid that controls shell Mg/Ca, we might expect both factors to exert an additive effect. Our model predicts that pH is more important, because foraminifera modify vacuole pH to a value substantially exceeding that of seawater, although further culturing experiments are required to confirm this.

An alternative biomineralisation model has been proposed, wherein ions predominantly arrive at the site of calcification through channels (the transmembrane transport model), with a small proportion arriving as a result of seawater leakage from vacuoles (Nehrke et al., 2013). However, there is no clear reason that a lower ambient pH would result in greater seawater leakage and therefore a higher $\text{Mg}/\text{Ca}_{\text{test}}$, and this model may not be able to explain the observed relationship between seawater carbonate chemistry and Mg incorporation. As is the case for the majority of the experiments described here, these should also be viewed with the caveat that observations derived from shallow-dwelling benthic foraminifera may not be directly applicable to planktic species.

4.2 Implications for reconstructing relative temperature shifts

Both transient and unidirectional temperature shifts are often associated with a change in ocean pH, as has been demonstrated for both the PETM (Penman et al., 2014) and the EOT (Pearson et al., 2009) based on $\delta^{11}\text{B}$ measurements of planktic foraminifera. It therefore follows that some portion of the observed foraminifera Mg/Ca shift over such events is not due to temperature, but changes in the pH of seawater. Figure 4 shows the extent to which relative temperature shifts would be overestimated as a result of assuming that temperature is the only dominant control on foraminifera Mg incorporation. The degree to which the associated temperature change may have been overestimated is dependent on the magnitude of the pH shift and the type of regression used to describe $\text{Mg}/\text{Ca}_{\text{test}}$ change with pH. Using a logisitic relationship, Mg/Ca is most sensitive to pH change between 7.9–8.4, as the regression slope becomes less steep outside of this range, which is not the case for a linear Mg/Ca-pH relationship.

Correcting PETM records for pH change is therefore sensitive to the type of regression used, if the pre-excursion pH was ≤ 7.8 , as indicated by the reconstructions of Panchuk et al. (2008) and Zeebe et al. (2009). If a linear fit is more appropriate, and assuming no other bias on deep-time Mg/Ca-derived relative temperature shifts, then a 0.3 unit pH drop would cause an apparent change in foraminifera Mg/Ca equivalent to a 1.6–1.8 °C temperature shift, implying that a reconstruction of relative temperature change would be overestimated by this amount if the effect of pH is not




Figure 4. Temperature overestimate across climatic events if it is assumed that pH change exerts no control on foraminifera Mg/Ca. Temperature overestimates based on both a logistic and linear relationship between Mg/Ca and pH are shown, for a 0.3 unit pH shift (solid lines) and a 0.15 unit pH shift (dashed lines). pH scenarios for both the PETM (Panchuk et al., 2008; Zeebe et al., 2009) and the EOT (Pearson et al., 2009) are overlain.

accounted for. If the logistic function is applied to the data then the contribution of pH would cause
 320 the temperature shift to be overestimated by $< 1^{\circ}\text{C}$.

For the EOT, the Mg/Ca shift associated with the pH reconstructions of Pearson et al. (2009)
 is less uncertain. This is because the maximum pH over this interval was probably higher than the
 background pH during the time of the early Eocene hyperthermals, and therefore there is a smaller
 difference between the pH-derived Mg shift between a logistic and linear function through the culture
 325 data (dashed lines in Fig. 4). If pH is unaccounted for in Mg/Ca-derived temperature reconstructions
 over the EOT (and again, assuming there are no other biases on Mg/Ca), then the relative
 temperature shift would be overestimated by 0.8–1.5 $^{\circ}\text{C}$. This bias is not strongly controlled by the
 assumption of $\delta^{11}\text{B}_{\text{sw}}$ (Pearson et al., 2009 report pH reconstructions based on $\delta^{11}\text{B}_{\text{sw}} = 38 \pm 1\%$),
 as the pH-related temperature shift is reasonably insensitive to both pH and the choice of Mg/Ca-pH
 330 regression over the seawater pH range 7.75–7.95 for a pH change of this magnitude.

Whilst it is beyond the scope of this paper to discuss the implications across all climatic events
 with an associated shift in carbonate chemistry, aside from the intervals discussed here, this finding
 has particular relevance to glacial-interglacial SST reconstructions based on foraminifera Mg/Ca,

which are typically viewed as being relatively robust as no Mg/Ca_{sw} correction is necessary during
335 the Pleistocene. Whilst glacial-interglacial surface ocean pH shifts were spatially variable (e.g. Fos-
ter et al., 2008; Martínez-Botí et al., 2015), a 0.1–0.15 unit change is typical, between ~ 8.1 – 8.25
which may result in a overestimation of glacial-interglacial SST change on the order of 1– 1.5 °C.

The relationship between pH and Mg/Ca means that relative temperature shifts in the past may
have been overestimated. However, a further consideration is the recent finding that the sensitivity of
340 the relationship between Mg/Ca and temperature may not be constant at below-modern Mg/Ca_{sw}
(Evans et al., 2016). Figure 5 shows calibrated and modelled variation in the exponential coefficient
(A) of a *G. ruber* Mg/Ca -temperature calibration with Mg/Ca_{sw} . The relationship between Mg/Ca
and temperature is assumed to take the form $Mg/Ca = B \exp^{AT}$, where B and A vary according to
an unknown function with respect to Mg/Ca_{sw} . At present, only the sensitivity in the modern ocean
345 (e.g. Anand et al., 2003; Kisakürek et al., 2008) and at $Mg/Ca_{sw} = 3.4 \text{ mol mol}^{-1}$ (Evans et al.,
2016) is known, indicating a reduction in sensitivity from 9 to $7.4 \text{ \% } ^\circ\text{C}^{-1}$. Further calibrations are
required to constrain this sensitivity at Paleogene Mg/Ca_{sw} ratios, which were reasonably invariant
between 2.0 – 2.5 mol mol^{-1} (Dickson, 2004; Coggon et al., 2010; Evans et al., 2013). Six different
model curves between the exponential coefficient (A) and Mg/Ca_{sw} are shown in Fig. 5. These were
350 derived by fitting both the observed sensitivity change as well as the calibrated nonlinear relationship
between seawater-test Mg/Ca reported by Evans et al. (2016), for a variety of assumptions regarding
the shape that this relationship should take (linear, power, exponential, quadratic). These curves
are better constrained than they appear from the two datapoints available, as only specific models
are capable of also capturing the upwards-convex seawater-test Mg/Ca relationship for *G. ruber*,
355 which is also the case for other foraminifera (Evans et al., 2015; Raitzsch et al., 2010). For example,
a linear regression through the two datapoints produces a seawater-test Mg/Ca curve that is convex
down, the opposite of that for all species studied so far (see Evans and Müller, 2012), and can be
immediately discounted. Nonetheless, the shift in sensitivity of Mg/Ca to temperature is not yet
precisely known for the Paleogene, and may lie anywhere in the range 5.0 – $7.5 \text{ \% } ^\circ\text{C}^{-1}$ based on the
360 data currently available.

The effect of the control exerted on Mg/Ca by pH and the shift in sensitivity with Mg/Ca_{sw} act
in opposite directions on reconstructed relative temperature shifts using foraminifera Mg/Ca , and
these effects may – to some extent – cancel each other in the Paleogene when Mg/Ca_{sw} was $\sim 50 \text{ \%}$
of modern. However, it should not be assumed that the two exactly balance, and at the very least
365 both of these factors should be propagated into error estimates. This may be the case to a lesser
extent for events that occurred within the last ~ 20 Ma, when seawater chemistry was closer to that
of today and therefore the effect of pH dominates the correction applied. Finally, whilst both factors
should be considered when reconstructing relative temperature shifts, the relationship between pH
and foraminifera Mg/Ca is such that the reporting of absolute temperatures will be highly sensitive
370 to differential carbonate chemistry throughout the Paleogene, as pH was probably broadly lower




Figure 5. Variation in the sensitivity of the *G. ruber* Mg/Ca thermometer with seawater Mg/Ca, for a range of assumptions regarding the relationship between Mg/Ca_{sw} and the coefficients of a Mg/Ca-temperature calibration of the form $Mg/Ca = B \exp^{AT}$. Datapoints are based on the cultures of Kisakürek et al. (2008) (modern seawater) and Evans et al. (2016) (Mg/Ca_{sw} = 3.4 mol mol⁻¹). Model lines are derived from least-squares modelling of these datasets as well as the seawater-shell Mg/Ca calibration of Evans et al. (2016). A range of model curves are shown based on different assumptions of the form the regression between the exponential coefficient of a Mg/Ca-temperature calibration and Mg/Ca_{sw} should take, shown in the legend. For example the bold red line assumes that both coefficients of a Mg/Ca-temperature equation vary quadratically with Mg/Ca_{sw}, the blue line assumes that the pre-exponential coefficient varies quadratically whilst the exponential coefficient varies with a power function with respect to Mg/Ca_{sw}. “quad. c=0” are quadratic equations forced through the origin, whilst “quad.+int.” are quadratic regressions with a *y* intercept. An estimate of early Eocene Mg/Ca_{sw} (Dickson, 2004) is shown for comparison. These models suggest the exponential coefficient of a Mg/Ca-temperature calibration may be reduced from 0.09 to 0.05–0.075 at this time.

throughout this Period (Tyrrell and Zeebe, 2004). Absolute Mg/Ca-derived temperatures should not be reported without an estimate of both the degree of nonlinearity of the seawater-test Mg/Ca relationship and the seawater pH for the interval of interest.

We now apply these findings to the PETM and the EOT in detail, in order to better constrain SST and hydrological shifts over the former, and to produce the first absolute Mg/Ca-derived temperatures across the EOT corrected for both Mg/Ca_{sw} and pH.

4.3 Implications for SST and hydrological shifts over the PETM

Coupled Mg/Ca- $\delta^{18}O$ measurements of foraminifera have been used to derive both SST and hydrological change across the PETM. Based on this technique, Zachos et al. (2003) reconstruct a 1.0–2.6‰ positive salinity shift at ODP Site 1209 (North Pacific), implying a change to a more evap-

orative regime during peak warming. Similarly, this technique has been applied to ODP Site 865 (equatorial Pacific) and DSDP Site 527 (subtropical South Atlantic) (Tripathi and Elderfield, 2004), with a more complex picture of hydrological change emerging. Uncorrected for pH, these data also imply enhanced evaporation in the Pacific (i.e. Mg/Ca shows a larger temperature shift than $\delta^{18}\text{O}$), whereas the Atlantic site is characterised by a shift towards more negative $\delta^{18}\text{O}_{\text{sw}}$ during the PETM. Finally, Zachos et al. (2006) reconstruct an negative $\delta^{18}\text{O}_{\text{sw}}$ shift at the coastal proximal Wilson Lake (New Jersey), which may be related to increased freshwater runoff, given that the PETM is known to be associated with seasonally increased precipitation in at least some regions (e.g. the Pyrenees; Schmitz and Pujalte, 2007). All of these datasets require revision in light of the control that both pH and Mg/Ca_{sw} exert on the reconstruction of relative temperature shifts. As an example, this is shown for ODP Site 1209 in Fig. 6.

The ~ 0.3 unit pH shift over the PETM (Penman et al., 2014) would result in a 15–20 % increase in foraminifera Mg/Ca using the pre-excursion pH scenario of 7.8. If this was the only previously unaccounted for source of error, this would result in a reduction of the reconstructed SST shift from 4.5 to 3 °C (Fig. 6). However, the potential shift in Mg/Ca-temperature sensitivity, or rather the large uncertainty in how this may have varied in the past means that the maximum SST excursion may lie anywhere between 3–6 °C. This full range of error includes the pH correction that we outline and assumes that the exponential constant of a Mg/Ca-temperature calibration may lie anywhere between the modern value (0.09) and the lowest modelled value shown in Fig. 5 (0.05). Until this is further constrained, it should be considered as a source of error for relative temperature shifts. It is not possible to assess at present whether reported temperature shifts are overestimates or underestimates, or fortuitously accurate.

The potential effect of seawater pH on foraminifera $\delta^{18}\text{O}$ has been previously recognised (Spero et al., 1997; Uchikawa and Zeebe, 2010) and has a mechanistic basis in oxygen isotope fractionation between carbonate species (Zeebe, 1999), i.e. this fractionation is also most appropriately causally ascribed to pH rather than $[\text{CO}_3^{2-}]$. *Orbulina universa* and *Globigerina bulloides* are characterised by cultured pH- $\delta^{18}\text{O}$ slopes of -0.89 and -2.51 per pH unit respectively, which means that a 0.3 unit pH shift over the PETM would result in underestimating the combined temperature and $\delta^{18}\text{O}_{\text{sw}}$ component of the foraminifera $\delta^{18}\text{O}$ shift by 0.27–0.75 ‰ (i.e. temperature and pH act in opposite directions on a $\delta^{18}\text{O}_{\text{shell}}$ record if the climatic event is characterised by a temperature increase and ocean pH decrease). This large interspecies difference in $\delta^{18}\text{O}$ -pH slopes is a further large source of uncertainty given that extinct species may be characterised by different slopes. Figure 6C shows a range of $\delta^{18}\text{O}_{\text{sw}}$ shifts over the PETM based on different assumptions regarding the control that Mg/Ca_{sw} and the relationship between pH, Mg/Ca and $\delta^{18}\text{O}$ exert on this record. The blue line shows $\Delta\delta^{18}\text{O}_{\text{sw}}$ ($\delta^{18}\text{O}_{\text{sw}}$ relative to the pre-CIE baseline) fully corrected for all of the effects described above, using our logistic Mg/Ca-pH relationship (Fig. 1), a reduced Mg/Ca-temperature sensitivity of 7.5 % °C⁻¹ for the Paleogene (Fig. 5) and the pH- $\delta^{18}\text{O}$ relationship for *G. bulloides*




Figure 6. (a) The fraction shift in Mg/Ca over the PETM based on the calibrations between Mg/Ca and pH shown in Fig. 1, and the pH reconstructions of Penman et al. (2014) (assuming initial pH = 7.8). All data are shown relative to Mg/Ca at modern pH and display the relative change as a result of pH only. (b) ODP Site 1209 *Morozovella* Mg/Ca-derived temperature shift across the PETM. The data are shown as originally reported, and with various corrections for both pH and Mg/Ca_{sw}, relative to the pre-carbon isotope excursion (CIE) baseline. (c) $\delta^{18}\text{O}_{\text{sw}}$ shift from coupled Mg/Ca- $\delta^{18}\text{O}$ data, based on the range of transient temperature shifts shown in panel (b). Our preferred record (blue line), corrected for the pH effect on both foraminifera Mg/Ca and $\delta^{18}\text{O}$, and accounting for the likely higher sensitivity of Mg/Ca to temperature in PETM seawater, implies little change in subtropical Pacific surface hydrology over the event. Data from Penman et al. (2014) and Zachos et al. (2003).

(Spero et al., 1997). This fully corrected record is characterised by $\Delta\delta^{18}\text{O}_{\text{sw}}$ within $\pm 0.25\%$ of zero, implying relatively little surface hydrology change at ODP Site 1209 over the PETM if this set
420 of corrections is indeed applicable for this extinct genus (*Morozovella*). For example, alternatively using the *O. universa* $\delta^{18}\text{O}$ -pH slope would result in a $\delta^{18}\text{O}_{\text{sw}}$ anomaly 0.4-0.5% more positive. Therefore, the blue line in Fig. 5C is a best estimate based on the available data and calibrations, and by no means definitively demonstrates no hydrology regime change across the PETM, but rather serves to highlight that (1) salinity anomalies were likely less extreme than previously suggested,
425 and (2) that such reconstructions are associated with large uncertainties.

Aside from those mentioned above, both $\delta^{18}\text{O}$ and Mg/Ca data are also available for DSDP Site 401 in the Bay of Biscay (Bornemann et al., 2014) and DSDP Site 277 in the southwest Pacific (Hollis et al., 2015). Furthermore, $\delta^{18}\text{O}$ and TEX_{86} data are available for Bass River (Sluijs et al., 2007; John et al., 2008), ~ 50 km to the East of Wilson Lake. Compiling the data for these sites
430 and applying the corrections that we detail above enables a more accurate picture of SST and hydrological shift over the PETM to be constructed. Figure 7 shows peak PETM SST and $\Delta\delta^{18}\text{O}_{\text{sw}}$ relative to the pre-CIE baseline, with error bars that incorporate the full range of uncertainties that we discuss above. Temperature shows a broad relationship with palaeolatitude, with $< 2^\circ\text{C}$ warming in the (sub)tropics and $2\text{--}7^\circ\text{C}$ at higher latitudes. The Mg/Ca data from DSDP Site 401 do not
435 conform to this, which show only a 0.3 mmol mol^{-1} raw Mg/Ca change and therefore a corrected temperature shift of $0.2 \pm 0.1^\circ\text{C}$. In contrast, the *Morozovella* $\delta^{18}\text{O}$ excursion at this site would imply a SST change of $5\text{--}7^\circ\text{C}$ if interpreted purely in terms of temperature. Conversely, the oxygen isotope data may simply reflect freshwater flux, given that there is also evidence for an abrupt shift in hydrological regime from the style of clay deposition (Bornemann et al., 2014), although
440 modelling results predict a substantial surface ocean warming in the northeast Atlantic (Jones et al., 2013). It is possible that the small Mg/Ca shift at DSDP Site 401 is a result of poor foraminifera preservation and/or drilling disturbance or dissolution. However, the raw ratios are not abnormal for this time interval, and the high clay content and relatively shallow palaeo-water depth of $1.5\text{--}2$ km (D'haenens et al., 2012) means that differential preservation or dissolution across the PETM is un-
445 likely (Bornemann et al., 2014). Nonetheless, with the exception of this site, the broad relationship between the magnitude of the SST excursion and latitude is consistent with enhanced high latitude warming compared to the tropics and a reduction in latitudinal temperature gradients with global warming (Bijl et al., 2009).

Compiled $\Delta\delta^{18}\text{O}_{\text{sw}}$ shifts over the PETM (Fig. 7b) are all within error of zero, with the exception
450 of the northeast Atlantic (which is a result of the small Mg/Ca shift discussed above) and Wilson Lake. The difference between reconstructed hydrology at Wilson Lake and Bass River is not a GDGT calibration artefact, as Wilson Lake shows a larger raw TEX_{86} shift across the PETM (Sluijs et al., 2007). It is more likely that $\delta^{18}\text{O}$ at Wilson Lake is affected to a greater extent from terrestrial freshwater input as it was located closer to the palaeoshoreline, whilst Bass River essentially shows




Figure 7. Compiled sea surface temperature and $\delta^{18}\text{O}_{\text{sw}}$ peak PETM shifts relative to pre-CIE values. **(a)** GDGT and Mg/Ca-derived warming as a function of latitude. Mg/Ca estimates are adjusted for both pH and the sensitivity of the Mg/Ca-temperature relationship in PETM seawater (see text for details), and are the average of estimates derived from the two surface-dwelling foraminifera genera *Morozovella* and *Acarinina*. Labels refer to ODP/DSDP site numbers or locality names. **(b)** $\Delta\delta^{18}\text{O}_{\text{sw}}$ reconstructions based on foraminifera $\delta^{18}\text{O}$ coupled with either Mg/Ca or GDGT-derived sea surface temperature shifts. $\delta^{18}\text{O}$ was adjusted for pH following Uchikawa and Zeebe (2010). Inner error bars are the uncertainty relating to the $\delta^{18}\text{O}$ -pH slope (datapoints use the $\delta^{18}\text{O}$ -pH slope of *G. bulloides* whilst the inner positive error bar shows where the data would lie if the slope of *O. universa* were to be used), outer error bars also include the uncertainty relating to the pH and sensitivity corrections applied to the Mg/Ca data. Data not specifically discussed in the main text are from Sluijs et al. (2006) and Sluijs et al. (2011). **(c)** The magnitude of the carbon isotope excursion at each site in *Acarinina*, with the exception of DSDP Site 401 (*Morozovella*).

455 no shift in surface hydrology. Alternatively, because the temperature record at these sites is derived
from different material, it is possible that a seasonal bias in foraminifera or GDGT production could
impact on the accuracy of hydrology reconstructions. Furthermore, microanalytical approaches to
proxy extraction from foraminifera shells is becoming more common (e.g. Kozdon et al., 2013;
Evans et al., 2015). As a result, the impact of diagenesis on planktic species from sites previously
460 thought to be well-preserved is becoming more apparent. The relatively large negative $\Delta\delta^{18}\text{O}_{\text{sw}}$ shift
that we reconstruct for the equatorial Pacific (ODP Site 865, Fig. 7) is in part because we use the
new spatially-resolved $\delta^{18}\text{O}$ data of Kozdon et al. (2013), which are characterised by a larger shift
in raw $\delta^{18}\text{O}$ than the multiple whole-shell analyses of Tripathi and Elderfield (2004). If other sites
are similarly diagenetically compromised then it is possible that current datasets do not accurately
465 portray temperature and hydrology change during the CIE.

Although some sites may be biased in these respects, or are likely influenced by near-coastal pro-
cesses, applying these corrections overall implies a slight freshening of surface waters during the
peak CIE. In distal sites, this can only result from a change in hydrology towards relatively more
precipitation. Although the detail of hydrological change across the full PETM interval is evidently
470 more temporally complex, particularly when differences between surface and thermocline water are
considered (e.g. Tripathi and Elderfield, 2004), this suggests that most areas for which records are
available were characterised by a salinity decrease. A salinity increase associated with the peak-
CIE PETM interval is only possible at sites other than ODP Site 1209 if the pH- $\delta^{18}\text{O}$ slope of
Morozovella and *Acarinina* was substantially lower than that of the extant species for which this
475 relationship has been calibrated (Spero et al., 1997) and/or (1) if the reduction in the exponential
coefficient of the Mg/Ca-temperature calibration was greater than shown in Fig. 5, and/or (2) the
pH-Mg/Ca relationship was steeper for these extinct species than shown in Fig. 1. Whilst some of
these issues will be challenging to address, we highlight that all of these factors should be consid-
ered as sources of error, and applying corrections for both pH and Mg/Ca_{sw} gives a current best
480 estimate of no surface hydrological change to a slight freshening at any site distal from the coast,
with the possible exception of ODP Site 1209. Further calibration of the pH- $\delta^{18}\text{O}$ relationship in
both symbiont bearing and asymbiotic foraminifera would enable the likely inter-species variability
of this relationship for extinct foraminifer with more confidence. An additional consideration is the
impact of diagenesis on these geochemical records.

485 Finally, because a drilling disturbance at DSDP Site 401 and dissolution at other sites cannot
be ruled out, we also show the magnitude of the $\delta^{13}\text{C}$ excursion in Fig. 7C (based on the same
foraminifera as the $\Delta\delta^{18}\text{O}_{\text{sw}}$ data) as an indicator of sites where this may be an issue, and therefore at
which the reconstructions should be regarded as not necessarily representing peak-PETM conditions.
This should be viewed with the caveat that other processes may affect the magnitude of the $\delta^{13}\text{C}$ shift
490 at a given site as recorded in planktic foraminifera. Within a given site all reconstructions are based
on coeval material, however, this does leave the possibility that some sites record post-maximum

CIE conditions (e.g. DSDP Site 401), whilst those with the largest CIEs inform us of peak PETM hydrology.

We do not apply a salinity correction to the Mg/Ca data as most sites are within error of no
495 salinity change and because such a correction requires some assumption regarding the salinity– $\delta^{18}\text{O}_{\text{sw}}$ relationship at the time. However, if robust evidence for salinity change is shown then this additional factor should be accounted for. Using the Mg/Ca-salinity slopes of Hönisch et al. (2013), a PETM salinity change of 1 ‰ would impact ΔT reconstructions by 0.15–0.5 °C.

4.4 Towards accurate absolute Mg/Ca-derived temperatures: the EOT as an example

500 Mg/Ca-derived reconstructions of the thermal evolution of the surface ocean across the Eocene–Oligocene climate transition is known from only a handful of sites: the Tanzania Drilling Project (Lear et al., 2008), St. Stephens Quarry Alabama (Wade et al., 2012), and four locations in the Southern Ocean (Bohaty et al., 2012). Planktic foraminifera boron isotope measurements indicate that the transition was associated with a 0.1–0.15 unit positive pH shift, from a pre-event pH of
505 7.8–8.0 (Pearson et al., 2009). As for the PETM, the Mg/Ca shift over the EOT resulting from this pH change is counteracted to some extent by the reduced sensitivity of the relationship between Mg/Ca-temperature, as Mg/Ca_{sw} reconstructions based on three independent proxies within a few million years of the transition are 2.0–2.4 mol mol⁻¹ (Horita et al., 2002; Coggon et al., 2010; Evans et al., 2013). However, the pH change over this interval is characterised by a reversal to more acidic
510 conditions between 33.6–33.4 Ma (Pearson et al., 2009), which means that this will impact relative temperature reconstructions across the EOT irrespective of the reduced sensitivity in Mg/Ca-temperature. As the pH record is far lower resolution than the Mg/Ca data, it is possible that there are unavoidable inaccuracies derived from linearly interpolating between datapoints.

In order to reconstruct absolute temperature (as well as relative shifts) from Mg/Ca before the
515 Pleistocene, two additional pieces of information are required: (1) a Mg/Ca_{sw} reconstruction for the time interval of interest, in order to update the coefficients of the Mg/Ca-temperature calibration, and (2) a pH record for the time of interest in order to adjust the Mg/Ca data according to Eqs. (1) or (2). As before, this assumes that the pH-Mg/Ca relationship that we present is broadly characteristic of all foraminifera, and that the *G. ruber* calibrations of the nonlinearity of the Mg/Ca_{test}-
520 Mg/Ca_{sw}-temperature surface of Evans et al. (2016) are applicable to other species. Whilst not all foraminifera are characterised by the same pre-exponential Mg/Ca-temperature coefficient in the modern ocean (e.g. Anand et al., 2003), this may be difficult to assess for extinct species. The curvature of a seawater-test Mg/Ca calibration and the reduced sensitivity of the relationship between Mg/Ca and temperature, although now known for *G. ruber* (Evans et al., 2016), should be cali-
525 brated in other extant foraminifera in order to see how ubiquitously these equations may be applied in deep-time. Nonetheless, now that the necessary calibrations are available for at least one species it is possible to reconstruct appropriately corrected absolute Mg/Ca temperatures for the Paleogene

for the first time, with the caveat that doing so assumes that these calibrations are applicable to the extinct species on which the existing records are based. This is shown in Fig. 8 for St. Stephens
530 Quarry (see Wade et al., 2012). We use this EOT dataset as an example of how these corrections should be applied, noting that the absolute temperatures are likely to shift as further pH reconstructions, seawater Mg/Ca estimates, and foraminifera seawater-shell Mg/Ca calibrations become available. For example, a pH record is currently only available from Tanzania, which although based on the same species as the St. Stephens Quarry Mg/Ca record (*T. ampliapertura*) may not necessarily
535 reflect the detail of carbonate chemistry changes at this site. Realistic Mg/Ca errors for this time are $\pm 4^\circ\text{C}$, of which 2°C is derived from the uncertainty in $\text{Mg}/\text{Ca}_{\text{sw}}$ ($\pm 0.4 \text{ mol mol}^{-1}$), 1.5°C results from a 1‰ uncertainty in $\delta^{11}\text{B}_{\text{sw}}$ at this temperature, and $\sim 0.5^\circ\text{C}$ from the uncertainty in extrapolating the seawater-shell Mg/Ca calibrations to Paleogene $\text{Mg}/\text{Ca}_{\text{sw}}$ (Fig. 5). Within the assumptions stated above (i.e. that these calibrations are applicable to extinct foraminifera), reducing
540 uncertainty in $\text{Mg}/\text{Ca}_{\text{sw}}$ reconstructions is clearly a priority in order to improve the precision of absolute Mg/Ca-derived ocean temperatures before the Pleistocene.

Although the absolute temperatures shown in Fig. 8 could shift in either direction by $\sim 4^\circ\text{C}$ as a result of these uncertainties, the Mg/Ca record from SSQ remains in good agreement with TEX_{86} from the same site Wade et al. (2012); the two proxies do not deviate by more than 2°C where (ap-
545 proximately) coeval samples have been measured. TEX_{86} temperatures are associated with an error of a similar magnitude to Mg/Ca (Tierney and Tingley, 2014). Therefore it would be coincidental if not impossible if both proxies were offset from the true value by approximately the same amount and in the same direction. However, because the effect of pH and temperature act in the opposite direction on $\text{Mg}/\text{Ca}_{\text{test}}$, it is possible to arrive at the “right” values by any combination of incorrect
550 assumptions regarding pH and seawater chemistry. For this reason, apparent proxy agreement with Mg/Ca should be viewed with caution (including agreement in relative temperature shifts), unless both a precise and accurate $\text{Mg}/\text{Ca}_{\text{sw}}$ and pH record are available.

At St. Stephens Quarry (Fig. 8b), the magnitude of the precursor cooling described by Wade et al. (2012) is sensitive to the corrections that we detail. Because it was not known otherwise at
555 the time, the published 3.9°C temperature shift used 0.09 as the exponential coefficient (A) of a Mg/Ca-temperature calibration and is uncorrected for pH change. Correcting for the difference in modern-EOT Mg/Ca-temperature sensitivity, shown by the orange dashed line on Fig. 8b, results in temperatures far higher than those from TEX_{86} and a precursor cooling of 6.3°C . Finally, applying a pH correction brings the absolute temperatures back in good agreement with those derived from
560 TEX_{86} and results in a slightly smaller precursor cooling of 5.9°C (albeit higher than that originally reported), as 0.4°C of apparent cooling can be ascribed to the ~ 0.05 pH change over this 200 ka interval. This further highlights how these factors may have a significant effect on both absolute and relative Mg/Ca temperature shifts. Given the stability of tropical SST at one site throughout the Eocene (Pearson et al., 2007), this event may represent unprecedented subtropical cooling compared




Figure 8. (a) EOT pH reconstruction from Pearson et al. (2009) based on Tanzania Drilling Project samples. The error bar is derived from an uncertainty in $\delta^{11}\text{B}_{\text{sw}}$ of $\pm 1\%$. (b) Subtropical sea surface temperature from Mg/Ca and TEX_{86} . These absolute Mg/Ca temperatures are the first to be adjusted for both the pH effect on Mg incorporation and the nonlinear relationship between seawater-shell Mg/Ca and temperature. St. Stephens Quarry Mg/Ca data from both solution ICPMS Wade et al. (2012) and laser ablation (this study) are shown. The combined Mg/Ca error estimate is that due to uncertainty in the $\text{Mg}/\text{Ca}_{\text{sw}}$ ratio ($\pm 0.5 \text{ mol mol}^{-1}$), $\delta^{11}\text{B}_{\text{sw}}$ and in the calibration regressions of Evans et al. (2016). The corrected Mg/Ca data imply 6°C of precursor cooling at $\sim 33.9 \text{ Ma}$. (c) ODP Site 744 and DSDP Site 522 benthic foraminifera $\delta^{18}\text{O}$ (Zachos et al., 1996). Note that the inter-interval Mg/Ca and $\delta^{11}\text{B}_{\text{sw}}$ error (i.e. the error in relative changes) is far smaller than shown by the bars.

565 to the earlier Paleogene. The corrected records that we present highlight that global cooling markedly
greater than previously thought is likely to have occurred several 100 ka prior to the onset of major
Antarctic glaciation, given that it is observed at several widely distributed sites. A subtropical SST
shift of almost 6 °C is greater than the change reconstructed for most of the subtropics between
present day and the Last Glacial Maximum (LGM), although it is interesting to note that the LGM-
570 Recent temperature difference for this region is of the same magnitude (Waelbroeck et al., 2009),
which may imply that this region was similarly sensitive to global climate change in the Paleogene.

The excellent agreement between the published solution data and our new laser-ablation analyses
(Figs. 3b and 8b) not only demonstrates that the SSQ Mg/Ca data of Wade et al. (2012) are robust,
but also provides good evidence that laser-ablation and solution ICPMS results are directly compa-
575 rable. It is not possible to assess whether the laser-ablation data we present here are more accurate
than the solution analyses, as the two are within error of each other (Fig. 3). It is possible that the
laser-ablation data are offset to higher Mg/Ca because it is more challenging to remove clay miner-
als from whole specimens, although given that we exclude all areas of the test with elevated Al/Ca
this is unlikely. The offset could alternatively have its roots in the necessity to standardise laser-
580 ablation data to glasses in the absence of a homogeneous carbonate material (e.g. Hathorne et al.,
2008). However, we note that inter-laboratory comparison studies of solution data are themselves
not always within 10 % agreement (Rosenthal et al., 2004); these data do not necessarily highlight
an issue with standardising carbonate Mg using the NIST glasses. Given the recent proliferation
of Paleogene laser-ablation derived trace element data (e.g. Creech et al., 2010; Evans et al., 2013;
585 Hollis et al., 2015), the consistency that we observe between datasets (Fig. 3b) is encouraging.

5 Conclusions

Using a combination of new and compiled data on the relationship between foraminifera Mg/Ca
and seawater carbonate chemistry, we show that pH exerts a large control on Mg incorporation
into planktic foraminiferal calcite. Using these data, we present new calibrations of how differential
590 modern and ancient pH may be corrected for. This correction is on the order of 5-20 % per 0.1 unit
pH shift between pH 7.7-8.3 (total scale), and has the implication that both absolute and relative
Mg/Ca-derived temperature changes are likely to be inaccurate and imprecise if secular variation in
carbonate chemistry is unaccounted for. We apply these findings to the Paleocene–Eocene Thermal
Maximum (PETM) and Eocene–Oligocene Transition (EOT). Over the PETM, the ~ 0.3 unit pH
595 shift (Penman et al., 2014) may be counteracted to an extent by the recent finding that the sensitiv-
ity of the relationship between Mg/Ca and temperature was likely lower in the Paleogene (Evans
et al., 2016), although the uncertainty in relative temperature and hydrology change is substantially
greater than previously thought. However, coupled with the effect that pH exerts on foraminifera
 $\delta^{18}\text{O}$ (Spero et al., 1997), we show that at most sites peak PETM sea surface hydrology was likely

600 characterised by a shift to less saline conditions relative to the pre-onset baseline. In contrast, the
broadly unidirectional pH shift over the EOT means that relative temperature shifts have been pre-
viously underestimated, as the reduced sensitivity of this thermometer at this time dominates the
correction. We report the first pH-corrected absolute Mg/Ca temperatures for the Paleogene that
take into account palaeo-Mg/Ca_{sw}, the differential modern-deep time Mg/Ca-temperature sensitiv-
605 ity, and pH.

Acknowledgements. D. Evans would like to thank Shai Oron (IUI, Eilat) for assistance with plankton tows.
M. Henehan would like to thank Gavin Foster for his part in facilitating Mg/Ca experiments contained in this
work, and express appreciation for the invaluable contributions of Katy Prentice, James Rae, Eleni Anagnostou,
Tom Chalk, Joe Stewart and Claudia Alt during culture experiments. This work was supported by Natural
610 Environment Research Council (NERC) PhD studentships granted to D. Evans and M. Henehan, NERC grant
NE/G014817 to B. Wade, ISF grant 551/10 awarded to J. Erez and EU FP7 ASSEMBLE grant to G. Foster and
M. Henehan. The authors would like to thank Ellen Thomas and two anonymous reviewers whose constructive
comments greatly improved this contribution.

References

- 615 Anand, P., Elderfield, H., and Conte, M.: Calibration of Mg/Ca thermometry in planktonic foraminifera from a sediment trap time series, *Paleoceanography*, 18, 28–31, 2003.
- Bentov, S. and Erez, J.: Impact of biomineralization processes on the Mg content of foraminiferal shells: a biological perspective, *Geochem. Geophys. Geosy.*, 7, Q01P08, doi:10.1029/2005GC001015, 2006.
- Bentov, S., Brownlee, C., and Erez, J.: The role of seawater endocytosis in the biomineralization process in calcareous foraminifera, *P. Natl. Acad. Sci. USA*, 106, 21500–21504, 2009.
- 620 Bijl, P., Schouten, S., Sluijs, A., Reichert, G., Zachos, J., and Brinkhuis, H.: Early Palaeogene temperature evolution of the southwest Pacific Ocean, *Nature*, 461, 776–779, 2009.
- Bohaty, S. M., Zachos, J. C., and Delaney, M. L.: Foraminiferal Mg/Ca evidence for Southern Ocean cooling across the Eocene–Oligocene transition, *Earth Planet. Sc. Lett.*, 317, 251–261, 2012.
- 625 Bornemann, A., Norris, R. D., Lyman, J. A., D’haenens, S., Groeneveld, J., Röhl, U., Farley, K. A., and Speijer, R. P.: Persistent environmental change after the Paleocene–Eocene Thermal Maximum in the eastern North Atlantic, *Earth Planet. Sc. Lett.*, 394, 70–81, 2014.
- Burton, E. and Walter, L.: The effects of $p\text{CO}_2$ and temperature on magnesium incorporation in calcite in seawater and MgCl_2 - CaCl_2 solutions, *Geochim. Cosmochim. Ac.*, 55, 777–785, 1991.
- 630 Coggon, R., Teagle, D., Smith-Duque, C., Alt, J., and Cooper, M.: Reconstructing past seawater Mg/Ca and Sr/Ca from mid-ocean ridge flank calcium carbonate veins, *Science*, 327, 1114–1117, 2010.
- Coxall, H., Wilson, P., Pälike, H., Lear, C., and Backman, J.: Rapid stepwise onset of Antarctic glaciation and deeper calcite compensation in the Pacific Ocean, *Nature*, 433, 53–57, 2005.
- Creech, J., Baker, J., Hollis, C., Morgans, H., and Smith, E.: Eocene sea temperatures for the mid-latitude southwest Pacific from Mg/Ca ratios in planktonic and benthic foraminifera, *Earth Planet. Sc. Lett.*, 299, 483–495, 2010.
- 635 de Nooijer, L. J., Toyofuku, T., and Kitazato, H.: Foraminifera promote calcification by elevating their intracellular pH, *P. Natl. Acad. Sci. USA*, 106, 15374–15378, 2009.
- D’haenens, S., Bornemann, A., Stassen, P., and Speijer, R. P.: Multiple early Eocene benthic foraminiferal assemblage and $\delta^{13}\text{C}$ fluctuations at DSDP Site 401 (Bay of Biscay – NE Atlantic), *Mar. Micropaleontol.*, 88, 15–35, 2012.
- Dickson, J.: Echinoderm skeletal preservation: calcite-aragonite seas and the Mg/Ca ratio of Phanerozoic oceans, *J. Sediment. Res.*, 74, 355–365, 2004.
- Erez, J.: The source of ions for biomineralization in foraminifera and their implications for paleoceanographic proxies, *Rev. Mineral. Geochem.*, 54, 115–149, 2003.
- 645 Evans, D. and Müller, W.: Deep time foraminifera Mg/Ca paleothermometry: nonlinear correction for secular change in seawater Mg/Ca, *Paleoceanography*, 27, PA4205, doi:10.1029/2012PA002315, 2012.
- Evans, D., Müller, W., Oron, S., and Renema, W.: Eocene seasonality and seawater alkaline earth reconstruction using shallow-dwelling large benthic foraminifera, *Earth Planet. Sc. Lett.*, 381, 104–115, doi:10.1016/j.epsl.2013.08.035, 2013.
- 650 Evans, D., Erez, J., Oron, S., and Müller, W.: Mg/Ca-temperature and seawater-test chemistry relationships in the shallow-dwelling large benthic foraminifer *Operculina ammonoides*, *Geochim. Cosmochim. Ac.*, 148, 325–342, 2015a.

- Evans, D., Brierley, C., Raymo, M. E., Erez, J., and Müller, W.: Planktic foraminifera shell chemistry response to seawater chemistry: Pliocene–Pleistocene seawater Mg/Ca, temperature and sea level change, *Earth Planet. Sc. Lett.*, accepted, 2016.
- Foster, L., Finch, A., Allison, N., Andersson, C., and Clarke, L.: Mg in aragonitic bivalve shells: seasonal variations and mode of incorporation in *Arctica islandica*, *Chem. Geol.*, 254, 113–119, 2008.
- Hathorne, E., James, R., Savage, P., and Alard, O.: Physical and chemical characteristics of particles produced by laser ablation of biogenic calcium carbonate, *J. Anal. Atom. Spectrom.*, 23, 240–243, 2008.
- Henehan, M. J., Rae, J. W., Foster, G. L., Erez, J., Prentice, K. C., Kucera, M., Bostock, H. C., Martínez-Botí, M. A., Milton, J. A., Wilson, P. A., Marshall, B. J., and Elliott, T.: Calibration of the boron isotope proxy in the planktonic foraminifera *Globigerinoides ruber* for use in palaeo-CO₂ reconstruction, *Earth Planet. Sc. Lett.*, 364, 111–122, 2013.
- Hollis, C. J., Hines, B. R., Littler, K., Villasante-Marcos, V., Kulhanek, D. K., Strong, C. P., Zachos, J. C., Eggins, S. M., Northcote, L., and Phillips, A.: Onset of the Paleocene–Eocene Thermal Maximum in the southern Pacific Ocean (DSDP Site 277, Campbell Plateau), *Clim. Past Discuss.*, 11, 243–278, doi:10.5194/cpd-11-243-2015, 2015.
- Hönisch, B., Allen, K. A., Lea, D. W., Spero, H. J., Eggins, S. M., Arbuszewski, J., deMenocal, P., Rosenthal, Y., Russell, A. D., and Elderfield, H.: The influence of salinity on Mg/Ca in planktic foraminifers—Evidence from cultures, core-top sediments and complementary $\delta^{18}\text{O}$, *Geochim. Cosmochim. Ac.*, 121, 196–213, 2013.
- Horita, J., Zimmermann, H., and Holland, H.: Chemical evolution of seawater during the Phanerozoic: implications from the record of marine evaporites, *Geochim. Cosmochim. Ac.*, 66, 3733–3756, 2002.
- Jochum, K., Stoll, B., Herwig, K., Willbold, M., Hofmann, A., Amini, M., Aarburg, S., Abouchami, W., Hellebrand, E., Mocek, B., Raczek, I., Stracke, A., Alard, O., Bouman, C., Becker, S., Dücking, M., Brátz, H., Klemd, R., de Bruin, D., Canil, D., Cornell, D., de Hoog, C-J., Dalpé, C., Danyushevsky, L., Eisenhauer, A., Gao, Y., Snow, J. E., Groschopf, N., Günther, D., Latkoczy, C., Guillong, M., Hauri, E. H., Hófer, H. E., Lahaye, Y., Horz, K., Jacob, D. E., Kasemann, S. A., Kent, A. J. R., Ludwig, T., Zack, T., Mason, P. R. D., Meixner, A., Rosner, M., Misawa, K., Nash, B. P., Pfänder, J., Premo, W. R., Sun, W. D., Tiepolo, M., Vannucci, R., Vennemann, T., Wayne, D., and Woodhead, J. D.: MPI-DING reference glasses for in situ microanalysis: new reference values for element concentrations and isotope ratios, *Geochem. Geophys. Geosy.*, 7, Q02008, doi:10.1029/2005GC001060, 2006.
- Jochum, K. P., Scholz, D., Stoll, B., Weis, U., Wilson, S. A., Yang, Q., Schwalb, A., Börner, N., Jacob, D. E., and Andreae, M. O.: Accurate trace element analysis of speleothems and biogenic calcium carbonates by LA-ICP-MS, *Chem. Geol.*, 318, 31–44, 2012.
- John, C. M., Bohaty, S. M., Zachos, J. C., Sluijs, A., Gibbs, S., Brinkhuis, H., and Bralower, T. J.: North American continental margin records of the Paleocene–Eocene thermal maximum: implications for global carbon and hydrological cycling, *Paleoceanography*, 23, PA2217, doi:10.1029/2007PA001465, 2008.
- Jones, T. D., Lunt, D. J., Schmidt, D. N., Ridgwell, A., Sluijs, A., Valdes, P. J., and Maslin, M.: Climate model and proxy data constraints on ocean warming across the Paleocene–Eocene Thermal Maximum, *Earth-Sci. Rev.*, 125, 123–145, 2013.

- Katz, M. E., Miller, K. G., Wright, J. D., Wade, B. S., Browning, J. V., Cramer, B. S., and Rosenthal, Y.: Stepwise transition from the Eocene greenhouse to the Oligocene icehouse, *Nat. Geosci.*, 1, 329–334, 2008.
- 695 Kennett, J. and Stott, L.: Abrupt deep sea warming, paleoceanographic changes and benthic extinctions at the end of the Paleocene, *Nature*, 353, 225–229, 1991.
- Kisakürek, B., Eisenhauer, A., Bohm, F., Garbe-Schönberg, D., and Erez, J.: Controls on shell Mg/Ca and Sr/Ca in cultured planktonic foraminiferan, *Globigerinoides ruber* (white), *Earth Planet. Sc. Lett.*, 273, 260–269, 2008.
- 700 Kozdon, R., Kelly, D., Kitajima, K., Strickland, A., Fournelle, J., and Valley, J.: In situ $\delta^{18}\text{O}$ and Mg/Ca analyses of diagenetic and planktic foraminiferal calcite preserved in a deep-sea record of the Paleocene–Eocene thermal maximum, *Paleoceanography*, 28, 517–528, 2013.
- Lea, D., Mashiotta, T., and Spero, H.: Controls on magnesium and strontium uptake in planktonic foraminifera determined by live culturing, *Geochim. Cosmochim. Ac.*, 63, 2369–2379, 1999.
- 705 Lear, C., Bailey, T., Pearson, P., Coxall, H., and Rosenthal, Y.: Cooling and ice growth across the Eocene–Oligocene transition, *Geology*, 36, 251–254, 2008.
- Lear, C. H., Elderfield, H., and Wilson, P. A.: Cenozoic deep-sea temperatures and global ice volumes from Mg/Ca in benthic foraminiferal calcite, *Science*, 287, 269–272, 2000.
- Lewis, E. and Wallace, D.: Program developed for CO₂ system calculations, ORNL/CDIAC-105, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, 2006.
- 710 Liu, Z., Pagani, M., Zinniker, D., DeConto, R., Huber, M., Brinkhuis, H., Shah, S., Leckie, R., and Pearson, A.: Global cooling during the Eocene–Oligocene climate transition, *Science*, 323, 1187–1190, 2009.
- Longerich, H., Jackson, S., and Günther, D.: Laser ablation inductively coupled plasma mass spectrometric transient signal data acquisition and analyte concentration calculation, *J. Anal. Atom. Spectrom.*, 11, 899–904, 1996.
- 715 Martínez-Botí, M., Marino, G., Foster, G., Ziveri, P., Henehan, M., Rae, J., Mortyn, P., and Vance, D.: Boron isotope evidence for oceanic carbon dioxide leakage during the last deglaciation, *Nature*, 518, 219–222, 2015.
- 720 Mashiotta, T. A., Lea, D. W., and Spero, H. J.: Glacial–interglacial changes in Subantarctic sea surface temperature and $\delta^{18}\text{O}$ -water using foraminiferal Mg, *Earth Planet. Sc. Lett.*, 170, 417–432, 1999.
- Miller, K. G., Browning, J. V., Aubry, M.-P., Wade, B. S., Katz, M. E., Kulpecz, A. A., and Wright, J. D.: Eocene–Oligocene global climate and sea-level changes: St. Stephens Quarry, Alabama, *Geol. Soc. Am. Bull.*, 120, 34–53, 2008.
- 725 Müller, W., Shelley, M., Miller, P., and Broude, S.: Initial performance metrics of a new custom–designed ArF excimer LA–ICPMS system coupled to a two-volume laser–ablation cell, *J. Anal. Atom. Spectrom.*, 24, 209–214, 2009.
- Nehrke, G., Keul, N., Langer, G., de Nooijer, L. J., Bijma, J., and Meibom, A.: A new model for biomineralization and trace-element signatures of Foraminifera tests, *Biogeosciences*, 10, 6759–6767, doi:10.5194/bg-10-6759-2013, 2013.
- 730 Palmer, M.: Rare earth elements in foraminifera tests, *Earth Planet. Sc. Lett.*, 73, 285–298, 1985.

- Panchuk, K., Ridgwell, A., and Kump, L.: Sedimentary response to Paleocene–Eocene Thermal Maximum carbon release: a model-data comparison, *Geology*, 36, 315–318, 2008.
- 735 Pearce, N. J., Perkins, W. T., Westgate, J. A., Gorton, M. P., Jackson, S. E., Neal, C. R., and Chenery, S. P.:
A compilation of new and published major and trace element data for NIST SRM 610 and NIST SRM 612
glass reference materials, *Geostandard. Newslett.*, 21, 115–144, 1997.
- Pearson, P. N., Van Dongen, B. E., Nicholas, C. J., Pancost, R. D., Schouten, S., Singano, J. M., and Wade, B. S.:
Stable warm tropical climate through the Eocene Epoch, *Geology*, 35, 211–214, 2007.
- 740 Pearson, P. N., Foster, G. L., and Wade, B. S.: Atmospheric carbon dioxide through the Eocene–Oligocene
climate transition, *Nature*, 461, 1110–1113, 2009.
- Penman, D. E., Hönisch, B., Zeebe, R. E., Thomas, E., and Zachos, J. C.: Rapid and sustained surface ocean
acidification during the Paleocene–Eocene Thermal Maximum, *Paleoceanography*, 29, 357–369, 2014.
- Politi, Y., Arad, T., Klein, E., Weiner, S., and Addadi, L.: Sea urchin spine calcite forms via a transient amor-
phous calcium carbonate phase, *Science*, 306, 1161–1164, 2004.
- 745 Raitzsch, M., Dueñas-Bohórquez, A., Reichart, G.-J., de Nooijer, L. J., and Bickert, T.: Incorporation of Mg
and Sr in calcite of cultured benthic foraminifera: impact of calcium concentration and associated calcite
saturation state, *Biogeosciences*, 7, 869–881, doi:10.5194/bg-7-869-2010, 2010.
- Regenberg, M., Regenberg, A., Garbe-Schönberg, D., and Lea, D. W.: Global dissolution effects on planktonic
foraminiferal Mg/Ca ratios controlled by the calcite-saturation state of bottom waters, *Paleoceanography*,
750 29, 127–142, 2014.
- Ries, J. B., Cohen, A. L., and McCorkle, D. C.: Marine calcifiers exhibit mixed responses to CO₂-induced
ocean acidification, *Geology*, 37, 1131–1134, 2009.
- Rosenthal, Y., Perron-Cashman, S., Lear, C. H., Bard, E., Barker, S., Billups, K., Bryan, M., Delaney, M. L.,
Demenocal, P. B., Dwyer, G. S., Elderfield, H., German, C. R., Greaves, M., Lea, D. W., Marchitto, T. M.,
755 Pak, D. K., Paradis, G. L., Russell, A. D., Schneider, R. R., Scheiderich, K., Stott, L., Tachikawa, K.,
Tappa, E., Thunell, R., Wara, M., Weldeab, S., and Wilson, P. A.: Interlaboratory comparison study of Mg/Ca
and Sr/Ca measurements in planktonic foraminifera for paleoceanographic research, *Geochem. Geophys.
Geosy.*, 5, Q04D09, doi:10.1029/2003GC000650, 2004.
- Russell, A., Hönisch, B., Spero, H., and Lea, D.: Effects of seawater carbonate ion concentration and tempera-
760 ture on shell U, Mg, and Sr in cultured planktonic foraminifera, *Geochim. Cosmochim. Ac.*, 68, 4347–4361,
2004.
- Schmitz, B. and Pujalte, V.: Abrupt increase in seasonal extreme precipitation at the Paleocene–Eocene bound-
ary, *Geology*, 35, 215–218, 2007.
- 765 Sluijs, A., Schouten, S., Pagani, M., Woltering, M., Brinkhuis, H., Damsté, J., Dickens, G., Huber, M., Re-
ichart, G., Stein, R., Matthiessen, J., Lourens, L. J., Pedentchouk, N., Backman, J., Moran, K., and the
Expedition 302 Scientists: Subtropical Arctic Ocean temperatures during the Palaeocene/Eocene thermal
maximum, *Nature*, 441, 610–613, 2006.
- 770 Sluijs, A., Brinkhuis, H., Schouten, S., Bohaty, S. M., John, C. M., Zachos, J. C., Reichart, G.-J.,
Damsté, J. S. S., Crouch, E. M., and Dickens, G. R.: Environmental precursors to rapid light carbon injection
at the Palaeocene/Eocene boundary, *Nature*, 450, 1218–1221, 2007.

- Sluijs, A., Bijl, P. K., Schouten, S., Röhl, U., Reichart, G.-J., and Brinkhuis, H.: Southern ocean warming, sea level and hydrological change during the Paleocene-Eocene thermal maximum, *Clim. Past*, 7, 47–61, doi:10.5194/cp-7-47-2011, 2011.
- Sosdian, S. and Rosenthal, Y.: Deep-sea temperature and ice volume changes across the Pliocene–Pleistocene climate transitions, *Science*, 325, 306–310, 2009.
- 775 Spanswick, R.: Evidence for an electrogenic ion pump in *Nitella translucens*. I. The effects of pH, K⁺, Na⁺, light and temperature on the membrane potential and resistance, *BBA-Biomembranes*, 288, 73–89, 1972.
- Spero, H. J., Bijma, J., Lea, D. W., and Bemis, B. E.: Effect of seawater carbonate concentration on foraminiferal carbon and oxygen isotopes, *Nature*, 390, 497–500, 1997.
- 780 ter Kuile, B. and Erez, J.: Uptake of inorganic carbon and internal carbon cycling in symbiont-bearing benthonic foraminifera, *Mar. Biol.*, 94, 499–509, 1987.
- ter Kuile, B., Erez, J., and Padan, E.: Mechanisms for the uptake of inorganic carbon by two species of symbiont-bearing foraminifera, *Mar. Biol.*, 103, 241–251, 1989.
- Tierney, J. E. and Tingley, M. P.: A Bayesian, spatially-varying calibration model for the TEX₈₆ proxy, *Geochim. Cosmochim. Ac.*, 127, 83–106, 2014.
- 785 Tripathi, A. and Elderfield, H.: Abrupt hydrographic changes in the equatorial Pacific and subtropical Atlantic from foraminiferal Mg/Ca indicate greenhouse origin for the thermal maximum at the Paleocene–Eocene Boundary, *Geochem. Geophys. Geosy.*, 5, Q02006, doi:10.1029/2003GC000631, 2004.
- Tyrrell, T. and Zeebe, R. E.: History of carbonate ion concentration over the last 100 million years, *Geochim. Cosmochim. Ac.*, 68, 3521–3530, 2004.
- 790 Uchikawa, J. and Zeebe, R. E.: Examining possible effects of seawater pH decline on foraminiferal stable isotopes during the Paleocene–Eocene Thermal Maximum, *Paleoceanography*, 25, PA2216, doi:10.1029/2009PA001864, 2010.
- Wade, B. S., Houben, A. J., Quaijtaal, W., Schouten, S., Rosenthal, Y., Miller, K. G., Katz, M. E., Wright, J. D., and Brinkhuis, H.: Multiproxy record of abrupt sea-surface cooling across the Eocene–Oligocene transition in the Gulf of Mexico, *Geology*, 40, 159–162, 2012.
- 795 Waelbroeck, C., Paul, A., Kucera, M., Rosell-Melé, A., Weinelt, M., Schneider, R., Mix, A., Abelmann, A., Armand, L., Bard, E., Barker, S., Barrows, T. T., Benway, H., Cacho, I., Chen, M.-T., Cortijo, E., Crosta, X., de Vernal, A., Dokken, T., Duprat, J., Elderfield, H., Eynaud, F., Gersonde, R., Hayes, A., Henry, M., Hillaire-Marcel, C., Huang, C.-C., Jansen, E., Juggins, S., Kallel, N., Kiefer, T., Kienast, M., Labeyrie, L., Leclaire, H., Londeix, L., Mangin, S., Matthiessen, J., Marret, F., Meland, M., Morey, A. E., Mulitza, S., Pflaumann, U., Pisias, N. G., Radi, T., Rochon, A., Rohling, E. J., Saffi, L., Schäfer-Neth, C., Solignac, S., Spero, H., Tachikawa, K., Turon, J.-L.: Constraints on the magnitude and patterns of ocean cooling at the Last Glacial Maximum, *Nat. Geosci.*, 2, 127–132, 2009.
- 800 Winguth, A., Shellito, C., Shields, C., and Winguth, C.: Climate response at the Paleocene–Eocene thermal maximum to greenhouse gas forcing—A model study with CCSM3, *J. Climate*, 23, 2562–2584, 2010.
- Yu, J. and Broecker, W.: Comment on “Deep-Sea Temperature and Ice Volume Changes Across the Pliocene–Pleistocene Climate Transitions”, *Science*, 328, 1480, 2010.

- Yu, J., Day, J., Greaves, M., and Elderfield, H.: Determination of multiple element/calcium ratios in foraminiferal calcite by quadrupole ICP-MS, *Geochem. Geophys. Geosy.*, 6, Q08P01, doi:10.1029/2005GC000964, 2005.
- Zachos, J. C., Quinn, T. M., and Salamy, K. A.: High-resolution (104 years) deep-sea foraminiferal stable isotope records of the Eocene–Oligocene climate transition, *Paleoceanography*, 11, 251–266, 1996.
- Zachos, J. C., Wara, M. W., Bohaty, S., Delaney, M. L., Petrizzo, M. R., Brill, A., Bralower, T. J., and Premoli-Silva, I.: A transient rise in tropical sea surface temperature during the Paleocene–Eocene thermal maximum, *Science*, 302, 1551–1554, 2003.
- Zachos, J. C., Schouten, S., Bohaty, S., Quattlebaum, T., Sluijs, A., Brinkhuis, H., Gibbs, S., and Bralower, T.: Extreme warming of mid-latitude coastal ocean during the Paleocene–Eocene Thermal Maximum: inferences from TEX₈₆ and isotope data, *Geology*, 34, 737–740, 2006.
- Zeebe, R. E.: An explanation of the effect of seawater carbonate concentration on foraminiferal oxygen isotopes, *Geochim. Cosmochim. Ac.*, 63, 2001–2007, 1999.
- Zeebe, R. E., Zachos, J. C., and Dickens, G. R.: Carbon dioxide forcing alone insufficient to explain Palaeocene–Eocene Thermal Maximum warming, *Nat. Geosci.*, 2, 576–580, 2009.