

1 **Author's response**

2 ***Anonymous Referee #1***

3
4 Received and published: 24 September 2015

5
6 **General comments**

7
8 This paper by E. Gautier and co-authors presents an interesting study of local scale
9 variability of sulfate records achieved in a low accumulation site (Dome C, Antarctica), in order to
10 assess the representativeness of a single ice core record for such reconstruction. One of the main
11 outcome of this study is an intra-site variability larger than the one reported in literature for inter-site
12 studies for most of the largest volcanic eruptions of the last 2 kyr. The most surprising result is the
13 absence of the Tambora signature in 2/3 cores out of the 5 drilled and analysed in this work. The
14 increasing interest in the last years in extracting information about climate forcing induced by volcanic
15 eruptions recorded in ice cores makes this paper a good piece of science that deserves publication in
16 "Climate of the Past" after few minor revisions.

17
18 **From a methodological point of view, the authors use a new method with respect to recent
19 literature to identify the volcanic spikes along each sulfate profile. The method is based on the
20 calculation of a background non-volcanic level above which volcanic spikes are detected using a
21 "moving window" in the depth profile. In my opinion it would be better to calculate the running
22 mean in a constant temporal range (and not a constant depth range) but I think that to the
23 purpose of this study it should not make a big difference in the obtained results.**

24
25 We agree with the reviewer that a time window should be used in general to treat time series but on
26 the field it was decided to use a constant depth window for simplicity (no datation was available at the
27 time of the drilling) in selecting the ice core sections to be retrograded to France (for isotopic analysis).
28 As mentioned by the reviewer, the difference between the two approaches should not produce a bias in
29 the analysis, as one sample is equivalent to approx. 4 months for top and 7 months for bottom samples.

30
31 **Minor comments.**

32 **As concerning the Tambora eruption, in the text you write that 2 out of 5 cores don't show the
33 sulfate peak while in the caption of figure 8 you write that 3 cores out of 5 don't show this
34 signature. Correct the text according to what we can see from figure 8 (it seems to me that just 2
35 of the 5 cores show the sulfate peak and that there is no "intermediate" peak as written in the
36 text).**

37
38 The correct statement is that 2 cores out of 5 do not show the sulphate peak. The caption of the figure
39 8 is corrected accordingly.

40 The peak was detected in core 1, 4 and 5, with peaks of 455, 188 and 307 ppb respectively. Even if the
41 peak in core 4 is not obvious in figure 8 (especially compared with the high concentrations in core 1),
42 it was detected by the algorithm.

43
44
45 **P. 3985 line 19 and following : :.Change "Maximums" in maxima.**

46
47 Thank you, the correction was made.

48
49 **It would be interesting to have a new table 2 showing two more columns: the mean volcanic flux
50 and the corresponding SD; this would allow a direct comparison with the fluxes and
51 uncertainties calculated in other papers dealing with this topic.**

53 That is right, these two columns are added in the revised version, caption is modified accordingly. We
54 also added Castellano's data for similar volcanic peaks, (Castellano et al., 2005) for comparison.
55

56 **There is no mention in the paper to the uncertainty of the IC measurements, but I believe that**
57 **part of the differences in the maximum concentration of sulfate when a volcanic event is**
58 **detected can be ascribed to the error associated to the measurement.**
59

60 The uncertainty (relative standard deviation) of the IC measurement is below 4%, (based on standards
61 runs). Therefore the uncertainty associated with the quantification represents only a small portion of
62 the variability recorded and commented below.
63

64 **Can you give an estimate of how big is this uncertainty with respect to the "real" uncertainty in**
65 **the amount of sulfate deposition?**
66

67 The relative error on the flux (estimated as 10%) takes into account the IC measurement relative
68 standard deviation (below 4% based on standards runs), the error on firn density (relative error
69 estimated as 2%) and the error on samples time length (10%) (Information added in table 2 caption).
70

71 **For future works it would be important to know a few details of the sampling site (i.e. the**
72 **approx. distance of the 5 cores from the FIRETRACC ice core and, above all, from the EDC96**
73 **and EDC99 drilling sites).**
74

75 The drilling site was located between Concordia station and EDC drilling tent, 300m west of the EDC
76 drilling tent (added in the text)
77

78 **P.3990 line 8. Check the reference Sigl et al. that seems to be not correct.**
79

80 Right, thank you, the correction was made.
81
82

83 **Anonymous Referee #2**


84
85 Received and published: 24 September 2015

86
87 The manuscript discusses the issue of multiple ice coring for extraction of a volcanic record at the
88 Antarctic Dome C location. The manuscript represents a substantial amount of dedicated and careful
89 work and the results are of interest to a large community and relevant in the context of climate change,
90 constraining of volcanic forcing, IPCC, etc. The manuscript is generally well structured and written,
91 the figures are relevant and referencing is appropriate, except as mentioned below.

92
93 General comments:

94
95 I urge the authors to study a recent publication by Gfeller et al., that is also concerned with multiple
96 ice coring at a single site, although at a higher accumulation site in Greenland. That study is concerned
97 with both seasonal and inter-annual variability of the cores. Whereas seasonality is probably irrelevant
98 for the present study, it may be of interest to try out the approach of Gfeller et al. for the longer term
99 variability, i.e. the volcanic record. In particular, the representativeness parameter as introduced in
100 Gfeller et al. would be interesting to derive for the Antarctic cores. The requirement for applying the
101 the Gfeller approach is that the sulfate concentrations are similar to log normal distributed (Gfeller et
102 al., figure 3). I am uncertain about if that is the case for the Antarctic sulfate records with their
103 volcanic spikes, but in the Gfeller et al. study the method works for conductivity that is often similar
104 to sulfate, so it should be worth investigating.

105
106 The sulphate concentrations do have indeed a log normal distribution (see figure above, based on core
107 1 concentrations), and Gfeller approach seems appropriate.



108
109
110 **You already have a common timescale for your five cores based on your synchronization, so the**
111 **analysis should be fairly straightforward.**

112
113 Gfeller approach is quite different from the approach we adopted because the variability assessment is
114 based on the entire record, while we base our study on isolated peaks. Indeed one of the major
115 differences resides in the fact that in Gfeller all data are equivalent where in our case we used the a
116 priori information that a peak is considered as volcanic if it is detected at least in two cores. Using the
117 Gfeller approach, where all data are used, including background data, delivers the following results on
118 time period of -570 to 1952, common to the 5 cores (4825 concentration values per cores):

	$\check{R}^2_{1,\infty}$	$\check{R}^2_{2,\infty}$	$\check{R}^2_{3,\infty}$	$\check{R}^2_{4,\infty}$	$\check{R}^2_{5,\infty}$
SO ₄ ²⁻	0.72	0.84	0.89	0.91	0.93

120
121

122 However, these representativeness coefficients aggregate the background + volcanoes and thus cannot
123 be directly compared with our approach. Nevertheless the same trends are observed, with a decreasing
124 noise as the number of cores increases (our Figure 6). Because the Gfeller's approach is not
125 compatible with discret signal, we have decided to leave our approach unchanged but add the above
126 table and the associated figure in the supplement material, to give a comparison with the Gfeller
127 « scale » when the full dataset is taken into account.

128
129

130 **It is important that you provide a table or a column in table 2 showing your best estimate of the**
131 **volcanic flux and sulfur deposition for each eruption, i.e. that you somehow provide the mean of**
132 **the five cores including the error/uncertainty estimate. This is the number that is important for**
133 **geographical deposition interpolations, databases, and modelers. In other words, your main**
134 **result for a larger community.**

135
136
137

We thank the reviewer for this comment and this information is now added in our Table 2.

138 **Are there no existing datasets you can compare your results to? What about the EDC volcanic**
139 **record of Severi et al., 2007? It would make much sense to see how the sulfur fluxes of an**
140 **independent study compare to your results. Do they fall within your error estimates? One could**
141 **even discuss the effect of the EDC deep core being drilled further away from your closely spaced**
142 **cores (again following the approach of Gfeller et al.)**

143

144 Sulfate flux of identified volcanic eruption are not provided in Severi et al. 2007, but they are
145 calculated in Castellano et al. 2005. For similar peak, Castellano's flux generally falls into the average
146 flux + 40% uncertainty, but it sometimes exceed this value. Castellano's concentrations and flux are
147 generally higher than our result (Castellano's data are now displayed in Table 2)

148

149 Regarding the comparison with EDC cores, it would indeed be very interesting to follow Gfeller
150 approach but first all the cores will need to be synchronized and interpolated to reach the same
151 sampling resolution. As mentioned before this will lead to a comparison of the whole time series
152 profiles and not only of the volcanoes peak similarity. This could be the scope of a future work that
153 wants to measure the representativeness of a time series at DC but we don't think that Gfeller's
154 approach is well fit for discret events like volcanoes, e.g. deciding what should be tagged as volcano is
155 not included in the Gfeller's approach.

156

157

158 **Specific comments:**

159

160 **Peak discrimination method:**

161 **1) I wonder why you determine the background based on 1m long sections when you sometimes**
162 **have volcanic spikes covering almost half of that interval length? In figures 3b and 8 this**
163 **approach appears to result in too high background determinations for core 1? I would suggest to**
164 **work on longer sections. C1782**

165

166 We are actually working on a 1m-moving window, therefore the background corresponding to one
167 even is calculated a large number of time (each point is considered in at least 50 runs). The 1m-
168 window was also chosen because ice cores were treated, logged and decontaminated by 1m section.

169

170 **2) To determine the background, why do you use the mean across 1m intervals rather than the**
171 **median? The median is much more efficiently discriminating outliers (in your case volcanic**
172 **spikes).**


173

174 Correct, median could have been a better criteria but the difference between median and mean is not
175 expected to be fundamental, as the difference will only play at the margin, for very small events.
176 Looping based on the mean until no peak is further detected will reduce the difference between mean

177 and median. If the background is assumed to be a noise controlled by surface processes, then a close to
178 normal distribution is expected for the background which will result in the median equals the mean.
179 As an example, on the first ten meters in core 1, the median of the background values is 79.67 ppb,
180 while the mean is 81.87 ppb.

181
182
183
184
185
186

3) It would be good to show the derived background together with the data over a longer section of the ice core, so we can better visually judge how well the background determination works.



187
188

Here is illustrated the variation of the background along depth in core 1, red dots are detected peaks, the dark line stands for the background concentration. If this is what the reviewer suggests, this figure can be added in the supplementary online materials.

191
192

Section 2.1: Please sketch/explain the lateral pattern of the five drill locations. Are the 5 cores drilled along a straight line on the snow surface? In that case, the distance between cores 1 and 5 would be 4m and not 1m?

194
195

Correct, we change the text as “drilled along a 5 m straight line, and spaced approximately 1 m apart” corrected on line 110.

199
200

P. 3981, l. 6: I suggest to replace ‘global’ with ‘local’ as global has a different meaning in the context of volcanism.

202
203

Correct, we have corrected the text.

204
205

Figures 3 and 8: Many coloured straight lines are shown close to the background level. If those represent the background level estimates then please mention in caption.

206
207

They actually don't. The different colors stand for different core profiles, none of them represents the background in itself.

209
210

Figure 4: The depth scale is wrong. In ice cores you rarely have both linear depth and age scales.

211
212

Correct, we made a poor manipulation to have both scales on the graph, which does not seem feasible with the program we use. We kept sulfate vs. age on the figure 4.

214
215

In figure 6, I am somewhat puzzled by the logarithmic fit to the data points. The fit suggests that the more ice cores you drill, the more volcanic events you will find. With no upper limit. That is

217
218

219 **not convincing. Instead, I would expect something similar to the representativeness parameter of**
220 **Gfeller et al., with an upper limit for (infinitely) many cores.**

221
222 The reason is simple. While Gfeller used an hypothetical upper limit to scale his coefficient, there is a
223 priori not known upper limit for the number of volcanic peak to be detected. Again this is another
224 illustration of the limit of Gfeller approach for discret event. We obviously agree that there should be
225 a fix number of volcanic peaks at the end and that an asymptotic value should be reached. However, as
226 our criteria is based on the detection of a common peak at least in two ice cores, multiplying the
227 number of cores increases the probability of such criteria to be verified, but as the number of core
228 increases our criteria of common detection should also become more stringent (e.g. with 20 cores, a
229 detection in 5 cores could be used), finally resulting in an asymptotic value as the number of core
230 increases. All the difficulty resides in the number of occurrence that should be taken to label an outlier
231 as a event (and therefore the level of confidence).
232 We think that the log fit is actually an approximation (resulting from a poor statistic) of a more general
233 law that should level off with more cores. To avoid such confusion, we decided to remove the
234 equation.
235
236

237 **Authors answer to:**

238 ***Interactive comment on “Variability of sulfate signal in ice-***
239 ***core records based on five replicate cores” by E. Gautier et al.***

240 **EW Wolff (Editor)**

241 ew428@cam.ac.uk
242 Received and published: 27 October 2015

243 **I will be asked to give a formal editorial comment after you post your replies to review- ers.**
244 **However meanwhile it is obvious that both reviewers are generally favourable to your paper,**
245 **and I will therefore be encouraging you to submit a revised version for CP, taking account of**
246 **their comments. I also have a few comments of my own.**

247 **There are a few typos which you already have from me.**

248 **Page 3980, line 6. If a peak has to pass the threshold in 3 consecutive points that means it has to**
249 **be most probably 6 cm wide. At the bottom (of the studied section this would mean the peak**
250 **must span more than 2 years. Such a threshold is likley to exclude some genuine peaks. Please**
251 **comment on this. I wonder also if some of the cases where you see a peak in only 2,3,or 4 of the**
252 **cores are ones whwere a peak is present but not across 3 samples. While this is technically a "no**
253 **peak detected" it is probably not what the reader imagines when they read this. Please comment.**

254 This choice of 3 consecutive data points is a compromise to avoid detecting noises instead of volcanic
255 peaks. Volcanic peaks detected in ice cores tend to be wider than expected if a typical 1-3 years-long
256 fallout is considered, especially at high depth (Wolff et al. 2005). The widening has been attributed to
257 diffusional effects on sulfate in the ice, by Barnes et al. (2003). Following their assessment, Castellano
258 et al. (2004) estimated that the peak broadening during the holocene was close to 2 cm. In the bottom
259 of the core, 6 cm wide represents more than two years, but considering the typical fallout time as well
260 as the peak widening, it seems improbable that a volcanic eruption will be imprinted in less than 3
261 consecutive data points for any of the 5 cores. .

262 Regarding the second comment, the algorithm disregards peaks not made of 3 consecutive samples in
263 any given ice core. These “sharp” peaks are simply not treated and not retained. It is therefore possible
264 that a volcanic peak is found in less than 5 cores because of such selection criteria.

265 However, for both comments above, the reader should understand that to build a more reliable
266 volcanic record, peaks shape must also be considered. As a result, after the algorithm treatment, the
267 last step is a visual inspection across all the profiles. For the sake of the objectivity of the statistical
268 assessment, no visual sorting was applied in the present paper.

269 In the main text it is now clearly mentioned that a final visual inspection must be performed to build a
270 more reliable volcanic record, also based on peaks shape.

271 **I found the mathematical description from lines 3-12 very hard to follow. Could you also explain**
272 **it in simpler terms.**


273 We agree with this comment and have simplified the text as follows, which summarizes the procedure
274 with the same rigor as the discussion paper:

275 After correcting the depth shift between cores, a composite profile was built by summing all the peaks
276 identified in the 5 cores. In this composite, sulfate peaks from different cores are associated to a same
277 event as soon as their respective depth (corresponding to the maximum concentration) are included in
278 a 20cm depth window. This level of tolerance is consistent with the dispersion in width and shape of
279 peaks observed. A number of occurrences is then attributed to each sulfate peak, reflecting the number
280 of time it has been detected in the 5 cores dataset (Figure 4).

281 **Fig 3. These are both examples where the peak is seen in all cores. I would like also to see some**
282 **examples where the peak is only seen in fewer cores. I know there is one in Fig 8 but I suggest to**
283 **expand Fig 3 to include 2 such events.**

284 We agree with this comment, the figure 3 was modified as follows:

285



286

287 **Fig 4 and elsewhere. I am not sure I know how you made the average when the peak is, for**
288 **example, seen only in 3 cores. Is the value shown for sulfate the sum of peak heights divided by 3**
289 **or by 5? Or is it something different?**

290 If detected in 3 cores, the sum is divided by 3. The average is calculated on detected peaks. The paper
291 first comments the fact that peaks are not always detected, and that even when they are, there is still a
292 variability in sulfate concentration. (Table 2 caption was modified accordingly)

293

294 **List of relevant modifications:**

295 Table 1: Dates have been modified in the deeper part, which showed discrepancies with Sigl
296 et al. 2015. Core dating was revised accordingly.

297 Table 2: figure + caption Table modified following reviews. In this new version of the paper,
298 background was calculated individually, for each volcanic events (while it was considered to
299 be around 85ppb all the time in the previous version). That must lead to more accurate flux
300 estimations in the table (although the variation with previous results is not very significant)

301 Figure 4: Scale corrected

302 Fig 6: Equation fit removed

303 Fig 8: bottom graph added

304 Paragraph “Composite building from the 5 ice cores”: simplified explanation

305 SOM added: including Gfeller approach results, and an example on core 1 of the background
306 detection

307

308
309
310
311
312

313 **Variability of sulfate signal in ice-core records based on five replicate cores**

314

315 Gautier Elsa^{1,2}, Savarino Joël^{1,2}, Erbland Joseph^{1,2}, Lanciki Alyson^{1,2}, Possenti Philippe^{1,2}

316

317 ¹Univ. Grenoble Alpes, LGGE, F-38000 Grenoble, France

318 ²CNRS, LGGE, F-38000 Grenoble, France

319

320 **Abstract**

321 Current volcanic reconstructions based on ice core analysis have significantly improved over

322 the past few decades by incorporating multiple core analysis with high temporal resolution

323 from different parts of the Polar Regions. Regional patterns of volcanic deposition are based

324 on composite records, built from cores taken at both poles. However, in many cases only a

325 single record at a given site is used for these reconstructions. This assumes that transport and

326 regional meteorological patterns are the only source of the dispersion of the volcanic-products.

327 Here we evaluate the local scale variability of a sulfate profile in a low accumulation site

328 (Dome C, Antarctica), in order to assess the representativeness of one core for such

329 reconstruction. We evaluate the variability with depth, statistical occurrence, and sulfate flux

330 deposition variability of volcanic eruptions detected on 5 ice cores, drilled 1 meter away from

331 each other. Local scale variability, essentially attributed to snow drift and surface roughness

332 at Dome C, can lead to a non-exhaustive record of volcanic events when a single core is used

333 as the site reference with a bulk probability of 30 % of missing volcanic events and almost

334 60 % uncertainty on the volcanic flux estimation. Averaging multiple records almost erases

335 the probability of missing volcanic events and can reduce by half the uncertainty pertaining to

336 the deposition flux.

337 **Introduction**

GAUTIER Elsa 24/11/y 18:49

Supprimé: last

GAUTIER Elsa 24/11/y 18:49

Supprimé: . Relying on limited and disparate sulfate profiles at first, they have progressively

GAUTIER Elsa 24/11/y 18:50

Supprimé: ed

GAUTIER Elsa 24/11/y 18:50

Supprimé: s

GAUTIER Elsa 24/11/y 18:50

Supprimé: flux

GAUTIER Elsa 24/11/y 18:50

Supprimé: now

GAUTIER Elsa 24/11/y 18:50

Supprimé: several

GAUTIER Elsa 24/11/y 18:51

Supprimé: it is worth mentioning that most of the time

GAUTIER Elsa 24/11/y 18:52

Supprimé: such

GAUTIER Elsa 24/11/y 18:52

Supprimé: implicitly

GAUTIER Elsa 24/11/y 18:53

Supprimé: In the present work,

GAUTIER Elsa 24/11/y 18:53

Supprimé: variability

353 When a large and powerful volcanic eruption occurs, the energy of the blast is sufficient to inject
354 megatons of material directly into the upper atmosphere [Robock, 2000]. While ashes and pyroclastic
355 materials fall rapidly to the ground because of gravity, gases remain over longer time scales. Among
356 gases, SO₂ is of a particular interest due to its conversion to tiny sulfuric acid aerosols, which can
357 potentially impact the radiative budget of the atmosphere [Rampino and Self, 1982; Timmreck, 2012].
358 In the troposphere a combination of turbulence, cloud formation, rainout and downward transport are
359 efficient processes that clean the atmosphere of sulfuric acid, and volcanic sulfuric acid layers rarely
360 survive more than a few weeks, limiting their impact on climate. The story is different when volcanic
361 SO₂ is injected into the stratosphere. There, the dry, cold and stratified atmosphere allows sulfuric acid
362 layers to remain for years, slowly spreading an aerosols blanket around the globe. The tiny aerosols
363 then act as efficient reflectors and absorbers of incoming solar radiations, significantly modifying the
364 energy balance of the atmosphere [Kiehl and Briegleb, 1993] and the ocean [Gleckler et al., 2006;
365 Miller et al., 2012; Ortega et al., 2015]. With a lifetime of 2 to 4 years, these aerosols of sulfuric acid
366 ultimately fall into the troposphere where they are removed within weeks.
367 In Polar Regions, the deposition of the sulfuric acid particles on pristine snow will generate an acidic
368 snow layer, enriched in sulfate. The continuous falling of snow, the absence of melting and the ice
369 thickness make the polar snowpack the best records of the Earth's volcanic eruptions. Hammer [1977]
370 was the first to recognize the polar ice propensity to record such volcanic history. Built on the seminal
371 work of Hammer et al., a paleo-volcanism science developed around this discovery with two aims.
372 The first relies on the idea that the ice record can reveal past volcanic activity and, to a greater extent,
373 its impact on Earth's climate history [Robock, 2000; Timmreck, 2012]. Indeed, at millennium time
374 scale, volcanoes and the solar activity are the only two recognized natural climate forcings [Stocker et
375 al., 2013]. Based on ice records, many attempts are made to extract the climate forcing induced by a
376 volcanic eruption [Crowley and Unterman, 2013 ; Gao et al., 2008; Gao et al., 2007; Sigl et al., 2013;
377 Sigl et al., 2014; Zielinski, 1995]. However, such an approach is inevitably prone to large uncertainty
378 pertaining to the quality of the ice record and non-linear effects between deposition fluxes and source
379 emissions [Pfeiffer et al., 2006].

GAUTIER Elsa 24/11/y 18:55

Supprimé: on

GAUTIER Elsa 24/11/y 18:55

Supprimé: y

GAUTIER Elsa 24/11/y 18:55

Supprimé: for

GAUTIER Elsa 24/11/y 18:56

Supprimé: them

GAUTIER Elsa 24/11/y 18:57

Supprimé: where

GAUTIER Elsa 24/11/y 18:57

Supprimé: s

GAUTIER Elsa 24/11/y 18:58

Supprimé: s

GAUTIER Elsa 24/11/y 18:58

Supprimé: o

GAUTIER Elsa 24/11/y 18:59

Supprimé: the

GAUTIER Elsa 24/11/y 19:00

Supprimé: and thus have a limited action on the Earth's

GAUTIER Elsa 24/11/y 19:00

Supprimé: the

GAUTIER Elsa 24/11/y 19:00

Supprimé: reaches

GAUTIER Elsa 24/11/y 19:01

Supprimé: the

GAUTIER Elsa 24/11/y 19:02

Supprimé: in the atmosphere

GAUTIER Elsa 24/11/y 19:03

Supprimé: efficiently

GAUTIER Elsa 24/11/y 19:04

Supprimé: the

GAUTIER Elsa 24/11/y 19:04

Supprimé: to reach the ground

GAUTIER Elsa 24/11/y 19:05

Supprimé: fold

GAUTIER Elsa 24/11/y 19:06

Supprimé:

GAUTIER Elsa 25/11/y 09:09

Supprimé: t

401 The second aim of the paleo-volcanism is to provide an absolute dating scale when clear volcanic
402 events in differently located ice cores can be unambiguously attributed to the same dated event [Severi
403 *et al.*, 2007]. The time synchronization of different proxy records is possible, allowing study of the
404 phasing response of different environmental parameters to climate perturbation [Ortega *et al.*, 2015;
405 *Sigl et al.*, 2015] or estimating the snow deposition over time [Parrenin *et al.*, 2007]. Whatever the
406 intent, paleo-volcanism should rely on robust and statistically relevant ice core records.

407 Work to establish a volcanic index, undertaken to date, has assumed volcanic event are clearly
408 identified, without any false signal from background variations induced by other sulfur sources (eg
409 marine, anthropogenic, etc). Seasonal layer counting is used whenever possible, bi-polar comparison
410 of ice sulfate records has become the method of choice to establish an absolute dated volcanic index
411 [Langway *et al.*, 1988]. Both known or unknown events can be used to synchronize different cores.
412 However, only a limited number of peaks, with characteristic shape or intensity, and known to be
413 associated with a dated eruption, can be used to set a reliable time scale [Parrenin *et al.*, 2007]. This
414 restriction is partly fueled by the poor and/or unknown representativeness of most volcanic events
415 found in ice cores. Most of the time, a single core is drilled at a given site and used for cross
416 comparison with other sites. This approach is clearly insufficient for ambiguous events.

417 At a large scale, sulfate deposition is highly variable in space and mainly associated with atmospheric
418 transport and precipitation patterns. At a local scale (ca. 1m), variability can emerge from post-
419 deposition processes. While sulfate is a non-volatile species supposed to be well preserved in snow,
420 spatial variability is induced by drifted snow, wind erosion leading to surface roughness
421 heterogeneities [Libois *et al.*, 2014]. These effects are amplified in low accumulation sites where most
422 of the deep drilling sites are performed [EPICA-community-members, 2004; Jouzel, 2013; Lorius *et al.*,
423 1985]). To the best of our knowledge, one single study has used multiple drillings at a given site to
424 analyze the representativeness of the ice core record [Wolff *et al.*, 2005]. This study took advantage of
425 the two EPICA cores drilled at Dome C, 10 m apart (Antarctica, 75°06'S, 123°21'E, elevation 3220 m,
426 mean annual temperature -54.5°C) [EPICA-community-members, 2004] to compare the dielectric
427 profile (DEP) along the 788 m common length of the two cores. For the two replicate cores, statistical
428 analysis showed that up to 50 % variability in the pattern of any given peak was encountered as a

GAUTIER Elsa 24/11/y 19:07

Supprimé: Per se,

GAUTIER Elsa 24/11/y 19:07

Supprimé: t

GAUTIER Elsa 24/11/y 19:07

Supprimé: ing

GAUTIER Elsa 24/11/y 19:07

Supprimé: a

GAUTIER Elsa 24/11/y 19:08

Supprimé: is

GAUTIER Elsa 24/11/y 19:08

Supprimé: the

GAUTIER Elsa 24/11/y 19:09

Supprimé: To date

GAUTIER Elsa 24/11/y 19:10

Supprimé: a core or

GAUTIER Elsa 24/11/y 19:10

Supprimé: the work

GAUTIER Elsa 24/11/y 19:10

Supprimé: s

GAUTIER Elsa 24/11/y 19:11

Supprimé: a clear identification of a

GAUTIER Elsa 24/11/y 19:12

Supprimé: confusion with

GAUTIER Elsa 24/11/y 19:12

Supprimé: Besides s

GAUTIER Elsa 24/11/y 19:13

Supprimé: is

GAUTIER Elsa 24/11/y 19:13

Supprimé: K

GAUTIER Elsa 24/11/y 19:13

Supprimé: both

GAUTIER Elsa 24/11/y 19:14

Supprimé: of the

GAUTIER Elsa 24/11/y 19:14

Supprimé:

GAUTIER Elsa 24/11/y 19:14

Supprimé: ,

GAUTIER Elsa 24/11/y 19:14

Supprimé: which

GAUTIER Elsa 24/11/y 19:15

Supprimé: particularly

450 consequence of the spatial variability of the snow deposition. The authors concluded that ice-core
451 volcanic indices from single cores at such low-accumulation sites couldn't be reliable and what was
452 required was a network of close-spaced records. However, as mentioned in Wolff's conclusion, this
453 statistical study relied only on two records. Additionally, DEP signals are known to be less sensitive
454 than sulfate signals for volcanic identification, and more accuracy is expected by comparing sulfate
455 profiles. The authors thus encouraged conducting a similar study on multiple ice cores to see if the
456 uncertainty could be reduced.

457 In the present study we took advantage of the drilling of 5 ice cores at Dome C, initially intended for
458 the analysis of sulfur isotopes of the volcanic sulfate. Putting aside the number of records, our
459 approach is similar in many points to Wolff's work. However, it has the advantage of relying on highly
460 resolved sulfate profiles. In addition, the spatial scale is slightly smaller as the 5 cores were drilled 1-
461 meter apart. The comparison of 5 identically processed cores is a chance to approach the
462 representativeness of a single core reconstruction at a low accumulation site, the most prone to spatial
463 variability. Therefore new constraints on variability of sulfate deposition, recorded by spatial
464 heterogeneity in such sites are expected from the present work. Even if recent publications [Sigl et al.,
465 2014], underline the need of using multiple records in low accumulation sites, to overcome the spatial
466 variability issue, such records are not always available. This lack of records adds uncertainty in the
467 volcanic flux reconstruction based on polar depositional pattern. Our study should help to better
468 constrain the error associated with local scale variability, and ultimately, the statistical significance of
469 volcanic reconstructions. The present study discusses the depth shift, occurrence of events and
470 deposition flux variability observed in the 5 cores drilled.

471

472 **Experimental setup and Methods**

473 **Core drilling**

474 The project "VolSol", initiated in 2009, aimed at constraining the estimation of the natural part of
475 radiative forcing, composed of both volcanic and solar contributions using ice core records of sulfate
476 and Beryllium-10. In order to build a robust volcanic index including a discrimination of stratospheric
477 events based on sulfur isotopic ratios [Baroni et al., 2008; Savarino et al., 2003], 5 x 100 m-firm cores

GAUTIER Elsa 24/11/y 19:17

Supprimé: variability

GAUTIER Elsa 24/11/y 19:17

Supprimé: induced

480 | (dia. 10 cm) were drilled in 2010/2011 [along a 5 m straight line, and spaced approximately 1 m apart.](#)
481 | The drilling took place at the French-Italian station Concordia (Dome C, Antarctica, 75°06'S,
482 | 123°21'E, elevation 3220 m, mean annual temperature -54.5°C) more precisely between Concordia
483 | station and EDC drilling tent (300m west of the EDC drilling tent). At this site, the mean annual snow
484 | accumulation rate is about 25 kg m⁻² y⁻¹, leading to an estimated time-period covered by the cores of
485 | 2500 years. Cores were logged and bagged in the field, and temporarily stored in the underground core
486 | buffer (- 50 °C) before analysis. The unusual number of ice core drilled at the same place was driven
487 | by the amount of sulfate necessary to conduct the isotopic analysis. However, this number of replicate
488 | cores drilled 1m apart offers the opportunity to question the [representativeness](#) of a volcanic signal
489 | extracted from a single core per site.

490

491 **Sampling, Resolution and IC Analyses**

492 | Analyses were directly performed on the field during two consecutive summer campaigns. Thirty
493 | meters were analyzed in 2011, the rest was processed the following year. The protocol was identical
494 | for each core and the steps followed were:

- 495 | - Decontamination of the external layer by scalpel scrapping
- 496 | - Longitudinal cut with a band saw of a 2 cm stick of the most external layer
- 497 | - Sampling of the ice stick at a 2 cm-resolution (ca. 23 600 samples)
- 498 | - Thawing the samples in 50 ml centrifuge tubes, and transfer in 15 ml centrifuge tubes positioned in
499 | an autosampler
- 500 | - Automatic analysis with a Metrohm IC 850 in suppressed mode (NaOH at 7 mM, suppressor H₂SO₄
501 | at 50 mM, Dionex AG11 column), in a fast IC configuration (2 min run) with regular calibration
502 | (every 60 samples) using certified sulfate reference solution (Fisher brand, 1000 ppm certified).

503 | Due to the fragility of snow cores, the first 4 m were only analyzed on a single core (Figure 1). We
504 | will thus not discuss the variability of the Pinatubo and Agung eruptions present in these first 4 meters.
505 | Concentration data are deposited in the public domain and made freely available in NOAA National
506 | Climatic Data center.

507

508 **Peaks discrimination method**

509 | As with most algorithms used for peak detection, the principle is to detect anomalous sulfate
510 concentration peaks from a background noise (stationary or not), which could potentially indicate a
511 volcanic event. The estimation of the background value should therefore be as accurate as possible.
512 Using core 2 as our reference core, we observed a background average value stationary and close to 85
513 ppb \pm 30 ppb (1σ) at Dome C during the 2,500 years of the record. However, the variability is
514 | sufficient enough to induce potential confusion on detection of small peaks. Therefore, a stringent
515 algorithm using PYTHON language (accessible on demand) was developed to isolate each possible
516 peak. The algorithm treats the full ice record by 1-meter section (ca. 45-50 samples). For each meter, a
517 mean concentration (m) and standard deviation (σ) is calculated regardless of the presence or not of
518 peaks in the section. Then, every value above the $m + 2\sigma$ is removed from the 1-meter dataset. A new
519 mean and standard deviation is calculated and the same filtration is applied. Iteration runs until no
520 more data above $m + 2\sigma$ is found. At that point, m represents the background mean concentration.
521 The process runs for each 1-m section, starting from the surface sample and until the end of the core.
522 Then, each 1-meter dataset is shifted by one sample; the process is reset and the peak detection run
523 again on each new 1-m dataset. Sample shift is applied until the last sample of the first 1-meter section
524 is reached so that no bias is introduced by the sampling scheme. Every concentration data point is thus
525 compared approximately with its 100 neighbor data (50 of each side). Each data point isolated by the
526 algorithm is further tested. To be considered as a point belonging to a potential volcanic peak, the data
527 should be detected in a given core (i.e. for being above the $m + 2\sigma$ final threshold) in at least 50 % of
528 the 50 runs. Additionally, the point has to be part of at least three consecutive points passing the same
529 50 % threshold detection. This algorithm was applied individually on each core, giving 5 different
530 lists of peak. In total, 54, 51, 47, 50 and 47 peaks were detected on core 1, 2, 3, 4 and 5, respectively.
531 A manual detection is then required if one wants to build a more accomplished volcanic record from
532 several profiles, which must be based on shape criteria, and not only statistical criteria. However, in
533 the scope of this paper, no manual sorting was applied, so that the statistical assessment could rely on
534 more objective criteria (the number of occurrences).

535

GAUTIER Elsa 24/11/y 19:18

Supprimé: detection

537 **Core synchronization and dating**

538 Core 1 was entirely dated with respect to the recently published volcanic ice core database [Sigl *et al.*,
539 2015] using *Analyseries* 2.0.8 software (<http://www.lsce.ipsl.fr/Phocea/Page/index.php?id=3>), and
540 covers the time period of -588 to 2010 CE. Figure 2 shows the age-depth profile obtained for this core.
541 A total of 13 major volcanic eruptions well dated were used as time markers to set a time scale (bold
542 date in Table 1). Core 1 was entirely dated through linear interpolation between those tie points. Dated
543 core 1 was then used as a reference to synchronize the remaining 4 cores, using the same tie points and
544 10 additional peaks (non-bold date in Table 1), presenting characteristic patterns common to each core.
545 In total, 23 points were therefore used to synchronize the cores.

546

547 **Composite building from the 5 ice cores**

548 Through the routine described above, the five cores are depth-synchronized using the 23 tie points and
549 other potential volcanic events in each core cores are detected independently. Therefore, the number
550 of peaks detected in each core is different (between 47 and 54) and their depth (with the exception of
551 the tie points used) is slightly different to each other cores due to sampling scheme and position of the
552 maximum concentration. After correcting the depth shift between cores, a composite profile was built
553 by summing all the peaks identified in the 5 cores. In this composite, sulfate peaks from different
554 cores are associated to a same event as soon as their respective depth (corresponding to the maximum
555 concentration) are included in a 20cm depth window. This level of tolerance is consistent with the
556 dispersion in width and shape of peaks observed. A number of occurrences is then attributed to each
557 sulfate peak, reflecting the number of time it has been detected in the 5 cores dataset (Figure 4).

558

559 **Results and Discussions**

560 **Depth offset between cores**

561 Depth offsets between cores are the result of the surface roughness at the time of drilling, variability in
562 snow accumulation, heterogeneous compaction during the burying of snow layers and logging

GAUTIER Elsa 24/11/y 19:19

Supprimé: s

GAUTIER Elsa 25/11/y 10:02

Commentaire [1]: Simplified explanation

564 uncertainty. This aspect has been discussed previously, over a similar time-scale (Wolff et al. 2005),
565 and over a longer time-scale (Barnes et al. 2006) in Dome C. Surface roughness, attributed to wind
566 speed, temperatures and accumulation rate, is highly variable in time and space. These small features
567 hardly contribute to the depth offset on a larger spatial scale, in which case glacial flow can control the
568 offset between synchronized peaks, as it seems to be the case in South pole site (Bay et al. 2010).
569 However, in Dome C, and at the very local spatial scale we are considering in the present work,
570 roughness is significant regarding to the accumulation rate. It is therefore expected that synchronized
571 peaks should be found at different depths. The offset trend fluctuates with depth, due to a variable
572 wind speed (Barnes et al. 2006). To estimate the variability in the depth shift for identical volcanic
573 events, we used the tie points listed in Table 1. For each peak maximum, we evaluate the depth offset
574 of core 1, 3, 4 and 5, with respect to core 2. To avoid logging uncertainty due to poor snow
575 compaction in the first meters of the cores and surface roughness at the time of the drilling, we used
576 the UE 1809 depth in core 2 (13.30 m) as a depth reference horizon from which all other depth cores
577 were anchored using the same 1809 event. For this reason, only eruptions prior to 1809 were used to
578 evaluate the offset variability, that is 18 eruptions instead of the 23 used for the core synchronization.
579 Figure 5, shows the distribution of depth shift of the cores with respect to core 2. While the first 40 m
580 appear to be stochastic in nature, a feature consistent with the random local accumulation variations
581 associated with snow drift in Dome C site, it is surprising that at greater depth, offset increases (note
582 that the positive or negative trends are purely arbitrary and depends only on the reference used, here
583 core 2). The maximum offset, obtained between core 3 and 5 is about 40 cm. Such accrued offsets
584 with depth were also observed by Wolff et al., [2005] and were attributed to the process of logging
585 despite the stringent guidelines used during EPICA drilling. Similarly, discontinuities in the depth
586 offset, observed by Barnes et al., [2006] were interpreted as resulting from logging errors. As no
587 physical processes can explain a trend in the offsets, we should also admit that the accrued offset is
588 certainly the result of the logging process. In the field, different operators were involved but a
589 common procedure was used for the logging. Two successive cores extracted from the drill were
590 reassembled on a bench to match the non uniform drill cut and then hand sawed meter by meter to get
591 the best precise depth core, as neither the drill depth recorder nor the length of the drilled core section

GAUTIER Elsa 24/11/y 19:21

Supprimé: s

593 can be used for establishing the depth scale. This methodology involving different operators should
594 have randomized systematic errors but obviously this was not the case. Despite the systematic depth
595 offset observed, synchronization did not pose fundamental issues as the maximum offset in rescaled
596 profiles never exceeds the peak width (ca. 20 cm) thank to the 10 possible comparisons when pair of
597 core are compared. Confusion of events or missing of events are thus very limited in our analysis (see
598 next section).

GAUTIER Elsa 25/11/y 09:19
Supprimé: width

600 **Variability in events occurrence**

601 The variability in events occurrence in the 5 ice cores has been evaluated through the construction of a
602 composite record (Figure 4) and the counting of events in each core as described in the method. By
603 combining the five ice cores, we listed a total amount of 91 sulfate peaks (Pinatubo and Agung not
604 included), which are not necessarily from volcanic sources. Some peaks can be due to post deposition
605 effects affecting the background deposition, or even contamination. When it comes to defining a
606 robust volcanic index, peak detection issues emerge. Chances to misinterpret a sulfate peak and assign
607 it, by mistake, to a volcanic eruption, as well as chances to miss a volcanic peak, can be discussed
608 through a statistic analysis conducted on our five cores.

609 We try to evaluate to what extent multiple cores comparison facilitates the identification of volcanic
610 peaks, among all sulfate peaks that can be detected in a core. To do so, we assumed that a peak is of
611 volcanic origin as soon as it is detected at least in two cores. In other words, the probability to have
612 two non-volcanic peaks synchronized in two different cores is nil. It is expected that combining an
613 increasing number of cores will increasingly reveal the real pattern of the volcanic events. All possible
614 combinations from 2 to 5 cores comparison were analyzed, totalizing 26 possibilities for the entire
615 population. The results for each comparison were averaged, giving a statistic on the average number of
616 volcanic peaks identified per number of cores compared. The results of the statistical analysis are
617 presented in Figure 6. As expected, in a composite made of 1 to 5 cores, the number of sulfate peaks
618 identified as volcanic peaks (for being detected at least twice) increases with the number of cores
619 combined in the composite. Thus, while only 30 peaks can be identified as volcanic from a two cores
620 study, a study based on 5 cores can yields 62 such peaks. The 5-cores comparison results in the

GAUTIER Elsa 25/11/y 09:10
Supprimé: d

623 composite profile given in Figure 4a. The initial composite of 93 peaks is reduced to 64 volcanic
624 peaks (Pinatubo and Agung included) after removing the single peaks (Figure 4b). Each characteristic
625 of the retained peaks is given in Table 2. The main conclusion observing the final composite record is
626 that only 17 of the 64 peaks were detected in all of the 5 cores and 68 % of all peaks were at least
627 present in two cores. At the other side of the spectrum, 2-cores analysis reveals that only 33 % (30
628 peaks on average) of the peaks are identified as possible eruptions. Two cores comparison presents
629 still a high risk of not extracting the most robust volcanic profile at low accumulation sites, a
630 conclusion similar to *Wolff et al.*, [2005]. Surprisingly, it can also be noticed that this 5-core
631 comparison doesn't results in an asymptotic ratio of identified volcanic peaks, suggesting that 5 cores
632 are not sufficient either to produce a full picture. High accumulation sites should be prone to less
633 uncertainty; however, this conclusion remains an a priori that still requires a confirmation.

634 Large and small events are not equally concerned by those statistics. Figure 7 shows that the
635 probability of presence is highly dependent on peak flux and the chance to miss a small peak
636 (maximum flux in the window $[f + 2\sigma : f + 5\sigma]$, f being the background average flux) is much higher
637 than the chance to miss a large one (maximum flux above $f + 8\sigma$). However, it is worth noticing that
638 major eruptions can also be missing from the record, as it has already been observed in other studies
639 [*Castellano et al.*, 2005; *Delmas et al.*, 1992]. The most obvious example in our case is the Tambora
640 peak (1815 AD), absent in 2 of our 5 drillings, while presenting an intermediate to strong signal in the
641 others (Figure 8). The reason for the variability in event occurrence has been discussed already by
642 *Castellano et al.*, [2005]. In the present case of close drillings, long-range transport and large-scale
643 meteorological conditions can be disregarded due to the small spatial scale of our study; the snow drift
644 and surface roughness is certainly the main reasons for missing peaks. The fact that two close events
645 as UE 1809 and Tambora are so differently recorded indicates how punctual, in time and space post-
646 depositional effects can affect the recording of eruptions.

647

648 **Variability in signal strength**

649 To compare peak height variability, detected peaks were corrected by subtracting the background from
650 peak maximum. We considered C_i/C_{mean} variations, C_i being the SO_4^{2-} maximum concentration in core

GAUTIER Elsa 24/11/y 19:24

Supprimé: s

652 i (1 to 5), and C_{mean} being the mean of those concentration for the event i . For concentration values,
653 positive by definition, the log-normal distribution is more appropriate; geometric means and geometric
654 standard deviations were used, as described by *Wolff et al.*, [2005] (Table 3). In our calculation, the
655 geometric standard deviation based on 2 cores is 1.35; in other words, maximum concentrations are
656 uncertain by a factor 1.35. This factor is slightly lower than the one obtained in *Wolff et al.*, [2005]
657 (1.5). Our cores are drilled closer (one meter from each others, instead of 10 m for *Wolff et al.*), which
658 might slightly reduce the uncertainty. The peaks height variability obtained by averaging 5 cores
659 (1.21), matches *Wolff et al.* forecast. Based on a 50 % uncertainty on 2 cores, *Wolff et al.* predicted a
660 20 % uncertainty on a 5 cores study (consistent with a reduction of the standard deviation by a factor
661 of $1/\sqrt{n}$, by averaging n values). Comparing the peaks maximum enables us to compare our study with
662 *Wolff's* study, also based on peaks maximum. However, in our case, comparing maximum induces a bias
663 related to the sampling method: with a two centimeters resolution on average, peak's height is directly
664 impacted by the cutting, which tends to smooth the maximum. Comparing the total sulfate deposited
665 during the event is more appropriate. Proceeding on a similar approach, but reasoning on mass of
666 deposited sulfate rather than maximum concentration, the obtained variability is higher than
667 previously: 41 % uncertainty on volcanic deposited sulfate mass, on a 5-cores study (F_i/F_{mean} , F_i being
668 the mass flux of peak i), and 56 % uncertainty on a 2-cores comparison (F_i/F_1). The difference in the
669 signal dispersion between the two approaches rests on the fact that peak maximum has a tendency to
670 smooth the concentration profile as a consequence of the sampling strategy. This artifact is suppressed
671 when the total mass deposited is considered. In any case, uncertainty seems to be significantly reduced
672 when comparing 5 cores instead of 2.

673

674 **Conclusion:**


675 This study confirms in many ways previous work on multiple drilling variability [*Wolff et al.*, 2005].
676 As already discussed, peaks flux uncertainty can be significantly reduced (56 % to 41 %) by averaging
677 5 ice-cores signals instead of 2. A 5-cores composite profile has been built using the criteria that a
678 peak is considered as volcanic if present at least in two cores. We observed that the number of
679 volcanic peaks listed in a composite profile increases with the number of cores considered. With 2

680 | cores, only 33 % of the peaks present in the composite profile are tagged as volcanoes. [This](#)
681 | [percentage](#) increases to 68 % with 5 cores. However, we did not observe an asymptotic value, even
682 | with 5 cores drilled. A record based on a single record in a low accumulation site is therefore very
683 | unlikely to be a robust volcanic record. Of course, peaks presenting the largest flux are more likely to
684 | be detected in any drilling, but the example of the Tambora shows that surface topography is variable
685 | enough to erase even the most significant signal, although rarely. This variability in snow surface is
686 | evidenced in the depth offset between two cores drilled less than 5 meters from each other, as peaks
687 | can easily be situated 40 cm apart.

688 | In low accumulation sites such as Dome C, where surface roughness can be on the order of the snow
689 | accumulation and highly variable, indices based on chemical records should be considered with
690 | respect to the time-scale of the proxy studied. Large time-scale trends are faintly sensitive to this effect.
691 | On the contrary, a study on episodic events like volcanic eruptions or biomass burning, with a
692 | deposition time in the order of magnitude of the surface variability scale should be based on a
693 | multiple-drilling analysis. A network of several cores is needed to obtain a representative record, at
694 | least in terms of recorded events. However, although lowered by the number of cores, the flux remains
695 | highly variable, and still uncertain by a factor of 1.4 with 5 cores. This point is particularly critical in
696 | volcanic reconstructions that [rely](#) on the deposited flux to estimate the mass of aerosols loaded in the
697 | stratosphere, and to a larger extent, the climatic forcing induced. Recent reconstructions largely take
698 | into account flux variability associated with regional pattern of deposition, but this study underlines
699 | the necessity of not neglecting local scale variability in low accumulation sites. Less variability is
700 | expected with higher accumulation rate, but this still has to be demonstrated. Sulfate flux is clearly
701 | one of the indicators of the eruption strength, but due to transport, deposition and post-deposition
702 | effects, such direct link should not be taken for granted.

703 | With such statistical analysis performed systematically at other sites, we should be able to reveal even
704 | the smallest imprinted volcanoes in ice cores, extending the absolute ice core dating, the
705 | teleconnection between climate and [volcanic events](#) and improving the time-resolution of mass
706 | balance calculation of ice sheets.

707



709 **Acknowledgments**

710 Part of this work would not have been possible without the technical support from the C2FN (French
711 National Center for Coring and Drilling, handled by INSU. Financial supports were provided by
712 LEFE-IMAGO, a scientific program of the Institut National des Sciences de l'Univers (INSU/CNRS),
713 the Agence Nationale de la Recherche (ANR) via contract NT09-431976- VOLSOL and by a grant
714 from Labex OSUG@2020 (Investissements d'avenir – ANR10 LABX56). E. G. deeply thanks the
715 Fulbright commission for providing the PhD Fulbright fellowship. The Institute Polaire Paul-Emile
716 Victor (IPEV) supported the research and polar logistics through the program SUNITEDC No. 1011.
717 We would also like to thank all the field team members present during the VOLSOL campaign and
718 who help us. Data are available at the World Data Center for paleoclimatology
719 (<http://www.ncdc.noaa.gov/paleo/wdc-paleo.html>).

720

721

722

723

724 **References**

725

726 Baroni, M., J. Savarino, J. Cole-Dai, V. K. Rai, and M. H. Thiemens (2008), Anomalous sulfur isotope
727 compositions of volcanic sulfate over the last millennium in Antarctic ice cores, *J Geophys Res*,
728 *113*(D20), D20112, doi: 10.1029/2008jd010185.

729 Barnes, P. R. F., E. W. Wolff, and R. Mulvaney (2006), A 44 kyr paleoroughness record of the
730 Antarctic surface, *J. Geophys. Res.*, 111, D03102, doi:10.1029/2005JD006349.

731 Bay, R. C., R. A. Rohde, P. B. Price, and N. E. Bramall (2010), South Pole paleowind from automated
732 synthesis of ice core records, *J. Geophys. Res.-Atmos.*, 115, doi:10.1029/2009jd013741.

733 Castellano, E., S. Becagli, M. Hansson, M. Hutterli, J. R. Petit, M. R. Rampino, M. Severi, J. P.
734 Steffensen, R. Traversi, and R. Udisti (2005), Holocene volcanic history as recorded in the sulfate

735 stratigraphy of the European Project for Ice Coring in Antarctica Dome C (EDC96) ice core, *J*
736 *Geophys Res*, *110*(D6), D06114, doi: 10.1029/2004jd005259.

737 Crowley, T. J., and M. B. Unterman (2013), Technical details concerning development of a 1200 yr
738 proxy index for global volcanism, *Earth Syst. Sci. Data*, *5*(1), 187-197, doi: 10.5194/essd-5-187-2013.

739 Delmas, R. J., S. Kirchner, J. M. Palais, and J. R. Petit (1992), 1000 years of explosive volcanism
740 recorded at the South-Pole, *Tellus Ser. B-Chem. Phys. Meteorol.*, *44*(4), 335-350.

741 EPICA-community-members (2004), Eight glacial cycles from an Antarctic ice core, *Nature*, *429*,
742 623-628, doi: 10.1038/nature02599.

743 Gao, C., A. Robock, and C. Ammann (2008), Volcanic forcing of climate over the past 1500 years: An
744 improved ice core-based index for climate models, *J Geophys Res*, *113*(D23), D23111, doi:
745 10.1029/2008jd010239.

746 Gao, C., L. Oman, A. Robock, and G. L. Stenchikov (2007), Atmospheric volcanic loading derived
747 from bipolar ice cores: Accounting for the spatial distribution of volcanic deposition, *J Geophys Res*,
748 *112*(D9), D09109, doi: 10.1029/2006jd007461.

749 Gleckler, P. J., K. AchutaRao, J. M. Gregory, B. D. Santer, K. E. Taylor, and T. M. L. Wigley (2006),
750 Krakatoa lives: The effect of volcanic eruptions on ocean heat content and thermal expansion,
751 *Geophys Res Lett*, *33*(17), L17702, doi: 10.1029/2006gl026771.

752 Hammer, C. U. (1977), Past Volcanism Revealed by Greenland Ice Sheet Impurities, *Nature*,
753 *270*(5637), 482-486.

754 Jouzel, J. (2013), A brief history of ice core science over the last 50 yr, *Climate of the Past*, *9*(6),
755 2525-2547, doi: 10.5194/cp-9-2525-2013.

756 Kiehl, J. T., and B. P. Briegleb (1993), The Relative Roles of Sulfate Aerosols and Greenhouse Gases
757 in Climate Forcing, *Science*, *260*(5106), 311-314, doi: 10.1126/science.260.5106.311.

758 Langway, C. C., H. B. Clausen, and C. U. Hammer (1988), An inter-hemispheric volcanic time-
759 marker in ice cores from Greenland and Antarctica, *Annals of Glaciology*, *10*, 102-108.

760 Libois, Q., G. Picard, L. Arnaud, S. Morin, and E. Brun (2014), Modeling the impact of snow drift on
761 the decameter-scale variability of snow properties on the Antarctic Plateau, *Journal of Geophysical*
762 *Research: Atmospheres*, *119*(20), 11,662-611,681, doi: 10.1002/2014jd022361.

763 Lorius, C., J. Jouzel, C. Ritz, L. Merlivat, N. I. Barkov, Y. S. Korotkevich, and V. M. Kotlyakov
764 (1985), A 150,000-year climatic record from Antarctic ice, *Nature*, 316(6029), 591-596, doi:
765 10.1038/316591a0.

766 [Miller, G. H., Geirsdóttir, Á., Zhong, Y., Larsen, D. J., Otto-Bliesner, B. L., Holland, M. M., Bailey,](#)
767 [D. a., Refsnider, K. a., Lehman, S. J., Southon, J. R., Anderson, C., Björnsson, H. and Thordarson, T.](#)
768 (2012), Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean
769 feedbacks, *Geophys. Res. Lett.*, 39(2), L02708, doi: 10.1029/2011gl050168.

770 Ortega, P., F. Lehner, D. Swingedouw, V. Masson-Delmotte, C. C. Raible, M. Casado, and P. Yiou
771 (2015), A model-tested North Atlantic Oscillation reconstruction for the past millennium, *Nature*,
772 523(7558), 71-74, doi: 10.1038/nature14518.

773 [Parrenin, F., Barnola, J.-M., Beer, J., Blunier, T., Castellano, E., Chappellaz, J., Dreyfus, G., Fischer,](#)
774 [H., Fujita, S., Jouzel, J., Kawamura, K., Lemieux-Dudon, B., Loulergue, L., Masson-Delmotte, V.,](#)
775 [Narcisi, B., Petit, J.-R., Raisbeck, G., Raynaud, D., Ruth, U., Schwander, J., Severi, M., Spahni, R.,](#)
776 [Steffensen, J. P., Svensson, a., Udisti, R., Waelbroeck, C. and Wolff, E. W. \(2007\), The EDC3](#)
777 [chronology for the EPICA dome C ice core, *Climate of the Past*, 3\(3\), 485-497, doi: 10.5194/cp-3-](#)
778 [485-2007.](#)

779 Pfeiffer, M. A., B. Langmann, and H. F. Graf (2006), Atmospheric transport and deposition of
780 Indonesian volcanic emissions, *Atmos Chem Phys*, 6, 2525-2537, doi:10.5194/acp-6-2525-2006.

781 Rampino, M. R., and S. Self (1982), Historic eruptions of Tambora (1815), Krakatau (1883), and
782 Agung (1963), their stratospheric aerosols, and climatic impact, *Quat. Res.*, 18(2), 127-143, doi:
783 10.1016/0033-5894(82)90065-5.

784 Robock, A. (2000), Volcanic eruptions and climate, *Reviews of Geophysics*, 38(2), 191-219.

785 Savarino, J., A. Romero, J. Cole-Dai, S. Bekki, and M. H. Thiemens (2003), UV induced mass-
786 independent sulfur isotope fractionation in stratospheric volcanic sulfate, *Geophys Res Lett*, 30(21),
787 2131, doi: 10.1029/2003gl018134.

788 [Severi, M., Becagli, S., Castellano, E., Morganti, a., Traversi, R., Udisti, R., Ruth, U., Fischer, H.,](#)
789 [Huybrechts, P., Wolff, E. W., Parrenin, F., Kaufmann, P., Lambert, F. and Steffensen, J. P. \(2007\),](#)

790 Synchronisation of the EDML and EDC ice cores for the last 52 kyr by volcanic signature matching,
791 *Clim. Past*, 3(3), 367-374, doi: 10.5194/cp-3-367-2007.

792 [Sigl, M., McConnell, J. R., Layman, L., Maselli, O., Mcgwire, K., Pasteris, D., Dahl-jensen, D.,](#)
793 [Steffensen, J. P., Vinther, B., Edwards, R., Mulvaney, R. and Kipfstuhl, S.](#) (2013), A new bipolar ice
794 core record of volcanism from WAIS Divide and NEEM and implications for climate forcing of the
795 last 2000 years, *Journal of Geophysical Research: Atmospheres*, 118(3), 1151-1169, doi:
796 10.1029/2012jd018603.

797 [Sigl, M., McConnell, J. R., Toohey, M., Curran, M., Das, S. B., Edwards, R., Isaksson, E., Kawamura,](#)
798 [K., Kipfstuhl, S., Krüger, K., Layman, L., Maselli, O. J., Motizuki, Y., Motoyama, H. and Pasteris, D.](#)
799 [R.](#) (2014), Insights from Antarctica on volcanic forcing during the Common Era, *Nature Clim. Change*,
800 4(8), 693-697, doi: 10.1038/nclimate2293.

801 [Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee,](#)
802 [M., Chellman, N., Dahl-Jensen, D., Fischer, H., Kipfstuhl, S., Kostick, C., Maselli, O. J., Mekhaldi, F.,](#)
803 [Mulvaney, R., Muscheler, R., Pasteris, D. R., Pilcher, J. R., Salzer, M., Schüpbach, S., Steffensen, J.](#)
804 [P., Vinther, B. M. and Woodruff, T. E.](#) (2015), Timing and climate forcing of volcanic eruptions for
805 the past 2,500 years, *Nature*, doi: 10.1038/nature14565.

806 Stocker, T. F., D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, B. V.,
807 and M. P. M. (2013), IPCC, 2013: The Physical Science Basis, Fifth Assessment Report of the
808 Intergovernmental Panel on Climate Change, Intergovernmental Panel on Climate Change 2013,
809 United Kingdom and New York, NY, USA.

810 Timmreck, C. (2012), Modeling the climatic effects of large explosive volcanic eruptions, *Wiley*
811 *Interdiscip. Rev.-Clim. Chang.*, 3(6), 545-564, doi: 10.1002/wcc.192.

812 Wolff, E. W., E. Cook, P. R. F. Barnes, and R. Mulvaney (2005), Signal variability in replicate ice
813 cores, *Journal of Glaciology*, 51(174), 462-468, doi: 10.3189/172756505781829197.

814 Zielinski, G. A. (1995), Stratospheric loading and optical depth estimates of explosive volcanism over
815 the last 2100 years derived from the Greenland- Ice-Sheet-Project-2 ice core, *J Geophys Res*,
816 100(D10), 20937-20955.

817

818 **Table 1** – Tie points used to set the time scale and synchronize the cores. Volcanic events are
 819 named "Ev x" if they are not assigned to a well-known eruption. Dating of the events is based
 820 on *Sigl et al.*, [2015].
 821

GAUTIER Elsa 25/11/y 10:00
Commentaire [2]: Dates have been modified in the deeper part, which showed discrepancies with Sigl et al. 2015. Cores dating was revised accordingly.

Eruption	core 1	core 2	core 3	core 4	core 5	date of deposition
Surface	0	0	0	0	0	2010
Pinatubo	1.53					1992
Krakatoa	8.82	8.92	8.67	8.71	8.63	1884
Cosiguina	11.98	11.83	11.65	11.62	11.46	1835
Tambora	12.85			12.6	12.57	1816
UE 1809	13.33	13.3	13.04	13.08	12.98	1809
ev 7	15.98	15.93	15.66	15.67	15.52	1762
Serua/UE	19.29	19.22	18.93	18.94	18.78	1695
<u>Ev 10</u>	21.87	21.74	21.53	21.48	21.4	1646
kuwae	30.18	30.04	29.92	29.85	29.73	1459
ev 16 - A	37.35	37.29	37.17	37.04	36.91	1286
ev 16 - B	37.77	37.77	37.62	37.52	37.4	1276
ev 16 - C	38.1	38.04		37.78		1271
Samalas	38.49	38.46	38.28	38.2	38.09	1259
ev 17	39.59	39.56	39.46	39.36	39.2	1230
ev 18	41.87	41.83	41.7	41.6	41.41	1172
ev 22	50.26	50.3	50.2	50.11	49.87	95 99
ev 27	60.77	60.72	60.66		60.27	68 4
ev 31	65.72	65.74	65.68	65.6	65.25	54 1
ev 35	76.06	76.13	76	75.94	75.64	23 5
ev 46	90.42	90.53	90.36	90.41	89.95	-2 14
ev 49	97.15	97.16	97.19	97.22	96.74	-4 26
ev 51	100.16	100.19		100.22	99.7	-5 29

822

823

824 **Table 2** – Sulfate peak (maximum concentration, in ng.g⁻¹, and flux of volcanic sulfate
825 deposited, in kg.km⁻²) considered as volcanic eruptions based on the statistical analysis of the
826 5 cores. Flux is calculated by integrating the peak, using the density profile obtained during
827 the logging process. Volcanic flux values are corrected from background sulfate (calculated
828 separately for each sulfate peak). 0 stands for non-detected events in the cores. Agung
829 (3.77m) and Pinatubo (1.52m) were not included in the statistical analysis because they were
830 analyzed only in core one and thus are marked as not applicable (N/A). The estimation of the
831 average volcanic flux is calculated considering detected peaks only (non detected peaks are
832 not included in this estimation). The relative error on the flux (estimated as 10%) takes into
833 account the IC measurement relative standard deviation (below 4% based on standards runs),
834 the error on firm density (relative error estimated as 2%) and the error on samples time length
835 (10%). The last column displays data obtained from Castellano *et al.* (2005), for identical
836 volcanic peaks. For similar peaks Castellano's flux generally falls into the average flux + 40%
837 uncertainty, sometimes exceeding this value.

GAUTIER Elsa 25/11/y 09:52

Commentaire [3]: Table modified following reviews. In this new version of the paper, background was calculated individually, for each volcanic events (while it was considered to be around 85ppb all the time in the previous version). That must lead to more accurate flux estimations in the table (although the variation with previous results is not very significant)

Peak depth (m)	date (year)	core 1		core 2		core 3		core 4		core 5		average*			Castellano et al., 2005	
		[SO ₄ ²⁻]	Volc. flux	[SO ₄ ²⁻]	Volc. flux	[SO ₄ ²⁻]	Volc. flux	[SO ₄ ²⁻]	Volc. flux	[SO ₄ ²⁻]	Volc. flux	[SO ₄ ²⁻]	Volc. flux	1σ (flux)	[SO ₄ ²⁻]	Volc. flux
1.52	1992	188	5.0	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	188	5.0	0.5	313	11
3.77	1964	207	5	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	362	8
6.24	1929	0	0.0	164	1.3	0	0.0	132	1.1	0	0.0	148	1.2	0.1		
8.59	1891	0	0.0	0	0.0	0	0.0	134	1.3	117	0.9	126	1.1	0.1	140	3.1
8.92	1885	232	8.1	262	8.8	236	10.5	240	10.2	216	7.7	237	9.1	0.9	289	9.3
11.83	1839	220	7.7	173	5.4	190	4.9	177	5.5	173	4.0	187	5.5	0.6		
12.08	1834	0	0.0	0	0.0	144	2.5	0	0.0	137	1.3	140	1.9	0.2		
12.91	1816	455	13.1	0	0.0	0	0.0	188	1.8	307	6.0	317	7.0	0.7	606	39.3
13.3	1809	436	16.6	291	10.5	392	12.7	408	16.3	461	13.4	398	13.9	1.4	271	10.2
15.93	1762	176	2.7	248	6.7	201	3.4	0	0.0	0	0.0	208	4.2	0.4	174	4.5
19.29	1695	287	13.4	0	0.0	168	9.2	194	7.3	0	0.0	217	10.0	1.0	185	8.8
20.3	1674	261	7.8	0	0.0	0	0.0	196	4.3	178	2.3	212	4.8	0.5	142	5.3
20.7	1666	0	0.0	0	0.0	0	0.0	123	1.6	149	2.4	136	2.0	0.2		
21.74	1646	257	10.1	249	10.3	259	13.2	282	17.5	257	13.2	261	12.8	1.3		
22.72	1625	181	4.8	146	2.7	141	2.9	0	0.0	0	0.0	156	3.5	0.3	175	8.0
23.77	1600	225	10.6	0	0.0	170	2.5	0	0.0	0	0.0	197	6.5	0.7	194	13.4
25.78	1557	144	2.1	0	0.0	0	0.0	148	2.2	0	0.0	146	2.1	0.2		
30	1459	496	33.2	442	31.1	422	31.6	543	37.2	559	36.9	493	34.0	3.4	399	31.7
30.56	1449	0	0.0	143	1.8	131	2.8	0	0.0	0	0.0	137	2.3	0.2		
31.83	1417	0	0.0	0	0.0	0	0.0	155	2.6	148	2.6	151	2.6	0.3		
33.51	1377	0	0.0	0	0.0	140	2.3	0	0.0	162	5.4	151	3.9	0.4		
34.85	1348	273	12.4	288	14.2	209	7.9	303	18.3	269	13.2	268	13.2	1.3	211	10.4
37.29	1286	325	18.3	324	16.1	373	17.1	347	14.8	458	30.7	365	19.4	1.9	258	22.4
37.77	1276	563	28.9	605	40.4	570	28.8	525	26.3	497	21.6	552	29.2	2.9	304	20.5
38.04	1271	205	4.1	180	3.1	0	0.0	235	5.1	0	0.0	206	4.1	0.4		
38.46	1259	1086	59.7	1022	63.8	928	61.4	1030	78.5	1428	104.8	1099	73.6	7.4	637	60.4

39.25	1239	0	0.0	0	0.0	132	2.6	147	2.4	151	2.7	143	2.5	0.3		
39.56	1230	268	17.8	260	16.8	279	15.6	315	18.7	320	16.7	288	17.1	1.7	337	25.2
41.17	1191	0	0.0	216	4.2	247	12.9	0	0.0	241	7.3	235	8.1	0.8	227	18.0
41.83	1172	437	30.9	401	29.4	377	25.2	378	23.3	433	29.4	405	27.6	2.8	311	20.8
44.4	1111	186	5.3	0	0.0	243	5.4	225	9.7	195	6.2	212	6.7	0.7		
44.87	1099	174	2.5	0	0.0	0	0.0	153	2.4	0	0.0	163	2.5	0.2		
45.81	1075	129	1.6	144	2.3	0	0.0	0	0.0	0	0.0	137	2.0	0.2		
47.15	1041	187	3.6	193	3.6	217	4.4	0	0.0	203	6.2	200	4.5	0.4		
47.5	1031	192	7.0	163	5.0	166	3.1	0	0.0	198	4.5	180	4.9	0.5		
48	1018	0	0.0	155	3.2	168	2.8	0	0.0	0	0.0	161	3.0	0.3		
49.63	976	132	2.0	0	0.0	139	2.5	0	0.0	0	0.0	135	2.2	0.2		
50.3	959	209	8.2	256	15.6	236	12.6	220	11.9	227	12.1	230	12.1	1.2		
52.49	902	254	3.9	0	0.0	215	4.8	184	5.9	233	7.7	222	5.6	0.6		
54.35	852	0	0.0	0	0.0	0	0.0	155	2.3	249	5.2	202	3.7	0.4		
55.65	819	184	8.8	193	7.3	191	6.7	181	7.1	249	5.2	200	7.0	0.7		
58.26	749	155	3.2	202	3.4	0	0.0	201	6.6	0	0.0	186	4.4	0.4		
60.72	684	287	12.9	216	14.0	243	7.8	0	0.0	230	4.9	244	9.9	1.0		
64.49	577	528	36.0	0	0.0	430	25.8	367	21.4	393	23.3	430	26.6	2.7		
65.74	541	287	19.1	274	12.7	283	20.5	306	21.5	304	16.3	291	18.0	1.8		
68.41	465	132	2.9	0	0.0	182	4.4	0	0.0	0	0.0	157	3.7	0.4		
69.41	436	194	10.7	168	3.8	0	0.0	207	11.1	233	9.1	201	8.7	0.9		
72.38	352	0	0.0	172	4.7	203	5.3	0	0.0	188	5.8	188	5.3	0.5		
73.13	331	0	0.0	169	4.1	152	2.8	0	0.0	0	0.0	160	3.5	0.3		
73.95	304	0	0.0	0	0.0	171	3.7	190	5.7	0	0.0	180	4.7	0.5		
76.13	235	205	12.1	258	20.0	237	21.7	287	23.8	262	13.0	250	18.1	1.8		
77.17	206	179	5.4	206	15.4	211	12.5	219	13.2	272	13.5	217	12.0	1.2		
78.31	172	250	15.3	0	0.0	156	4.3	203	5.4	219	7.7	207	8.2	0.8		
79.98	125	165	4.4	187	3.7	0	0.0	162	3.2	167	3.3	170	3.7	0.4		
84.5	-4	202	9.8	199	7.7	222	5.0	0	0.0	188	7.9	203	7.6	0.8		
85.44	-37	0	0.0	155	4.4	0	0.0	0	0.0	240	8.6	197	6.5	0.7		
87.89	-128	236	11.2	212	9.6	270	12.9	244	12.1	0	0.0	241	11.4	1.1		
89.28	-173	0	0.0	0	0.0	0	0.0	190	5.6	164	3.7	177	4.7	0.5		
90.53	-214	276	18.8	286	26.1	278	16.5	296	18.1	241	6.9	275	17.3	1.7		
91.72	-251	0	0.0	0	0.0	0	0.0	227	10.4	244	12.5	236	11.4	1.1		
94.83	-347	0	0.0	191	4.6	198	5.9	216	8.7	0	0.0	201	6.4	0.6		
97.16	-426	331	22.6	228	15.4	403	35.2	436	48.5	675	75.0	414	39.3	3.9		
97.31	-431	0	0.0	131	2.9	0	0.0	0	0.0	0	0.0	65	1.5	0.1		
100.19	-529	219	12.1	224	6.6	0	0.0	247	15.9	235	7.7	231	10.6	1.1		

838

839

840

841

842

843

844

845

846 **Table 3** – Statistics on sulfate signal for identical peaks in core 1, 2, 3, 4 and 5. Geometric
 847 standard deviations are calculated on peaks heights (i.e maximum concentration reached, in
 848 ng.g⁻¹) and on peaks sulfate flux (i.e total mass of volcanic sulfate deposited after the
 849 eruption). Background corrections are based on background values calculated separately for
 850 each volcanic event.

Study	Number of compared cores	Geom. std deviation based on maximum concentration	Geom std deviation based on deposition flux
Wolff and others	2	1.5	
This study	2*	1.3 <u>5</u>	1. <u>56</u>
This study	5	1.21	1. <u>41</u>

851 * : C_x/C_1 , with x=2,3,4,5

852
 853
 854
 855
 856
 857
 858
 859
 860
 861
 862
 863
 864
 865
 866

867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884






Figure 1 - Sulfate profiles on the 5 replicate cores obtained during a drilling operation at Dome C – Antarctica in 2011.



885
886
887
888
889


Figure 2 - Age versus depth in core 1 drilled in 2011 CE, Dome C – Antarctica



891
 892 **Figure 3** –Kuwae (a, top), Krakatoa (b, middle) and Tambora (c, bottom) sulfate
 893 concentration profiles after depth synchronization. All peaks are within a 20 cm uncertainty,
 894 enabling to clearly attribute each occurrence to a single event.

GAUTIER Elsa 25/11/y 10:05
Commentaire [4]: The bottom graph has been added in this version

895
 896
 897
 898
 899
 900
 901
 902
 903
 904




906

907 **Figure 4** – a) Composite sulfate peak profile deduced from our statistical analysis of the 5
 908 cores using our detection peak and synchronization algorithms (see text). The numbers
 909 indicate the number of time a common peak is found in the cores. Unnumbered peaks, peaks
 910 found only in single core. b) same as a) without the single detected peaks. All the remaining
 911 peaks are considered as volcanic eruptions. See Table 2 for details.

912

913



914

915

916 **Figure 5** – Depth offset of 18 common and well-identified volcanic events in cores 1, 3, 4 and


917 5 relatively to core 2. To overcome offset due to the drilling process and poor core quality on

918 the first meters, UE 1809 (depth ca. 13 m) is taken as the origin and horizon reference.

919

920

921




922

923

924 | **Figure 6** – Black dots with red line (left axis) represent the number of sulfate peaks that can
925 | be identified as volcanic peaks in a composite profile, made of n cores (with n ranging from 1
926 | to 5). A sulfate peak appearing simultaneously in at least two cores is considered to be a
927 | volcanic peak. Blue diamonds represent the ratio of identified volcanic peaks, i.e the number
928 | of identified volcanic peaks (plotted on the left axis), relatively to the total number of sulfate
929 | peaks (no discrimination criteria) in a composite made of 5 cores. In our case, the 5 ice-cores
930 | composite comprises 91 sulfate peaks (Agung and Pinatubo excluded). With two cores, only
931 | 33% of them would be identified as being volcanic peaks (detected in both cores), while 68%
932 | of them can be identified as volcanic events using 5 cores.

933

934
935
936




937
938 **Figure 7** - Peaks probability to be detected in 2, 3, 4 or 5 cores, as function of their flux. The
939 three categories of flux are defined by peaks flux value, relatively to the average flux, and
940 quantified by x time (2, 5 and 8) the flux standard deviation, calculated for a 30 ppb standard
941 deviation in concentrations.

942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957

958

959



960

961 **Figure 8:** Close look at UE 1809 and Tambora (1815) events showing the absence of the
962 Tambora event in 2 out of the 5 cores. This figure illustrates the possibility of missing major
963 volcanic eruptions when a single core is used.


964

965 | **SOM**

966
 967 1. Gfeller *et al.* (2014) approach on Dome C 5 cores: calculation of the representativeness
 968

n (number of cores)	1	2	3	4	5
$\tilde{R}_{n,\infty}^2 \text{SO}_4^{2-}$	0.72	0.84	0.89	0.91	0.93


969
 970
 971



972
 973
 974

975 [Figure S1: Representativeness of sulfate in the cores depending on the number of cores n \(based on](#)
 976 [Gfeller et al., 2014 approach\).](#)

977
 978



979

980 | Figure S2 - Variation of the background along depth in core 1, red dots are detected peaks, the dark
981 | line stands for the background concentration.
982 |
983 |
984 |
985 |
986 |
987 |