

## Minor Revision

1. The description of studied area, some precisions are important: the annual precipitation and temperature averages are not sufficient as you discuss in your paper about January temperature (winter) and about summer and winter precipitation. You must add the present-day ones as reference in your paper (with values) in the text and also in the figures showing the past variations (by a little star for example).

**Done:**

**p. 5 l.15-17.**

**Figure 3.**

in the same paragraph, you have to describe more in the text the present day vegetation, not only through Quercus and Cedar as component of the forest. A rapid description of the components of the different types of vegetation cited in the figure 1 is important.

**Done :**

**p.5 l. 25-27.**

Please name Quercus rotundifolia by Quercus. ilex subsp rotundifolia (Tela botanica - I think that it is this species after searching on internet). The reader could see that you have deciduous and evergreen Quercus (Q. Canariensis being deciduous) in the vegetation, it is not evident for him (her).

You mentioned the different species of Pine in response to reviewer 1. If they are a component of the vegetation you have to mention that in your description.

**Done**

**p. 5 l. 22-24.**

2. In material and method paragraph.

a. You have to note that the Quercus (deciduous and evergreen) have been gathered in a unique group (this was a precision asked by reviewer 1) and please do not call it “evergreen Quercus” (as in your figure) if it is not only evergreen ones. So note this curve only “Quercus sp” to be more rigorous. All the three reviewers ask you questions about that.

**Done:**

**Figure 4.**

b. Your discussion is mainly centered on pollen-based climate reconstructions. Then this part of the methods has to be more developed. In fact, you refer to a paper published in 2014. This paper is focused on South Africa. I would like to believe that the method goes well in Mediterranean area too. Nevertheless it remains necessary to bring several precisions. How the Mediterranean taxa had been included in the method? And Cedar which is not a component of the African vegetation)? Have you done tests before application on Mediterranean? This has to be added in the paper or at least a reference that state on the methodology in the Mediterranean. In any case, it is important to detail which taxa have been taking into account for the reconstructions (all taxa?? what about Mediterranean??? All are taken or only some of them are taken??? Same for other groups of taxa - temperate, Cedar,...). it is important to detailed the method as the discussion is based on the reconstructions.

**Done:**

p. 7 l. 12-22.

Table 2 is used to detail which taxa have been taken into account for the reconstructions. We think that this table is not necessary to be included in the manuscript, however, if you think it is useful then we agree to keep it. We let you decide.

c. You have to say in the text that the dates have been done on bulk sediment.  
**Done:**

p. 6 l. 33-34.

3. Results: thanks to add a table to describe more the vegetation changes. However, I would like to see the pollen zones drawn on the diagram even if you do not use them in the discussion. It will make easier the reader for the interpretation of diagram and table.

**Done:**

**Figures 3 and 4.**

4. Discussion: I have seen nothing in the discussion that takes into account the fact that it will be interesting to compare your results with the previous pollen-based climate reconstructions that has been done on Alboran sea cores (Dormoy et al., 2009; Combourieu-Nebout et al., 2009; and in Morocco (Cheddadi et al., 2009) even it had been done with other methods. Reviewer 1 requested such discussion and you respond that you will do that in the revised manuscript. Please do that. Concerning the SI changes, there are papers that already states that the Mediterranean seasonal contrast takes place after 4.2 (see for example Combourieu-Nebout et al. 2013 on Adriatic Sea in which the summer and winter precipitation curves are represented).

**Actually, we believe that there is a misunderstanding concerning this point. As we have stated in our reply to reviewer #1, in Cheddadi et al., (2009) (as stated by the reviewer) there are three climate records among which 2 are from Morocco. Lake Ifrah which chronology does not encompass the late Holocene and therefore it is not realistic to compare it with our record which covers the last 6000 years BP. Then, the second record is the one published by Cheddadi et al., (1998) based on Lake Tigalmamine record. The latter, which indeed covers the last 6000 years, is not in conflict with our climate data since both reconstructions show a decreasing trend of the annual precipitation. Thus, we do insist that (1) there is no conflict between the two climate reconstructions and (2) if a comparison should be made then it should be with Cheddadi et al. (1998) and not with Cheddadi et al. (2009).**

**Having made this statement we have added a comparison with the climate reconstruction from the Alboran Sea record (Combourieu-Nebout et al., 2009), lake levels (Magny et al., 2013) and climate data from Italian pollen records (Peyron et al., 2009). We did not include Dormoy et al. (2009) because the upper age of their record is 4000 which does make the comparison not appropriate with our record which encompasses the last 6000 years. (p. 10 l. 5-14).**

**Another sentence was added for comparing the reconstructed Pann from Hachlaf (which shows an aridity trend after 5ka) with that obtained from Lake Tigalmamine (Cheddadi et al., 1998) has also been added to the discussion (p. 9 l. 7-9).**

5. Reviewer 3 asked you to enlarge the discussion to all the western Mediterranean. In fact, it will be very interesting to replace the data of the lake Hachlah in a broader context. You prefer to stay in a short area in the western part of the west Mediterranean, that's a choice. Nevertheless, in that case, you have to take into account and compare with all the references

from the restricted area you chose: continental, marine (Alboran Sea), done on one site or several and compilation of sites which includes sites from the studied area (ex: Magny et al., several papers; Fletcher and Zielhofer, 2013). Do not forget that and add all the references (for example) that covered the time slice studied in the discussion and really discuss your results face to them.

**Indeed, we prefer to keep this study in a more regional context (Western Mediterranean). As stated above (in reply to point 4 above) we added a section for a comparison the records you have cited.**

6. other corrections to be done.

a. Some French expressions remain in the text (p.5 l.31 “varie de” has to be changed in “varies” please check in all text of other ones are still remaining.

**Done:**

**p. 6 l. 8.**

b. Please correct d13C in  $\delta^{13}C$  in all text (p.7 l. 30 for example)

**Done:**

**p. 6 l. 31.**

**p. 8 l. 18.**

c. The line in red has not been removed in the diagrams as proposed in the response to reviewer. Either you explain what it means in the or you remove it from the figures.

**Done:**

**The presence of the red line is now explained in the captions of both figures 3 and 4.**

d. Please complete the caption of figure 4 as requested by the reviewer 1

**Done.**

e. You added the pollen sums in the fig.4. Please use bigger characters as they are really difficult to read in the proposed figure.

**Done.**

**MARKED-UP MANUSCRIPT VERSION :**

1 **Climate change and ecosystems dynamics over the last 6000 years**  
2 **in the Middle Atlas, Morocco**  
3  
4

5 M. Nourelbait<sup>1,2,3</sup>, A. Rhoujjati<sup>2</sup>, A. Benkaddour<sup>2</sup>, M. Carré<sup>3</sup>, F. Eynaud<sup>4</sup>, P. Martinez<sup>4</sup>, and R.  
6 Cheddadi<sup>3</sup>  
7

8 <sup>1</sup>Université Chouaib Doukkali, Laboratoire Géosciences Marines et Sciences des Sols, unité  
9 associée CNRST (URAC 45), El Jadida, Morocco.

10 <sup>2</sup>Université Cadi Ayyad, Faculté des Sciences et Techniques, unité associée CNRST (URAC  
11 42), Gueliz Marrakech, Morocco.

12 <sup>3</sup>Université Montpellier 2, Institut des Sciences de l'Evolution, UMR UM2-CNRS-IRD 5554,  
13 Montpellier, France.

14 <sup>4</sup>University of Bordeaux, UMR EPOC 5805, CS 50023, 33615 Pessac, Bordeaux, France.  
15  
16  
17

18 Received: 25 July 2015 – Accepted: 28 July 2015 – Published: 1 September 2015  
19

20 Correspondence to: M. Nourelbait ([nourelbait.m@gmail.com](mailto:nourelbait.m@gmail.com))  
21

22 Published by Copernicus Publications on behalf of the European Geosciences Union.

1 | **Abstract**

2 | The present study aims at reconstructing past climate changes and their environmental  
3 | impacts on plant ecosystems during the last 6000 years in the Middle Atlas, Morocco. Mean  
4 | January temperature ( $T_{jan}$ ), annual precipitation ( $P_{ann}$ ), winter ( $P_w$ ) and summer ( $P_s$ )  
5 | precipitation and a seasonal index (SI) have all been quantified from a fossil pollen record.  
6 | Several bio and geo-chemical elements have also been analyzed to evaluate the links between  
7 | past climate, landscape and ecosystem changes.

8 | Over the last 6000 years, climate has changed within a low temperature and precipitation  
9 | range with a trend of aridity and warming towards the present.  $T_{jan}$  has varied within a ca.  
10 |  $2^{\circ}\text{C}$  range and  $P_{ann}$  within less than  $100\text{ mmyr}^{-1}$ . The long-term changes reconstructed in our  
11 | record between 6ka cal BP and today are consistent with the aridity trend observed in the  
12 | Mediterranean basin. Despite the overall limited range of climate fluctuation, we observe  
13 | major changes in the ecosystem composition, the carbon isotopic contents of organic matter  
14 | ( $\delta^{13}\text{C}$ ), the total organic carbon and nitrogen amount, and the carbon to nitrogen ratio (C/N)  
15 | after ca. 3750 cal BP. The main ecosystem changes correspond to a noticeable transition in  
16 | the conifer forest between the Atlas cedar, which expanded after 3750 cal BP, and the pine  
17 | forest. These vegetation changes impacted the sedimentation type and its composition in the  
18 | lake.

19 | Between 5500 and 5000 cal BP, we observe an abrupt change in all proxies which is coherent  
20 | with a decrease in  $T_{jan}$  without a significant change in the overall amount of precipitation.

# 1 Introduction

The amplitude of climate change during the Holocene (11,700 cal BP to the present) is known to be globally less extreme than during the post-glacial period (Bianchi and McCave, 1999; Bond et al., 2001; Debret et al., 2007). However, several studies have shown that there were climate fluctuations (Alley et al., 1997; Wanner et al., 2008) related to the internal variability of the climate system, solar activity, albedo (Ruddiman, 2003; Eddy, 1982; Stuiver *et al.*, 1991), volcanic eruptions (Kelly & Sear, 1984; Sear et al., 1987; Bryson, 1989; Mann et al., 2005), ocean circulation (Manabe & Stouffer, 1988; Dansgaard et al., 1989; Lascaratos et al., 1999; Rohling, 2002), etc. which all have a direct impact on the terrestrial ecosystems (Davis, 1963; Emmanuel *et al.*, 1985). Although climate changes were less pronounced during the Holocene (Andersen et al., 2004; Mayewski et al., 2004; Witt and Schumann, 2005; Frigola et al., 2007; Cheddadi & Bar-Hen, 2009) than during the last post-glacial period, they have still been noticeable enough to be recorded by different proxies (Dorale et al., 1992; Williams et al., 2002; Geiss et al., 2003, 2004). At the global scale, the Holocene climate stability allowed a sustainable vegetation dynamics with long-term ecosystems changes, plant species expansions and migrations, and an increase of species diversity over all latitudes (Rohde, 1992). However, the Holocene period has also recorded some abrupt and cold events such as the one at 8.2 ka cal BP (e.g. Alley and Agustsdottir, 2005) which recorded a depletion of about 4°C in winter temperature in the Eastern Mediterranean (Weninger et al., 2009).

In Morocco, climate changes during the Holocene have also been quantified and they show significant fluctuations (Cheddadi et al., 1998). As a matter of fact, the climate variability of the Holocene is less known than that of the post-glacial (Mayewski et al., 2004) because it has a lower amplitude and is less abrupt. This statement is even more acute in the Mediterranean region where high resolution and chronologically well-constrained Holocene records are much less numerous than in Europe or North America. The Mediterranean area is currently a hotspot of biodiversity (Myers et al., 2000) and it is one of the largest regions in the world that undergo long-lasting and pronounced droughts during the summer season (Roberts et al., 2004; Milano et al., 2013). The southern rim of the Mediterranean region is even more arid than the northern one because of the influence of the Azores high and the Saharan winds which increase the impact of the drought effect during the summer season. Most of the winter precipitation (Pw) originates from the trade winds which carry moisture from the Mediterranean Sea (Martin, 1981). The amount of Pw has a strong impact on the persistence of water bodies and on the lake levels in the Mediterranean area. Strong lake level

1 fluctuations during the Holocene were observed in Lake Van, Turkey (Lemcke and Sturm,  
2 1997), Lago Dell'Accesa and Lago di Mezzano, Italy (Magny et al., 2006), lake Kinneret,  
3 (Hazan et al., 2005) and the Dead Sea, Israel (Migowski et al., 2006), lake Siles, Spain  
4 (Carrion, 2002), and lakes Sidi Ali and Tigalmamine in Morocco (Lamb and Van der kaars,  
5 1995; Märsche-Soulie et al., 2008).

6 The analysis of marine and continental records from the central part of the Mediterranean  
7 shows that the lake levels were high between 10,300 and 4500 cal BP due to an enhanced  
8 moisture availability during both summer and winter (Magny et al., 2013). After 5000 cal BP,  
9 pollen data from southwestern Europe show that drought increased and led to a sustained  
10 reduction of the forest cover (Roberts et al., 2001; Jalut et al., 2009; Jiménez-Moreno et al.,  
11 2015). These environmental changes show that within the long-term climate trend there were  
12 humid-arid episodes that are related to internal forcings of the climate system such as, in the  
13 case of these westernmost Mediterranean ecosystems, the centennial changes in the North  
14 Atlantic Oscillation modes (Jiménez-Moreno et al., 2015), the enhancement/weakening of the  
15 trade winds, or the increase in the coastal upwelling off northwestern Africa (McGregor et al.,  
16 2007).

17 Climate reconstructions from marine pollen records suggest that the Mediterranean  
18 environments may react with a reduced time lag to rapid climate changes (Fletcher et al.,  
19 2010). The response of the western Mediterranean ecosystems has even been synchronous  
20 with the North Atlantic variability during the post-glacial period and the Holocene  
21 (Combourieu Nebout et al., 2009). Changes in the pollen assemblages of a marine record from  
22 the Alboran Sea also show very synchronous fluctuations between the surrounding land  
23 ecosystems changes and the sea surface temperature fluctuations (Fletcher and Sánchez Goñi,  
24 2008; [Combourieu Nebout et al., 2009](#)). Pollen records from the Middle Atlas (Reille, 1976;  
25 Lamb and Van der kaars, 1995; Cheddadi et al., 2009; Rhoujjati et al., 2010; Nour el Bait et  
26 al., 2014; Tabel et al., 2016) and the Rif mountains (Cheddadi et al., 2016) show that the  
27 Holocene climate change had a major impact on the ecosystems composition with a clear  
28 succession of different species sensitive to winter frost, strong rainfall seasonality and/or the  
29 total amount of annual rainfall throughout the year.

30 The aim of the present study is to evaluate the impacts of the climate changes on the  
31 ecosystems and the landscape of the Middle Atlas during the last six millennia. Our approach  
32 is multidisciplinary and based on the analysis of pollen grains, elemental and isotopic  
33 geochemistry and grain size from a fossil record collected in [Lake](#) Hachlaf, Middle Atlas.  
34 Temperature and precipitation variables have been quantified. They show a moderate change



1 which is superimposed by an aridity trend that is combined with an increase in winter  
2 temperature over the past 6000 years. We also observed some noticeable ecosystem and  
3 landscape changes with one rapid and quite abrupt climate fluctuation between 5500 and 5000  
4 detectable in all the proxies used.

## 5 2 Study area

6 The Middle Atlas Mountains, lying in northwestern Morocco, consist of two geological sets  
7 called Pleated and Tabular Middle Atlas (Fig. 1a). The latter is formed by a Paleozoic  
8 basement covered by a Mesozoic thick layer and Cenozoic and Quaternary volcanic flows  
9 (Texier et al., 1985; Herbig, 1988; Harmand and Moukadiri, 1986). The Liasic limestone and  
10 dolostone are shaped by karstic mechanisms (Martin, 1981; Baali, 1998; Hinaje and Ait  
11 Brahim, 2002; Chillasse and Dakki, 2004). In this geomorphological and structural  
12 composition, there exist nowadays about twenty permanent or semi-permanent natural lakes  
13 (Chillasse and Dakki, 2004) among which we can find the studied site, Dayet (lake) Hachlaf  
14 (33°33'20" N; 5°0'0" W; 1700m a.s.l.). This small water body is located about ten kilometers  
15 North-East of Ifrane national park (Fig. 1b). Available meteorological data (HCEFLCD,  
16 20041980–2008) at Dayet HachlafIfrane station show an average annual rainfall of ca.  
17 600885 mm with Pw and Ps ca. 150 and ca. 70 mm, respectively. The,–a mean  
18 Januarymaximum temperature is ca 4of 18 °C and a mean minimum temperature of about 6.3  
19 °C with ca. 90 rainy days per year, and ca. 70 frosty days among which ca. 17 with snow  
20 precipitation. The surface area and depth of the lake change throughout the year reaching up  
21 to respectively 14 ha and 4 m during late spring. The lake is fed by rainwater, snow, surface  
22 runoff and groundwater and has no river inflow.

23 The forest cover around the site (Fig. 1c) is composed of holm oak (*Q. illex subsp.*  
24 *rotundifolia*) which is evergreen and}, zeen oak (*Q. canariensis*}), which is deciduousare both  
25 evergreen, and Atlas cedar (*Cedrus atlantica*) with occurrences of Pinus halepensis.}-  
26 Nowadays, there are some degraded populations of *Cedrus atlantica* with cultivated lands  
27 around the lake. At higher altitude (1700 to 2500 m, Fig. 1c) an herbaceous/shrubby  
28 vegetation (Artemisia herba-alba and Poaceae) dominates the landscapesite.

## 29 3 Materials and methods

30

1 In April 2008, a 2.5m core (33°33'2.49" N, 4°59'41.57" W) was collected using a Russian  
2 corer. Each section of the core was then sub-sampled for the analysis of pollen content (30  
3 samples), grain size (39 samples), organic matter (43 samples) and its isotopic composition  
4 ( $\delta^{13}\text{C}_{(\text{org})}$ ; 46 samples), and total nitrogen and carbonates (43 samples).

5 Pollen grains were extracted using a standard laboratory procedure: HCl (20 %), KOH (10  
6 %),  $\text{ZnCl}_2$ , acetolysis ( $\text{CH}_3\text{CO}_2\text{O}$  and  $\text{H}_2\text{SO}_4$ ), KOH (10 %), ethanol and glycerine. The  
7 identification and counting of pollen grains were performed with an optical microscope (Leica  
8 DM750) using a  $\times 40$  magnification ( $\times 63$  for accurate identifications). The pollen percentages  
9 were calculated on the total sum of pollen grains originating from vascular terrestrial plants.

10 The total pollen grains counted ~~varies~~variede between ca. 200 and 1300. Aquatic plants  
11 percentages (including Cyperaceae and Juncaceae) were excluded from the total pollen sum.  
12 Cyperaceae were considered as aquatic plants since there are *Juncus* and *Cyperus* genera  
13 growing around the lake today.

14 The particle size analysis was carried out at the “*Laboratoire Marocain d’Agriculture*  
15 (LABOMAG)” and was only performed on the sediment fraction  $< 2$  mm. The proportions of  
16 five fractions were identified as follows: coarse sand (2000–200  $\mu\text{m}$ ), fine sand (200–50  $\mu\text{m}$ ),  
17 coarse silt (50–20  $\mu\text{m}$ ), fine silt (20–2  $\mu\text{m}$ ) and clay (below 2  $\mu\text{m}$ ).

18 Organic matter amount (OM) was estimated based on the content of the organic carbon in  
19 lacustrine sediments (OC), elaborated by spectrometry (NF ISO 14235). Sediment OC was  
20 oxidized in a sulfochromic environment with an excess of potassium dichromate at 135 °C.  
21 Subsequently, the determination of chromate ions  $\text{Cr}^{3+}$  formed was analysed by spectrometry.

22 For total nitrogen (TN), the method used was based on the Kjeldahl mineralization (ISO  
23 11464: 1994), but the catalyst used was the titanium dioxide ( $\text{TiO}_2$ ). The technique consists in  
24 assaying the total nitrogen content in the sediment as ammonium, nitrate, nitrite and organic  
25 form.

26 Carbonates were measured by adding HCl to the bulk sediment to decompose all carbonates  
27 (NF ISO 10693: Juin, 1995). The volume of the carbonic gas produced was measured using a  
28 Scheibler apparatus.

29 Stable isotope ratios measurements of carbon were performed on a Thermo Fischer Flash  
30 2000 Elemental Analyzer in line with a VG Isoprime Mass Spectrometer at the University of  
31 Bordeaux. All samples were pretreated with 1N HCl to remove inorganic carbon. The  
32 analytical precision of 0.15‰ was estimated from several calibrated laboratory standards  
33 analyzed along the samples. Stable isotopic ratios were reported as:  $\delta^{13}\text{C}\text{‰} = [(\frac{^{13}\text{C}}{^{12}\text{C}})_{\text{sample}} / \frac{^{13}\text{C}}{^{12}\text{C}}_{\text{std}} - 1] * 1,000$ , where the standard used is Vienna Pee Dee Belemnite (PDB)

34

1 Besides the multi-proxy analysis, four organic samples were dated. All this dates have been  
2 done on bulk sediment. We used the BACON software (Blaauw and Christen, 2011) to  
3 compute the age/depth model (Fig. 2). The default  $^{14}\text{C}$  calibration curve used by BACON for  
4 terrestrial northern hemisphere samples is IntCal13. The AMS  $^{14}\text{C}$  dates were also calibrated  
5 using the “CALIB 7.1” program (Stuiver and Reimer, 1986; table 1). The fossil record  
6 continuously encompassed the last 6000 years.

7 Annual precipitation ( $P_{ann}$ ), mean January temperature ( $T_{jan}$ ) and precipitation seasonal  
8 index (SI) assessment (Fig. 3) were based on pollen data as follows:

$$PSI_{(s)} = (\sum P_w - \sum P_s) / \sqrt{P_{ann}}$$

11 Where  $PSI_{(s)}$  is the seasonal index quantified for sample  $s$ ;  $P_w$  is the sum of December,  
12 January and February precipitation;  $P_s$  is the sum of June, July and August precipitation;  
13  $P_{ann}$  is the total annual precipitation.

14 The monthly mean precipitation  $P_{ann}$  and  $T_{jan}$  were obtained using the probability density  
15 function of modern plant species (pdf-method). This method is described in Chevalier et al.  
16 (2014). In order to apply) and it to a fossil pollen record collected in the Mediterranean area it  
17 requiredrequires a modern database of Mediterranean plant species distributions and their  
18 corresponding modern climate variables. We used a georeferenced database of plant species  
19 that have been georeferenced from *Flora Europaea* ((Hulten and Fries 1986; Jalas et al. 1972,  
20 1973, 1976, 1979, 1980, 1983, 1986, 1989, 1991, 1994) and Hulten and Fries (1986).  
21 Additional geographical distributions were obtained from GBIF (2012) and personal field  
22 observations using GPS in Morocco. In order to use plant species distributions for the pollen-  
23 based climate reconstruction we assigned pollen taxa to the most probable plant species in our  
24 plant database (table 2). The modern climate variables were extracted from the  
25 WORLDCLIM database (Hijmans et al., 2005) and interpolated onto the species occurrences  
26 for inferring their pdfs.)-

## 27 4 Results

28 During the last 6000 years, the main change in the forest cover is marked by a decline of the  
29 pine populations, the expansion of Atlas cedars after 3750 cal BP and the persistence of the  
30 evergreen oaks. Although the latter dominate today the landscape around Lakelake Hachlaf,  
31 the microscope identification of the fossil pollen grains that originate from deciduous or  
32 evergreen plants may often be dubious and therefore, may not be reproducible by another

1 pollen analyst. We have assigned all oak pollen grains to the evergreen *Quercus ilex* in the  
2 climate reconstruction. All other taxa, including trees, shrubs and herbs, also show some  
3 changes but within a much lower range than that of the two conifer taxa, Atlas cedar and pine  
4 (Fig. 4). We have applied a constrained cluster analysis to depict the main changes in the  
5 pollen fossil record. There are four main clusters summarizing the main changes in the  
6 ecosystem composition around ~~Lake~~ Lake Hachlaf over the last 6000 years (table ~~32~~).

7 The grain size analysis revealed the presence of three fractions (Fig. 3) with the following  
8 proportions: clay (22.87%), silt (60.46% with 41.9% of fine silt) and sand (16.67 %). The  
9 dominant silty fraction tends to increase from the bottom to the top of the core after a brief  
10 decline between ca. 5600 and 5200 cal BP. The sandy fraction follows the same pattern. Clay  
11 shows an opposite trend to both the sandy and silty fractions.

12 Carbonates ( $\text{CaCO}_3$ ) content is high throughout the record except around 5200 cal BP (Fig.  
13 3). They are positively correlated with silt and sand. The total organic carbon (TOC) content  
14 is also high and varies significantly between 4 and 27.4% (Fig. 3). The total nitrogen (TN)  
15 remains low throughout the record. The carbon to nitrogen ratio (C/N) varies between 9 and  
16 17.4, and the  $\delta^{13}\text{C}_{\text{org}}$  between  $-21$  and  $-27\text{‰}$  (Fig. 3). Two origins of the organic matter are  
17 thus identified, with lake algae characterized by  $\text{C/N} < 11$  and very depleted  $\delta^{13}\text{C}_{\text{org}}$  and  
18 terrestrial plants characterized by  $\text{C/N} > 11$  and less depleted  ~~$\delta^{13}\text{C}_{\text{org}}$~~   $\delta^{13}\text{C}_{\text{org}}$  (Fig. 5).  
19  $\delta^{13}\text{C}_{\text{org}}$  and C/N are positively correlated (Fig. 3). TOC and TN are highly correlated (0.99,  
20 Figs. 3 and 6) as well.

21 In order to interpret the different bio and geo-chemical proxies within a climatic frame, a  
22 pairwise correlation was performed between the three climate variables and  $\delta^{13}\text{C}$ , C/N, TN  
23 and TOC (Fig. 6). Although there could be no causal relationship, SI and Tjan are well  
24 correlated together. They are both correlated negatively with  $\delta^{13}\text{C}$  and C/N and positively  
25 with TN and TOC (Fig. 6).

## 26 **5 Discussion**

27 The Holocene climate around the Mediterranean Sea was suitable for the expansion of human  
28 populations and their organization towards true civilizations (Kaniewski et al., 2012). The  
29 persistence and longevity of many Mediterranean populations may be linked to the relative  
30 suitability and also to an overall stability of the Holocene climate. However, climatic events  
31 have been recorded within the Holocene (e.g. Rohling and Pälike, 2005) and a causal  
32 relationship has been made between some abrupt climatic events and societal changes in the  
33 Mediterranean (Berger and Guilaine, 2009; Kaniewski et al., 2008).

1 In the present study, we have focused on the environmental and climate changes that occurred  
2 during the last 6 millennia in the northern part of the Moroccan Middle Atlas Mountains. We  
3 have evaluated the vegetation dynamics using the palynological content of a fossil sequence  
4 and analyzed its bio- and geo-chemical content to reconstruct the overall landscape changes.  
5 The reconstructed Tjan and Pann show a relatively low amplitude of change over the last  
6 6000 years (Fig. 3). Pann decreases progressively by ca. 100mm which is in line with the  
7 aridity trend that has been observed in other fossil records (Risacher and Fritz, 1992; Brooks,  
8 2006; Hastenrath, 1991; Anderson and Leng, 2004; Umbanhowar et al., 2006) and  
9 particularly in the Mediterranean area (Pons and Reille, 1988; Julià et al., 2001; Burjachs et  
10 al., 1997; Yll et al., 1997; Roberts et al., 2001; Valino et al., 2002, Jalut et al., 2009) and  
11 northern Africa (Ritchie, 1984; Ballouche et al., 1986; Lamb et al., 1989). At a more regional  
12 scale, reconstructed Pann is coherent with that obtained from Lake Tigalmamine (Cheddadi et  
13 al., 1998) which shows a decreasing trend over the last ca. 5000 cal BP. The arid trend  
14 observed after ca. 5ka cal BP is marked by a spread of Poaceae and a progressive replacement  
15 of pines by Atlas cedars which better stand the high seasonal contrast of precipitation at the  
16 altitude of Hachlaf ~~Lake~~lake. SI increases from 3 to 7 times over the last 6000 years (Fig. 3).  
17 A study of drought thresholds influencing the growth and photosynthesis was performed on  
18 different cedar stands and species (*C. atlantica*, *C. libani*, *C. brevifolia* & *C. deodora*) of  
19 different origins (Aussenac & Finkelstein, 1983). This study showed that among many  
20 conifers, cedar trees may keep a sustained photosynthesis activity even when drought is very  
21 high. Thus, a strong precipitation contrast between Ps and Pw (Fig. 3) may not affect the Atlas  
22 cedar overall growth as long as the total amount of rainfall is sufficient (higher than 600  
23 mm/year) and the winter temperature is low enough (below 6°C) for the vegetative cycle  
24 (Aussenac et al., 1981). The Mediterranean climate is known for its strong seasonal  
25 distribution of precipitation throughout the year. Summers are fairly dry and most of the  
26 annual precipitation occurs during the cold months (end of autumn and beginning of winter).  
27 Currently, 75% of the Moroccan territory with a grassy or wooded vegetation (thus excluding  
28 the desert) records between 500 and 800mm of annual rainfall with an SI between 5 and 8  
29 (Fig. 7). The whole range of SI in Morocco is between -1 in areas where Pann is less than  
30 100mm with a random distribution as for instance in the South of Morocco, and 15 in areas  
31 where the annual rainfall is quite high (over 800 mm) and occurs mainly in the winter season  
32 such as in the Rif mountains today (Fig. 7). SI is higher in mountainous areas. Nowadays, in  
33 the areas surrounding Hachlaf lake (located at ca. 1600m elevation) SI is around 5. Such SI  
34 has changed over the past thousand years as confirmed, at least between 6000 cal BP and

1 today, by the studied fossil archive (Fig. 3). The amplitude between Pw and Ps precipitation  
2 has increased 2 to 3 times towards the present (Fig. 3). Since Pann has a decreasing trend, the  
3 opposite increased seasonality is related to a significant reduction in the amount of rainfall  
4 during the months of June, July and August (Fig. 3). This strengthening of the contrast  
5 between Pw and Ps had a rather limited impact on the dominating taxa because they can  
6 withstand the summer drought and the overall amount of Pw remained sufficient for their  
7 persistence. However, a change in the amplitude of SI has probably favoured those species  
8 best adapted to the length of the dry season, as for instance evergreen oaks rather than  
9 deciduous. Pollen-based climate reconstructions from records collected in the Alboran Sea  
10 (Combourieu-Nebout et al., 2009) and Italy (Magny et al., 2013; Peyron et al., 2013) suggest  
11 a rather steady and low seasonal contrast between Pw and Ps (about two times) over the past  
12 6000 years cal BP. Such discrepancy between the reconstructed SI from Hachlaf and the  
13 marine record may potentially be related to the fact that marine records collect pollen grains  
14 from a much wider geographical source area than continental (mountainous) records which  
15 probably tends to smooth the local/regional changes. The reconstructed seasonality from the  
16 Italian records (Magny et al., 2013; Peyron et al., 2013) is buffered by the less abrupt  
17 precipitation seasonal contrast at the European temperate latitude than at the arid  
18 Mediterranean one.

19 SI was lower than 5 before 3750 cal BP despite an amount of precipitation between 600 and  
20 700 mm $\text{yr}^{-1}$  (Fig. 3). During that period, water probably persisted in the lake all throughout  
21 the year which allowed the presence of aquatic plants (Fig. 4) flowering during late spring and  
22 summer, and algae identified in the pollen data, through the low values of  $\delta^{13}\text{C}_{\text{org}}$  and the  
23 C/N ratio being greater than 11 (Figs. 3 and 5). The proportion of aquatic plants cannot be  
24 directly related to a high lake level and may not be used to state the lake level changes but  
25 only the presence of water in the site. The  $\delta^{13}\text{C}_{\text{org}}$  and C/N (Fig. 5) provide information  
26 concerning the origin of the organic matter (*in situ* production versus input from the  
27 catchment area) but not on the lake level changes. Thus, high  $\delta^{13}\text{C}_{\text{org}}$  and C/N ratios (Fig. 3)  
28 with low presence of aquatic plants (Fig. 4) may not be inconsistent in cases where there is a  
29 low terrestrial input (low Sand/Silt, Fig. 3) during a period when the lake level is high.

30 The relationship between  $\delta^{13}\text{C}_{\text{org}}$  and the C/N ratio indicates the occurrence of two main  
31 types of organic matter mainly originating from a C3 metabolism. Lacustrine algae can be  
32 considered as dominantly autochthonous; in the lower part of the record, the organic matter,  
33 with higher C/N ratios and less depleted  $\delta^{13}\text{C}_{\text{org}}$  corresponds to a terrestrial input. Indeed,  
34 Fresh organic matter from lake algae is known to be protein-rich and cellulose-poor with

1 molar C/N values commonly between 4 and 10, whereas vascular land plants, are protein-poor  
2 and cellulose-rich, creating organic matter usually with C/N ratios of 20 and greater (Meyers,  
3 1994, 2003). However, a C/N ratio > 11 may correspond to a mixture of both local and  
4 terrestrial organic matter (Fig. 5).

5 After 3750 cal BP, Atlas cedars noticeably spread around the site while the pine populations  
6 strongly regress. A series of fossil pollen records in the Middle Atlas show that Atlas cedar  
7 populations expanded after ca. 6 ka cal BP. The sustained expansion of Atlas cedar after ca.  
8 3750 cal BP around Hachlaf ~~Lake~~ expresses its late occurrence at higher altitude. Around  
9 lake Tigalmamine (Lamb et al., 1995), the Ras El Ma marsh (Nour El Bait et al., 2014) and  
10 the Ait Ichou marsh (Tabel et al., 2016) which are all located at about 100 to 200 meters  
11 altitude below Hachlaf lake (ca. 1700m asl), Atlas cedar occurs much earlier. The expansion  
12 of Atlas cedar around the lake is probably related to both an upslope spread and a south-north  
13 migration.

14 During this ecosystem transition we observe a major change in both Pann and Tjan. The  
15 increase of SI after 3750 cal BP is due to a combined increase of Pw and decrease of Ps (Fig.  
16 3). The expansion of cedar forests in the studied area may be related to their better adaptation  
17 to strong SI than pines at higher altitude.

18 Competition is another parameter that might be worth considering. After 3750 cal BP, the  
19 C/N ratio is below 11 and the  $\delta^{13}\text{C}$  remains below  $-26\text{‰}$  which suggest the important primary  
20 productivity of the lake associated with low input of land plant derived organic matter. Atlas  
21 cedar forests have a more important growth in both height and diameter than pines which  
22 leads to a higher biomass production. This is linked to the genetic model of growth that is  
23 very distinct between the two taxa (Kaushal et al., 1989). Thus, the expansion of Atlas cedar  
24 population around the site may explain the high input of OM into the lake.

25 Over the last six millennia, superimposed to the overall climate trend, we observe one  
26 relatively abrupt event between 5500 and 5000 cal BP during which Tjan declined by about  
27  $2^{\circ}\text{C}$  compared to its average over 6000 years. A climatic transition between 6 and 5 ka cal BP  
28 at the end of the Holocene thermal maximum has been globally identified (Steig, 1999;  
29 Mayewski et al., 2004; Wanner et al., 2008; Brooks, 2012). This transition has been recorded  
30 by a wide range of climate proxies (e.g. Kaufman et al., 2004; Jansen et al., 2009; Seppä et  
31 al., 2009; Bartlein et al., 2011) and has been related to different biosphere feedbacks and  
32 potentially to a decay of the remaining Laurentide ice sheet (Renssen et al., 2009). All proxies  
33 from the Hachlaf sequence as well as the reconstructed climate variables have recorded  
34 marked changes during that period of time. SI has the lowest value of the record and a

1 succession of abrupt changes are recorded in the C/N ratio, the grain size fractions, the  $\delta^{13}\text{C}$ ,  
2 TN, TOC and  $\text{CaCO}_3$  (Fig. 3). Carbonates, considered as a “paleo-thermometer” (Meyers,  
3 1994, 2003), also decrease abruptly around 5200 cal BP (Fig. 3). The latter may be linked to a  
4 low evaporation of the lake which may have been favored by low winter temperature around  
5 5200 cal BP. The fine grain size sediment also increased as a consequence of low seasonal  
6 precipitation contrast and/or a continuous sediment input to the lake. Such sustained input of  
7 clay and decreasing carbonate content suggest a higher lake level between 5500 and 5000 cal  
8 BP (Fig. 3). Thus, the Tjan and SI decrease may have contributed to the higher lake level or at  
9 least to the presence of water throughout the year (Fig. 3). At the same time, the sand to silt  
10 ratio is very low which confirms a low energy during the sedimentation process. The major  
11 change in the ecosystem composition around the lake is the rapid collapse of the pine forest  
12 which has inevitably released an important amount of terrestrial carbon (biomass) into the  
13 lake (positive peaks in  $\delta^{13}\text{C}$  and C/N, Fig. 3).

## 14 **6 Conclusions**

15 This study marks a new contribution to the knowledge of past climates and environmental  
16 history in North Africa mountainous areas. The range of climate change in the Middle Atlas,  
17 Morocco, was rather minor between 6000 cal BP and the present. Annual precipitation and  
18 January mean temperature have respectively varied within a range of ~~100 mm~~~~400mm~~ and 2 to  
19 3°C. However, they both show a trend towards a more arid and warmer climate as well as a  
20 higher rainfall seasonality. Pann became as contrasted as today after 3750 cal BP. The aridity  
21 trend observed in Hachlaf over the last 6000 years is consistent with other climate  
22 reconstructions available from other Mediterranean fossil records. Besides these overall  
23 climatic trends, we also observe an abrupt cold event between 5500 and 5000 cal BP which is  
24 well marked in all environmental proxies from our studied fossil record. The  $\delta^{13}\text{C}$  and C/N  
25 ratios, which are well correlated together, suggest an increase in the organic matter input from  
26 the catchment area. Concomitantly, the pollen record indicates a decline of the pine forest  
27 which may have contributed to the organic matter input into the lake too. The marked change  
28 in both the carbonates content and clay composition of the record were probably related to a  
29 perennial presence of water throughout the year. Synchronously, seasonality index and  
30 January mean temperature were the lowest of the record which has contributed to a reduction  
31 of the evaporation.



1 The increase in rainfall seasonality has probably favored the expansion of Atlas cedars around  
2 the studied site at the expense of the pine forest.

3  
4 **Acknowledgements.** This work was supported by the Volubilis Program (Programme mixte  
5 Interuniversitaire Franco-marocain, MA/11/251), 2011, by CNRS-CNRST Convention, 2009  
6 (ScVie07/09) and the French national program EC2CO-Biohefect, “Variabilité  
7 paléoclimatique et impact sur les forêts de conifères au Maroc depuis la période glaciaire”.  
8 MN received a postdoc grant from the EU Framework Programme Erasmus Mundus EU  
9 METALIC II (2013-2442 / 001-001 – EMA2) for completing this study at ISEM. We thank  
10 Claire Grandchamps for her revision of the English version of the manuscript. This is an  
11 ISEM contribution n° 2016-020.

## 1 | **References**

- 2 Alley, R. B., Mayewski, P. A., Sowers, T., Stuiver, M., Taylor, K. C., and Clark, P. U.:  
3 Holocene climatic instability: a prominent, widespread event 8200 yr ago, *Geology*, 25, 483–  
4 486, 1997.
- 5 Alley, R.B., Agustsdottir, A.M.: The 8k event: cause and consequences of a major Holocene  
6 abrupt climate change, *Quat. Sci. Rev.*, 24, 1123–1149, 2005.
- 7 Andersen, C., Koç, N., Jennings, A., and Andrews, J. T.: Non uniform response of the major  
8 surface currents in the Nordic Sea to insolation forcing: implications for the Holocene climate  
9 variability, *Paleoceanography*, 19, PA2003, doi:10.1029/2002PA000873, 2004.
- 10 Anderson, N. J. and Leng, M. J.: Increased aridity during the early Holocene in West  
11 Greenland inferred from stable isotopes in laminated-lake sediments, *Quat. Sci. Rev.*, 23,  
12 841–849, 2004.
- 13 Aussenac, G. and Finkelstein, D.: Influence de la sécheresse sur la croissance et la  
14 photosynthèse du cèdre, *Ann. Sci. for.*, 40, 67–77, 1983.
- 15 Aussenac, G., Granier, A., and Gross, P.: Etude de la croissance en hauteur du Cèdre (*Cedrus*  
16 *atlantica* Manetti) Utilisation d'un appareillage de mesure automatique, *Ann. Sci. for.*, 38,  
17 301–316, 1981.
- 18 Baali, A.: Genèse et évolution au Plio-Quaternaire de deux bassins intramontagneux en  
19 domaine carbonaté méditerranéen. Les bassins versants des dayets Afourgagh et Agoulmam  
20 (Moyen Atlas, Maroc), PhD thesis, University of Rabat, 326 pp., 1998.
- 21 Ballouche, A. : Paléoenvironnements de l'homme fossile holocène au Maroc. Apports de la  
22 palynologie, PhD thesis, University of bordeaux, 134 pp., 1986.
- 23 Bartlein, P.J., Harrison, S.P., Brewer, S., Connor, S., Davis, B.A.S., Gajewski, K., Guiot, J.,  
24 Harrison-Prentice, T.I., Henderson, A., Peyron, O., Prentice, I.C., Scholze, M., Seppä, H.,  
25 Shuman, B., Sugita, S., Thompson, R.S., Viau, a. E., Williams, J., Wu, H.: Pollen-based  
26 continental climate reconstructions at 6 and 21 ka: a global synthesis. *Clim. Dyn.*, 37, 775–  
27 802, 2011.

- 1 Berger, J. F. and Guilaine, J.: The 8200 cal BP abrupt environmental change and the  
2 Neolithic- transition: a Mediterranean perspective, *Quat. Int.*, 200, 31–49, 2009.
- 3 Bianchi, G. G. and McCave, N.: Holocene periodicity in North Atlantic climate and deep-  
4 ocean flow south of Iceland, *Nature*, 397, 515–517, 1999.
- 5 Blaauw, M. and Christen, J.A.: Flexible Paleoclimate Age-Depth Models Using an  
6 Autoregressive Gamma Process, *Bayesian Analysis*, 6, 457–474, 2011.
- 7 Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M., Showers, W., Hoffmann, S., Lotti-  
8 Bond, R., Hajdas, I., and Bonani, G.: Persistent solar influence on North Atlantic climate  
9 during the Holocene, *Science*, 294, 2130–2136, 2001. Brooks, N.: Beyond collapse: climate  
10 change and causality during the Middle Holocene Climatic Transition, 6400–5000 years  
11 before present, *Geogr. Tidsskr. Geogr.*, 112, 93–104, doi:10.1080/00167223.2012.741881,  
12 | 2012.
- 13 Brooks, N.: Cultural responses to aridity in the middle Holocene and increased social  
14 complexity, *Quat. Int.*, 151, 29–49, 2006.
- 15 Bryson, R.A.: Late Quaternary volcanic modulation of Milankovitch climate forcing. *Theor.*  
16 *Appl. Climatol.*, 39, 115–125, 1989.
- 17 Burjachs, F., Giralt, S., Roca, J.R., Seret, G., and Julia, R.: Palinologia holocenica y  
18 desertizacion en el Mediterraneo occidental. *El Paisaje Mediterraneo a Traves del Espacio y*  
19 *del Tiempo. Implicaciones en la Desertificacion* (eds J.J. Ibanez, B.L. Valero & C. Machado),  
20 379–394. Geoforma Editores, Logrono, Spain, 1997.
- 21 Carrion, J. S.: Patterns and processes of Late Quaternary environmental change in a montane  
22 region of Southwestern Europe, *Quat. Sci. Rev.*, 21, 2130–2136, 2002.
- 23 Cheddadi, R., [Lamb, H.F., Guiot, J., and van der Kaars, S.: Holocene climatic change in](#)  
24 [Morocco: a quantitative reconstruction from pollen data. 14, 883–890, 1998.](#)
- 25 [Cheddadi, R.](#) and Bar-Hen, A.: Spatial gradient of temperature and potential vegetation  
26 feedback across Europe during the late Quaternary, *Clim. Din.*, 32, 371–379, 2009.
- 27 Cheddadi, R., Bouaissa, O., Rhoujjati, A. and Dezileau, L.: Holocene Environmental changes  
28 in the Rif Mountains, Morocco. *Quat.*, 27, 15–25, 2016.

- 1 Cheddadi, R., Fady, B., François, L., Hajar, L., Suc, J. P., Huang, K., Demarteau, M.,  
2 Vendramin, G. G., and Ortu, E.: Putative glacial refugia of *Cedrus atlantica* from Quaternary  
3 pollen records and modern genetic diversity, *J. Biogeogr.*, 36, 1361–1371, 2009.
- 4 Chevalier, M., Cheddadi, R. and Chase, B. M.: CREST (Climate REconstruction SofTware):  
5 a probability density function (PDF)-based quantitative climate reconstruction method, *Clim.*  
6 *Past*, 10, 2081–2098, doi:10.5194/cp-10-2081-2014, 2014.
- 7 Chillasse, L. and Dakki, M.: Potentialités et statuts de conservation des zones humides du  
8 Moyen-Atlas (Maroc), avec référence aux influences de la sécheresse, *Sécheresse*, 15, 337–  
9 45, 2004.
- 10 Combourieu Nebout, N., Peyron, O., Dormoy, I., Desprat, S., Beaudouin, C., Kotthoff, U.,  
11 and Marret, F.: Rapid climatic variability in the west Mediterranean during the last 25 000  
12 years from high resolution pollen data, *Clim. Past*, 5, 503–521, doi:10.5194/cp-5-503-2009,  
13 2009.
- 14 Dansgaard, W., White, J.W.C., and Johnsen, S.J.: The abrupt termination of the Younger  
15 Dryas climatic event, *Nature*, 339, 532–534, 1989.
- 16 Davis, M.B.: On the theory of pollen analysis. *Am. J. Sci.*, 261, 899–912, 1963.
- 17 Debret, M., Bout-Roumazeilles, V., Grousset, F., Desmet, M., McManus, J. F., Massei, N.,  
18 Sebag, D., Petit, J.-R., Copard, Y., and Trentesaux, A.: The origin of the 1500-year climate  
19 cycles in Holocene North-Atlantic records, *Clim. Past*, 3, 569–575, doi:10.5194/cp-3-569-  
20 2007, 2007.
- 21 Dorale, J. A., Gonzalez, L. A., Reagan, M. K., Pickett, D. A., Murrell, M. T., and Baker, R.  
22 G.: A high resolution record of Holocene climate change in speleothem calcite from Cold  
23 Water Cave, northeast Iowa, *Science*, 258, 1626–1630, 1992.
- 24 Eddy, J.A.: The solar constant and surface temperature. *AIP Conf. Proc.*, La Jolla, CA, USA,  
25 9-11 March 1981, 82, 247, 1982.
- 26 Emmanuel, W.R., Shugart, H.H. and Stevenson, M.P.: Climate change and the broad-scale  
27 distribution of terrestrial ecosystem complexes. *Clim. Chang.*, 7, 29–43, 1985.

- 1 Fletcher, W., J., Sánchez Goñi, M., F., Allen, J. R. M., Cheddadi, R., Combourieu-Nebout,  
2 N. and Huntley, B.: Millennial-scale variability during the last glacial in vegetation records  
3 from Europe. *Quat. Sci. Rev.*, 29, 2839–2864, 2010.
- 4 Fletcher, W.J. and Sánchez Goñi, M.F.: Orbital- and sub-orbital-scale climate impacts on  
5 vegetation of the western Mediterranean basin over the last 48,000 yr, *Quat. Res.*, 70, 451–  
6 464, 2008.
- 7 Frigola, J., Moreno, A., Cacho, I., Canals, M., Sierro, F. J., Flores, J. A., Grimalt, O., Hodell,  
8 D., and Curtis, J. H.: Holocene climate variability in the western Mediterranean region from a  
9 deepwater sediment record, *Paleoceanography*, 22, PA2209, doi:10.1029/2006PA001307,  
10 2007.
- 11 [GBIF: Recommended practices for citation of the data published through the GBIF Network.](#)  
12 [Version 1.0 \(Authored by Vishwas Chavan\), Copenhagen: Global Biodiversity Information](#)  
13 [Facility, 12, ISBN: 87-92020-36-4,](#)  
14 [http://links.gbif.org/gbif\\_best\\_practice\\_data\\_citation\\_en\\_v,2012.](http://links.gbif.org/gbif_best_practice_data_citation_en_v,2012)
- 15 Geiss, C. E., Banerjee, S. K., Camill, P., and Umbanhowar, J. C. E.: Sediment-magnetic  
16 signature of land-use and drought as recorded in lake sediment from south-central Minnesota,  
17 USA, *Quat. Res.*, 62, 117–125, 2004.
- 18 Geiss, C. E., Umbanhowar, C. E. J., Camill, P., and Banerjee, S. K.: Sediment magnetic  
19 properties reveal Holocene climate change along the Minnesota prairie-forest ecotone,  
20 *Paleolimnol.*, 30, 151–166, 2003.
- 21 Harmand, C. and Moukadiri, A.: Synchronisme entre tectonique compressive et volcanisme  
22 alcalin: exemple de la province quaternaire du Moyen Atlas (Maroc), *B. Soc. Geol. Fr.*, 8,  
23 595–603, 1986.
- 24 Hastenrath, S.: *Climate Dynamics of the Tropics*, Kluwer Academic Publishers, 1383-8601,  
25 Springer Netherlands, 463–488 pp., 1991.
- 26 Hazan, N., Stein, M., Agnon, A., Marco, S., Nadel, D., Negendank, J. F. W., Schwab, M., and  
27 Neev, D.: The late Pleistocene–Holocene limnological history of Lake Kinneret (Sea of  
28 Galilee), Israel, *Quat. Res.*, 63, 60–77, 2005.

- 1 [HCEFLCD : Haut-Commissariat aux Eaux et Forêts et Lutte Contre la Désertification. Bilan](#)  
2 [annuel, Santé des Forêts au Maroc. Etudes d'aménagement concerté des forêts et des parcours](#)  
3 [collectifs de la province d'Ifrane. Composante III : études forestières. Rapports 9 et 10, 2004.](#)
- 4 Herbig, H. G.: Synsedimentary tectonics in the Northern Middle Atlas (Morocco) during the  
5 Late Cretaceous and Tertiary, in: *The Atlas System of Morocco*, edited by: Jacobshagen, V.,  
6 Springer-Verlag, Berlin, 321–337, 1988.
- 7 Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., and Jarvis, A.: Very high resolution  
8 interpolated climate surfaces for global land areas, *Int. J. Climatol.*, 25, 1965–1978, 2005.
- 9 Hinaje, S. and Ait Brahim, L.: Les Bassins Lacustres du Moyen Atlas (Maroc): un exemple  
10 d'Activité Tectonique Polyphasée Associée à des Structures d'Effondrement, *Com. Instituto*  
11 *Geológico e Mineiro*, 89, 283–294, 2002.
- 12 Hulten, E. and Fries, M.: *Atlas of North European vascular plants: north of the Tropic of*  
13 *Cancer I-III*. Koeltz Scientific Books, Königstein, DE, 1986.
- 14 ISO 11464: 1994. Qualité du sol-Prétraitement des échantillons pour analyses physico-  
15 chimiques. Norme révisée par ISO 11464 : 2006, 11, 2006.
- 16 Jalas, J. and Suominen, J.: (eds) *Atlas florae Europaeae. Distribution of vascular plants in*  
17 *Europe*. The Committee for Mapping the Flora of Europe and Societas Biologica Fennica  
18 Vanamo, Helsinki, vols 1–10, 1972, 1973, 1976, 1979, 1980, 1983, 1986, 1989, 1991, 1994.
- 19 Jalut, G., Dedoubat, J. J., Fontugne, M. and Otto, T. : Holocene circum-Mediterranean  
20 vegetation changes: Climate forcing and human impact. *Quat. Int.*, 200, 4–18, 2009.
- 21 Jansen, E., Andersson, C., Moros, M., Nisancioglu, K. H., Nyland, B. F. and Telford, R. J.:  
22 The Early to Mid-Holocene Thermal Optimum in the North Atlantic, in *Natural Climate*  
23 *Variability and Global Warming: A Holocene Perspective* (eds R. W. Battarbee and H. A.  
24 Binney), Wiley Blackwell, Oxford, UK. doi: 10.1002/9781444300932.ch5, 2008
- 25 Jiménez-Moreno, G., Rodríguez-Ramírez, A., Pérez-Asensio, J. N., Carrión, J. S., López-  
26 Sáez, J. A., Villarías-Robles, J. J. R., Celestino-Pérez, S., Cerrillo-Cuenca, E., León, A., and  
27 Contreras, C.: Impact of late-Holocene aridification trend, climate variability and geodynamic

1 control on the environment from a coastal area in SW Spain, *Holocene*, 25, 607–627,  
2 | doi:10.1177/0959683614565955, 2015.

3 Julià, R., Riera, S., and Wansard, G.: Advances in Mediterranean lacustrine studies and future  
4 prospects: the Southern European group of ELDP project (1999– 2001) contribution, *Terra*  
5 | *Nostra*, 3, 43–51, 2001.

6 Kaniewski, D., Paulissen, E., Van Campo, E., Al-Maqdissi, M., Bretschneider, J., and Van  
7 Lerberghe, K.: Middle East coastal ecosystem response to middle-to-late Holocene abrupt  
8 climate changes, *Proc. Natl. Acad. Sci. U.S.A.*, 16, 13941–13946,  
9 doi:10.1073/pnas.0803533105, 2008.

10 Kaniewski, D., Van Campo, E., and Weiss, H.: Drought is a recurring challenge in the Middle  
11 East, *Proc. Natl. Acad. Sci. U.S.A.*, 109, 3862–3867, 2012.

12 Kaufman, D. S., Ager, T. A., Anderson, N. J., Anderson, P. M., Andrews, J. T., Bartlein, P. J.,  
13 Brubaker, L. B., Coats, L.L., Cwynar, L. C., Duvall, M. L., Dyke, a. S., Edwards, M.E.,  
14 Eisner, W.R., Gajewski, K., Geirsdóttir, a., Hu, F.S., Jennings, A.E., Kaplan, M.R., Kerwin,  
15 M.W., Lozhkin, a. V., MacDonald, G.M., Miller, G.H., Mock, C.J., Oswald, W.W., Otto-  
16 Bliesner, B.L., Porinchu, D.F., Rühland, K., Smol, J.P., Steig, E.J., and Wolfe, B.B.:  
17 Holocene thermal maximum in the western Arctic (0–180°W), *Quat. Sci. Rev.*, 23, 529–560,  
18 doi:10.1016/j.quascirev.2003.09.007, 2004.

19 Kaushal, P., Guehl, J. M., and Aussenac, G.: Differential growth response to atmospheric  
20 carbon dioxide enrichment in seedlings of *Cedrus atlantica* and *Pinus nigra* ssp. *Laricio* var.  
21 *Corsicana*, *Can. J. For. Res.*, 19, 1351–1358, 1989.

22 Kelly, P.M. and Sear, C.B.: Climatic impact of explosive volcanic eruptions. *Nature*, 311,  
23 740–743, 1984.

24 Lamb, C.J., Lawton, M.A., Dron, M., and Dixont, R.A.: Signals and Transduction  
25 Mechanisms for Activation of Plant Defenses against Microbial Attack, *Cell*, 56, 215–224,  
26 1989.

27 Lamb, H. F. and Van der Kaars, S.: Vegetational response to Holocene climatic change:  
28 pollen and palaeolimnological data from the Middle Atlas, *Holocene*, 5, 400–408, 1995.

- 1 Lamb, H.F., Gasse, F., Benkaddour, A., El Hamouti, N., van der Kaars, S., Perkins, W.  
2 T., Pearce, N. J., and Roberts, C. N.: Relation between century-scale Holocene arid intervals  
3 in tropical and temperate zones, *Nature*, 373, 134–137, doi:10.1038/373134a0, 1995.
- 4 Lascaratos, A., Roether, W., Nittis, K., and Klein, B.: Recent changes in deep water formation  
5 and spreading in the eastern Mediterranean Sea: a review, *Prog. Oceanography*, 44, 5–36,  
6 1999.
- 7 Lecompte, M.: La végétation du moyen atlas central. Esquisse phyto-écologique et carte des  
8 séries de végétation au 1/200 000, *Rev. geogr. Maroc.*, 16, 1–31, 1969.
- 9 Lemcke, G. and Sturm, M.:  $^{18}\text{O}$  and trace element measurements as proxy for the  
10 reconstruction of climate changes at Lake Van (Turkey): preliminary results, in: Third  
11 Millenium BC Climate Change and Old World Collapse (Proceedings of the NATO  
12 Advanced Research Workshop on Third Millenium BC Abrupt Climate Change and Old  
13 World Social Collapse, held at Kemer, Turkey, 19–24 September 1994), edited by: Nüzhet  
14 Dalfes, H., Kukla, G., and Weiss, H., Springer, Berlin, Heidelberg, New York, 653–678,  
15 1997.
- 16 Magny, M., Combourieu Nebout, N., de Beaulieu, J. L., Bout-Roumazeilles, V., Colombaroli,  
17 D., Desprat, S., Francke, A., Joannin, S., Peyron, O., Revel, M., Sadori, L., Siani, G., Sicre,  
18 M. A., Samartin, S., Simonneau, A., Tinner, W., Vanni`ere, B., Wagner, B., Zanchetta, G.,  
19 Anselmetti, F., Brugiapaglia, E., Chapron, E., Debret, M., Desmet, M., Didier, J., Essallami,  
20 L., Galop, D., Gilli, A., Haas, J. N., Kallel, N., Millet, L., Stock, A., Turon, J. L., and Wirth,  
21 S.: North–south palaeohydrological contrasts in the central Mediterranean during the  
22 Holocene: tentative synthesis and working hypotheses, *Clim. Past.*, 9, 1901–1967,  
23 doi:10.5194/cpd-9-1901-2013, 2013.
- 24 Magny, M., De Beaulieu, J. L., Drescher-Schneider, R., Vanniere, B., Waltersimonnet, A. V.,  
25 Millet, L., Bossuet, G., and Peyron, O.: Climatic oscillations in central Italy during the Last  
26 Glacial–Holocene transition: the record from Lake Accesa, *J. Quat. Sci.*, 21, 311–320, 2006.
- 27 Manabe, S. and Stouffer, R. J.: Two stable equilibria of a coupled ocean-atmosphere model. *J.*  
28 *Clim.*, 1, 841–866, 1988.
- 29 Mann, M. E., Cane, M. A., Zebiak, S. E. and Clement, A.: Volcanic and solar forcing of the  
30 tropical Pacific over the past 1000 years. *J. Clim.*, 18, 447–456, 2005.



- 1 Märsche-Soulié, I., Benkaddour, A., Elkhiaï, N., Gemayel, P., and Ramdani, M.:  
2 Charophytes, indicateurs de paléo-bathymétrie du lac Tigalmamine (Moyen Atlas, Maroc),  
3 *Geobios*, 41, 435–444, 2008.
- 4 Martin, J.: Carte géomorphologique du Moyen Atlas central au 1/100.000, Notes & Mém.  
5 *Serv. géol. Maroc*, 258 bis, 445 pp., 1973.
- 6 Martin, J.: Le Moyen Atlas central étude géomorphologique, Notes et Mémoires du service  
7 Géologique N° 258 bis Rabat Maroc, 447 pp, 1981.
- 8 Mayewski, P. A., Rohling, E., Stager, C., Karlén, K., Maasch, K., Meeker, L. D., Meyerson,  
9 E., Gasse, F., Van kreveld, S., Holmgren, K., Lee-thorp, J., Rosqvist, G., Rack, F.,  
10 Staubwasser, M., Schneider, R., and Steig, E. J.: Holocene climate variability, *Quat. Res.*, 62,  
11 243–255, 2004.
- 12 McGregor, H. V, Dima, M., Fischer, H. W., and Mulitza, S.: Rapid 20th-century increase in  
13 coastal upwelling off northwest Africa. *Science*, 315, 637–9, 2007.
- 14 Meyers, P. A.: Applications of organic geochemistry to paleolimnological reconstructions: a  
15 summary of examples from the Laurentian Great Lakes, *Org. Geochem.*, 34, 261–289, 2003.
- 16 Meyers, P. A.: Preservation of elemental and isotopic source identification of sedimentary  
17 organic matter, *Chem. Geol.*, 144, 289–302, 1994.
- 18 Migowski, C., Stein, M., Prasad, S., Negendank, J. F. W., and Agnonc, A.: Holocene climate  
19 variability and cultural evolution in the Near East from the Dead Sea sedimentary record,  
20 *Quat. Res.*, 66, 421–431, 2006.
- 21 Milano, M., Ruelland, D., Fernandez, S., Dezetter, A., Fabre, J. Servat, E., Fritsch, J. M.,  
22 Ardoin–Bardin, S., and Thivet, G.: Current state of Mediterranean water resources and future  
23 trends under global changes, *Hydrolog. Sci. J.*, 58, 498–518, doi:  
24 10.1080/02626667.2013.774458, 2013.
- 25 Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B., and Kent, J.:  
26 Biodiversity hotspots for conservation priorities, *Nature*, 403, 853–858, 2000.
- 27 NF ISO 10693: Juin, 1995. Qualité du sol-Détermination de la teneur en carbonate - Méthode  
28 volumétrique, 7, 1995.

- 1 NF ISO 14235. Qualité du sol–Dosage du carbone organique par oxydation sulfochimique,  
2 11, 1998.
- 3 Nour El Bait, M., Rhoujjati, A., Eynaud, F., Benkaddour, A., Dezileau, L., Wainer, K.,  
4 Goslar, T., Khater, C., Tabel, J., and Cheddadi, R.: An 18,000 year pollen and sedimentary  
5 record from the cedar forests of the Middle Atlas, Morocco, *J. Quat. Sci.*, 29, 423–432, 2014.
- 6 [Peyron, O., Magny, M., Goring, S., Joannin, S., de Beaulieu, J.-L., Brugiapaglia, E., Sadori,](#)  
7 [L., Garfi, G., Kouli, K., Ioakim, C., and Combourieu-Nebout, N.: Contrasting patterns of](#)  
8 [climatic changes during the Holocene across the Italian Peninsula reconstructed from pollen](#)  
9 [data. \*Clim. Past\*, 9, 1233-1252, doi:10.5194/cp-9-1233-2013, 2013.](#)
- 10 Pons, A. and Reille, M.: The holocene- and upper pleistocene pollen record from Padul  
11 (Granada, Spain): A new study, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 66, 255–249–263,  
12 1988
- 13 Reille, M.: Analyse pollinique de sédiments postglaciaires dans le Moyen Atlas et le Haut  
14 Atlas marocains: premiers résultats, *Ecol. Mediterr.*, 2, 153–170, 1976.
- 15 Renssen, H., Seppä, H., Heiri, O., Roche, D.M., Goosse, H., and Fichefet, T.: The spatial and  
16 temporal complexity of the Holocene thermal maximum. *Nat. Geosci.*, 2, 411–414, 2009.
- 17 Rhoujjati, A., Ortu, E., Baali, A., Taïeb, M., and Cheddadi, R.: Environmental changes over  
18 the past 29,000 years in the Middle Atlas (Morocco): a record from Lake Ifrah, *J. Arid*  
19 *Environ.*, 74, 737–745, 2010.
- 20 Risacher, F. and Fritz, B.: Mise en évidence d’une phase climatique holocène extrêmement  
21 aride dans l’Altiplano central, par la présence de la polyhalite dans le salar de Uyuni  
22 (Bolivie), *Paleoclimatology, CR. Acad. Sci. Paris*, 314, 1371–1377, 1992.
- 23 Ritchie, M. : Analyses polliniques de sédiments holocènes supérieurs des Hauts-Plateaux du  
24 Maghreb oriental, *Pollen Spores, Paris*, 16, 489–496, 1984.
- 25 Roberts, N., Reed, J. M., Leng, M. J., Kuzucuoglu, C., Fontugne, M., Bertaux, J., Woldring,  
26 H., Bottemaj, S., Black S., Hunt, E., and Karabiyikoglu, M.: The tempo of Holocene climatic  
27 change in the eastern Mediterranean region: new high-resolution crater-lake sediment data  
28 from central Turkey, *Holocene*, 11, 721–736, 2001.

- 1 Roberts, N., Stevenson, T., Davis, B., Cheddadi, R., Brewer, S., and Rosen, A.: Holocene  
2 climate, environment and cultural change in the circum Mediterranean region, in: Past  
3 Climate Variability through Europe and Africa, edited by: Battarbee, R. W., Gasse, F., and  
4 Stickley, C. E., Kluwer Academic Press, Dordrecht, 343–362, 2004.
- 5 Rohde, K.: Latitudinal gradients in species diversity: The search for the primary cause, *Oikos*,  
6 65, 514–527, 1992.
- 7 Rohling, E. J. and Pälike, H.: Centennial-scale climate cooling with a sudden cold event  
8 around 8,200 years ago, *Nature*, 434, 975–979, doi:10.1038, 2005.
- 9 Rohling, E., Mayewski, P., Abu-Zied, R., Casford, J., and Hayes, A.: Holocene atmosphere-  
10 ocean interactions: records from Greenland and the Aegean Sea, *Clim. Dyn.*, 18, 587–593,  
11 2002.
- 12 Ruddiman, W. F.: Orbital insolation, ice volume, and greenhouse gases, *Quat. Sci. Rev.*, 22,  
13 1597–1629, 2003.
- 14 Sear, C.B., Kelly, P.M., Jones, P.D. and Goodess, C. M.: Global surface-temperature  
15 responses to major volcanic eruptions, *Nature*, 330, 365–367, 1987.
- 16 Seppä, H., Bjune, A.E., Telford, R.J., Birks, H.J.B., and Veski, S.: Last nine-thousand years  
17 of temperature variability in Northern Europe, *Clim. Past.*, 5, 523–535, 2009.
- 18 Steig, E.J.: Mid-Holocene climate change. *Science*, 286, 6–8, 1999.
- 19 Stuiver, M. and Reimer, P. J.: Extended  $^{14}\text{C}$  data base and revised calib 3.0  $^{14}\text{C}$  age calibration  
20 program, *Radiocarbon*, 35, 215–230, 1986.
- 21 Stuiver, M., Braziunas, T.F., Becker, B. and Kromer, B., Climatic, solar, oceanic and  
22 geomagnetic influences on late-glacial and Holocene atmosphere  $^{14}\text{C}/^{12}\text{C}$  change, *Quat. Res.*,  
23 35, 1–24, 1991.
- 24 Tabel, J., Khater, K., Rhoujjati, A., Dezileau, L., Bouimetarhan, I., Carré, C., Vidal, L.,  
25 Benkaddour, A., Nour El Bait, M., and Cheddadi, R. : Environmental changes over the past  
26 25,000 years in the southern Middle Atlas, Morocco, *J. Quat. Sci.*, in press, DOI:  
27 10.1002/jqs.2841, 2016.

- 1 Texier, J. P., Raynal, J. P., and Lefevre, D.: Nouvelles positions pour un cadre chronologiques  
2 raisonné du Quaternaire marocain, CR. Acad. Sc. Paris, 301, 183–188, 1985.
- 3 Umbanhowar, C. E. J., Camill, P., Geiss, C. E., and Teed, R.: Asymmetric vegetation  
4 responses to mid-Holocene aridity at the prairie-forest ecotone in south-central Minnesota,  
5 Quat. Res., 66, 53–66, 2006.
- 6 Valino, M.D., Rodríguez, A.V., Zapata, M.B.R., Garcia, M.J.G., and Gutiérrez, I.B.:  
7 Climatic changes since the Late-glacial/Holocene transition in La Mancha Plain (South-  
8 central Iberian Peninsula, Spain) and their incidence on Las Tablas de Daimiel marshlands,  
9 Quat. Int., 73–84, 2002.
- 10 Wanner, H., Beer, J., Bütikofer, J., Crowley, T. J., Cubasch, U., Flückiger, J., Goosse, H.,  
11 Grosjean, M., Joos, F., Kaplan, J. O., Küttel, M., Müller, S. A., Prentice, I. C., Solomina, O.,  
12 Stocker, T. F., Tarasov, P., Wagner, M., and Widmann, M.: Mid- to Late Holocene climate  
13 change: an overview, Quat. Sci. Rev., 27, 1791–1828, 2008.
- 14 Weninger B., Clare L., Rohling E. J., Bar-Yosef O., Böhner U., Budja M., Bundschuh M.,  
15 Feurdean A., Gebel H.-G., Jöris O., Linstädter J., Mayewski P., Mühlenbruch T., Reingruber  
16 A., Rollefson G., Schyle D., Thissen L., Todorova H., and Zielhofer C.: The Impact of Rapid  
17 Climate Change on prehistoric societies during the Holocene in the Eastern Mediterranean,  
18 Documenta Praehistorica, 36, 7–59, 2009.
- 19 Williams, J. T., Post, D. M., Cwyner, L. C., Lotter, A. F., and Levesque, A. J.: Rapid and  
20 widespread vegetation responses to past climate change in the North Atlantic region,  
21 Geology, 11, 971–974, 2002.
- 22 Witt, A. and Schumann, A. Y.: Holocene climate variability on millennial scales recorded in  
23 Greenland ice cores, Nonlinear Process. Geophys., 12, 345–352, 2005.
- 24 Yll, E.I., Perez-Obiol, R., Pantaleon-Cano, J., and Roure, J.M.: Palynological Evidence for  
25 Climatic Change and Human Activity during the Holocene on Minorca (Balearic Islands),  
26 Quat. Res., 48, 339–347, 1997.

1

Depth (cm)	Material dated	<sup>14</sup> C age yr BP	95,4 % (2σ) cal age ranges (BP)	Relative area under probability distribution	Median probability cal BP
60	Bulk	2535 ± 30	2494 – 2746	0,447	2624
120	Bulk	3220 ± 35	3371 – 3509	0,936	3436
170	Bulk	4390 ± 35	4859 – 5047	0,991	4949
240	Bulk	5200 ± 40	5897 – 6021	0,943	5958

2

3

Zones	Depth (cm)	Age (cal BP)	Pollen taxa data description		Plant species	
<u>Alisma</u> Zone I		<u>Alisma plantago-aquatica</u> 250–190	6227–5171	AP	27–60%	————— Mainly <i>Quercus</i> ——— and <i>Olea</i> . — Peak of <i>Pinus</i> (47%) — at 6100 cal BP — then decreasing. ————— Low percentages — of <i>Cedrus atlantica</i> with — initial spread — around 5800 cal BP.
		<u>Alnus</u>	<u>Alnus glutinosa</u>			
		<u>Berberis</u>	<u>Berberis hispanica</u>			
		<u>Brassica</u>	<u>Brassica</u>			
		<u>Campanula</u>	<u>Campanula afra</u>			
		<u>Caryophyllaceae</u>	<u>Caryophyllaceae</u>			
		<u>Centaurea</u>	<u>Centaurea cyanus</u>			
		<u>Chenopodiaceae</u>		NAP	39–72%	<u>Chenopodiaceae</u> — Herbs dominated by Poaceae (11–48%), <i>Illecebrum</i> (3–19%), Apiaceae (2–5%), Brassicaceae (1–5%), Asteraceae (0–5%), Cichorioideae (1–6%), Chenopodiaceae (0.5–2%) and Cereals (0–1%).
		<u>Asteroidaeae</u>	<u>Compositae Subfam. Asteroidaeae</u>	DT	18–26	————— Rapid fluctuations
		<u>Cichorioideae</u>	<u>Compositae Subfam. Cichorioideae</u>			

<u>Corylus</u>	<u>Corylus avellana</u>				
<u>Cupressaceae</u>	<u>Cupressaceae</u>				
<u>Ephedra</u>	<u>Ephedra fragilis</u>				
<u>Euphorbia</u>	<u>Euphorbia characias</u>				
<u>Geranium</u>	<u>Geranium macrorrhizum</u>				
<u>Helianthemum</u>	<u>Helianthemum canariense</u>				
<u>Ilex</u>	<u>Ilex aquifolium</u>				
<u>Juglans</u>	<u>Juglans regia</u>				
<u>Myriophyllum</u>	<u>Myriophyllum aquaticum</u>				
<u>Plantago</u>	<u>Plantago lanceolata</u>				
<u>Polygonaceae</u>	<u>Polygonaceae</u>				
<u>Ranunculaceae</u>	<u>Ranunculaceae</u>				
<u>Salix</u>	<u>Salix pedicellata</u>				
<u>Saxifraga</u>	<u>Saxifraga</u>				
<u>Taxus</u>	<u>Taxus baccata</u>				
<u>Urtica</u>	<u>Urtica dioica</u>				
<u>Papaveraceae</u>	<u>Papaveraceae</u>				
<u>Pinus</u>	<u>Pinus halepensis</u>				
<u>Olea</u>	<u>Olea europaea</u>				
<u>Paronychia</u>	<u>Paronychia argentea</u>				
<u>Erica</u>	<u>Erica arborea</u>				
<u>Quercus</u>	<u>Quercus ilex</u>				
<u>Cedrus</u> Zone H	190—111	5171—3651	AP	28—56 %	<p>————— <i>Pinus</i> dominates — the pollen record but regresses at 5500 cal BP (from 44 to less than 2 %).</p> <p>————— <i>Cedrus atlantica</i> continues — to expand (0—5 %).</p> <p>— We observe a peak of Rosaceae (6 %).</p>
<u>Artemisia</u>	<u>Artemisia herba-alba</u>				

1  
2

Zones	Depth (cm)	Age (cal BP)	Pollen data description		
Zone I	250 – 190	6227 – 5171	AP	27 – 60%	- Mainly <i>Quercus</i> and <i>Olea</i> . - Peak of <i>Pinus</i> (47%) at 6100 cal BP then decreasing. - Low percentages of <i>Cedrus atlantica</i> with initial spread around 5800 cal BP.
			NAP	39 – 72 %	- Herbs dominated by Poaceae (11 – 48 %),

					<i>Illecebrum</i> (3 – 19 %), <i>Apiaceae</i> (2 – 5 %), <i>Brassicaceae</i> (1 – 5 %), <i>Asteraceae</i> (0 – 5 %), <i>Cichorioideae</i> (1 – 6 %), <i>Chenopodiaceae</i> (0.5 – 2 %) and <i>Cereals</i> (0 – 1 %).
			DT	18 – 26	- Rapid fluctuations
Zone II	190 – 111	5171 – 3651	AP	28 – 56 %	- <i>Pinus</i> dominates the pollen record but regresses at 5500 cal BP (from 44 to less than 2 %). - <i>Cedrus atlantica</i> continues to expand (0 – 5 %). - We observe a peak of <i>Rosaceae</i> (6 %).
			NAP	43 – 72 %	- Herbs are dominated by <i>Poaceae</i> , <i>Illecebrum</i> and <i>Asteraceae</i> which reach their maximum (53, 20 and 10 %, respectively). - <i>Cereals</i> disappear.
			DT	19 – 29	- Moderate to high with two peaks.
Zone III	111 – 60	3651 – 2351	AP	23 – 58 %	- Strong expansion of <i>Cedrus atlantica</i> and <i>Quercus</i> . - An abrupt decline of <i>Cedrus atlantica</i> around 2653 cal BP is recorded. - <i>Pinus</i> regresses as well but shows a peak of 20% at 3300 cal BP.
			NAP	41 – 76 %	- Herbs dominate the pollen record. - Sharp decline in <i>Poaceae</i> , <i>Asteraceae</i> , <i>Chenopodiaceae</i> and <i>Caryophyllaceae</i> at 5600 cal BP. - Appearance of <i>Cereals</i> around 2653 cal BP.
			DT	20 – 31	- High.
Zone IV	60 – 5	2351 – 173	AP	23 – 43 %	- Abundance of <i>Cedrus atlantica</i> , <i>Quercus</i> , <i>Olea</i> and <i>Rosaceae</i> . - Sharp decline and disappearance of <i>Pinus</i> .
			NAP	56 – 76 %	- Herbs continue to dominate the pollen record with <i>Poaceae</i> , <i>Cereals</i> , <i>Brassicaceae</i> , <i>Chenopodiaceae</i> and <i>Caryophyllaceae</i> which are most abundant. - <i>Asteraceae</i> , <i>Illecebrum</i> and <i>Apiaceae</i> decline. - <i>Centaurea</i> and <i>Cichorioideae</i> disappear.
			DT	21 – 32	- High.

1  
2

1 **Table 1.** Radiocarbon ages for the Hach-I core. Calibrations were performed using Calib 7.1  
2 (Stuiver and Reimer, 1986).

3 **Table 2.** Pollen taxa assigned to the most probable plant species in our plant database.

4 **Table 3.** Pollen zones identified in the fossil record using a constrained cluster analysis. AP:  
5 arboreal pollen taxa, NAP: non-arboreal pollen taxa, DT: taxa diversity.

6 **Figure 1.** The study area. (a) Geographical location of the tabular and pleated Middle Atlas  
7 (MA); (b) sketch of the geological and geomorphological characteristics of the Hachlaf area  
8 (from Martin, 1973); (c) phytoecological map showing the main ecosystems and the location  
9 of the Hachlaf Lake (Dayet Hachlaf) within an oak forest (from Lecompte, 1969).

10 **Figure 2.** (a) Lithology of the core Hach-I and radiocarbon  $^{14}\text{C}$  dates; (b) age/depth model  
11 from BACON software (Blaauw and Christen, 2011).

12 **Figure 3.** Diagram showing the sediment fractions (clay, silt and Sand/Silt ratio), the pollen  
13 percentages of *Cedrus atlantica* and *Pinus*, geochemical elements ( $\delta^{13}\text{C}$  [ $\delta^{13}\text{C}\%$ ],  
14 nitrogen to carbon ratio [C/N], total organic carbon [TOC], Total Nitrogen [NT]) and  
15 carbonates concentrations ( $\text{CaCO}_3$ ), January mean temperature (Tjan), Annual precipitation  
16 (Pann), winter and summer precipitations (Pw and Ps) and precipitation seasonality index  
17 (SI). The red rectangles are pointing the values of present-day Tjan, Pann, Pw and Ps  
18 (HCEFLCD, 2004). The red line shows the limit 3.7 ka cal BP.

19 **Figure 4.** Diagram showing the percentages of the main pollen taxa identified in the Hach-I  
20 core. Cyperaceae and Juncaceae are included within aquatic taxa. The dashed black curves  
21 shows an exaggeration ( $\times 7$ ) of the percentages of some taxa. On the right, pollen zones with their  
22 boundaries are set up using a constrained hierarchical clustering (R Development Core Team, 2013).  
23 The taxonomic diversity is computed using a rarefaction analysis. The red line shows the limit 3.7 ka  
24 cal BP.

25 **Figure 5.**  $\delta^{13}\text{C}$  and C/N bi-plot (from Meyers, 1994).

26 **Figure 6.** Pairwise correlation between the three climatic variables (Tjan, Pann and SI) and  
27 the chemical elements.

28 **Figure 7.** Modern SI (upper panel) and Pann (middle panel) from the gridded WorldClim  
29 dataset (Hijmans et al., 2005) over Morocco. The lower panel shows the distribution of Pann  
30 vs. SI: the lowest index occurs in southern Morocco where Pann is lower than  $200 \text{ mm.y}^{-1}$   
31 and the highest index occurs in the high altitudinal areas (Middle Atlas and Rif mountains).