

**The biogeophysical climatic impacts of anthropogenic land use**

M. C. Smith et al.

# The biogeophysical climatic impacts of anthropogenic land use change during the Holocene

M. C. Smith<sup>1</sup>, J. S. Singarayer<sup>1</sup>, P. J. Valdes<sup>2</sup>, J. O. Kaplan<sup>3</sup>, and N. P. Branch<sup>4</sup>

<sup>1</sup>Centre for Past Climate Change and Department of Meteorology, University of Reading, Reading, UK

<sup>2</sup>School of Geographical Sciences, University of Bristol, Bristol, UK

<sup>3</sup>Institute of Earth Surface Dynamics, University of Lausanne, Lausanne, Switzerland

<sup>4</sup>School of Archaeology, Geography and Environmental Science, University of Reading, Reading, UK

Received: 29 August 2015 – Accepted: 31 August 2015 – Published: 1 October 2015

Correspondence to: M. C. Smith (mcsmith@pgr.reading.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

The first agricultural societies were established around 10 ka BP and had spread across much of Europe and southern Asia by 5.5 ka BP with resultant anthropogenic deforestation for crop and pasture land. Various studies have attempted to assess the biogeochemical implications for Holocene climate in terms of increased carbon dioxide and methane emissions. However, less work has been done to examine the biogeophysical impacts of this early land use change. In this study, global climate model simulations with HadCM3 were used to examine the biogeophysical effects of Holocene land cover change on climate, both globally and regionally, from the early Holocene (8 ka BP) to the early industrial era (1850 CE).

Two experiments were performed with alternative descriptions of past vegetation: (i) potential natural vegetation simulated by TRIFFID but no land-use changes, and (ii) where the anthropogenic land use model, KK10 (Kaplan et al., 2009, 2011) has been used to set the HadCM3 crop regions. Snapshot simulations have been run at 1000 year intervals to examine when the first signature of anthropogenic climate change can be detected both regionally, in the areas of land use change, and globally. Results indicate that in regions of early land disturbance such as Europe and S.E. Asia detectable temperature changes, outside the normal range of variability, are encountered in the model as early as 7 ka BP in the June/July/August (JJA) season and throughout the entire annual cycle by 2–3 ka BP. Areas outside the regions of land disturbance are also affected, with virtually the whole globe experiencing significant temperature changes (predominantly cooling) by the early industrial period. Large-scale precipitation features such as the Indian monsoon, the intertropical convergence zone (ITCZ), and the North Atlantic storm track are also impacted by local land use and remote teleconnections. We investigated how advection by surface winds, mean sea level pressure (MSLP) anomalies, and tropospheric stationary wave train disturbances in the mid- to high-latitudes led to remote teleconnections.

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## 1 Introduction

The first agricultural societies were established in the Near East around 10 kaBP and had spread across most of Europe by 5.7 kaBP (Zohary et al., 2012) and to India by 9 kaBP (Tauger, 2013). In China domestication of millet and rice began about 8.5 kaBP initially spreading more slowly than in Europe but reaching S.E. Asia by 5.5 kaBP (Roberts, 2013; Tauger, 2013). Agriculture was also independently developed in Mesoamerica with maize possibly being cultivated as far back as 9 kaBP (Piperno et al., 2009) but, as in China, it spread slowly to other areas.

The most important anthropogenic alteration of the natural environment was the clearing of forests to establish cropland and pasture, and the exploitation of forests for fuel and construction materials (Darby, 1956). This long history of anthropogenic land use change (ALCC) has implications for regional hydrology and climate, and possibly for global climate. Deforestation results in both biogeochemical and biogeophysical changes. The biogeochemical changes tend to increase temperature by the emission of greenhouse gases such as CO<sub>2</sub> and CH<sub>4</sub> (CH<sub>4</sub> emissions are influenced not just directly by deforestation but by irrigation in rice agriculture and by emissions from livestock and humans). The impacts of biogeophysical changes are many and varied, being dependent on the local climate, soil, and the natural vegetation that is being replaced, e.g. if natural savannah or grassland is replaced by crops the impact will not be as great as if woodland is replaced.

There are several mechanisms by which biogeophysical changes due to deforestation can affect regional climate. A combination of reduction in aerodynamic roughness, in the root extraction of moisture and in the capture of precipitation on the canopy leads to reduced evaporation and thus decreases the fluxes of moisture and latent heat from the surface to the atmosphere. These changes work to increase the local surface temperature (Lean and Rowntree, 1993). Conversely, the increase in surface albedo due to deforestation acts to decrease surface temperature by increasing the reflection of shortwave radiation. This is particularly true at high latitudes where ly-

CPD

11, 4601–4641, 2015

### The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ing snow is a factor for some of the year and the snow covered ground is no longer masked by the canopy of the forest. Generally, in mid to high latitudes the albedo increase is considered to be the dominant effect; leading to a net cooling of the regional surface temperature, whereas in the moist tropics the evaporation is more important and, therefore, a localised overall warming may result (Betts et al., 2006).

The extent and timing of the early anthropogenic land surface changes is the subject of much debate, as is their role in changing Holocene climate. Ruddiman (2003) proposed the idea that anthropogenic impacts on greenhouse gases, and consequently climate change, began thousands of years ago as a consequence of early agriculture and have been increasing in amplitude ever since, which he termed “the early anthropogenic hypothesis”. The idea has been hotly debated in the literature (e.g. Broecker and Stocker, 2006; Joos et al., 2004; Singarayer et al., 2011; Mitchell et al., 2013; Kaplan et al., 2011). Whilst the early anthropogenic hypothesis may likely not account entirely for the pre-industrial rises in CO<sub>2</sub> and CH<sub>4</sub> there is no doubt that land use changes do have climatic impact on both regional and global scales. The real debate is the scale of these effects of early agriculture.

Whilst paleoecological and archaeological evidence of anthropogenic land use changes exists, there are not enough sites to comprehensively determine continental scale impacts of deforestation (Kaplan, 2009). Therefore, in order to better estimate impacts of anthropogenic land use, several databases of land use change have been developed. Examples of these include the HYDE 3.1 (History Database of the Global Environment, Goldewijk et al., 2011), KK10 (Kaplan et al., 2009 and 2011) and Pongratz et al. (2008) models. Although the methodologies differ in the details, the basic premise of these models is that from an estimated database of historical population trends, anthropogenic deforestation is calculated based on population density and the suitability of land for crops or pasture.

To quantify the impact of ALCC on climate, datasets of past ALCC can be used in conjunction with climate models. Several studies have estimated the influence of pre-industrial ALCC on global climate (He et al., 2014; Kutzbach, 2011; Pongratz et al.,

CPD

11, 4601–4641, 2015

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2010). Globally, the biogeophysical effects of anthropogenic land use change have been estimated to cause a slight cooling that is offset by the biogeochemical warming, giving a net global warming (He et al., 2014; Pongratz et al., 2010). At the local to regional scale, in the most intensively altered landscapes of Europe, Asia, and North America, the biogeophysical effects can be comparable with the biogeochemical (He et al., 2014; Pongratz et al., 2010). In addition, Strandberg et al. (2014) used a regional climate model to evaluate the climatic effect of anthropogenic deforestation in Europe at 6 and 0.2 kaBP with both the HYDE 3.1 and KK10 ALCC scenarios. For the KK10 scenario at 6 kaBP small but significant temperature differences were found in summer and, at 0.2 kaBP, changes up to  $\pm 1^\circ\text{C}$  were found over widespread areas in both summer and winter. Other authors (e.g. Oglesby et al., 2010; Cook et al., 2012) have modelled a decrease in precipitation in response to deforestation in Mesoamerica.

These existing studies are, however, limited in either temporal or spatial extent and do not address the question of when anthropogenically induced climate change first occurs. In this study global climate model simulations are used to provide a comprehensive evaluation of the influence that the biogeophysical effects of regional human-induced land cover change have had on the climate both globally and regionally throughout much of the Holocene. As described in detail in Sect. 2, the period under consideration is from 8 kaBP to pre-industrial (1850) and snapshot simulations with HadCM3 were run at 1000 year intervals. The results are highlighted in Sect. 3. For evaluation purposes palaeoclimate reconstructions from Bartlein et al. (2011) and Marcott et al. (2013) have been compared with the results from the model runs (Sect. 4). The implications are discussed in Sect. 5.

## 2 Methodology

### 2.1 Model description

The climate simulations in this study were performed with the UK Hadley Centre coupled global climate model, HadCM3 (Gordon et al., 2000; Pope et al., 2000) with the Met Office Surface Exchange Scheme (MOSES2.1) (Essery, 2003) and TRIFFID (Top-down Representation of Interactive Foliage and Flora Including Dynamics) (Cox, 2001) dynamic vegetation.

HadCM3, is a coupled atmospheric, ocean and sea ice model. The atmospheric component has a horizontal resolution of 2.5° latitude and 3.75° longitude with 19 unequally spaced levels in the vertical and a 30 min time step. It has an Eulerian advection scheme and includes effects of CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub>, CFC11 and CFC12. The spatial resolution over the ocean is 1.25° by 1.25° with 20 unequally spaced layers extending to a depth of 5200 m. It also includes a 1.25° by 1.25° resolution model for the formation of sea-ice with simple dynamics whereby the sea-ice drifts on the ocean currents (Cattle and Crossley, 1995).

TRIFFID is coupled to the GCM (General Circulation Model) via MOSES every 10 days of the model run. Within TRIFFID nine surface types are specified: 5 plant functional types (PFTs) and 4 non-vegetation types.

HadCM3 was widely used in both the third and fourth assessment reports of the Intergovernmental Panel on Climate Change (IPCC, 2001, 2007) and still performs well in a number of tests relative to other global GCMs (Covey et al., 2003; IPCC, 2007). For the fifth IPCC assessment it has been superseded by HadGEM2 (Collins et al., 2011), but being relatively computationally efficient HadCM3 can be the better choice for some palaeoclimate modelling applications as it allows more and/or longer runs to be conducted than would be possible with a higher-resolution model.

CPD

11, 4601–4641, 2015

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2 Project-specific model configuration

The version of HadCM3 used does not include interactive ice, carbon cycle, or methane and thus must be forced with prescribed changes in orbit, greenhouse gases and ice-sheet evolution. Orbital parameters are taken from Berger and Loutre (1991), atmospheric concentrations of gases are determined from ice cores (CO<sub>2</sub> from Vostok, Petit et al., 1999; Loulergue et al., 2008, and CH<sub>4</sub>, and N<sub>2</sub>O from EPICA, Spahni et al., 2005) and the ice-sheet evolution is estimated using the ICE5G model of Peltier (2004). For further details of these natural forcings of the climate model readers are referred to Singarayer et al. (2011).

To prescribe Holocene ALCC the KK10 dataset of Kaplan et al. (2009 and 2011) was used. The original dataset is on a 5' spatial resolution and has modelled crop and pasture land use for every year from 8 kaBP to present. For this study the data at 1000 year intervals were taken (8, 7 kaBP, etc.) for both crop and pasture combined and upscaled to the spatial resolution of HadCM3 to formulate a time series of cropland masks (Fig. 1). Within TRIFFID the global crop area is designated by a cropland mask, which can only be occupied by agricultural-type vegetation (i.e. C<sub>3</sub> and C<sub>4</sub> grasses) or bare soil (Betts et al., 2007). Hence, the actual cropland is equivalent to the mask area, less inland water, urban and ice tiles, and less the area covered by non-grassland vegetation or bare soil. The crop mask area is not dynamically updated by climate data. The cropland area incorporates the natural C<sub>3</sub>/C<sub>4</sub> grass fractional areas before converting tree fractions.

For each simulation, the boundary condition forcings (orbit, greenhouse gases and ice sheets) were specified, and in all simulations the initial conditions were the same, based on a spun-up early industrial simulation. Simulations were run for 1000 years. By the final 500 years of the simulation the climate system has adjusted to a new surface equilibrium and thus these final 500 years were averaged to result in the mean altered climatic conditions. The relatively long averaging period increases the signal-to-noise ratio between the modified and control climates, and thus distinguishes differences that

### The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



are statistically significant, but which can be hidden by decadal/multidecadal variability in shorter averaging periods. This is especially important for assessing the impact of agriculture in the earlier time slices of the Holocene when the land use change is small and localised.

## 3 Results

### 3.1 Surface air temperature

#### 3.1.1 Local impacts of land use

In the regions where ALCC was significant, surface air temperature changes can be seen in all the time slice simulations (Figs. 2 and 3) and for all time slices except 8 ka BP (not shown) the temperature anomalies in most regions are outside the normal range of variability, which is considered to be within 2 standard deviations of the mean. The anomalies are more pronounced in the JJA season (Fig. 2) than DJF (Fig. 3). This is due to a combination of the land imbalance between the Northern and Southern Hemispheres, the lack of land surface changes in the extra tropical Southern Hemisphere and the enhanced effect of land surface changes during the season of greatest solar insolation and plant growth (which is JJA in the Northern Hemisphere).

The direct temperature response to ALCC varies with the degree of latitude but the relationship is not straightforward as it depends on local climate, soil, and the natural vegetation. In the extratropics, where the albedo effect is generally dominant, there is a trend towards increasing (negative) anomalies with an increase in disturbance fraction (Fig. 4a for P.I.). At 7 kaBP the range of extra tropical temperature anomalies within the areas of land disturbance is  $+0.1/-1.2^{\circ}\text{C}$  (JJA) and  $+0.6/-0.5^{\circ}\text{C}$  (DJF), by 4 kaBP it has increased to  $+0.4/-2^{\circ}\text{C}$  (JJA) and  $+0.9/-1^{\circ}\text{C}$  (DJF) and by the pre-industrial period (PI; 1850) it had reached  $+0.7/-4^{\circ}\text{C}$  (JJA) and  $+0.3/-2^{\circ}\text{C}$  (DJF). Although there are some positive temperature anomalies, the vast majority of grid points

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





show a negative temperature trend. Regions with the highest ALCC intensity show the largest negative temperature anomalies, in particular Europe and E. Asia/China, where the agricultural land use occurs earliest and has the highest concentration of land conversion (Figs. 3 and 4).

5 The tropical response shows less of a trend because the impact of reduced evaporation is more significant and thus there are conflicting signals between the cooling effect of increased albedo and the warming effect of reduced evaporation. In some tropical areas ALCC leads to net cooling while in other areas net warming is simulated (Figs. 2, 3, and 4a), partly dependent on the availability of moisture at the surface and partly on cloud cover changes (not shown). The main areas that show a warm anomaly response are Southern Africa and India in JJA from 5 kaBP and in DJF the area bordering the Bay of Bengal where E. India is warmer from 6 kaBP extending to the east coast of the Bay of Bengal by 2 kaBP. The Indian JJA warming is enhanced by cloud feedbacks; a decrease in monsoon circulation leads to decreased cloudiness, thus increasing the shortwave radiation reaching the surface and warming the lower atmosphere. In contrast, tropical South America generally shows a net cooling in response to ALCC. Around the mid to late Holocene the tropical temperature anomaly range within the areas of land disturbance is  $\pm 0.5^\circ\text{C}$  and by the early industrial era (1850 CE) it had reached  $\pm 1^\circ\text{C}$  (Fig. 4).

20 Analysis of the standard deviation of both the KK10 and Control simulations indicated no significant changes in the amplitude of interannual variability of surface temperature or precipitation (using the *F* test statistic).

### 3.1.2 Remote impacts of land use

25 In addition to the local temperature changes described above, cooling can also be observed in regions remote from the areas of major ALCC, particularly in the Northern Hemisphere. The most intense cooling is always in the regions of ALCC but even as early as 7 kaBP in the JJA season our model simulations show a band of cooling that stretches across much of the extra-tropical Northern Hemisphere and the North

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Atlantic (Fig. 2). This cooling starts influencing the northern Pacific regions by 5 kaBP and by the early industrial era the surface air temperature over most of the world's land masses and much of the ocean is cooler due to the effects of ALCC in remote areas.

In the DJF season in the Northern Hemisphere ALCC leads to cooling both locally and regionally starting at 7 kaBP (Fig. 3). The model simulations also show cooling in the Arctic and warming in Siberia. Cooling remote from the areas of major ALCC becomes more extensive by 3 ka BP and most landmasses of the Northern Hemisphere are cooler than the control simulation by 2 kaBP. The Siberian warm anomaly has ceased by 3 kaBP but it remains less affected by the cooling than other regions. In the Southern Hemisphere, cooling remains more localised until 2 ka BP by which time the majority of the land surface is cooler than the control.

There is an increased temperature anomaly response for the same level of disturbance fraction in the later timeslices (Fig. 4b) implying that the responses to the land use changes are not just due to the local effects. Some possible mechanisms for these remote impacts are large-scale circulation changes (such as stationary waves in the upper troposphere at mid-high latitudes and monsoonal circulation changes), near-surface advection, and the amplifying factors of snow cover. These mechanisms will be discussed in more detail in Sect. 3.2 for atmospheric dynamics and Sect. 3.3.2 for snow cover changes. Changes to the natural vegetation cover (outside the regions of land use) due to the climatic impacts of land use were also investigated as a potential mechanism of further feedbacks but were not found to be significant.

## 3.2 Atmospheric dynamics

### 3.2.1 Surface advection

Some of the cooling in regions remote from the areas of land disturbance is due to advection by low-level winds. In regions with a prevailing wind direction an advection pattern can be clearly seen. Figure 5 shows advection of cold air from the areas of land use change in East Asia to the east across the West Pacific and from the region

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of land disturbance in Mexico to the west across the East Pacific, this also affects the sea surface temperatures (SSTs) in these regions of the Pacific (not shown). In other areas where the surface wind direction is more variable the effect is more difficult to detect but probably does contribute to the spread of the cold anomaly outward from the region of land disturbance.

### 3.2.2 Mean sea level pressure

Changes to Mean Sea Level Pressure (MSLP) can also have an effect on the climate system. The colder surface air temperature in the region of disturbance means reduced ascent in those regions and thus higher MSLP. This can be seen quite clearly in China from 6 ka and in Europe by 4 kaBP (Fig. 5 for PI JJA), although the DJF situation in Europe is less coherent probably due to more remote influences such as the North Atlantic storm track. These MSLP changes could play a part in steering weather systems and thus influencing the climate in remote regions. For example, the JJA MSLP anomaly pattern (Fig. 6a and b) over the North and Central Atlantic is indicative of a negative phase of the North Atlantic Oscillation (NAO) with above-normal pressure over the North Atlantic and below-normal pressure over the central Atlantic. This pattern is apparent from 5 kaBP. The NAO alters the intensity and location of the North Atlantic jet stream and storm track and thus the patterns of heat and moisture transport (Hurrell, 1995). A negative NAO would contribute to a tendency to wetter summers in all but the southernmost regions of Europe (Folland et al., 2009) and this was seen in the results described in Sect. 3.3.1. The DJF NAO shows a trend towards a positive NAO which could result in drier winters over the Mediterranean region ([http://www.cpc.ncep.noaa.gov/data/teledoc/nao\\_pmap.shtml](http://www.cpc.ncep.noaa.gov/data/teledoc/nao_pmap.shtml)) but the pattern is not as consistent as JJA.

CPD

11, 4601–4641, 2015

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.2.3 Upper tropospheric dynamics

Cooler surface air temperature means that the density of the air is greater and, therefore, the geopotential height in cooler regions will be lower (See Fig. 6c and d). In the JJA season from 7 kaBP there is a reduction of the 500 hPa geopotential height over the extra-tropical Northern Hemisphere in a pattern similar to the temperature pattern but more extensive, completely encircling the globe. From 3 kaBP onwards the height reduction expands southwards so the geopotential height is lowered almost everywhere by pre-industrial times. The most intense reduction is always in a zonal belt across Europe and E. Asia. The Southern Hemisphere response from 5 kaBP appears to show a standing wave pattern affecting the subtropical highs.

In the DJF season a stationary wave in the anomaly field in both the Northern and Southern Hemispheres is apparent at all timeslices but most pronounced at 4 kaBP (Fig. 6d). This is a recognised response to surface temperature anomalies as described in Hoskins and Karoly (1981) although, in this case, there are multiple thermal anomalies caused by ALCC. By 2 kaBP there is a reduction in 500 hPa geopotential height over most of the globe. Note that there are several areas that are not statistically significant in Fig. 6c and d implying a large amount of variability. These geopotential height changes contribute to the simulated remote temperature changes by altering the regions of vorticity, which in turn influence the regions of ascent and descent and thus the surface climatic conditions. The geopotential height anomalies can also alter the pattern of the upper level winds thus influencing surface storm tracks.

In particular, the positioning of the geopotential height anomalies in the earlier timeslices (up to 4 kaBP, Fig. 6d) indicate an increased tendency towards a positive Tropical/Northern Hemisphere (TNH) pattern (<http://www.cpc.ncep.noaa.gov/data/teledoc/tnh.shtml>) with above average heights over the Bering Strait/Gulf of Alaska and north-eastward of the Gulf of Mexico and below average heights over eastern Canada. This would be expected to cause cooler temperatures over the continental United States by increasing the transport of cold polar air into the United States. In several time

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



slices it can be seen that DJF temperature anomalies over Bering Strait/Alaska (e.g. Fig. 3) show a warming pattern where there are positive geopotential height anomalies in Fig. 6c and d. The DJF temperature anomalies (warming) over Siberia up to 4 ka (Fig. 3) also relate to the stationary wave pattern. The decreased heights over the polar regions in most of the time slices are indicative of a positive Arctic Oscillation (AO) (Fig. 6b) which has been shown to be correlated with milder winters in Siberia (Tubi and Dayan, 2012).

### 3.3 Hydroclimate

Precipitation responses to ALCC (Fig. 7 for JJA and Fig. 8 for DJF) tend to be caused by a response to large-scale circulation changes rather than being directly attributable to local land use. The European precipitation response in the DJF season is not entirely consistent throughout the time slices but the general response is a slight decrease in precipitation around the western and Mediterranean coasts with this dryness extending further into the continent by 1 ka BP. The simulations show that Europe in the JJA season has an increase in precipitation compared to the control from 7 ka BP onwards, which gradually increases in extent, possibly influenced by the increased tendency to a negative North Atlantic Oscillation (NAO). Positive anomalies begin in the warm pool of the Gulf of Mexico and extend across the North Atlantic following the track of positive anomalies to the 850 hPa wind field (Fig. 9). In addition, as the cooler temperature anomalies extend quite high in the troposphere over Europe this increases relative humidity throughout the low to mid troposphere (not shown) and thus the likelihood of large-scale precipitation.

In India there is a decrease in monsoon precipitation from 5 ka BP, which then gradually increases in intensity. This is partly driven by the slightly cooler Indian sub-continent temperatures (Fig. 10) in the critical months for monsoon development and also by cooler temperatures in Europe and East Asia and increased snow cover on the Tibetan plateau. This leads to decreased monsoonal circulation and decreased cloudiness. There are also changes to the East Asian monsoon with wetter conditions to the

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



north and south of the region and drier conditions in the centre. This pattern is seen reasonably consistently from 4 kaBP onwards.

### 3.3.1 Inter Tropical Convergence Zone (ITCZ)

Analysis of the precipitation fields (Figs. 8 and 9) shows an overall southward migration of the ITCZ. These changes are most obvious in the Atlantic and Pacific Oceans and over the continent of Africa. In the DJF season there are changes to the ITCZ in the Western Pacific but a consistent pattern is not seen until 3 ka when there is southward shift in the ITCZ over the Atlantic and Atlantic coasts leading by 2 ka BP to a decrease in precipitation in the interior of southern Africa and wetter conditions on the coasts.

There is generally increased precipitation over the Indian Ocean and the Amazonian region of South America and a reduction over the Bay of Bengal but this pattern is not entirely consistent throughout the timeslices. Similarly, in the JJA season there are changes to the ITCZ throughout all the timeslices but these do not all show a consistent pattern. The most persistent changes are increased precipitation over the Pacific from 5 kaBP, over Central America from 7 kaBP and a southward shift from 2 kaBP. This southward shift in the ITCZ in the JJA season impacts the West African monsoon with lower precipitation in a belt across the monsoon region by 2 kaBP although the west coast of North Africa is wetter.

The generally cooler temperatures in the Northern Hemisphere may influence the latitudinal position of the Hadley cell and thus the location of the ITCZ via the influence on the inter-hemispheric temperature gradient resulting in the strengthening of the northward cross-equatorial energy transport (e.g. Kang et al., 2008). This shift south of the ITCZ to transport heat to the cooler Northern Hemisphere is seen in both the DJF and JJA seasons.

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.3.2 Snow cover

Lower surface air temperatures in the ALCC scenario relative to the control lead to an increase in winter snow accumulation (Fig. 11). This increase is seen by 5 kaBP mostly in northern and mountainous regions. The areas affected gradually increase so that by 3 kaBP more temperate and lower lying areas see increases in snow depth. The effects are most pronounced in North America and Europe. In regions outside the areas of permanent snow cover increases in snow depth will delay the melting of the snow pack and thus result in a longer period of snow cover. The increased snow cover due to the cold temperature anomalies will cause additional cooling due to the increased albedo. This will be greatest in regions of deforestation where the snow-covered ground is no longer masked by the canopy of the forest. This increased snow cover would also lead to decreases in precipitation due to lower rates of moisture recycling over land.

## 4 Temporal evolution of Holocene climate

The inclusion of land use changes through the Holocene has a significant impact on the progression of global average temperatures, such as to alter the direction of the multi-millennial trend. In the control experiment the changes in orbital configuration, greenhouse gases, and icesheets/sea level lead to monotonically increasing global temperatures through the Holocene (Fig. 12a). Analysis of previous experiments to assess the sensitivity to different natural forcings (data from Singarayer and Valdes, 2010; Singarayer et al., 2011) suggest that while the changes to orbital configuration effect a cooling in global temperature over the Holocene, this is outweighed by increases in greenhouse gases ( $\sim 17$  ppm  $\text{CO}_2$  from 8 kaBP to late pre-industrial time), which result in overall warming. This contrasts with recent data compilations that suggest a general decline in global temperatures since the mid-Holocene (Marcott et al., 2013). The increasing magnitude and spread of ALCC through the Holocene reconstructed in KK10

CPD

11, 4601–4641, 2015

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



counteracts the influence of increasing greenhouse gases, so that temperatures are effectively steady from 3 kaBP in HadCM3 (Fig. 13a).

These global trends are composed of considerable heterogeneity at the regional scale. When broken down into zonal regions it can be clearly seen that the difference in trends with/without land use is greatest in the Northern extratropics (Fig. 12b), where the Holocene trend is modified from increasing temperatures to decreasing temperatures in the late Holocene by the addition of ALCC. There are impacts on mean temperatures in the tropics (Fig. 12c) and southern extratropics (Fig. 12d) but not sufficient to influence the direction of the Holocene trend.

Mid-Holocene (MH) minus late Pre-Industrial (PI) anomalies of annual mean surface air temperature (Fig. 13a) show near global distribution of cooling, except over high latitude sea-ice regions, which are particularly influenced by changes in obliquity (higher in the MH than PI). The cooling is most dramatic over the tropics and monsoonal regions, where changes in the seasonality of insolation (due to orbital precession variation) intensify monsoon circulation in the early and mid-Holocene and the resulting additional cloud cover reduces incoming shortwave radiation as well as increased surface water altering the balance of sensible to latent heat fluxes. The inclusion of land use change in the KK10 experiment reduces the magnitude of MH cooling, especially in the mid-latitudes. Over Europe and eastern North America the anomaly is reversed to a warming (i.e. over these regions the cooling from deforestation shown in Fig. 2, increases and outstrips the warming from greenhouse gases). These are also regions where there is the highest concentration of pollen reconstructions (Fig. 13c; Bartlein et al., 2011; Mauri et al., 2014). The influence of land use change improves the data-model comparison with the reconstruction by Bartlein et al. (2011) over these key areas (Fig. 13a–c). Likewise, when simulated top-level ocean temperatures are compared with SST data used within the Marcott et al. (2013) compilation, the inclusion of land use improves the data-model comparison (Fig. 13d and e). However, the largest MH warming in the model is in the summer months, whereas, recent seasonal temperature

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)







changes found in this study were greater e.g. a pre-industrial global annual mean temperature anomaly of  $-0.23^{\circ}\text{C}$  as opposed to the  $-0.17^{\circ}\text{C}$  estimated by He et al. (2014). Running similar simulations with a greater number of models would improve the robustness of the results as the variability in the results from different models can be greater than the variability of the property that is being assessed (Pitman et al., 2009; de Noblet-Ducoudré et al., 2012; Brovkin et al., 2013). These inconsistencies have been attributed to disagreements in how land use change is implemented, the parameterisation of albedo, the representation of crop phenology and evapotranspiration and the partitioning of available energy between latent and sensible heat fluxes (Pitman et al., 2009; de Noblet-Ducoudré et al., 2012). The results could be further improved by the running of transient simulations that could capture events such as the Maunder minimum. This was included in the 0.2 kaBP simulations of Strandberg et al. (2014).

From late preindustrial era simulations, one using observed atmospheric greenhouse gas concentrations and the other using greenhouse gas concentrations in a world with no anthropogenic emissions (based on linear projection from earlier Holocene trends from Kutzbach et al., 2011), He et al. (2014) estimated a net global warming of  $0.9^{\circ}\text{C}$  due to the biogeochemical effects of ALCC, with between  $0.5$  and  $1.5^{\circ}\text{C}$  warming in the areas of most intense land use changes. Incorporating this degree of warming into our early industrial era (1850 CE) simulations there would still be a net cooling in Europe, E. Asia and N.E. America with, e.g., a net cooling of up to  $2^{\circ}\text{C}$  in parts of Europe. To put this in perspective the IPCC (IPCC WG II, 2014a) consider a temperature rise of more than  $2^{\circ}\text{C}$  to be undesirable and that changes of  $1^{\circ}\text{C}$  could have an impact on vulnerable ecosystems. However, the temperature changes in this study took place over a much longer period than the timeframe considered in the IPCC and ecosystems and human societies would have had more time to adapt. The consequences of these changes for agricultural societies would vary depending on the pre-existing conditions. For example, in drier regions, where crops are more likely to be water limited, cooler, wetter summer conditions may have been beneficial to the agricultural output although the risk of erosion would be increased. The generally lower temperatures might also

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



make societies more vulnerable to further transient cooling effects such as volcanic activity.

There are discrepancies between simulations of the mid-Holocene climate and the independent data-based reconstructions. Both the simulations from this study and virtually all the PMIP3 (Palaeoclimate Model Intercomparison 3; <https://pmip3.lsce.ipsl.fr>) models show a temperature increase from the mid-Holocene to the PI whereas the Marcott et al. (2013) (and Mann et al. (2008) on a shorter timescale) reconstructions show a decrease. It is interesting to note that ALCC reduces this mismatch for HadCM3, especially in key areas such as Europe. Other factors that could lead to this discrepancy are uncertainties in the proxy reconstructions and deficiencies in climate models. These climate model deficiencies include low resolution and sensitivity and, importantly, their dependence on soil moisture whereby energy is utilised for evaporation rather than for temperature increase.

This study shows a significant increase in precipitation over Europe with increasing land use which means that the PI becomes wetter than the mid-Holocene, which leads to increases in soil moisture and changes in the sensible to latent heat flux balance and, in combination with increased albedo caused by deforestation, this results in cooler temperatures for PI than MH. If the land–atmosphere coupling strength was different and soil moisture was strongly reduced with deforestation it is likely that the cooling effect would be smaller (cf. Strandberg, 2014).

Further uncertainties arise from the robustness of the land use reconstructions, which is difficult to evaluate due to the lack of global-scale evidence for human impact on the Earth’s land surface. Much of the uncertainty comes from the lack of knowledge about the magnitude and distribution of the global human population and the rate of technological evolution and intensification through time. As part of our initial investigations simulations were also run using an alternative land use scenario (the HYDE 3.1 dataset; Goldewijk et al., 2011). The HYDE 3.1 reconstruction has substantially lower levels of land use early in the Holocene (as compared with KK10), which resulted in a later development of consistent temperature anomalies at 4 ka BP

CPD

11, 4601–4641, 2015

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

⏪

⏩

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(not shown) in comparison with the KK10 land use scenario. The decision was taken to proceed with the KK10 data due to its assumptions of a larger per capita land use earlier in the Holocene. Several ongoing international initiatives that aim to synthesise palaeoecological and archaeological data promise to lead to more robust reconstructions of Holocene ALCC in the future (e.g. PAGES LandCover6k project; <http://www.pages-igbp.org/workinggroups/landcover6k/intro>).

By the early industrial period simulated biogeophysical temperature changes in the regions of land disturbance are of the same order of magnitude (e.g. 0.83 °C annual anomaly in the main agricultural areas of Europe) as the changes seen due to CO<sub>2</sub> increases during the industrial period (0.85 °C, IPCC synthesis report, 2014b). Part of Ruddiman's original hypothesis (Ruddiman, 2003) is that pre-industrial global warming caused by anthropogenic CO<sub>2</sub>/CH<sub>4</sub> emissions should have been ~ 2 °C at higher latitudes, but there was no evidence for this warming. Ruddiman (2003) attributed this to a natural cooling trend caused by decreasing summer insolation. This study suggests that biogeophysical effects of the land use changes may also have played a part in counteracting the warming due to anthropogenic greenhouse gas emissions as acknowledged in Ruddiman (2013). The precipitation changes might also have an impact on the availability of water for rice irrigation and on natural wetlands thus affecting the production of methane.

## 6 Conclusions

In our global model simulations that use a Holocene ALCC scenario as a boundary condition, a surface temperature response to the biogeophysical effects of ALCC is seen in regions of early land use such as Europe and S.E. Asia as early as 7 ka BP in the JJA season and throughout the entire annual cycle by 2–3 ka BP. Areas outside the major regions of ALCC are also affected, with virtually the whole globe experiencing significant temperature changes with a net global cooling of 0.23 °C by the pre-industrial period. Although the temperature changes are predominantly cooling some regions

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



such as India, Southern Africa and Siberia show warming as a response to ALCC. The greatest changes are generally seen in the JJA season with a mean regional cooling of 1.4°C experienced in Europe and 1°C in E. Asia in the early industrial period (1850 CE). Much of the precipitation response to the land use tends to be due to large-scale circulation changes such as a decrease in the intensity of the Indian monsoon, the southward movement of the ITCZ and changes to the North Atlantic storm track. In Europe there is a slight decrease in precipitation in the DJF season and a more substantial increase in the JJA season. Some causal factors for the teleconnections are advection by surface winds, MSLP anomalies, and tropospheric stationary wave train disturbances in the mid- to high-latitudes.

The potential for an early global impact of ALCC on climate strongly implied by this study suggests that due consideration of this should be taken in simulations covering the Holocene. The inclusion of ALCC in the model improves the model comparison for surface air temperature with the data-driven palaeoclimate reconstructions especially in key areas such as Europe. The remote teleconnections seen in this study have implications for the regional modelling of land use change due to circulation changes that occur outside the domain of the regional model.

Overall, our model simulations indicate an increase in global surface air temperatures through the Holocene. Globally, the inclusion of ALCC data reduces the magnitude of this warming especially in the late Holocene when the temperatures remain relatively constant. Regionally, in the Northern extratropics, this warming is reversed in the late Holocene. It should be noted that in this study it is not possible to distinguish the anthropogenic component of the biogeochemical changes as the same atmospheric CO<sub>2</sub> and CH<sub>4</sub> concentrations (from ice core measurements) are prescribed for both the KK10 and control simulations. However, the level of early industrial warming due to the biogeochemical impacts of ALCC predicted by He et al. (2014) would negate the early industrial biogeophysical cooling seen in this study in all regions except for the most intensively altered landscapes of Europe, E. Asia and N.E. America.

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Other caveats are the large uncertainties in the land use data and, therefore, in our understanding of the Holocene evolution of land surface-climate interactions as well as our ability to evaluate climate models. To reduce these uncertainties there is an urgent need to extend land cover reconstructions and prehistory of land use globally (cf. LandCover6k PAGES initiative).

### Data availability

Data is available from the Bristol Research Initiative for the Dynamic Global Environment website: <http://www.bridge.bris.ac.uk/resources/simulations>.

*Acknowledgements.* M. C. Smith is supported by a University of Reading doctoral scholarship. M. C. Smith and J. S. Singarayer would like to thank Sandy Harrison and Beni Stocker for useful discussions on land cover reconstructions. J. O. Kaplan was supported by the European Research Council (313797 COEVOLVE).

### References

- Bartlein, P. J., Harrison, S. P., Brewer, S., Connor, S., Davis, B. A. S., Gajewski, K., Guiot, J., Harrison-Prentice, T. I., Henderson, A., Peyron, O., Prentice, I. C., Scholze, M., Seppä, H., Shuman, B., Sugita, S., Thompson, R. S., Viau, A. E., Williams, J., and Wu, H.: Pollen-based continental climate reconstructions at 6 and 21 ka: a global synthesis, *Clim. Dynam.*, 37, 775–802, doi:10.1007/s00382-010-0904-1, 2011.
- Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years, *Quaternary Sci. Rev.*, 10, 297–317, doi:10.1016/0277-3791(91)90033-Q, 1991.
- Betts, R. A., Falloon, P. D., Goldewijk, K. K., and Ramankutty, N.: Biogeophysical effects of land use on climate: model simulations of radiative forcing and large-scale temperature change, *Agr. Forest Meteorol.*, 142, 216–233, doi:10.1016/j.agrformet.2006.08.021, 2007.
- Broecker, W. S. and Stocker, T. F.: The Holocene CO<sub>2</sub> rise: anthropogenic or natural?, *EOS Trans. AGU*, 87, 27–27, doi:10.1029/2006EO030002, 2006.

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Brovkin, V., Boysen, L., Arora, V. K., Boisier, J. P., Cadule, P., Chini, L., Claussen, M., Friedlingstein, P., Gayler, V., v. d. Hurk, B. J. J. M., Hurtt, G. C., Jones, C. D., Kato, E., de Noblet-Ducoudré, N., Pacifico, F., Pongratz, J., and Weiss, M.: Effect of anthropogenic land-use and land-cover changes on climate and land carbon storage in CMIP5 projections for the twenty-first century, *J. Climate*, 26, 6859–6881, 2013.

Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System model – HadGEM2, *Geosci. Model Dev.*, 4, 1051–1075, doi:10.5194/gmd-4-1051-2011, 2011.

Cook, B. I., Anchukaitis, K. J., Kaplan, J. O., Puma, M. J., Kelley, M., and Gueyffier, D.: Pre-Columbian deforestation as an amplifier of drought in Mesoamerica, *Geophys. Res. Lett.*, 39, L16706, doi:10.1029/2012GL052565, 2012.

Covey, C., AchutaRao, K. M., Cubasch, U., Jones, P., Lambert, S. J., Mann, M. E., Phillips, T. J., and Taylor, K. E.: An overview of results from the Coupled Model Intercomparison Project, *Global Planet. Change*, 37, 103–133, doi:10.1016/S0921-8181(02)00193-5, 2003.

Cox, P. M.: Description of the TRIFFID Dynamic Global Vegetation Model, Tech. Note 24, Hadley Centre, Met Office, 16 pp., available at: <http://www.metoffice.co.uk> (last access: 25 September 2015), 2001.

Darby, H. C.: The clearing of the woodland in Europe. In: *Man's Role in Changing the Face of the Earth*, edited by: Thomas Jr., W. L., University of Chicago Press, Chicago, 183–216, 1956.

de Noblet-Ducoudré, N., Pitman, A., Delire, C., v. d. Hurk, B. J. J. M., Boisier, J. P., Brovkin, V., Cruz, F., Voldoire, A., v. d. Molen, M. K., Müller, C., Bonan, G. B., Gayler, V., Reick, C. H., Strengers, B. J., and Lawrence, P. J.: Determining robust impacts of land-use induced land-cover changes on surface climate over North America and Eurasia; results from the first set of LUCID experiments, *J. Climate*, 25, 3261–3281, 2012.

Essery, R. L. H., Best, M. J., Betts, R. A., Cox, P. M., and Taylor, C. M.: Explicit representation of subgrid heterogeneity in a GCM land surface scheme, *J. Hydrometeorol.*, 4, 530–543, doi:10.1175/1525-7541(2003)004<0530:EROSHI>2.0.CO;2, 2003.

Folland, C. K., Knight, J., Linderholm, H. W., Fereday, D., Ineson, S., and Hurrell, J. W.: The summer North Atlantic oscillation: past, present, and future, *J. Climate*, 22, 1082–1103, doi:10.1175/2008JCLI2459.1, 2009.



Goldewijk, K. K., Beusen, A., Van Drecht, G., and De Vos, M.: The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12 000 years, *Global Ecol. Biogeogr.*, 20, 73–86, doi:10.1111/j.1466-8238.2010.00587, 2011.

Gordon, C., Cooper, C., Senior, C. A., Banks, H., Gregory, J. M., Johns, T. C., Mitchell, J. F. B., and Wood, R. A.: The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments, *Clim. Dynam.*, 16, 147–168, doi:10.1007/s003820050010, 2000.

He, F., Vavrus, S. J., Kutzbach, J. E., Ruddiman, W. F., Kaplan, J. O., and Krumhardt, K. M.: Simulating global and local surface temperature changes due to Holocene anthropogenic land cover change, *Geophys. Res. Lett.*, 41, 623–631, 2014.

Hoskins, B. J. and Karoly, D. J.: The steady linear response of a spherical atmosphere to thermal and orographic forcing, *J. Atmos. Sci.*, 38, 1179–1196, 1981.

Hurrell, J. W.: Decadal trends in the North Atlantic Oscillation: regional temperatures and precipitation, *Science*, 269, 676–679, doi:10.1126/science.269.5224.676, 1995.

Intergovernmental Panel on Climate Change (IPCC): Impacts, Adaptation and Vulnerability – Contribution of Working Group II to the Third Assessment Report of IPCC, Cambridge University Press, 2001.

Intergovernmental Panel on Climate Change (IPCC): The Physical Science Basis – Contribution of Working Group I to the Fourth Assessment Report of the IPCC, Cambridge University Press, 2007.

IPCC: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change edited by: Field, C. B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., Genova, R. C., Girma, B., Kissel, E. S., Levy, A. N., MacCracken, S., Mastrandrea, P. R., and White, L. L., Cambridge University Press, Cambridge, UK and New York, NY, USA, 1132 pp., 2014a.

IPCC: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Core Writing Team, Pachauri, R. K., and Meyer, L. A., IPCC, Geneva, Switzerland, 151 pp., 2014b.

**The biogeophysical climatic impacts of anthropogenic land use**

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Joos, F., Gerber, S., Prentice, I. C., Otto-Bliesner, B. L., and Valdes, P. J.: Transient simulations of Holocene atmospheric carbon dioxide and terrestrial carbon since the last glacial maximum, *Global Biogeochem. Cy.*, 18, GB2002, doi:10.1029/2003GB002156, 2004.

Kang, S. M., Held, I. M., Frierson, D. M., and Zhao, M.: The response of the ITCZ to extratropical thermal forcing: idealized slab-ocean experiments with a GCM, *J. Climate*, 21, 3521–3532, 2008.

Kaplan, J. O., Krumhardt, K. M., and Zimmermann, N.: The prehistoric and preindustrial deforestation of Europe, *Quaternary Sci. Rev.*, 28, 3016–3034, doi:10.1016/j.quascirev.2009.09.028, 2009.

Kaplan, J. O., Krumhardt, K. M., Ellis, E. C., Ruddiman, W. F., Lemmen, C., and Goldewijk, K. K.: Holocene carbon emissions as a result of anthropogenic land cover change, *Holocene*, 21, 775–791, doi:10.1177/0959683610386983, 2011.

Kutzbach, J. E., Vavrus, S. J., Ruddiman, W. F., and Philippon-Berthier, G.: Comparisons of atmosphere–ocean simulations of greenhouse gas-induced climate change for pre-industrial and hypothetical “no-anthropogenic” radiative forcing, relative to present day, *Holocene*, 21, 793–801, doi:10.1177/0959683611400200, 2011.

Lean, J. and Rowntree, P. R.: A GCM simulation of the impact of Amazonian deforestation on climate using an improved canopy representation, *Q. J. Roy. Meteor. Soc.*, 119, 509–530, doi:10.1002/qj.49711951109, 1993.

Louergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola, J., Raynaud, D., Stocker, T. F., and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH<sub>4</sub> over the past 800 000 years, *Nature*, 453, 383–386, doi:10.1038/nature06950, 2008.

Mann, M. E., Zhang, Z., Hughes, M. K., Bradley, R. S., Miller, S. K., Rutherford, S., and Ni, F.: Proxy-based reconstructions of hemispheric and global surface temperature variations over the past two millennia, *P. Natl. Acad. Sci. USA*, 105, 13252–13257, doi:10.1073/pnas.0805721105, 2008.

Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A reconstruction of regional and global temperature for the past 11 300 years, *Science*, 339, 1198–1201, doi:10.1126/science.1228026, 2013.

Mauri, A., Davis, B. A. S., Collins, P. M., and Kaplan, J. O.: The climate of Europe during the Holocene: a gridded pollen-based reconstruction and its multi-proxy evaluation, *Quaternary Sci. Rev.*, 112, 109–127, doi:10.1016/j.quascirev.2015.01.013, 2015.

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mitchell, L., Brook, E., Lee, J., Buizert, C., and Sowers, T.: Constraints on the late Holocene anthropogenic contribution to the atmospheric methane budget, *Science*, 342, 964–966, doi:10.1126/science.1238920, 2013.

Oglesby, R. J., Sever, T. L., Saturno, W., Erickson III, D. J., and Srikishen, J.: Collapse of the Maya: could deforestation have contributed?, *J. Geophys. Res.*, 115, D12106, doi:10.1029/2009JD011942, 2010.

Peltier, W. R.: Global glacial isostasy and the surface of the ice-age Earth: the ICE-5G (VM2) model and GRACE, *Annu. Rev. Earth Pl. Sc.*, 32, 111–149, doi:10.1146/annurev.earth.32.082503.144359, 2004.

Petit, J. R., Lipenkov, V. Y., Kotlyakov, V. M., Barkov, N. I., Bender, M., Davis, M., Delaygue, G., Barnola, J., Stievenard, M., Lorius, C., Delmotte, M., Pepin, L., Chappellaz, J., Saltzman, E., Legrand, M., Basile, I., Jouzel, J., Raynaud, D., and Ritz, C.: Climate and atmospheric history of the past 420 000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429–436, doi:10.1038/20859, 1999.

Piperno, D. R., Ranere, A. J., Holst, I., Iriarte, J., and Dickau, R.: Starch grain and phytolith evidence for early ninth millennium BP maize from the Central Balsas River Valley, Mexico, *P. Natl. Acad. Sci. USA*, 106, 5019–5024, doi:10.1073/pnas.0812525106, 2009.

Pitman, A. J., de Noblet-Ducoudré, N., Cruz, F. T., Davin, E. L., Bonan, G. B., Brovkin, V., Claussen, M., Delire, C., Ganzeveld, L., Gayler, V., v. d. Hurk, B. J. J. M., Lawrence, P. J., v. d. Molen, M. K., Müller, C., Reick, C. H., Seneviratne, S. I., Strengers, B. J., and Voldoire, A.: Uncertainties in climate responses to past land cover change: first results from the LUCID intercomparison study, *Geophys. Res. Lett.*, 36, L14814, doi:10.1029/2009GL039076, 2009.

Pongratz, J., Reick, C., Raddatz, T., and Claussen, M.: A reconstruction of global agricultural areas and land cover for the last millennium, *Global Biogeochem. Cy.*, 22, GB3018, doi:10.1029/2007GB003153, 2008.

Pongratz, J., Reick, C. H., Raddatz, T., and Claussen, M.: Biogeophysical vs. biogeochemical climate response to historical anthropogenic land cover change, *Geophys. Res. Lett.*, 37, L08702, doi:10.1029/2010GL043010, 2010.

Pope, V. D., Gallani, M. L., Rowntree, P. R., and Stratton, R. A.: The impact of new physical parametrizations in the Hadley centre climate model: HadAM3, *Clim. Dynam.*, 16, 123–146, doi:10.1007/s003820050009, 2000.

Roberts, N.: *The Holocene: an Environmental History*, John Wiley and Sons, 2013.

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ruddiman, W. F.: The anthropogenic greenhouse era began thousands of years ago, *Climatic Change*, 61, 261–293, doi:10.1023/B:CLIM.0000004577.17928.fa, 2003.
- Ruddiman, W. F.: The early anthropogenic hypothesis: challenges and responses, *Rev. Geophys.*, 45, RG4001, doi:10.1029/2006RG000207, 2007.
- 5 Ruddiman, W. F.: The Anthropocene, *Annu. Rev. Earth Pl. Sc.*, 41, 45–68, doi:10.1146/annurev-earth-050212-123944, 2013.
- Schmidt, G. A., Shindell, D. T., and Harder, S.: A note on the relationship between ice core methane concentrations and insolation, *Geophys. Res. Lett.*, 31, L23206, doi:10.1029/2004GL021083, 2004.
- 10 Singarayer, J. S. and Valdes, P. J.: High-latitude climate sensitivity to ice-sheet forcing over the last 120 kyr, *Quaternary Sci. Rev.*, 29, 43–55, doi:10.1016/j.quascirev.2009.10.011, 2010.
- Singarayer, J. S., Valdes, P. J., Friedlingstein, P., Nelson, S., and Beerling, D. J.: Late Holocene methane rise caused by orbitally controlled increase in tropical sources, *Nature*, 470, 82–85, doi:10.1038/nature09739, 2011.
- 15 Spahni, R., Chappellaz, J., Stocker, T. F., Louergue, L., Hausammann, G., Kawamura, K., Flückiger, J., Schwander, J., Raynaud, D., Masson-Delmotte, V., and Jouzel, J.: Atmospheric methane and nitrous oxide of the late Pleistocene from Antarctic ice cores, *Science*, 310, 1317–1321, doi:10.1126/science.1120132, 2005.
- Strandberg, G., Kjellström, E., Poska, A., Wagner, S., Gaillard, M.-J., Trondman, A.-K., Mauri, A., Davis, B. A. S., Kaplan, J. O., Birks, H. J. B., Bjune, A. E., Fyfe, R., Giesecke, T., Kalnina, L., Kangur, M., van der Knaap, W. O., Kokfelt, U., Kuneš, P., Latałowa, M., Marquer, L., Mazier, F., Nielsen, A. B., Smith, B., Seppä, H., and Sugita, S.: Regional climate model simulations for Europe at 6 and 0.2 k BP: sensitivity to changes in anthropogenic deforestation, *Clim. Past*, 10, 661–680, doi:10.5194/cp-10-661-2014, 2014.
- 20 Tauger, M. B.: *Agriculture in World History*, Routledge, 2013.
- Tubi, A. and Dayan, U.: The Siberian high: teleconnections, extremes and association with the Icelandic low, *Int. J. Climatol.*, 33, 1357–1366, doi:10.1002/joc.3517, 2013.
- Zohary, D., Hopf, M., and Weiss, E.: *Domestication of Plants in the Old World: the Origin and Spread of Domesticated Plants in South-West Asia, Europe, and the Mediterranean Basin*, 4th edn., Oxford University Press, Oxford, 2012.
- 30

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Figure 1.** Fraction of anthropogenically disturbed land at 1000 year intervals from the late pre-industrial (1850 CE) period to 7 ka BP. The land disturbance data is based on the anthropogenic land-use scenario KK10 (Kaplan et al., 2009, 2011).

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.



**Figure 2.** JJA Temperature Anomalies (°C) for KK10 minus Control at timeslices from the late pre-industrial period (1850 CE) to 7 kaBP. The stippling indicates grid boxes where the anomalies are significant at the 95% level using Wilcoxon ranksum statistical analysis.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.



**Figure 3.** DJF Temperature Anomalies (°C) for KK10 minus Control at timeslices from the late pre-industrial period (1850 CE) to 7 ka BP. The stippling indicates grid boxes where the anomalies are significant at the 95 % level using Wilcoxon ranksum statistical analysis.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)


[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)







**Figure 4.** Relationship between the fraction of anthropogenically disturbed land (from KK10) and the resultant JJA temperature anomaly ( $^{\circ}\text{C}$ ). **(a)** For the late pre-industrial (1850 CE) period for extratropical and tropical grid cells. **(b)** For 7, 4, 2 ka BP and late pre-industrial (1850 CE) timeslices.

**The biogeophysical climatic impacts of anthropogenic land use**

M. C. Smith et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.




**Figure 5.** DJF surface temperature anomalies for KK10 – Control and KK10 surface winds for the late pre-industrial (1850 CE) and 2 ka BP.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.



**Figure 6.** Modifiers of climate in regions outside the areas of anthropogenic land use change. All anomalies are for KK10 – Control: **(a)** JJA MSLP changes at 1850 CE, the stippling indicates grid boxes where the anomalies are significant at the 95% level using Wilcoxon ranksum statistical analysis; **(b)** as **(a)** but for 4 ka BP; **(c)** DJF 500 Pa geopotential height anomalies at 1850 CE demonstrating stationary wave pattern. The stippling indicates grid boxes where the anomalies are significant at the 95% level using Wilcoxon ranksum statistical analysis. **(d)** as **(c)** but for 4 ka BP.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.



**Figure 7.** JJA Precipitation Anomalies ( $\text{mm day}^{-1}$ ) for KK10 minus Control at timeslices from the late pre-industrial period (1850 CE) to 7 ka BP. The stippling indicates grid boxes where the anomalies are significant at the 95 % level using Wilcoxon ranksum statistical analysis.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



**Figure 8.** DJF Precipitation Anomalies ( $\text{mm day}^{-1}$ ) for KK10 minus Control at timeslices from the late pre-industrial period (1850 CE) to 7 ka BP. The stippling indicates grid boxes where the anomalies are significant at the 95% level using Wilcoxon ranksum statistical analysis.

The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 9.** JJA 850 hPa Wind Strength Anomalies ( $\text{m s}^{-1}$ ) for KK10 minus Control for the late pre-industrial period (1850).



**Figure 10.** Indian temperature seasonality; KK10 – Control surface temperature anomalies for the late pre-industrial (1850 CE) simulation; the vertical bars indicate the normal range of variability which is considered to be within 2 standard deviations of the mean.

**The biogeophysical climatic impacts of anthropogenic land use**

M. C. Smith et al.

[Title Page](#)

[Abstract](#)   [Introduction](#)

[Conclusions](#)   [References](#)

[Tables](#)   [Figures](#)

[◀](#)   [▶](#)

[◀](#)   [▶](#)

[Back](#)   [Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.




**Figure 11.** DJF Snowdepth Anomalies (cm) for KK10 minus Control for the late pre-industrial period (1850 CE). The stippling indicates grid boxes where the anomalies are significant at the 95% level using Wilcoxon ranksum statistical analysis.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.



**Figure 12.** Time series plots for the Holocene simulations from HadCM3. **(a)** annual mean global Surface Air Temperature (SAT) for the Control simulation in grey, and KK10 simulation in black; **(b)** anomaly in northern extratropical (30–90° N) annual mean temperature from the equivalent simulation at 8 ka BP; **(c)** same as **(b)** but for the tropics (30° N–30° S); **(d)** same as **(b)** but for the southern extratropics (30–90° S).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.



**Figure 13.** Mid-Holocene minus Pre-Industrial temperature anomalies. **(a)** Annual mean surface air temperature anomalies from the control experiment; **(b)** annual mean surface air temperature anomalies from the KK10 experiment; **(c)** Bartlein et al. (2011) pollen-based reconstructions of the mean annual air temperature anomaly; **(d)** ocean annual mean temperature for the top two layers (25 m) from the control experiment with palaeo-proxy reconstructions (Marcott et al., 2013) overlain in coloured circles; **(e)** as in **(d)** but for the KK10 experiment.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## The biogeophysical climatic impacts of anthropogenic land use

M. C. Smith et al.



**Figure 14.** Mid-Holocene minus Pre-Industrial precipitation anomalies. **(a)** annual mean precipitation anomalies from the control experiment; **(b)** annual precipitation anomalies from the KK10 experiment; **(c)** Bartlein et al. (2011) pollen-based reconstructions of the mean annual precipitation anomaly.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)