

Interactive comment on “An astronomical correspondence to the 1470 year cycle of abrupt climate change” by A. M. Kelsey et al.

S. Obrochta

obrochta@gipc.akita-u.ac.jp

Received and published: 2 December 2015

This paper presents an interesting investigation of astronomical factors that could produce a 1,470-year cycle. However, similar to previous comments, I think that the relevance of a 1,470-year cycle to millennial-scale variability must be better explained. The authors are focused on a body of work that is primarily 10–20 years old. The field has greatly evolved since that time, with the current consensus being that there is little evidence for an intrinsic “1,500-year cycle” in climate. Furthermore, the authors’ response to Professor Wolff’s comment leaves the impression that they would benefit from a more detailed explanation on the historical background of the 1,470-year cycle and the current state of research, which motivates me to comment.

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



In the early 1990s, both the North Atlantic sediment and the Greenland ice archives were shown to vary coherently (Bond et al., 1993). A sharp 1,470-year spectral peak was later detected in the GISP2 ice core water isotope record (Grootes and Stuvier, 1997). Coincidentally, an enigmatic ice-rafting proxy, the percentage of hematite-stained quartz and feldspar (HSG), was reported to exhibit an average 1,470-year cycle length (Bond et al., 1997). As the authors are aware this resulted in tremendous interest. However, as I will explain below, work from the last decade indicates it is unlikely that either Greenland water isotopes or HSG contain evidence for a 1,470-year climate cycle.

Of the manuscripts discussing a 1,470-year climate cycle cited in the authors' response to Professor Wolff (1), all but one predate the publication of the Greenland Ice Core Chronology 2005 (GICC05) (Andersen et al., 2006; Svensson et al., 2008). [Regarding the Bond chapter in the AGU Monograph, please note that the publisher's "how to cite" section (2) indicates the correct citation is Bond et al. (1999).] GICC05 and the North GRIP (NGRIP) ice core strongly suggest that the GISP2 chronology contains significant inaccuracies, and thus NGRIP superseded GISP2 as the key last-glacial reference series. GISP2 on its original chronology (Meese-Sowers) exhibited an inconsistent climate/annual layer thickness relationship; Svensson et al. (2006) note that: *"the existence of the proposed 1,470yr cycle depends on the exact timing and phasing of the onset of D–O events, and ... this is exactly where we believe that the GISP2 time scale is inaccurate."* Furthermore, the GISP2 age scale places the Toba Eruption about 4,000 years too early (e.g., Svensson et al., 2013).

The GICC05 chronology has been transferred to GISP2 through high-precision volcanic synchronization with NGRIP (Seierstad et al., 2014). This largely corrects the inconsistent climate-layer thickness relationship of GISP2 (Obrochta et al., 2014a) and drastically changes the characteristics of the time series. The sharp 1,470-year peak that dominated the millennial band is removed, with power redistributed to a larger number of low-amplitude peaks (see figure on last page). While one of these is near 1,500

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

years, I personally see no *a priori* reason to focus attention on this one frequency when it is of no higher significance than the others.

Regarding HSG, it has been pointed out that $1,470 \pm 532$ years is obtained by averaging 1374 ± 502 and 1536 ± 563 years. At keystone North Atlantic Site DSDP 609, HSG cyclicity was reported to be 1476 ± 585 years (Bond et al., 1999). After updating the radiocarbon calibration, correcting errors in the event stratigraphy correlation, and importing GICC05, this becomes 1573 ± 669 years. More importantly, within the GICC05-dated interval of Site 609, HSG is bimodally distributed with rare occurrence of 1,500 years cycles. Modes at 1,000-years and 2,000-years simply average to 1500 ± 500 . That is the 1,500-year cycle in HSG – the nonexistent mean in a bimodal population. Thus, I interpret that HSG provides little support for 1,500-year intervals of climate change (Obrochta et al., 2012).

Finally, there is a growing number of records from the North Atlantic region that extend beyond 200 ka and contain Greenland-like variability during the last glaciation (MIS 4-2) but exhibit a drastically different pattern during the penultimate glaciation (MIS 6; fewer, longer, and less abrupt "events") (e.g., Kandiano et al., 2004; Martrat et al., 2004; Martrat et al., 2007; Margari et al., 2010; Margari et al., 2014; Obrochta et al., 2014b). That is, the nature of millennial variability is inconsistent between individual glaciations. This suggests that internal processes are more important to the actual expression of sub-orbital climate change than external forcing.

In the absence of Greenland ice, one may consider HSG variability over the previous two glaciations from the same location, DSDP 609/IODP U1308 (Obrochta et al., 2014b). The MIS 8 record is similar to the MIS 4-2 record, with no clustering of cycles at a 1,500-year period. Cycle lengths are more uniformly distributed, but there are two modes around 1,000 and 2,000. (It should be noted that suborbital age control is much less precise for MIS 8.) MIS 6 on the other hand is quite dissimilar to the records from the surrounding glaciations. high amplitude events are spaced 4-8 ky apart, with no indication of 1,500-year periodicity.

Therefore I reiterate that, while I find this work to be an interesting investigation of astronomical factors that could produce a 1,470-year cycle, I question the relevance to the paleoclimate community as the primary archives previously interpreted to exhibit this cycle are now interpreted much differently in the light of improved chronology.

Notes

(1) <http://www.clim-past-discuss.net/11/C2098/2015/cpd-11-C2098-2015.pdf> (2) <http://onlinelibrary.wiley.com/doi/10.1029/GM112p0035/summary>

References

Andersen, K.K., et al., 2006. The Greenland ice core chronology 2005, 15-42 ka. Part 1: constructing the time scale. *Quaternary Science Reviews* 25(23-24) 3246–3257.

Bond, G.C., et al., 1997. A pervasive millennial-scale cycle in North Atlantic Holocene and glacial climates. *Science* 278(5341) 1257–1266.

Bond, G.C., et al., 1999. The North Atlantic's 1-2 kyr climate rhythm: relation to Heinrich events, Dansgaard/Oeschger cycles and the little ice age. in: Clark, P.U., Webb, R.S., and Keigwin, L.d., (eds.), *Mechanisms of Global Climate Change at Millennial Time Scales*. AGU, Washington D.C., pp. 35–58.

Bond, G.C., et al., 1993. Correlations between climate records from North-Atlantic sediments and Greenland ice. *Nature* 365(6442) 143–147.

Grootes, P.M., and Stuvier, M., 1997. Oxygen 18/16 variability in Greenland snow and ice with 10-3- to 105-year time resolution. *Journal of Geophysical Research* 102(C12) 26455–66470.

Kandiano, E.S., Bauch, H.A., and Müller, A., 2004. Sea surface temperature variability in the North Atlantic during the last two glacial-interglacial cycles: comparison of faunal, oxygen isotopic, and Mg/Ca-derived records. *Palaeogeography, Palaeoclimatology, Palaeoecology* 204(1-2) 145–164.

Margari, V., et al., 2010. The nature of millennial-scale climate variability during the past two glacial periods. *Nature Geoscience* 3(2) 127–131.

Margari, V., et al., 2014. Land-ocean changes on orbital and millennial time scales and the penultimate glaciation. *Geology*, 10.1130/G35070.1.

Martrat, B., et al., 2004. Abrupt temperature changes in the Western Mediterranean over the past 250,000 years. *Science* 306(5702) 1762–1765.

Martrat, B., et al., 2007. Four Climate Cycles of Recurring Deep and Surface Water Destabilizations on the Iberian Margin. *Science* 317(5837) 502–507.

Obrochta, S.P., et al., 2012. A re-examination of evidence for the North Atlantic “1500-year cycle” at Site 609. *Quaternary Science Reviews* 55, 23–33.

Obrochta, S.P., et al., 2014a. Conversion of GISP2-based sediment core age models to the GICC05 extended chronology. *Quaternary Geochronology* 20(0) 1–7.

Obrochta, S.P., et al., 2014b. Climate variability and ice-sheet dynamics during the last three glaciations. *Earth and Planetary Science Letters* 406(0) 198–212.

Seierstad, I.K., et al., 2014. Consistently dated records from the Greenland GRIP, GISP2 and NGRIP ice cores for the past 104 Åka reveal regional millennial-scale $\delta^{18}\text{O}$ gradients with possible Heinrich event imprint. *Quaternary Science Reviews* 106(0) 29–46.

Svensson, A., et al., 2006. The Greenland Ice Core Chronology 2005, 15-42 ka. Part 2: comparison to other records. *Quaternary Science Reviews* 25(23-24) 3258–3267.

Svensson, A., et al., 2008. A 60 000 year Greenland stratigraphic ice core chronology. *Climate of the Past* 4(1) 47–57.

Svensson, A., et al., 2013. Direct linking of Greenland and Antarctic ice cores at the Toba eruption (74 ka BP). *Climate of the Past* 9(2) 749–766.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive comment on Clim. Past Discuss., 11, 4895, 2015.

CPD

11, C2548–C2554, 2015

Interactive
Comment

Full Screen / Esc


Printer-friendly Version

Interactive Discussion

Discussion Paper

C2553





Multitaper method spectral analysis of GISP2 $\delta^{18}\text{O}$ using original chronology (blue) and GICC05 (red).

Fig. 1.