

## ***Interactive comment on “Climatic and insolation control on the high-resolution total air content in the NGRIP ice core” by O. Eicher et al.***

### **Anonymous Referee #1**

Received and published: 24 December 2015

#### General comments

This paper considerably contributes to our understanding of the complicated nature of the variability of the air content of polar ice by providing and interpreting the first high-resolution air content record obtained along the deep Greenland ice core from the NGRIP drilling site. The local insolation effect on the air content is for the first time confirmed for the Northern Hemisphere and, what may be even more important - for a site with considerably (5-8 times) higher snow accumulation than at the Antarctic sites (Dome C, Vostok, Dome Fuji) where this effect was initially discovered. This finding may help to improve our (still very poor and only qualitative) understanding of the mechanisms by which the insolation signal is imprinted in the air content of ice (At present it is assumed that local integrated summer insolation (ISI), by controlling

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



---

[Interactive  
Comment](#)

temperature gradients in the near-surface snow during summertime, significantly affects the microstructure of snow. This implies that at sites with higher accumulation and therefore a shorter residence time of the snow near the surface, the probability that an insolation signal might be found in the properties of the ice should be lower. However the Greenland data shows that a longer and warmer summertime may well compensate for the negative effect of high accumulation.) In addition, the authors attempt to give the first possible (and quite realistic) explanation for the DO event-related variations of air content that have also been observed in the other Greenland ice cores (similar data from the GRIP ice core, as far as I remember, was presented at the PIRE meeting in Grenoble this year ). However the section entitled ‘Transient firnification model experiment’ in which they explain how a sharp increase in accumulation rate may cause an observed decrease in the air content of ice is rather short and somewhat unsubstantial. I would suggest that the authors devote a bit more space and consideration to this section, which contains a novel approach to air content interpretation, in order to make it more convincing to the reader and to improve the overall presentation of the work (see my specific comments below). I also think that the manuscript could still gain from more careful and precise writing.

#### Specific comments

P5510, L23-24. The empirical relationship between pore volume at close-off and snow temperature was for the first time discovered by Raynaud and Lebel (Nature, 1979, 281(5729), 289-291).

P5513 and Fig. 1. Descriptions of the apparatus and error estimate are not sufficient to judge whether the declared (very high!) accuracy of the measurements is correct. What are the volumes  $V_1$  and  $V_2$ ? How was the volume of the system measured? (The declared accuracy for  $V_1$  and  $V_2$  seems very high!) Where is the pressure gauge located in Fig. 1? What is the accuracy of pressure ( $p_{exp}$ ) measurements and what is the  $p_{exp}$  typical value during the measurements? Do you correct  $p_{exp}$  for the partial pressure of water vapor, and how do you estimate the latter? ... My feeling

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive  
Comment

is that the overall (absolute?) accuracy of the method involving the inhomogeneous temperature of the system is overestimated, but I cannot judge this based on the limited data available from the ms.

P5515, L4-6. From my point of view these systematic shifts between different sets of measurements give us the right impression about the real absolute accuracy of the method (i.e., of the order of 5%).

P5519, L8-9. ‘...where  $T_s$  is the snow temperature in Kelvin, here assumed to be the same as the temperature at bubble close-off depth...’ I don’t understand the meaning of this misleading sentence here. Then, in equation (8) for  $V_{cr}$  you distinguish between  $T_s$  and  $T_c$ , and this is correct. (Indeed, at present-day (stationary) conditions,  $T_s=T_c$  within uncertainties not exceeding 1.5 °C. This approximate equality was used by Martinerie et al.(1992, 1994) to derive empirical relationships  $V_c(T_s)$  (7) from data on air content in recent ice at different drilling sites using both  $T_s$  and  $T_c$  (whatever was available), but this information is not important for your consideration).

P5527, L21-24. Please explain (or provide reference) how did you estimate changes in temperature and accumulation rate during DO-event that you used in the modeling.

P5527, L24-27. These 4 lines of the text do not explain how was the resulting modeled TAC evolution shown by solid blue curve in Fig 11 actually obtained. Please give more details explaining the calculations, which will allow the reader to judge whether the proposed scenario is realistic or not.

P5529, L7-11. I agree with the authors’ conclusion, but I myself would not refrain from an attempt to calculate, e.g. using the CWT technique, the time delay between the filtered air content record on the AICC2012 ice age scale and the ISI curve on the astronomical scale. Such an exercise could help to quantify the effect of the climate-related variations on the precision of the air content-based chronology for the Greenland ice core.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Finally, I suggest that the authors mention in this paper, where it is appropriate, the work of Suwa and Bender, 2008 ('O<sub>2</sub>/N<sub>2</sub> ratios of occluded air in the GISP2 ice core') in which both the local insolation signal and the millennial scale signals that are in phase with the local temperature record of rapid climate change (DO events) are discussed in application to the O<sub>2</sub>/N<sub>2</sub> record from the GISP2 core. Provided the variations in the air content and the O<sub>2</sub>/N<sub>2</sub> ratio are both related to variations in the close-off porosity, as proposed in Lipenkov et al. (2011), it is relevant to compare the findings of the reviewed work with those of Suwa and Bender.

#### Technical comments

Please check and correct if needed the use of symbols in equations and their definitions. I give only few examples where corrections are needed: 1. To denote air content you use  $V$  in eqs. (1) and (8), but TAC in eqs. (2), (4), and (9). Why do you use different symbols for the same thing? 2. In eq. (2) you use for the first time  $R$  (gas constant) which is defined only after eq. (9). 3. In the middle part of eq. (9)  $T_s$  in the denominator should be replaced by  $T_c$ ,  $V_c(T)$  in the numerator should be written as  $V_c(T_s)$ , and  $P_a$  likely represent  $P_c$  (? the definition for this is not found).

---

Interactive comment on Clim. Past Discuss., 11, 5509, 2015.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)