

# Response to: anonymous reviewer one (cpd-11-C2845-2016)

## *General comments:*

**GC1:** *This is a well-written manuscript. The text is well structured and the figures are clear.*

**R1:** Thank you.

**GC2:** *No methods for reconstructing palaeoclimates from biotic proxies are perfect. It is important to be aware of the limitations of each method. This ms is an in-depth critical analysis of the Co-existence approach. It would have been useful to suggest alternatives to it whenever possible; otherwise we are left with nothing. The section on page 5742, lines 9 and foll. is therefore too short.*

**R2:** We argue that a method should not be used simply because there is no alternative. However, other methods do exist overcoming the most serious flaws of the Coexistence Approach and we have cited these in the text. We included a new (final) section outlining how we think the field of palaeoclimate reconstruction using biological proxies should test methods in future.

## *Specific comments*

**SC1:** *The first part of the title is slightly too confrontational.*

**R3:** This was purposefully done. The NECLIME project, which is the home of the Coexistence Approach (CA), consists of over 100 members (<http://www.neclime.de/members.html>). Previous manuscripts criticising the approach have failed to elicit any meaningful engagement from the group members and the method continues to be used essentially unchanged (and untested; contrary to what is said in Utescher et al., 2014). For instance, Hoorn et al. (2012: 28) disregard nearly all principal points raised by Grimm and Denk (2012) as being “methodological flawed” including such incorporated later in the Utescher et al. (2014) paper. In Utescher et al. (2014), the only reference to Grimm and Denk (2012) is on p. 67 pointing out “successful tests with modern floras (Mosbrugger and Utescher, 1997, Grimm and Denk, 2012, Fig. 7) [side note: there is no Figure 7 in either of the cited papers and Fig. 7 in the Utescher et

al. paper does not address this issue] ... *underline the suitability of Palaeoflora climate data for palaeoclimate reconstructions* [using the Coexistence Approach]". This is a clear miscitation of Grimm and Denk (2012) who demonstrated the poor, sometimes erratic, performance of CA/Palaeoflora using over 200 modern floras. Reconstructions resulted in many erroneous mean annual temperature intervals (as acknowledged by other researchers not using CA: e.g. Eldrett et al., 2014; Kotthoff et al., 2014). This practical issue alone would require full re-investigation using the substantially augmented database (Utescher et al., 2014) to understand to which extend these primary, data-related errors affected estimates in the past. The original paper (Mosbrugger and Utescher, 1997), the only other paper testing the method/main database on modern-day floras, used *four* modern validation floras: One North American (S.E. United States), two from the Upper Rhine Valley (Germany), and one from the Istrian Peninsula, the latter producing wrong precipitation values (see Grimm and Denk, 2012, for an in-depth discussion of the original validation). Even the new documentation guidelines proposed by Utescher et al. (2014), following earlier suggestions of Grimm and Denk (2012; but not cited as such, see Table 1 in Grimm et al., 2015), are not applied by the very same authors (e.g. Tang et al., 2015). The new strategy seems to be to acknowledge problems with the database and data-related issues (without providing actual details), but not the method (the Grimm and Denk, 2012, paper has been cited 30 times according GoogleScholar, apparently stirring up things a little), while, at the same time, all older results (using wrong database entries) are still treated as fully valid (see statistics presented in Utescher et al., 2014).

Table R1 lists the major fallacies and fantasies of the Coexistence Approach as proposed and applied (according Mosbrugger and Utescher, 1997; Utescher et al., 2014; and most palaeoclimate studies using the Coexistence Approach) and addressed in our ms. Nevertheless, Mosbrugger and Utescher (1997) conclude "*Taking the advantages and disadvantages into account, we consider the coexistence approach to represent a reasonable, robust and the most widely applicable technique for quantitative reconstructions of the terrestrial palaeoclimate in the Tertiary.*" Despite acknowledging the many surfacing, partly severe problems ("*uncertainties*"), Utescher et al. (2014) state "*With more than 15 years of experience of using the Coexistence Approach on Cenozoic micro- and macropalaeobotanical records, it is clear that the method provides a robust palaeoclimatic proxy.*" We feel a little confrontation is needed in this case, hoping it may crack cemented but unfounded beliefs.

**Technical corrections**

**TC1:** *Page 5728, line 2: used to*

**R4:** Correct. Two grammatical errors in the first sentence. Ooops!

**TC2:** *Page 5733, line 24: add call to table 1*

**R5:** Added.

**TC3:** *Page 5734, line 9: add call to table 1*

**R6:** Added.

**TC4:** *Page 5736, line 15: one climatic parameter*

**R7:** Added.

**TC5:** *Page 5741, line 15: species through*

**R8:** Corrected.

**TC6:** *Page 5743, line 23: some capital letters missing*

**R9:** Thank you for picking this up - it occurred during the typesetting process.

**TC7:** *Figure 2b: uniformitarianism [i missing]*

**R10:** Corrected.

**Table R1: Fallacies and fantasies of the Coexistence Approach**

---

<b>Fallacies</b>	<p>Violations of basic assumptions and errors will surface in the form of “climatic outliers” (e.g. Mosbrugger and Utescher, 1997: 73).</p> <p>Outliers can be recognised ignoring most community information.</p> <p>The climatic space of a taxon is rectangular (neither addressed nor argued for in Mosbrugger and Utescher, 1997; Utescher et al., 2014)</p> <p>Extinct climates can be reconstructed (considered a major advantage of the method, Utescher et al., 2014)</p> <p>Strict(!) niche conservatism can be considered to be the norm, at least for the all genera defining coexistence intervals as nearest living relatives. Note that Utescher et al. (2014) point out that even though highly precise intervals are “<i>usually considered beyond the resolution of the primary data</i>”, coexistence intervals of “<i>&lt;1 °C for MAT and &lt;100 mm for MAP, respectively, represent valid solutions in the sense of the CA algorithm</i>”.</p>
<b>Fantasies</b>	<p>Accuracy and precision are sufficient to resolve minute climate shifts in the order of few degree Celsius or few tenths of mm precipitation per month or year (see Coexistence Approach literature; e.g. examples/references given in <b>R9 in Response to Reviewer 2</b>).</p> <p>The pair of fossil species that defines the coexistence interval lived always close to the minimum and maximum tolerances of the according modern genus – this is an implicit necessary assumption of the Coexistence Approach never mentioned in any paper.</p>

---

## References

Eldrett, J. S., Greenwood, D. R., Polling, M., Brinkhuis, H., and Sluijs, A.: A seasonality trigger for carbon injection at the Paleocene–Eocene Thermal Maximum, *Clim. Past*, 10, 759–769, 2014.

Grimm, G. W., and Denk, T.: Reliability and resolution of the coexistence approach — A revalidation using modern-day data, *Rev. Palaeobot. Palynol.*, 172, 33–47, 2012.

Grimm, G. W., Bouchal, J. M., Denk, T., and Potts, A. J.: Fables and foibles: a critical analysis of the Palaeoflora database and the Coexistence Approach for palaeoclimate reconstruction., *bioRxiv*, doi, 10.1101/016378, 2015.

Hoorn, C., Straathof, J., Abels, H. A., Xu, Y., Utescher, T., and Dupont-Nivet, G.: A late Eocene palynological record of climate change and Tibetan Plateau uplift (Xining Basin, China). *Palaeogeogr. Palaeoclimat. Palaeoecol.*, 344–345, 16–38, 2012.

Kotthoff, U., Greenwood, D. R., McCarthy, F. M. G., Müller-Navarra, K., Prader, S., and Hesselbo, S. P.: Late Eocene to middle Miocene (33 to 13 million years ago) vegetation and

climate development on the North American Atlantic Coastal Plain (IODP Expedition 313, Site M0027), *Clim. Past*, 10, 1523–1539, 2014.

Mosbrugger, V., and Utescher, T.: The coexistence approach -- a method for quantitative reconstructions of Tertiary terrestrial palaeoclimate data using plant fossils, *Palaeogeogr. Palaeoclimat. Palaeoecol.*, 134, 61-86, 1997.

Tang, H., Eronen, J. T., Kaakinen, A., Utescher, T., Ahrens, B., and Fortelius, M.: Strong winter monsoon wind causes surface cooling over India and China in the Late Miocene, *Climates of the Past Discussions*, 11, 63–93, 2015.

Utescher, T., Bruch, A. A., Erdei, B., François, I., Ivanov, D., Jacques, F. M. B., Kern, A. K., Liu, Y.-S. C., Mosbrugger, V., and Spicer, R. A.: The Coexistence Approach—Theoretical background and practical considerations of using plant fossils for climate quantification, *Palaeogeogr. Palaeoclimat. Palaeoecol.*, 410, 58–73, 2014.

1 **Fallacies and Fantasies: the theoretical underpinnings of**  
2 **the Coexistence Approach for palaeoclimate**  
3 **reconstruction**

4

5 **Guido W. Grimm<sup>1</sup> and Alastair J. Potts<sup>2</sup>**

6 [1] University of Vienna, Department of Palaeontology, Vienna, Austria

7 [2] Nelson Mandela Metropolitan University, Centre of Coastal Palaeoscience, Port Elizabeth,  
8 South Africa

9 **Abstract**

10 The Coexistence Approach has been used [to](#) infer palaeoclimates for many Eurasian fossil  
11 plant assemblages. However, the theory that underpins the method has never been examined  
12 in detail. Here we discuss acknowledged and implicit assumptions, and assess the statistical  
13 nature and pseudo-logic of the method. We also compare the Coexistence Approach theory  
14 with the active field of species distribution modelling. We argue that the assumptions will  
15 inevitably be violated to some degree and that the method has no means to identify and  
16 quantify these violations. The lack of a statistical framework makes the method highly  
17 vulnerable to the vagaries of statistical outliers and exotic elements. In addition, we find  
18 numerous logical inconsistencies, such as how climate shifts are quantified (the use of a  
19 “center value” of a coexistence interval) and the ability to reconstruct “extinct” climates from  
20 modern plant distributions. Given the problems that have surfaced in species distribution  
21 modelling, accurate and precise quantitative reconstructions of palaeoclimates (or even  
22 climate shifts) using the nearest-living-relative principle and rectilinear niches (the basis of  
23 the method) will not be possible. The Coexistence Approach can be summarised as an  
24 exercise that shoe-horns a plant fossil assemblages into coexistence and then naively assumes  
25 that this must be the climate. Given the theoretical issues, and methodological issues  
26 highlighted elsewhere, we suggest that the method be discontinued and that all past  
27 reconstructions be disregarded and revisited using less fallacious methods.

1 **Keywords:** Cainozoic; Eurasia; mutual climate range; nearest-living-relative principle;  
2 physiological uniformitarianism; species distribution modelling; theoretical problems;  
3 univariate niche

4 **1 Introduction**

5 One of the most widely used methods to infer the palaeoclimates of Eurasia using fossil plant  
6 assemblages is the ‘Coexistence Approach’ (Utescher et al., 2014). Conceptually, this  
7 approach belongs to the family of mutual climate range techniques but also makes use of the  
8 ‘nearest-living-relative’ principle; a nearest-living-relative (NLR) is a modern taxon (species,  
9 group of species, genus, or higher) that is considered an analogue for the fossil taxon. Mutual  
10 climate range methods use the climatic preferences of modern species (a set of nearest-living-  
11 relatives), as defined by their current distribution, to infer the potential climatic niche for a  
12 fossil assemblage. In the case of the Coexistence Approach, the climate niche is defined using  
13 minimum and maximum climate values of an NLR, obtained from its present-day distribution.  
14 Pure mutual climate range techniques are usually restricted to reconstructing palaeoclimates  
15 of the recent past (i.e. Quaternary) where species in the fossil assemblages can be directly  
16 linked to modern species (e.g. Elias, 1997, 2001; Thompson et al., 2012; Harbert and Nixon,  
17 2015); the processes of extinction and speciation are ignored and niche conservatism is  
18 considered to be the norm. However, to apply these palaeoclimate reconstruction techniques  
19 to assemblages from older time periods requires the use of the nearest-living-relative  
20 principle, which is linked to the concept of physiological uniformitarianism (Tiffney and  
21 Manchester, 2001; Tiffney, 2008). The niche-space of an NLR is used to represent that of the  
22 fossil taxon. Thus, one assumes that the climate niche of the NLR (the modern species or  
23 species set) is identical to that of the associated fossil taxon (an extinct sister or ancestral  
24 species) and the mutually shared climate range of the NLRs enables the estimation of the  
25 climate conditions in which the fossil assemblage thrived (Fig. 1).

26 Despite the availability of alternative palaeoclimate reconstruction techniques using NLRs  
27 and the mutual climate range approach (e.g. Greenwood et al., 2005), the Coexistence  
28 Approach has become the *de facto* method for plant fossil assemblages of Eurasia for time  
29 periods spanning the Miocene to Late Cretaceous (Utescher et al., 2014). The cumulative  
30 citation count of studies using the Coexistence Approach is in excess of 10,000. On the  
31 surface, it reconstructs precise palaeoclimatic conditions (usually reported with a precision of  
32 0.1 °C and 1 mm precipitation per month or year) based on a series of acknowledged and

1 implicit basic assumptions (Table 1; Mosbrugger and Utescher, 1997; Utescher et al., 2014).  
2 These assumptions appear straightforward, but have theoretical and practical implications  
3 essentially ignored in the application of the Coexistence Approach (Mosbrugger and  
4 Utescher, 1997; Utescher et al., 2014; Grimm et al., 2015). Furthermore, the Coexistence  
5 Approach avoids any statistical processing (Mosbrugger and Utescher, 1997; Utescher et al.,  
6 2014). It relies to some degree on hard-to-grasp pseudo-logic, some of which is advocated as  
7 strengths of the method, e.g. the ability to reconstruct extinct climates (Utescher et al., 2014).  
8 The applicability of the nearest-living-relative principle for reconstructing past climates in a  
9 quantitative manner is never questioned. This is surprising in the light of ongoing discussions  
10 in the field of spatial distribution modelling, which shares a number of assumptions with  
11 mutual climate range and nearest-living-relatives methods. Below we discuss each of these  
12 issues in further detail.

## 13 **2 Theoretical background of the ‘Coexistence Approach’**

### 14 **2.1 Assumptions of the Coexistence Approach**

15 Mosbrugger and Utescher (1997) list four basic assumptions that need to be fulfilled (Table  
16 1). The first assumption has never been used in the application of the Coexistence Approach,  
17 and the three others superimpose additional uncertainty on the method and are easily violated,  
18 particularly if the aim is high accuracy *and* precision. Notably, none of the assumptions have  
19 been tested and verified for taxa commonly used in the Coexistence Approach.

20 The first assumption is anchored on the ability to define a “systematically close” NLR (Table  
21 1). However, Mosbrugger and Utescher (1997) or Utescher et al. (2014) do not provide a  
22 framework on how to quantify “systematically close” and in what respect systematic  
23 closeness should be relevant for the identification of the NLR. A focus on “systematic  
24 closeness” can lead to conflict with the nearest-living-relative principle. This principle is  
25 based on overall morphological similarity and not necessarily linked to phylogenetic  
26 relatedness, which is the current basis of systematics. Thus, a fossil may be “systematically  
27 close” to a modern species (or group) that has undergone significant shifts in morphology and  
28 fundamental niche, and the best modern analogue may be a more distantly related lineage that  
29 has been morphologically and ecologically stable (Fig. 2A). In addition, the degree of  
30 systematic relatedness of a fossil to an NLR requires the placement of fossils within a

1 phylogenetic framework (i.e. a tree or network) and this has never been explored in any  
2 Coexistence Approach study.

3 There are further issues with Assumption 1 when considering the taxonomic affiliation of an  
4 NLR. Given the timespan separating ancient assemblages and modern day taxa, it has been  
5 agreed that defining an NLR at the species level is highly problematic (Grimm and Denk,  
6 2012; Utescher et al., 2014). Thus, the Coexistence Approach usually defines a  
7 “systematically close” NLR as the genus or family to which the fossil can be assigned, with  
8 rare instances of an intrageneric lineage or a modern species (Grimm and Denk, 2012;  
9 Utescher et al., 2014; Grimm et al., 2015). For example, the NLR of a fossil oak leaf would be  
10 genus *Quercus*, the NLR of a deciduous, convexly lobed oak leaf would be *Quercus* Group  
11 *Quercus* (the white oak clade), and the NLRs of a fagaceous fossil of unknown generic  
12 affinity would be all Fagaceae. Hence “systematically close”, as used in the Coexistence  
13 Approach and other nearest-living-relative approaches, translates into simply being a member  
14 of the same taxonomic rank (e.g. genus or family), and the actual phylogenetic (= systematic)  
15 distances between fossils and their NLRs is never established. Under this implementation of  
16 assigning NLRs to higher taxonomic ranks (above species) includes the taxonomic problems  
17 linked to paraphyly (exclusive common origin; Fig. 2B). Fossils of a paraphyletic group will  
18 have different systematic distances to the modern members of the specified taxonomic group  
19 of NLRs. However, this is not a problem for the combination of mutual climate range  
20 approaches and nearest-living-relative principle as long as the assumption of physiological  
21 uniformitarianism is fulfilled (**Assumption 2**). Thus, shared ancestry remains important, but  
22 the ‘systematic closeness’ of **Assumption 1** is entirely superfluous for the application of  
23 mutual climate range techniques making use of the nearest-living-relative concept.

24 The second assumption (Table 1) is based upon the concept of physiological  
25 uniformitarianism (Tiffney and Manchester, 2001; Tiffney, 2008). Physiological  
26 uniformitarianism implies that as long as lineage stays within its environmental niche, it will  
27 not accumulate morphological changes. Hence, a modern species with the same, or very  
28 similar, morphological traits of a fossil of the same evolutionary lineage should share the  
29 same environmental niche. It also implies that members of the lineages that have undergone  
30 niche shifts also experienced morphological changes. **Assumption 2** is likely to be violated  
31 when morphological changes are evident between the fossil and modern members of an  
32 evolutionary lineage and an NLR of a fossil specimen should only be used if there is

1 morphological, not mere taxonomic, similarity (ideally identical) and have a common origin.  
2 This would exclude the use of most modern plant genera and all families as NLRs as they are  
3 typically composed of morphologically divergent species.

4 In addition, the use of morphologically diverse taxonomic groups to represent an NLR usually  
5 means that the environmental niche of the NLR is large, likely encompassing the niche of the  
6 fossil, but is not “climatically similar” to that of the fossil; thus, directly violating  
7 **Assumption 2.** Novel procedures and methods are required that take cognisance of the fact  
8 that the NLR niche is likely to be far broader than can be expected for that of the fossil. The  
9 actual assumption, as used by Coexistence Approach practitioners, is that *the climatic niche of*  
10 *a fossil taxon lies somewhere within the range of niches found within the species comprising*  
11 *the NLR.* This has two major implications for the setup and interpretation of reconstructed  
12 palaeoclimates using the Coexistence Approach (and other mutual climate range techniques  
13 that use NLRs): 1) a high resolution climate reconstruction should not be possible, especially  
14 when only minimum and maximum NLR tolerances are used (Fig. 3A), and 2) mixed floras  
15 may not be identified as mutually exclusive species (or communities) can have overlapping  
16 climate ranges at higher taxonomic levels (Fig. 3B). Thus, highly precise *and* accurate climate  
17 reconstructions can only be obtained using the Coexistence Approach if the critical species  
18 within a palaeoassemblage occupied niches close to the minimum **and** maximum tolerances  
19 of their corresponding modern genus- or family level NLRs.

20 The third assumption (Table 1), that the distributions of extant species are in equilibrium with  
21 their climate, is a topic rich with discussions in the ecological and species distribution  
22 modelling literature (Araújo and Pearson, 2005; Bond et al., 2005; Sexton et al., 2009;  
23 Franklin, 2010). Species are often not in equilibrium with their climate for abiotic (e.g. soil,  
24 fire) or biotic (e.g. competition) reasons, and thus their realised niches do not span their  
25 fundamental niches. Thus species will be plastic in their expression of the realised niche  
26 depending on external factors, which would exclude the reconstruction of palaeoclimate with  
27 high accuracy. Any change in the abiotic or biotic parameters can affect the distribution of a  
28 species (i.e. its realised niche) even if the fundamental niche remains unchanged.

29 The climatic niche is solely represented by minimum and maximum values in the Coexistence  
30 Approach, which are independently compiled for climate parameters in a univariate manner.  
31 However, it has been long established that biological climate niches are multi-dimensional  
32 (Köppen, 1936; Hutchinson, 1957; Walter, 1973; Walter and Breckle, 1983–1991; Schroeder,

1 1998). Using minimum-maximum tolerances along univariate axes can only roughly  
2 approximate the multidimensional climatic niche, and may be misleading (Klotz, 1999;  
3 Thompson et al., 2012). For example, two *mutually exclusive* taxa, for which **Assumption 3**  
4 applies, may still have an artificial mutual climate range regarding their minimum and  
5 maximum tolerances (Fig. 4A). In this context it is important to note that species distribution  
6 modelling started with algorithms that used minimum and maximum values, but quickly  
7 moved on to methods that better represented the bioclimatic niche of a species (discussed  
8 further below). Thus, the use of range values for climatic parameters does not accurately  
9 capture the climatic requirements or tolerances of an NLR ([Table 1](#)), which will affect the  
10 reconstructed palaeoclimate using the Coexistence Approach.

11 The fourth and last assumption has no apparent theoretical implications. Technical  
12 implications have been discussed in Grimm and Denk (2012), Thompson et al. (2012),  
13 Utescher et al. (2014), and Grimm et al. (2015). We do, however, wish to highlight that since  
14 local climate can substantially vary over short time scales, minimum and maximum tolerances  
15 may be unduly affected by the selected observation period of climate stations.

16 Not formally addressed by Mosbrugger and Utescher (1997) or Utescher et al. (2014) are two  
17 more fundamental assumptions [in the application](#) of the Coexistence Approach, which  
18 distinguish the method from mutual climate range techniques using modern-day species: 1)  
19 palaeoassemblages comprise only taxa that existed as actual communities (i.e. all fossil  
20 specimens are autochthonous and from the same point in time); 2) absence of a fossil taxon  
21 indicates true absence (i.e. each fossil plant assemblage comprehensively reflects the actual  
22 palaeocommunity; [Table 1](#)). The Coexistence Approach implicitly assumes that only an  
23 autochthonous and strictly coeval palaeoassemblage will result in a single coexistence  
24 interval. However, given that two mutually exclusive taxa can share a climate range of  
25 minimum and maximum along univariate climate parameters, so too can allochthonous taxa  
26 in a fossil assemblage. In addition, the expansion of the climate niche using higher-level  
27 NLRs automatically increases the probability of artificial coexistence. Thus, allochthonous  
28 assemblages (mixed floras) do not necessarily result in ambiguous intervals (Fig. 4B-D) and  
29 may very well be the reason for highly precise palaeoclimate estimates (< 1 °C for  
30 temperature parameters, < 100 mm precipitation per year, < 10 mm precipitation per month)  
31 observed in many studies using the Coexistence Approach (Denk et al., 2012; Grimm et al.,  
32 2015). Thompson et al. (2012) suggest that a benefit of mutual climate range techniques, in

1 comparison to indicator-species approaches, is that the reconstruction is only affected by the  
2 presence of taxa, not their absence. However, this does not apply to the Coexistence  
3 Approach, where the mere absence of a taxon can directly affect the outcome of the  
4 reconstruction (discussed further below). For instance, absence of a taxon may eliminate  
5 another NLR as “climatic outlier” rather than producing two “ambiguous” intervals.

6 We have outlined a range of probable and inevitable issues of the purported basic assumptions  
7 of the Coexistence Approach. These will all, to some unknown degree, decrease the precision  
8 and accuracy of any approach that attempts to reconstruct palaeoclimates. In this light, the  
9 Coexistence Approach is highly unlikely to reconstruct precise or accurate palaeoclimatic  
10 conditions. Utescher et al. (2014) state that it is impossible to test the accuracy of Coexistence  
11 Approach reconstructions (but see Grimm and Denk, 2012, for mean annual temperature  
12 estimates), but follow the original paper in assuming that violation of the basic assumptions  
13 will readily surface in the form of “*climatic outliers*”.

14

## 15 **2.2 The statistical nature of the Coexistence Approach**

16 According to Utescher et al. (2014) the “*Coexistence Approach by Mosbrugger and Utescher*  
17 (*1997*) is a nearest living relative method, which relies only on the presence/absence of a  
18 plant taxon within a fossil assemblage and the climatic requirements of its modern relatives.  
19 It avoids any statistical processing or further assumptions, except those given in Mosbrugger  
20 and Utescher (1997) [i.e. the four basic assumptions, see Table 1]”. In the original paper, no  
21 means of statistical processing were proposed, hence, the Coexistence Approach defines an  
22 interval for a past climate parameter assuming that statistical effects do not exist or are  
23 negligible. The Coexistence Approach ignores the majority of the community information  
24 because the reconstructed climate interval is always solely defined by the pair of the two most  
25 divergent, but putatively coexisting NLRs. Usually one member of the pair is an exotic  
26 element; here we define ‘exotic’ as any NLR whose niche is at odds with the majority of the  
27 assemblage (e.g. Fig. 5). The Coexistence Approach lacks a statistical framework to account  
28 for potential oddities, errors or violations of assumptions, the likelihood of which increases  
29 with assemblage size or depositional age. The approach naively relies on the presumption that  
30 any violation will readily surface in the form of so-called “*climatic outliers*” (Mosbrugger and  
31 Utescher, 1997; Utescher et al., 2014). This exposes palaeoclimate reconstructions using this

1 approach to the vagaries of statistical outliers and exotic elements (see Grimm and Denk,  
2 2012; Grimm et al., 2015, for real-world data examples).

3 A ‘climatic outlier’ is identified as an NLR or small number of NLRs that do not share the  
4 climate space for a given parameter with a slightly higher number of other NLRs (Fig. 5). In  
5 those cases where there are more than one interval that can be reconstructed using the same  
6 maximum possible number of NLRs, then alternative ‘*ambiguous*’ intervals are reported; each  
7 of these intervals recognises a different set of climatic outliers. Ambiguous intervals are  
8 interpreted by Utescher et al. (2014) as the only evidence for mixed floras rather than a  
9 violation of any the assumptions discussed above. Taxa identified as climatic outliers are  
10 typically removed from a Coexistence Approach analysis for a given palaeoassemblage and  
11 parameter. We wish to highlight that a climatic outlier is simply an NLR that is seemingly at  
12 odds with a few other NLRs and must not to be confused with a statistical outlier (Fig. 5).

13 There are two paramount problems with the current outlier elimination strategy used by the  
14 Coexistence Approach. First, *two* taxa *violating* the assumptions behind the Coexistence  
15 Approach may eliminate *one* taxon that is not. A typical situation is illustrated in Figure 5,  
16 where an NLR occupying a climate range that is in general agreement with the rest of the  
17 flora would be identified and eliminated as a climatic outlier because of presence of two  
18 deviant taxa that are at odds with the overall NLR community. Second, taxa identified as  
19 climatic outliers for one [climatic](#) parameter and therefore removed from the assemblage for  
20 estimating that parameter are still, in most cases, kept for analysing other parameters for the  
21 same assemblage. In some cases, these climatic outliers even define the coexistence interval  
22 in another parameter (Grimm et al., 2015). If we follow the logic that climatic outliers  
23 represent violations of the basic assumptions of the Coexistence Approach (Utescher et al.,  
24 2014), then it is imperative that they are removed from all reconstructions for a given  
25 assemblage or in general (Table 2). This has been rarely applied in any study that has  
26 identified climatic outliers in the Coexistence Approach, mainly to avoid wide, and thus  
27 uninformative, coexistence intervals (Grimm and Denk, 2012; Grimm et al., 2015). It could  
28 be argued that any palaeoassemblage represented by mutually exclusive NLRs should be  
29 ignored until the reason for the non-coexistence can be identified and corrected for.

30 Any mutual climate range approach needs a framework to identify statistical outliers as the  
31 assumptions will inevitably be violated, and establishing the degree of violation (e.g. degree  
32 of niche shifts) is not feasible based on current knowledge. Many palaeoassemblages will

1 comprise mixed floras with elements from different climate niches, and this would need to be  
2 explicitly addressed before reconstructing coexistence intervals. As stated above, the  
3 Coexistence Approach lacks any framework to identify exotic elements or allochthonous  
4 assemblages, unless they are sufficiently divergent to generate climatic outliers.  
5 Allochthonous assemblages comprising mutually exclusive species can share a climate  
6 interval (Fig. 4B), and this problem of pseudo-coexistence is exacerbated by the use of  
7 *higher-level taxa* (genera, families) as NLRs of a fossil *species/morphotypes*. Any slightly  
8 conflicting, but exotic, element in an assemblage will have a disproportionately high influence  
9 on the palaeoclimate estimates (Fig. 5). It is clear that not only ‘climatic outliers’ and  
10 ‘ambiguous intervals’ should be indicative of mixed floras, errors in the data, or violations in  
11 the assumptions, but also *any* narrow coexistence interval (see Grimm and Denk, 2012;  
12 Grimm et al., 2015, for real-world examples).

13 Mutual climate range techniques that apply simple statistics to filter exotic taxa, such as the  
14 Bioclimatic Approach (Greenwood et al., 2005), will be less susceptible to the presence and  
15 absence of a few exotic taxa, but will also usually fail to recognise mixed floras. The problem  
16 of mixed floras can only be overcome, to some degree, by using alternative mutual climate  
17 range techniques that make use of the full spectrum of distributional information, and thus  
18 include the climatic preference of all constituent elements of a palaeoassemblage (e.g. using  
19 the niche curves in Fig. 5). This includes methods such as the weighted mutual climate range  
20 approach (Thompson et al., 2012), the probability density function method (Chevalier et al.,  
21 2014), and the coexistence likelihood estimation method (Harbert and Nixon, 2015).  
22 However, these methods will probably begin to break down when the nearest-living-relative  
23 principle is needed to link fossils with extant lineages (Thompson et al., 2012; Harbert and  
24 Nixon, 2015), and may explain why their application has been limited to Quaternary  
25 assemblages.

## 26 **2.3 Pseudo-logic of the Coexistence Approach**

27 We wish to highlight four additional points regarding the use of the Coexistence Approach  
28 that lack any (bio)logical basis, specifically: 1) the use of the “center value” to identify and  
29 quantify climatic shifts, 2) that the reconstructed climate is based on only two nominally  
30 coexisting elements, 3) that the reconstructed climate is highly dependent on the presence or  
31 absence of a single or few taxa (the “Heisenberg effect”), and 4) the reconstruction of  
32 “extinct” climates. We elaborate on each of these points below.

1 The conclusions of most Coexistence Approach studies rely on shifts observed in the so-  
2 called “center value”. This value is simply the arithmetic mean of the upper and lower  
3 boundary of the coexistence interval. Practical tests have shown that there is little correlation  
4 between the actual climate and the “center value” (Klotz, 1999; Grimm and Denk, 2012). The  
5 use of this value highlights a fundamental misunderstanding of the niche concept. If we  
6 imagine the coexistence interval to be correct, then *all* values within the interval should be  
7 equally probable as no other information is incorporated regarding the probabilities of  
8 occurrence of the assemblage. Selecting the “center value” as an indicator of a shift in climate  
9 makes no statistical or biological sense. For example, Figure 6A shows two plant assemblages  
10 that differ only by the climatic preference of a single NLR. The replacement of one NLR by  
11 another with a preference towards *lower* values gives rise to a reconstructed climate shift  
12 towards *higher* values using the “center value”.

13 Many Coexistence Approach reconstructions rely on the presence of NLRs that nominally  
14 coexist, even if these elements have climate tolerances that are at odds with the rest of the  
15 assemblage (Fig. 5; cf. Grimm and Denk, 2012; Grimm et al., 2015). In extreme cases the  
16 same coexistence interval can be reconstructed based on plant assemblages with contrary  
17 climate tolerances (Fig. 6B). In Figure 6B, the elements of two plant assemblages have  
18 contrary climate tolerances and it is the two exotic taxa in each assemblage that ensure that  
19 the reconstructed coexistence intervals are the same. Thus, the precision of the reconstructed  
20 palaeoclimates is often entirely dependent on the presence or absence of specific, usually  
21 exotic NLRs. Across Coexistence Approach studies, a handful of NLRs that occur towards  
22 the tolerance margins over the *entirety* of all palaeofloras usually determine the coexistence  
23 intervals; it is these few NLRs that give rise to the praised precision of the technique (Grimm  
24 and Denk, 2012; Grimm et al., 2015).

25 The presence or absence of individual NLRs are generally at the base of reconstruction  
26 uncertainty in coexistence interval - we term this the “Heisenberg” effect. Figure 6C shows  
27 two very similar assemblages where the presence or absence of the two highlighted taxa  
28 changes the coexistence interval reconstructed by the Coexistence Approach in a degree that  
29 would be interpreted as a trend towards higher values. The Heisenberg effect renders  
30 palaeoclimate estimates obtained with the Coexistence Approach protocol highly susceptible  
31 to taxon-bias effects. The reconstructed climate is exceedingly dependent on what fraction of  
32 the actual vegetation has been captured by the fossil assemblages (note that in Fig. 6C all

1 NLRs have a mutually shared climate range). Thus, even if all assumptions needed for a  
2 mutual climate range approach that also uses the nearest-living-relative principle are fulfilled,  
3 the Heisenberg effect will lead to unstable, even random, climate reconstructions.

4 Uttescher et al. (2014) explicitly state that, as each parameter is independently reconstructed,  
5 the Coexistence Approach has the potential to reconstruct a climate that does not exist today:  
6 an “extinct climate”. It is hard to grasp how this can be logically accommodated with the  
7 basic assumptions of the Coexistence Approach and the actuo-palaeontological nearest-living-  
8 relative principle in general (Fig. 7). An extinct climate for a palaeoassemblage would  
9 indicate that the present-day niches of the NLRs are **not** representative of the fossils, and  
10 therefore would indicate direct violations of Assumptions 2 and 3 discussed above (Table 1).  
11 In addition, it is not possible to construct an extinct climate using species that are restricted to  
12 present-day climates if the principle of physiological uniformitarianism applies. Reasons why  
13 extinct climates are reconstructed using the Coexistence Approach include violations of basic  
14 assumptions, pseudo-coexistence, the inconsistent identification of climatic outliers within an  
15 assemblage across climate variables, and the single-dimension effect where climate  
16 parameters are analysed in isolation and are assumed to be unlinked. The reconstruction of an  
17 extinct climate should be seen as a direct indication of error, and not lauded as a benefit of the  
18 method.

19 Ignoring these logical inconsistencies in the conception and application of the method, the  
20 Coexistence Approach still cannot be expected to reproduce a robust quantitative  
21 reconstruction of the palaeoclimate, as 1) assumptions are likely to be violated but cannot be  
22 detected, 2) one cannot avoid using higher-level taxa to represent fossil species or  
23 morphotypes, and 3) the fossil record will always be incomplete to different degrees, and this  
24 will affect the calculated coexistence interval.

### 25 **3 Applicability of the nearest-living-relative principle: Lessons to be learnt** 26 **from species distribution modelling**

27 Species distribution modelling (SDM) is an exceptionally active field which aims to  
28 empirically model the species-environment relationships and thereby quantify the realised  
29 niche of a given taxon (Franklin, 2010; Peterson et al., 2011) or, in some cases, communities  
30 (e.g. Potts et al., 2013). The dawn of the field was the BIOCLIM software package (Nix,  
31 1986), which is comparable to the Coexistence Approach as it used the range (or percentile

1 range) of climatic variables in a rectilinear fashion. Booth et al. (2014) describe the roots of  
2 the field and highlight that one of the most active areas of SDM development has been of  
3 methods that trim the rectilinear climate envelopes of BIOCLIM. This was driven by the early  
4 realisation that the relationships between climate variables ~~was-were~~ poorly captured by the  
5 rectilinear approach; for example, a rectilinear niche may suggest that a species could survive  
6 in a situation where it is both hot and dry, but the actual climate niche is that it only occurs  
7 where it is hot and wet. More advanced methods have refined the  $n$ -dimensional hyperniche  
8 (Hutchinson, 1957) where response curves are used to capture the suitability of different  
9 conditions for species occurrence. ~~This revolution in the multidimensional quantification of  
10 the niche has completely bypassed the Coexistence Approach and many other nearest living  
11 relative methods.~~ BIOCLIM performed poorly in comparison to more recent methods in a  
12 comparison of more recent SDM methods (Elith et al., 2006) indicating that the simplistic use  
13 of range values for climatic variables, as used by the Coexistence Approach, is a poor  
14 representation of the realised niche of species or NLR.

15 The revolution in the multidimensional quantification of the niche has completely bypassed  
16 the Coexistence Approach. Measuring ecological niche overlap between species in  
17 multivariate space is an active area of investigation (Rödder and Engler, 2011; Broennimann  
18 et al., 2012), which can be used to determine the shared niche within a set of species.  
19 However, measuring the niche in such a manner also requires that all the variables selected  
20 are, in fact, significant in limiting the niche. Establishing the contribution and importance of  
21 different environmental variables (i.e. variable selection) in setting the bounds of a taxon's  
22 niche is a theoretical issue (Araújo and Guisan, 2006) where advances are also being made  
23 (Austin and Van Niel, 2011). In comparison, the Coexistence Approach blindly uses a wide  
24 range of environmental parameters in a univariate manner assuming that they are all important  
25 in determining a taxon's niche.

26

27 Furthermore, The assumption of niche conservatism (linked to the principle of physiological  
28 uniformitarianism) has generated considerable debate in the SDM literature as it has been  
29 used as justification for projecting models into altered climate states (past or future) and to  
30 predict the establishment and spread of invasive species (reviewed in Pearman et al., 2008a).  
31 These discussions have centred firstly on whether the current distribution for a given species,  
32 i.e. the realised niche, adequately represents the fundamental niche; and secondly, how

1 quickly the fundamental niche might be able to shift? Such concerns are absent in the  
2 theoretical underpinnings of Coexistence Approach (Mosbrugger and Utescher, 1997;  
3 Utescher et al., 2014). Unfortunately, niche shifts have been documented for a wide range of  
4 plant species [both](#) through space (Broennimann et al., 2007; Pearman et al., 2008a) and even  
5 over relatively short time scales (Pearman et al., 2008b; Veloz et al., 2011). Therefore the  
6 assumption of physiological uniformitarianism has limited applicability to reconstruct precise  
7 and accurate palaeoclimates, especially with increasing age of an assemblage.

## 8 **4 Conclusions**

9 Using best-possible climate data for modern North American woody plants, Thompson et al.  
10 (2012) were unable to reconstruct the climatic shifts from the Last Glacial Maximum to the  
11 present-day using an unweighted mutual climatic range method (which represents the niche  
12 using range values and is equivalent with the Coexistence Approach save for the use of NLRs  
13 [and recognition of climatic outliers](#)). This is in stark contrast to the beliefs of Coexistence  
14 Approach practitioners that the method can reconstruct climate shifts at high-precision,  
15 despite the additional error and uncertainty associated with the nearest-living-relative  
16 principle. The purported high precision in Coexistence Approach studies is dependent on  
17 phenomena such as pseudo-coexistence and the lack of a sound statistical framework.

18 We argue that the Coexistence Approach, as conceived by Mosbrugger and Utescher (1997),  
19 violates the basic concepts behind mutual climate range techniques and the nearest-living-  
20 relative principle. It imposes a number of assumptions that will inevitably be violated and has  
21 no ability to detect violations and generally lacks any safeguards against the reconstruction of  
22 artificial coexistence intervals and thus erroneous palaeoclimate estimates.

23 Given the theoretical problems outlined here, and the practical problems highlighted by  
24 Grimm et al. (2015) – for example, that any random real-world flora will eventually produce a  
25 “statistically significant” (according Mosbrugger and Utescher, 1997) [coexistence](#) interval  
26 with a high number (>20) of NLRs – we suggest that all palaeoclimate reconstruction studies  
27 using the Coexistence Approach be disregarded and that the palaeoassemblages be revisited  
28 with other methods and careful, well-documented, and well-investigated NLR-associations.

## 5 Where to go from here?

There are already a range of potential methods available for palaeoclimate reconstruction using plant fossils as proxies in a univariate manner that have been rarely used or recently proposed (e.g. Greenwood et al., 2005; Boyle et al., 2008; Thompson et al., 2012; Chevalier et al., 2014; Harbert and Nixon, 2015), and there are avenues ripe for exploration (e.g. Broennimann et al., 2012; Denk et al., 2013). However, all of these methods require (further) testing, and then careful, well-documented usage when reconstructing palaeoclimates. The development of the physiognomic approach (CLAMP) within the last two decades may serve as an example regarding validation, advancement, and, most importantly, documentation and transparency. The various publications demonstrate a constant thrive to reach higher precision and counter known problems (e.g. Kovach and Spicer, 1995; Herman and Spicer, 1997; Spicer et al., 2009; Yang et al., 2015; Li et al., 2016). All primary data is made freely accessible and means are implemented allowing for quick application (CLAMP online; Yang et al., 2011). CLAMP online does not only provide data, guidelines and templates for application, but also pinpoints shortcomings and ideas how to deal with them. No method is or will be 'perfect'. Nevertheless, it is crucial to define the principal accuracy and precision of any quantitative method. If this is not possible, as in the case of the Coexistence Approach (Utescher et al., 2014: 61), it must not be used. Therefore, we suggest that any current or future taxon-based method be:

1. Tested against the modern flora (e.g. Boyle et al., 2008; Thompson et al., 2012; Chevalier et al., 2014; Harbert and Nixon, 2015).
2. Tested with randomised and unlikely communities of modern flora. A robust (taxon-based) method that is to be applied to micro-, meso- and macrofossil assemblages must detect possible allochthonous elements/mixed floras.
3. First applied to the 'better-understood' palaeoclimates of the most recent past (e.g. present to the Last Glacial Maximum) and compared with available relevant proxies (e.g. Thompson et al., 2012).
4. Explored using both species-level and taxonomic levels of potential or probable nearest-living relatives (e.g. Boyle et al., 2008).
5. Examined using a jackknifing or similar procedure to ensure that results remain accurate and establish the actual precision that can be expected with fossil floras. Fossil floras will always only provide a fraction of the actual flora, and may include

1       incorrectly determined taxa. The accuracy of a result must not change due to the  
2       presence or absence of specific taxa in the assemblage, although precision can, and is  
3       likely to, decline.

4       6. Finally, tested in a stepwise fashion further and further into the past using available,  
5       well-studied and dated, more or less continuous records, such as the recently revised  
6       Icelandic record covering the last 15 million years ranging from subtropical lowland to  
7       ice age conditions (Denk et al., 2011; Denk et al., 2013)

8       After such a series of tests, the method can be considered an alternative means to reconstruct  
9       past climates for further exploration. However, the ultimate limitations of mutual climate  
10      range techniques or other nearest-living-relative methods for palaeoclimate reconstruction do  
11      not lie in the methodological framework to estimate, for example, the coexistence space, but  
12      rather the applicability of the nearest-living-relative principle. When it comes to application  
13      into the more distant past, the basic assumption of any method must be that the nearest-living-  
14      relative principle will be violated to an unknown degree. The degree of violation will likely  
15      increase with time, and may not necessarily surface during application or testing phase.  
16      Bivariate or multivariate approaches, which can tackle the problem of pseudo-coexistence  
17      (e.g. Fig. 4), will be more sensible in this respect. The capability to accurately and precisely  
18      predict palaeoclimate will not only deteriorate with increasing age, but also with  
19      compositional change of the fossil plant assemblages relative to the modern-day situation.  
20      Precise, highly sophisticated methods (e.g. Punyasena, 2008; Harbert and Nixon, 2015) or  
21      methods using few, overly precise, values to characterise the niche space of the NLR (e.g.  
22      Greenwood et al., 2005) run a higher risk of being affected by violations of the nearest-living-  
23      relative principle than methods that use semi-quantitative approximations of the niche (e.g.  
24      Thompson et al., 2012; Denk et al., 2013).

25      Taking into account all theoretical and practical issues involved, we suspect that quantitative  
26      palaeoclimate estimates at a high precision *and* accuracy is an impossible goal when the  
27      nearest-living-relative principle has to be applied. Therefore, our opinion is that method  
28      development should not focus on high (or higher) precision, as the basis of this precision is  
29      undermined as the temporal difference between fossil and NLR increases, but rather on  
30      establishing climate change trends in a robust and reproducible manner. Semi-quantitative  
31      approaches can detect such changes and may prove to be more robust (e.g. the Köppen  
32      signature approach proposed by Denk et al., 2013). Furthermore, the application of any

1 nearest-living-relative method on palaeofloras will always depend on thoughtful filtering a  
2 fossil assemblage for elements that have been shown to have a high likelihood of niche  
3 conservatism. Fossil-NLR associations must be carefully selected to ensure that the principle  
4 of physiological uniformitarianism applies, in contrast to the current practise of seemingly  
5 data-naive bulk analyses.

## 6 Acknowledgements

7 This study was funded by the Austrian Science Fund (FWF) with a grant to GWG, project  
8 number M1751-B16. AJP received support from the National Research Foundation  
9 (RCA13091944022).

## 10 References

11 Araújo, M. B., and Pearson, R. G.: Equilibrium of species' distributions with climate,  
12 Ecography, 28, 693–695, 2005.

13 Araújo, M. B., and Guisan, A.: Five (or so) challenges for species distribution modelling, J.  
14 Biogeogr., 33, 1677–1688, 2006.

15 Austin, M. P., and Van Niel, K. P.: Improving species distribution models for climate change  
16 studies: variable selection and scale, J. Biogeogr., 38, 1–8, 2011.

17 Bond, W. J., Woodward, F. I., and Midgley, G. F.: The global distribution of ecosystems in a  
18 world without fire, New Phytol., 165, 525–538, 2005.

19 Booth, T. H., Nix, H. A., Busby, J. R., and Hutchinson, M. F.: BIOCLIM: the first species  
20 distribution modelling package, its early applications and relevance to most current MaxEnt  
21 studies, Divers. Distr., 20, 1–9, 2014.

22 Boyle, B., Meyer, H. W., Enquist, B., and Salas, S.: Higher taxa as paleoecological and  
23 paleoclimatic indicators: A search for the modern analog of the Florissant fossil flora, Geol.  
24 Soc. Am. Sp. Paper, 435, 33–51, 2008.

25 Broennimann, O., Treier, U. A., Müller-Schärer, H., Thuiller, W., Peterson, A. T., and  
26 Guisan, A.: Evidence of climatic niche shift during biological invasion, Ecology Letters, 10,  
27 701–709, 2007.

28 Broennimann, O., Fitzpatrick, M. C., Pearman, P. B., Petitpierre, B., Pellissier, L., Yoccoz, N.  
29 G., Thuiller, W., Fortin, M.-J., Randin, C., Zimmermann, N. E., Graham, C. H., and Guisan,  
30 A.: Measuring ecological niche overlap from occurrence and spatial environmental data,  
31 Global Ecol. Biogeogr., 21, 481–497, 2012.

32 Chevalier, M., Cheddadi, R., and Chase, B. M.: CREST (Climate REconstruction SofTware):  
33 a probability density function (PDF)-based quantitative climate reconstruction method, Clim.  
34 Past, 10, 2081–2098, 2014.

1 Denk, T., Grímsson, F., Zetter, R., and Símonarson, L. A.: Late Cainozoic Floras of Iceland:  
2 15 Million Years of Vegetation and Climate History in the Northern North Atlantic., Topics  
3 in Geobiology, Springer, Heidelberg, New York, 854 pp., 2011.

4 Denk, T., Grimm, G. W., and Röseler, A.-K.: When field botany meets history: taxonomy of  
5 *Platanus mexicana* in Mexico, *Willdenowia*, 42, 99–115, 2012.

6 Denk, T., Grimm, G. W., Grímsson, F., and Zetter, R.: Evidence from "Köppen signatures" of  
7 fossil plant assemblages for effective heat transport of Gulf Stream to subarctic North Atlantic  
8 during Miocene cooling, *Biogeosci.*, 10, 7927–7942, 2013.

9 Elias, S. A.: Review of Palaeoentomology of the Kap København Formation, a Plio-  
10 Pleistocene Sequence in Peary Land, North Greenland by Jens Böcher, *Arctic and Alpine*  
11 *Research*, 29, 140, 1997.

12 Elias, S. A.: Mutual climatic range reconstructions of seasonal temperatures based on Late  
13 Pleistocene fossil beetle assemblages in Eastern Beringia, *Quaternary Sci. Rev.*, 20, 77–91,  
14 2001.

15 Elith, J., Graham, C. H., Anderson, R. P., Dudík, M., Ferrier, S., Guisan, A., Hijmans, R. J.,  
16 Huettmann, F., Leathwick, J. R., Lehmann, A., Li, J., Lohmann, L. G., Loiselle, B. A.,  
17 Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., Overton, J. M., Peterson, A. T.,  
18 Phillips, S. J., Richardson, K. S., Scachetti-Pereira, R., Schapire, R. E., Soberón, J., Williams,  
19 S., Wisz, M. S., and Zimmermann, N. E.: Novel methods improve prediction of species  
20 distributions from occurrence data, *Ecography*, 29, 129–151, 2006.

21 Franklin, J.: *Mapping Species Distributions: Spatial Inference and Prediction*, Cambridge  
22 University Press, Cambridge, UK, 2010.

23 Greenwood, D. R., Archibald, S. B., Mathewes, R. W., and Moss, P. T.: Fossil biotas from the  
24 Okanagan Highlands, southern British Columbia and northeastern Washington State: climates  
25 and ecosystems across an Eocene landscape, *Can. J. Earth Sci.*, 42, 167–185, 2005.

26 Grimm, G. W., and Denk, T.: Reliability and resolution of the coexistence approach — A  
27 revalidation using modern-day data, *Rev. Palaeobot. Palynol.*, 172, 33–47, 2012.

28 Grimm, G. W., Bouchal, J. M., Denk, T., and Potts, A. J.: Fables and foibles: a critical  
29 analysis of the Palaeoflora database and the Coexistence Approach for palaeoclimate  
30 reconstruction., *bioRxiv*, doi, 10.1101/016378, 2015.

31 Harbert, R. S., and Nixon, K. C.: Climate reconstruction analysis using coexistence likelihood  
32 estimation (CRACLE): A method for the estimation of climate using vegetation, *Am. J. Bot.*,  
33 102, 1277-1289, 2015.

34 Herman, A. B., and Spicer, R. A.: New quantitative palaeoclimate data for the Late  
35 Cretaceous Arctic: evidence for a warm polar ocean, *Palaeogeogr. Palaeoclimat. Palaeoecol.*,  
36 128, 227–251, 1997.

37 Hutchinson, G. E.: Concluding Remarks, *Cold Spring Harbor Symp. Quant. Biol.*, 22, 415–  
38 427, 1957.

39 Klotz, S.: Neue Methoden der Klimarekonstruktion - angewendet auf quartäre  
40 Pollensequenzen der französischen Alpen., *Tübinger Mikropaläontologische Mitteilungen*,  
41 Institut & Museum für Geologie & Paläontologie [now: Institute for Geosciences], Eberhard  
42 Karls University, Tübingen, 1999.

1 Köppen, W.: Das geographische System der Klimate., in: Handbuch der Klimatologie, Band  
2 1, Teil C., edited by: Köppen, W., and Geiger, R., Gebr. Borntraeger, Berlin, 1–44, 1936.

3 Kovach, W., and Spicer, R. A.: Canonical correspondence analysis of leaf physiognomy: a  
4 contribution to the development of a new palaeoclimatological tool, *Palaeoclimates: Data and*  
5 *Modelling*, 2, 125–138, 1995.

6 Li, S. F., Jacques, F. M. B., Spicer, R. A., Su, T., Spicer, T. E. V., Yang, J., and Zhou, Z.:  
7 Artificial neural networks reveal a high-resolution climatic signal in leaf physiognomy,  
8 *Palaeogeogr. Palaeoclimat. Palaeoecol.*, 442, 1–11, 2016.

9 Mosbrugger, V., and Utescher, T.: The coexistence approach -- a method for quantitative  
10 reconstructions of Tertiary terrestrial palaeoclimate data using plant fossils, *Palaeogeogr.*  
11 *Palaeoclimat. Palaeoecol.*, 134, 61–86, 1997.

12 Nix, H. A.: A biogeographic analysis of Australian elapid snakes, in: *Atlas of Elapid Snakes*  
13 of Australia, edited by: Longmore, R., Australian Government Publishing Service, Canberra,  
14 4–15, 1986.

15 Pearman, P. B., Guisan, A., Broennimann, O., and Randin, C. F.: Niche dynamics in space  
16 and time, *Trends in Ecology & Evolution*, 23, 149–158, 2008a.

17 Pearman, P. B., Randin, C. F., Broennimann, O., Vittoz, P., van der Knapp, W. O., Engler, R.,  
18 Le Lay, G., Zimmermann, N. E., and Guisan, A.: Prediction of plant species distributions  
19 across six millennia, *Ecology Letters*, 11, 357–369, 2008b.

20 Peterson, A. T., Soberón, J., Pearson, R. G., Anderson, R. P., Martínez-Meyer, E., Nakamura,  
21 M., and Araújo, M. B.: *Ecological Niches and Geographic Distributions*, Princeton University  
22 Press, Princeton, 2011.

23 Potts, A. J., Hedderson, T. A., Franklin, J., and Cowling, R. M.: The Last Glacial Maximum  
24 distribution of South African subtropical thicket inferred from community distribution  
25 modelling, *J. Biogeogr.*, 40, 310–322, 2013.

26 Punyasena, S. W.: Estimating Neotropical palaeotemperature and palaeoprecipitation using  
27 plant family climatic optima, *Palaeogeogr. Palaeoclimat. Palaeoecol.*, 265, 226–237, 2008.

28 Rödder, D., and Engler, J. O.: Quantitative metrics of overlaps in Grinnellian niches:  
29 advances and possible drawbacks, *Global Ecol. Biogeogr.*, 20, 915–927, 2011.

30 Schroeder, G.-F.: *Lehrbuch der Pflanzengeographie.*, Quelle & Meyer, Wiesbaden, 459 pp.,  
31 1998.

32 Sexton, J. P., McIntyre, P. J., Angert, A. L., and Rice, K. J.: Evolution and ecology of species  
33 range limits, *Ann. Rev. Ecol. Evol. Syst.*, 40, 415–436, 2009.

34 Spicer, R. A., Valdes, P. J., Spicer, T. E. V., Craggs, H. J., Srivastava, G., Mehrotra, R. C.,  
35 and Yang, J.: New development in CLAMP: calibration using global gridded meteorological  
36 data, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 283, 91–98, 2009.

37 Thompson, R. S., Anderson, K. H., Peltier, R. T., Strickland, L. E., Bartlein, P. J., and  
38 Shafer, S. L.: Quantitative estimation of climatic parameters from vegetation data in North  
39 America by the mutual climatic range technique, *Quaternary Sci. Rev.*, 51, 18–39, 2012.

40 Tiffney, B. H., and Manchester, S. R.: The use of geological and paleontological evidence in  
41 evaluating plant phylogeographic hypotheses in the Northern Hemisphere Tertiary., *Int. J.*  
42 *Plant Sci.*, 162, S3–S17, 2001.

1 Tiffney, B. H.: Phylogeography, fossils, and northern hemisphere biogeography: The role of  
2 physiological uniformitarianism., *Ann. Missouri Bot. Gard.*, 95, 135–143, 2008.

3 Utescher, T., Bruch, A. A., Erdei, B., François, I., Ivanov, D., Jacques, F. M. B., Kern, A. K.,  
4 Liu, Y.-S. C., Mosbrugger, V., and Spicer, R. A.: The Coexistence Approach—Theoretical  
5 background and practical considerations of using plant fossils for climate quantification,  
6 *Palaeogeogr. Palaeoclimat. Palaeoecol.*, 410, 58–73, 2014.

7 Veloz, S. D., Williams, J. W., Blois, J. L., He, F., Otto-Bliesner, B., and Liu, Z.: No-analog  
8 climates and shifting realized niches during the late Quaternary: implications for 21st-century  
9 predictions by species distribution models, *Global Change Biol.*, 18, 1698–1713, 2011.

10 Walter, H.: Vegetation of the Earth in relation to the climate and the eco-physiological  
11 conditions., Springer Verlag, New York, Heidelberg, Berlin, 237 pp., 1973.

12 Walter, H., and Breckle, S.-W.: *Ökologie der Erde*, Eugen Ulmer Verlag, Stuttgart, 1983–  
13 1991.

14 Yang, J., Spicer, R. A., Spicer, T. E. V., and Li, C.-S.: ‘CLAMP Online’: a new web-based  
15 palaeoclimate tool and its application to the terrestrial Paleogene and Neogene of North  
16 America, *Palaeobiodiversity and Palaeoenvironments*, 91, 163–183, 2011.

17 Yang, J., Spicer, R. A., Spicer, T. E. V., Arens, N. C., Jacques, F. M. B., Su, T., Kennedy, E.  
18 M., Herman, A. B., Steart, D. C., Srivastava, G., Mehrotra, R. C., Valdes, P. J., Mehrotra, N.  
19 C., Zhou, Z. K., and Lai, J. S.: Leaf Form-Climate Relationships on the Global Stage: An  
20 Ensemble of Characters, *Global Ecol. Biogeogr.*, 10, 1113–1125, 2015.

21

22 **Figure legends**

23 **Figure 1. The concept of the mutual climate range as used in the Coexistence Approach.**

24 **Figure 2. Difference between ‘systematically close’ and nearest-living-relative (NLR, i.e.  
25 best modern analogues).** Shown is a species phylogeny of a diversified ingroup; the  
26 outgroup in this example is a sister species of the ingroup. **A.** Standard definition of  
27 nearest-living relative (best modern analogue) vs. definition if Assumption 1 of the  
28 Coexistence Approach should be fulfilled. **B.** Same tree as in A, only that each species is  
29 categorised as a member of a distinct morphotaxon that can be distinguished in the fossil  
30 record. Note that all morphotaxa are mutually exclusive regarding their climatic niche,  
31 but there is no strict correlation between systematic closeness (phylogeny) and the  
32 climatic niche of the fossils and their nearest-living relatives (modern species of the same  
33 morphotypes as the fossils).

34 **Figure 3. Issues related to the use higher-level taxonomic classification (e.g. genus or  
35 family) as nearest-living relatives (NLRs) of fossil species.** In this example, two fossil  
36 species occupy a climate range within the modern climate range of their selected genus-  
37 level NLRs, fulfilling the principle of physiological uniformitarianism. **A.** The fossil  
38 species have a narrow shared climate range and coexisted in the past. The use of higher-  
39 level taxonomic ranks as NLRs will lead, in most cases, to a much broader and less  
40 precise reconstructed coexistence interval. **B.** The fossil species are mutually exclusive,

1 but the expansion of the niche space – due to the use of genera as NLRs – results in a  
2 coexistence interval (i.e. pseudo-coexistence).

3 **Figure 4. Pseudo-coexistence as a result of the representation of the climate niche using**  
4 **minimum and maximum tolerances.** **A.** Bivariate climate niches of two mutually  
5 exclusive species. These species have no overlapping climate space, but still reconstruct  
6 narrow coexistence intervals (orange bars) along univariate axes. **B.** Bivariate climate  
7 niches of NLRs of two floras growing under substantially different climates (indicated  
8 “x”s). Note that only the niches of three of the Community 1 species overlap with one or  
9 two of the Community 2 species. **C, D.** Univariate mutual climate ranges (MCR) of both  
10 communities; the overlap of the two MCR result in highly precise coexistence intervals  
11 for the artificially mixed communities including all elements from Community 1 and  
12 Community 2.

13 **Figure 5. “Climatic outliers” and the bias of the Coexistence Approach towards exotic**  
14 **nearest-living relatives (NLRs).** Shown are the niche response curves for 20 potential  
15 NLRs, of which 18 (grey and green) show a general overlap in their climatic preference.  
16 The two red NLRs are exotic elements with strongly differing climatic preferences. Bars  
17 indicate the minimum and maximum tolerances of each NLR, the dots highlight each  
18 NLR’s optimal climate value. Because the green NLR has no shared climate range with  
19 the two exotic NLRs (red), it would be excluded as a “climatic outlier” following the  
20 Coexistence Approach protocol. The resultant coexistence interval (orange bar) is highly  
21 precise but reflects neither the climatic preference of the non-exotic (grey and green) nor  
22 exotic group of NLRs (red).

23 **Figure 6. Logical inconsistencies in the application and theory of the Coexistence**  
24 **Approach.** Shown are coexistence intervals (orange bars) based on slightly (**A, C**) or  
25 extremely (**B**) different sets of nearest-living relatives (NLRs). **A.** Use of the “center  
26 value” to determine climate shifts. A single NLR (black) is replaced by a NLR tolerant to  
27 *lower* values (red), which would be eliminated as a “climatic outlier” by the two green  
28 NLRs; thus leading to a *higher* “center value”: **B.** All NLRs have contrasting climate  
29 tolerances, the exotic taxa in both floras (red) ensure that the reconstructed coexistence  
30 interval is the same. **C.** Two floras that only differ by the absence (white bars) or presence  
31 (black bars) of each a single taxon. The resulting coexistence intervals would be  
32 interpreted as a shift towards higher values. The green box shows the coexistence interval  
33 of a flora in which both taxa are represented.

34 **Figure 7. Impossibility of reconstructing extinct climates with the nearest-living-relative**  
35 **(NLR) principle.** Shown are the (realised) climate niches of five modern species, which,  
36 inevitably have to lie within the frame of the modern climate space. Any coexistence  
37 space (yellow square, showing the coexistence space of species 2, 3, and 4 using their  
38 minimum and maximum tolerances) must reflect a climate situation also found today.  
39 Any extinct climate (grey square) could only be defined by the coexistence of species  
40 with *different* climate niches than found in modern species, species with no living NLR or  
41 species belonging to lineages that underwent niche shift.

1 **Table 1. The assumptions of the Coexistence Approach (quotations from Utescher *et al.*,**  
 2 **2014)**

Description	Issues
<b>Assumption 1:</b> “ <i>For fossil taxa systematically closely related nearest living relatives (NLRs) can be identified.</i> ”	<ul style="list-style-type: none"> <li>a) Lack of a theoretical framework to define a systematically close relative</li> <li>b) Concept of physiological uniformitarianism assumes a common origin, but does not need quantification of phylogenetic closeness</li> </ul>
<b>Assumption 2:</b> “ <i>The climatic requirements of a fossil taxon are similar to those of its nearest living relative.</i> ”	<ul style="list-style-type: none"> <li>a) Physiological uniformitarianism cannot be generally assumed</li> <li>b) Different taxonomic ranks of fossils and their nearest-living relatives</li> </ul>
<b>Assumption 3:</b> “ <i>The climatic requirements or tolerances of a nearest living relative [i.e. minimum and maximum tolerances regarding single parameters that are considered per se to be independent from each other] can be derived from its [current] area of distribution</i> ”	<ul style="list-style-type: none"> <li>a) Distribution is not necessarily a function of climate, but also other biotic and abiotic parameters: the realised niche &lt; fundamental niche</li> <li>b) Minimum and maximum tolerances are poor estimates for the climatic niche of a taxon</li> <li>c) Climate parameters are not independent from each other</li> <li>d) There are no working frameworks to test if a potential nearest-living-relative fulfils <b>Assumption 3</b></li> </ul>
<b>Assumption 4:</b> “ <i>The modern climatic data used are reliable and of good quality</i> ”	More or less violated in all studies that applied the Coexistence Approach (see Grimm & Denk, 2012)
<b>Assumption 5:</b> Palaeoassemblages represent actual communities	<ul style="list-style-type: none"> <li>a) Fossils may be allochthonous, in particular microfossils (pollen).</li> <li>b) Fossils may not be strictly coeval (macrofossil lagerstätten usually cover substantial time periods)</li> </ul>
<b>Assumption 6:</b> Absence of a fossil in a palaeoassemblage is evidence of true absence	The fossil record is incomplete

1 **Table 2: The consequences of identifying a ‘climatic outlier’ in a palaeoassemblage**  
 2 **supposing that these represent violations of one or more of the four basic**  
 3 **assumptions.**

4

<b>Violation of basic assumption...</b>	<b>Consequence</b>
... 1: The nearest-living-relative (NLR) is not a close relative of a fossil taxon	There is no consequence as long as the NLR shares the same lineage and is a good physiological modern analogue.
... 2: The climatic requirements of the fossil taxon are different from the NLR	If different for one climate parameter, the NLR may be equally non-representative for other climate parameters of the fossil taxon. Any coexistence interval including this fossil taxon may be misinformed.
... 3: The NLR’s distribution is not representative for its climatic requirements (relict distribution)	Coexistence intervals delimited by the NLR are likely to be misinformed in any study using the NLR.
... 4: The modern climate data to estimate NLR minimum and maximum tolerances are unreliable.	If this is the case, then no coexistence interval is reliable and palaeoclimate reconstruction using modern analogues is impossible.

5