

References

- 110 Engelstaedter, S. and Washington, R.: Atmospheric controls on the annual cycle of North African dust, *Journal of Geophysical Research: Atmospheres*, 112, doi:10.1029/2006JD007195, <http://dx.doi.org/10.1029/2006JD007195>, 2007.
- Hagemann, S.: An improved land surface parameter dataset for global and regional climate models, MPI Report No. 289, Max Planck Institute for Meteorology, Hamburg, 2002.
- 115 Hoelzmann, P., Jolly, D., Harrison, S. P., Laarif, F., Bonnefille, R., and Pachur, H.-J.: Mid-Holocene land-surface conditions in northern Africa and the Arabian Peninsula: A data set for the analysis of biogeophysical feedbacks in the climate system, *Global Biogeochemical Cycles*, 12, 35–51, doi:10.1029/97GB02733, <http://dx.doi.org/10.1029/97GB02733>, 1998.
- Middleton, N. J. and Goudie, A. S.: Saharan dust: sources and trajectories, *Transactions of the Institute of British Geographers*, 26, 165–181, doi:10.1111/1475-5661.00013, <http://dx.doi.org/10.1111/1475-5661.00013>, 2001.
- 120 Stanelle, T., Bey, I., Raddatz, T., Reick, C., and Tegen, I.: Anthropogenically induced changes in twentieth century mineral dust burden and the associated impact on radiative forcing, *Journal of Geophysical Research: Atmospheres*, 119, 13,526–13,546, doi:10.1002/2014JD022062, <http://dx.doi.org/10.1002/2014JD022062>, 2014.
- 125 Tegen, I., Harrison, S. P., Kohfeld, K., Prentice, I. C., Coe, M., and Heimann, M.: Impact of vegetation and preferential source areas on global dust aerosol: Results from a model study, *Journal of Geophysical Research: Atmospheres*, 107, AAC 14–1–AAC 14–27, doi:10.1029/2001JD000963, <http://dx.doi.org/10.1029/2001JD000963>, 2002.