

Interactive comment on “A 60 000 year Greenland stratigraphic ice core chronology” by K. K. Andersen et al.

Anonymous Referee #2

Received and published: 19 December 2007

This paper presents an extension of the GIC005 ice core chronology from 42 to 60 kyr, based on layer counting of the NGRIP ice core. Since this chronology has been adopted as a reference for many other studies, the extension is a welcome addition, for which the authors are to be complimented.

While the layer counting is labelled as "multi-parameter", it appears to rely most heavily on the visual stratigraphic record. As a non-specialist in this area, my reaction to the examples shown in Figs 1 and 2 is very similar to that given in the comments of E. Wolff and D. Genty, namely it is very difficult to understand the objective basis on which layers are considered "uncertain". Since, as emphasized by E. Wolff, it is on this basis that the uncertainties on the time scale are estimated, this is obviously a critical point. The authors give two justifications for their uncertainty estimates. One,

Interactive
Comment

the agreement between two independent observers for the entire section (although, as the authors note, locally this difference can exceed the estimated uncertainty). As the authors also note, such a procedure cannot exclude systematic bias. To their credit, the authors are very open about these difficulties, and it is clear that they are not trying to hide anything. The question remains, nevertheless, as to whether such error estimates are quantitatively robust.

The second justification relies on the fact that the obtained chronology agrees with independently dated records within the estimated uncertainties. Since this is potentially a less subjective criteria, it is important to look closely at these comparisons, which are summarized in Table 2 and figs 4-7. When one does this, it can be noted that in fact the majority of the reference dates used for comparison are on layers younger than the time range of the extension reported in the present paper. While these comparisons are useful for supporting the general approach and reliability of the previously published range of GIC005, they are unfortunately not very helpful for evaluating the accuracy of the extended range. I therefore concentrate here on this extended time range. In Table 2, only one of the seven cited reference levels (the NAAZ II volcanic layer) falls in the extended portion of the time scale. The Ar/Ar date cited for this ash layer is 54.5 ± 1.0 ky, which is well within the uncertainty of the GIC005 age. However, when I looked at the cited reference, I found that it is in fact a secondary reference in which this date is given as a "private communication". In fact, it is not even clear to me whether this is actually a "recent" determination, as described, or the same as that given in a 1998 abstract (AGU Spring Meeting Suppl. 79, S377), having the same value, and one of the same authors as the "private communication".

The remaining comparisons are with radiometrically dated cave deposits. These involve not direct dating of events observed in the ice, but rather correlating climate records from the two reservoirs. The authors argue that, on the time scales considered, these climatic records can be assumed to be synchronous. I believe this is a reasonable assumption, although one can also find arguments why this might not be

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

strictly the case. Also, because speleotherms are not dated continuously, and have inferior resolution, there is some subjectivity about placing the control points. The most comprehensive comparison involves the Hulu Cave, where five well defined climatic features, fairly uniformly distributed over the extended time range, are compared. The agreement is within the estimated ice core uncertainties for all these features. Three climatic features from the Klee gruben Cave also appear to be in good agreement although, as the authors point out, agreement of features toward each end of that record are less obvious. With regard to this comparison, I think it would be useful for the authors to include the assumed correlation lines in Fig 6, in the same way as fig 5. Also shown is a comparison with a stalagmite profile from Moomi Cave, in which only one of three features falls within the estimated uncertainties. Although the climate records are not shown, the authors also claim satisfactory agreement with individual features from several other absolutely dated speleotherm records. Overall, if one accepts the critical assumption that the climate records from these two types of reservoirs are synchronous, the comparisons with the absolutely dated cave deposits are quite convincing.

Finally, there is one other important aspect of this chronology, and that is the accuracy of relatively short duration features. The details of such features is becoming more and more important, and one of the advantages of a continuous chronology, compared to one simply interpolated between reference levels, is that in principle it should give much better estimates of the duration of such features. This is nicely illustrated by the authors when they compare their chronology with that of Meese et al. As the authors note, although these two time scales agree within uncertainties at the beginning and end of the extended time region, they actually disagree by 20 % locally in between. In ice core studies themselves, paleo-accumulation rates deduced from continuous chronologies are often used to transform concentrations of trace species into fluxes. This local chronological information is in fact given implicitly by the annual layer thickness curve shown fig 3. If I understand correctly, the authors estimate that a maximum counting error (MCE) of 5 % (2 sigma) should be associated with this curve between 41.8-60

kyr. It should be noted that the method of comparison with independently dated control points, as shown in fig 4, does not really test this uncertainty. I believe it would be very useful to give a supplementary table with a numerical version of this curve.

There is in fact an aspect of this curve which I find puzzling. As pointed out by the authors, a widely used model chronology for GRIP and NGRIP is ss09sea. This model is based on an estimate of initial accumulation, coupled with a thinning function. The accumulation is related to the 18O record via a semi-empirical correlation. For nearby features, where the thinning is relatively constant, one expects changes in the layer thickness to be dominated by changes in the estimated accumulation. This implies that the layer thicknesses will be proportional to 18O. If one compares GI 10 and 11, however, one finds that GI 10 has larger amplitude in layer thickness than GI 11, but smaller amplitude in 18O. A similar situation occurs for GI 6 and 7. This suggests to me that either there is a problem with the model predictions of the layer thicknesses in these particular regions, or a problem with the layer counting. I think it might be quite useful to include the layer thicknesses of ss09sea in fig 3, so that one could see where the major differences between the modelled and observed thicknesses occur. Such information might be helpful in identifying the cause of the differences, as well as giving added confidence in regions where the two agree.

Interactive comment on Clim. Past Discuss., 3, 1235, 2007.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)