127

Clim. Past Discuss., 5, 127–151, 2009 www.clim-past-discuss.net/5/127/2009/ © Author(s) 2009. This work is distributed under the Creative Commons Attribution 3.0 License.

Climate of the Past Discussions is the access reviewed discussion forum of Climate of the Past

Late Glacial and Holocene changes in vegetation cover and climate in southern Siberia derived from a 15 kyr long pollen record from Lake Kotokel

P. E. Tarasov¹, E. V. Bezrukova^{2,3}, and S. K. Krivonogov⁴

¹Freie Universität Berlin, Geological Sciences, Palaeontology Section, Berlin, Germany ²Institute of Archaeology and Ethnography, Russian Academy of Sciences, Siberian Branch, Novosibirsk, Russia

³Institute of Geochemistry, Russian Academy of Sciences, Siberian Branch, Irkutsk, Russia ⁴United Institute of Geology, Geophysics and Mineralogy, Russian Academy of Sciences, Siberian Branch, Novosibirsk, Russia

Received: 21 October 2008 - Accepted: 28 October 2008 - Published: 14 January 2009

Correspondence to: P. E. Tarasov (ptarasov@zedat.fu-berlin.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.





Abstract

In this study a radiocarbon-dated pollen record from Lake Kotokel (52°47' N, 108°07' E. 458 m a.s.l.) located in southern Siberia east of Lake Baikal was used to derive quantitative characteristics of regional vegetation and climate since about 15 kyr BP (1 kyr=1000 cal. yr) until today. Quantitative reconstruction of the late glacial vegeta-5 tion and climate dynamics suggests that open steppe and tundra communities predominated in the study area prior to ca. 13.5 kyr BP and again during the Younger Dryas interval, between 12.8 and 11.6 kyr BP. The pollen-based climate reconstruction suggests lower-than-present mean January (~-38°C) and July (~12°C) temperatures and annual precipitation (~270-300 mm) values during these time intervals. Boreal 10 woodland replaced the primarily open landscape around Kotokel three times at about 14.8–14.7 kyr BP, during the Allerød Interstadial between 13.3–12.8 kyr BP and with the onset of the Holocene interglacial between 11.5 and 10.5 kyr BP, presumably in response to a noticeable increase in precipitation, and in July and January temperatures. The maximal spread of the boreal forest (taiga) communities in the region is 15 associated with a warmer and wetter-than-present climate ($T_w \sim 17-18^{\circ}$ C, $T_c \sim -19^{\circ}$ C, P_{ann} ~500–550 mm) occurred ca. 10.8–7.3 kyr BP. During this time interval woody vegetation covered more than 50% of the area within a 21×21 km window around the lake.

The pollen-based best modern analogue reconstruction suggests a decrease in woody cover percentages and in all climatic variables about 7–6.5 kyr BP. Since that time our results demonstrate gradual decrease in precipitation and mean January temperature towards their present-day values in the region around Lake Kotokel.

1 Introduction

The Lake Baikal region of northern Eurasia (Fig. 1a) experienced a boom of palaeoenvironmental studies during past decade (e.g. Colman et al., 1996; BDP-Members, 1997; 1998; 2005; Grachev et al., 1997; Williams et al., 1997; 2001; Demske et al., 5, 127–151, 2009

Vegetation and climate in southern Siberia since 15 kyr BP



2005; Oberhänsli and Mackay, 2005; Tarasov et al., 2007a and references therein). A rising interest to Lake Baikal – the world's largest, deepest and oldest freshwater reservoir – is easy to understand. The lake bottom sediments contain detailed and well preserved palaeoenvironmetal archives, which provide an excellent opportunity

for the reconstructions of the regional climate and environments (e.g. Bezrukova 1999; Bezrukova et al., 2005; Horiuchi et al., 2000; Khursevich et al., 2001; Prokopenko and Williams 2004; Tarasov et al. 2005).

Objective reconstruction of the past climate is one of the priority tasks for the scientists, working in the field of the past global changes and the Earth's system modeling (PAGES: http://www.pages.unibe.ch/). Since 1940s (e.g. Iversen, 1944) late Quater-

- (PAGES: http://www.pages.unibe.ch/). Since 1940s (e.g. Iversen, 1944) late Quaternary pollen records from the lake sediments became a frequently used proxy providing palaeoclimatic information at local and large regional scale (e.g. Grichuk, 1969; Bartlein et al., 1984; Guiot et al., 1989; Nakagawa et al., 2002; Frenzel et al., 1992; Seppä and Birks, 2001, 2002; Tarasov et al., 2007a and references therein).
- ¹⁵ Despite a relatively high number of the late glacial and Holocene pollen records generated for the Lake Baikal region, there have been very few attempts at their quantitative palaeoclimatic interpretation. The earlier pollen-based climate reconstructions were either solely qualitative (e.g. Khotinskii 1984; Bezrukova 1999; Prentice and Jolly, 2000; Demske et al., 2005) or concerned with the "Holocene optimum" time slice (e.g.
- ²⁰ Frenzel et al., 1992; Tarasov et al., 1999). For the first time Holocene changes in annual precipitation (P_{ann}), mean temperature of the warmest (T_w) and coldest (T_c) month and moisture index (α) were reconstructed from continuous and adequately dated pollen records recovered from the underwater Buguldeika site (Fig. 1b) in the southern part of Lake Baikal (Tarasov et al., 2007a). The latter results from Lake Baikal likely represent
- vegetation and climate dynamics in the relatively large region. However, pollen records from different parts of the lake and from surrounding coastal plains demonstrate spatial variations in the vegetation dynamics within the Lake Baikal region (e.g. Tarasov et al., 2002; Demske et al., 2005; Bezrukova et al., 2005), pointing out the need for more reconstructions from the region.

5, 127-151, 2009 Vegetation and climate in southern Siberia since 15 kyr BP P. E. Tarasov et al. **Title Page** Introduction Abstract Conclusions References Figures Tables Back Close Full Screen / Esc Printer-friendly Version Interactive Discussion



In the recent study we present the results of the quantitative reconstruction of the temperature and precipitation variables derived from the Lake Kotokel pollen record collected in 2004 (Bezrukova et al., 2008). The lake is situated at the eastern shore of Lake Baikal (Fig. 1) and is representative for the middle part of the Lake Baikal region. ⁵ The reconstructed climate variables since about 15 kyr BP (1 kyr=1000 cal. yr) until today are compared with the quantitative characteristics of vegetation around Kotokel, including dominant biome scores and woody cover percentages derived from the same pollen record. The regional environmental dynamics is then discussed together with the oxygen isotope records of the post glacial climate from the North Atlantic and North Pacific regions.

2 Regional setting

15

Kotokel (458 m a.s.l.) is a fresh-water lake occupying western part of the intermountain depression located at the eastern coast of Lake Baikal (Fig. 1b). A low-elevated mountain ridge (up to 729 m a.s.l.) separates the depression from Lake Baikal and the Ulan-Burgasy Ridge (up to 2033 m a.s.l.) bounds it from the east (Galaziy, 1993). The lake has an area of about 67 km², a maximum length of about 15 km and maximum width of about 5 km. An average water depth is 5–6 m and a maximum depth is about 15 m (see Tarasov et al., 1994 for more details and references).

The area has continental climate with long cold winters and relatively short cool summers and large seasonal variations in temperature and precipitation (e.g. Alpat'ev et al., 1976; Galaziy, 1993). Around Kotokel the mean January temperature is about -20°C and decreases to below -26°C with increase in elevation. The mean July temperature is about 16°C and annual precipitation sums vary from 400 mm in the coastal zone to above 500 mm in the upper elevation belt (Galaziy, 1993). Modern precipitation 25 distribution has well pronounced summer maximum. July and August are particularly

wet. During these months westerly winds dominating through the year become weak and south-eastern cyclones bring to the region warm and wet Pacific air and cause





heavy rainfalls at the eastern branch of the Polar Front (Bezrukova et al., 2008). The precipitation associated with the Atlantic air masses brought by the westerly winds is not abundant and mainly come during autumn and spring. Dry, cold and sunny weather typically occurs in winter, when the whole region is controlled by the stationary Siberian Anticyclone.

Modern vegetation at the eastern coast of Lake Baikal and around Kotokel is mainly composed of boreal coniferous and deciduous forests. Swampy vegetation is common in the Selenga River delta and south of Kotokel' Lake. Forests are composed of *Pinus sylvestris* (Scots pine), *Larix sibirica* (Siberian larch) and *Betula* (birch) species, with some admixture of *Populus tremula* (aspen) and shrubby alder *Alnus fruticosa* (Galaziy, 1993). Taiga forests composed of mainly boreal evergreen conifers (e.g. *Pinus sibirica* – Siberian pine, *Abies sibirica* – Siberian fir and *Picea obovata* – Siberian spruce) appear several kilometres eastward at the slopes of the Ulan-Burgasy Ridge. In the upper mountain belt open birch and larch forests coexist with *Pinus pumila* (shrubby

¹⁵ pine) and alpine tundra vegetation (Molozhnikov, 1986; Galaziy, 1993).

3 Material and methods

10

3.1 Coring, core lithology and age determination

In the early 1960s the Lake Kotokel Holocene sediments were first studied for pollen and non-pollen microfossils (Korde, 1968; Vipper and Smirnov, 1979; Tarasov et al., 1994). In 1997 a near-shore Cheremushka peat bog was drilled (Takahara et al., 2000). The KTK1 sediment core (52°47′ N, 108°07′ E) was recovered in July 2004 (Bezrukova et al., 2008; Shichi et al., 2008). The coring was performed from the water depth of about 3.5 m in the southern part of the lake (Fig. 1c) using Livingston piston core. A detail pollen analysis was performed on the upper 900 cm of the KTK1 core composed of the soft brownish black gyttja (0–810 cm) and slightly laminated gray clay (Bezrukova et al., 2008; Shichi et al., 2008).



Three radiocarbon dates (e.g. 6070±60, 10.680±40 and 11.670±60¹⁴C yr BP) were obtained from the brownish black gyttja unit (e.g. from 412–413 cm, 652–653 cm and 806–807 cm depth, respectively) using the AMS facility at Nagoya University in Japan suggesting that the analyzed 900 cm of sediments accumulated since about 15 kyr BP (see Bezrukova et al., 2008; Shichi et al., 2008 for details). Radiocarbon years were converted to calendar years using the CalPal program available online (Danzeglocke et al., 2008). The resulting age-depth model represents a linear regression, suggests rather constant sedimentation rate of 0.59 mm/yr along the profile, in line with the undisturbed character of the sediment. This model was applied to the KTK1 pollen record and used to date reconstructed changes in vegetation and climate discussed in the text.

3.2 Pollen analysis

25

The KTK1 core material stored at Institute of Geochemistry (Irkutsk) was sampled there for pollen analysis. In total 128 samples were taken as 1 cm slices, yielding an average temporal resolution of 118 years throughout the late glacial and Holocene period. Standard laboratory methods were used to extract pollen from the sediment samples, including HCl and KOH treatments, heavy-liquid separation and following acetolysis (Berglund and Ralska-Jasiewiczowa, 1986). Pollen and spores mounted in glycerin were counted under the light microscope with ×400–1000 magnification.
²⁰ Identification of the pollen and spores was performed using regional pollen atlases and the reference collection (Bezrukova et al., 2008).

Samples were generally rich in pollen and counting of 300 to 500 pollen grains per sample was easy to achieve (Fig. 2). However, the pollen content was lower (176–240 grains per sample) in the bottommost 3 samples. The quantitative climate reconstruction results from these samples need to be taken with caution.

The percentage diagram presented in Fig. 2 shows results of the pollen analysis from the KTK1 core. The diagram was constructed using the Tilia/Tilia-Graph/TGView

5, 127-151, 2009 Vegetation and climate in southern Siberia since 15 kyr BP P. E. Tarasov et al. **Title Page** Introduction Abstract Conclusions References Figures Tables Back Close Full Screen / Esc Printer-friendly Version Interactive Discussion



software (Grimm, 1993, 2004). Percentages for individual terrestrial pollen taxa at each level were calculated from the total sum of arboreal and terrestrial non-arboreal pollen taken as 100%. Spore percentages for cryptogam plants were calculated in relation to the total sum of counted pollen and spores. In order to facilitate discussion of the pollen record, the pollen diagram was subdivided into local pollen zones (PZ) based on square-root-transformation of the percentage data and stratigraphically constrained cluster analysis by the method of incremental sum of squares (Grimm, 1987).

3.3 Pollen-based vegetation and climate reconstruction

5

25

The quantitative method of biome reconstruction (Prentice et al., 1996) adapted for
reconstruction of vegetation in northern Eurasia (Tarasov et al., 1998) allows the objective assignment of pollen taxa to plant functional types (PFTs) and to biomes on the basis of the modern ecology, bioclimatic tolerance and geographical distribution of pollen producing plants. The method was successfully tested using an extensive surface pollen data set from northern Eurasia and used to reconstruct Last Glacial
Maximum (LGM) and mid-Holocene vegetation at the hemispheric (e.g. Prentice and Jolly, 2000) and large regional (e.g. Tarasov et al., 2000) scale. In the Lake Baikal region region (Fig. 1b) the method was applied to the Eemian pollen record from Continent (Tarasov et al., 2005) and to the Holocene record from Buguldeika (Tarasov et al., 2007a). In this study all identified terrestrial pollen taxa from the KTK1 record were assigned to the regional PFTs and biomes using the biome-PFT-taxon matrix published in the latter studies (see Tarasov et al., 2005; 2007a for further details).

The "biomization" provides no quantitative information about vegetation composition or structure and can mask temporal variations in the internal structure and composition of biomes (Williams et al., 2004). To get this important information out of the fossil pollen record we used a new approach for woody cover reconstruction (Tarasov et al., 2007b). The method combines extensive modern surface pollen and satellite-

based Advanced Very High Resolution Radiometer (AVHRR) datasets from northern Eurasia (DeFries et al., 1999; Tarasov et al., 2007b) with the best modern analogue



(BMA) approach (Overpeck et al., 1985; Guiot, 1990), allowing fossil pollen samples attribution to the vegetation characteristics associated with their closest modern pollen analogues. In this study the AVHRR-based estimates of woody cover percentages within a 21×21 km window around pollen sampling sites were attributed to the KTK1
 ⁵ pollen spectra (see Tarasov et al., 2007b for the method evaluation and design).

The BMA approach is frequently employed to infer past climates from fossil pollen assemblages (e.g. Guiot, 1990; Nakagawa et al., 2002). Recently it was used in the Lake Baikal region to reconstruct climate dynamics during the last and the recent interglacial (Tarasov et al., 2005, 2007a). In the present study we used the reference data set of 10 1173 modern pollen spectra from the large area of former Soviet Union and Mongolia with all main bioclimatic regions well represented (see Tarasov et al., 2005 for details) to reconstruct changes in annual precipitation, and in July and January (warmest and

coldest month) mean temperature (e.g. P_{ann} , T_w and T_c). All selected climatic variables are important to explain the spatial distribution of the main vegetation types in north-

- ern Eurasia and are commonly derived from fossil records and simulated with climate models (e.g. Kageyama et al., 2001; Battarbee et al., 2004). All terrestrial pollen taxa identified in the KTK1 record also appear in the list of 81 taxa presented in the reference pollen/climate data set. Modern climate values at each of the 1173 modern pollen sampling sites have been calculated from the high-resolution global climatol-
- ²⁰ ogy database that provides the 30-year average (1961–1990) of the monthly means of principal meteorological parameters on a 10 min grid (New et al., 2002).

4 Results

4.1 KTK1 pollen record and vegetation dynamics

Complete results of the pollen analysis and reconstructed vegetation and climate characteristics will be available in the PANGAEA data information system (http://www. pangaea.de). The simplified pollen percentage diagram is shown in Fig. 2. The re-



sults of the pollen analysis used in this study, including conventional description of the pollen zones and their qualitative palaeoenvironmental interpretation, were presented by Bezrukova et al. (2008). In order to avoid repetitions with the latter publication, we concentrate in this chapter on the quantitative interpretation of the KTK pollen data in terms of the late glacial and Holocene vegetation (Fig. 3) and climate dynamics (Fig. 4).

The pollen assemblages prior to 14.5 kyr BP (lower part of PZ KTK1-5) are characterised by highest percentages of herbaceous pollen taxa (e.g. *Artemisia*, Poaceae and Cyperaceae) and low percentages of tree pollen taxa (Fig. 2), pointing to an open character of vegetation around the lake. Pollen of shrubby birches and willows is relatively abundant. Tundra and steppe biomes have highest scores (Fig. 3b), suggesting that mainly these two vegetation types occupied the region about 15–14.5 kyr BP (Fig. 3c). The BMA reconstruction demonstrates very low woody cover percentages (Fig. 3d). However, woody cover rises to ca. 25% at one level dated to ca. 14.7–14.8 kyr BP, which is characterized by the relatively high amount of pine pollen.

Herbaceous pollen percentages show decline and shrubby taxa (e.g. birch, alder, willow and heath) predominate in the pollen assemblages between 14.5 and 13.4 kyr BP (upper part of PZ KTK1-5 and lower part of PZ KTK1-4). The biome reconstruction demonstrates that the tundra scores became higher than those of the steppe and taiga, suggesting reduction of the cold steppe communities and spread of the shrub tundra
 communities in the region. Low amounts of arboreal pollen (Fig. 2) and low woody

cover percentages (less than 10%) suggest that the landscape was open and scarce boreal trees could grow only in the locally favorable environments.

A sharp increase in spruce pollen percentages up to 54–68% is the characteristic feature of the pollen assemblages dated to ca. 13.3–12.8 kyr BP (middle part of

PZ KTK1-4). The biome reconstruction shows that numerical scores of taiga biome become slightly higher than those of tundra (Fig. 3b), suggesting that the vegetation around Kotokel turned to boreal woodland or forest-steppe (Fig. 3c). Consistently, vegetation cover reconstruction demonstrates increase in the total woody cover up to about 25% at that time (Fig. 3d).



Pollen of alder and birch shrubs once more become a dominant component of the pollen assemblages between ca. 12.7 and 11.5 kyr BP (upper part of PZ KTK1-4 and lower part of PZ KTK1-3). This time interval is also noticeable for highest scores of tundra biome. However, numerical scores of taiga remain higher than steppe in contrast to the time period prior to 14.5 kyr BP. The reconstructed total woody cover is about 10%,

suggesting that forests still occupied a limited area in the region.

After ca. 11.5 kyr BP pollen of boreal trees predominate in the pollen assemblages (Fig. 2). Results of the biome reconstruction demonstrate almost equally high scores for the tundra and taiga biomes (Fig. 3b), suggesting boreal woodland vegetation be-

- ¹⁰ tween 11.5 and 10.5 kyr BP (upper part of PZ KTK1-3). Since about 10.5 kyr BP the taiga biome scores are noticeably higher than the scores of non-arboreal biomes, suggesting well established boreal forest vegetation in the region. The pollen data show that birch tree pollen was relatively abundant in the pollen spectra between ca. 10.5 and 6.8 kyr BP (PZ KTK1-2) and that pine pollen went at the first place after that time (PZ KTK1-1). This abange in the pollen composition likely reflects abange in the form.
- (PZ KTK1-1). This change in the pollen composition likely reflects change in the forest communities around the lake and spread of Scots pine in the study region after 7 kyr BP. The results of the vegetation cover reconstruction suggest that total woody cover increase to above 50% after 10.5 kyr BP and decreased to present-day level of about 45% after 6.8 kyr BP.
- 20 4.2 Climate reconstruction

5

The results of the pollen-based quantitative climate reconstruction (Fig. 4a–c) suggest pronounced changes in the main climatic variables during the late glacial and Holocene, which can be summarized as follows. Prior to ca. 11.5 kyr BP climate conditions were rather unstable. The first relatively warm episode ($T_w \sim 13^\circ$ C, $T_c \sim -25^\circ$ C) occurred about 14.5 kyr BP followed by the cold episode ($T_w \sim 12^\circ$ C, $T_c \sim -38^\circ$ C) centered around ca. 14 kyr BP. The reconstructed precipitation sums were substantially lower-than-present (e.g. $P_{ann} \sim 270$ mm). The next warming occurred between 13.5 and 13 kyr BP. At that time both T_w and T_c reached modern levels, but P_{ann} remained slightly



lower-than-present, e.g. ~400 mm. The interval around 12.5–12 kyr BP was characterized by significantly lower-than-present temperatures and precipitation sums (e.g. $T_{W} \sim 12^{\circ}$ C, $T_{c} \sim -38^{\circ}$ C, $P_{ann} \sim 300$ mm). A shift towards warmer and wetter climate occurred after ca. 12 kyr BP. However, this gradual trend was interrupted by a smaller scale cold and dry oscillation dated to ca. 11.1 kyr BP. The onset of a warmer and wetter-than-present climate ($T_{W} \sim 17-18^{\circ}$ C, $T_{c} \sim -19^{\circ}$ C, $P_{ann} \sim 500-550$ mm) occurred ca. 10.8–7.3 kyr BP. The climate reconstruction shows a decrease in all climatic variables about 7–6.5 kyr BP (Fig. 4a–c). Since that time the climate reconstruction suggests gradual decrease in T_{c} , (Fig. 4c) and P_{ann} (Fig. 4a) towards present-day values in the region around Lake Kotokel. The T_{W} curve (Fig. 4b) does not show any significant changes after ca. 6.5 kyr BP.

5 Discussion

The recent study on the KTK1 core provides clear evidences for the noticeable changes in the pollen assemblages and, thus, in the pollen-producing vegetation communities around Lake Kotokel during the late glacial and Holocene. The quantitative interpretation of the regional vegetation history (Fig. 3b, d) is in concordance with that based on the qualitative interpretation of the pollen record (Fig. 2; Bezrukova et al., 2008), but provide more robust reconstruction of the regional vegetation dynamics.

Despite the biomization approach (Prentice et al., 1996) does not allow the reconstruction of transitional vegetation types (e.g. forest-steppe and tundra-steppe), this missing information can be obtained (Fig. 3c) by examining the relative values of the respective biome scores (e.g. Fig. 3b; Tarasov et al., 2000). Additional information concerning spread of the woody vegetation communities is provided by the quantitative reconstruction of the vegetation cover (Fig. 3d). Our reconstruction supports the earlier idea (e.g. Grichuk, 1984; Tarasov et al., 2007b; Bezrukova et al., 2008) that the late glacial landscape in southern Siberia even being dominated by the cold and drought resistant steppe and tundra communities (e.g. up to 95% of the total vegetation cover)

5, 127-151, 2009 Vegetation and climate in southern Siberia since 15 kyr BP P. E. Tarasov et al. **Title Page** Introduction Abstract Conclusions References Figures Tables Back Close Full Screen / Esc Printer-friendly Version Interactive Discussion



was not completely free of boreal coniferous and deciduous trees. Recorded peaks in the arboreal pollen percentages (Fig. 2) and corresponding peaks in the taiga biome scores (Fig. 3b) and in the woody cover percentages (Fig. 3d) suggest that trees could quickly spread out of their glacial refugia occupying up to ca. 25% of the area within the
 21 km window at least two times during the late glacial interval, e.g. 14.8–14.7 kyr and

- 5 21 km window at least two times during the late glacial interval, e.g. 14.6–14.7 kyr and 13.3–12.8 kyr BP. The latter aforestaion phase around Kotokel falls within the Allerød (AL) Interstadial in line with other palaeobotanical records from northern Asia (see Andreev and Tarasov, 2007 for the overview) and from the Lake Baikal region (e.g. Demske et I., 2005; Bezrukova et al., 2008; Shichi et al., 2008 and references therein).
- According to the applied age model the earlier short-term aforestation episode might be an analogue of the European Meiendorf or Bølling (e.g. Frenzel et al., 1992) in the middle Lake Baikal region. However, more definitive conclusion can not be adequately proved, because of the uncertainty in the age model, which is based on the extrapolation below the 13.6 kyr BP level. An attribution of the reconstructed increase
- ¹⁵ in the woody cover around Kotokel about 14.8–14.7 kyr BP to the Meiendorf raises further question concerning occurrence of the Bølling (BO) Interstadial in the region. The earlier publications presenting postglacial pollen records from the Lake Baikal region usually do not separate Bølling from Allerød (e.g. Bezrukova et al., 2005; Demske et al., 2005). The KTK1 pollen diagram (Fig. 2; Bezrukova et al., 2008) also does not pro-
- vide a clear evidence for their separation. The quantitative transformation of the pollen percentages using the AVHRR-pollen-based approach suggests a slight increase in the woody cover percentages about 14 kyr BP (Fig. 3d). The accuracy of the method tested with the pollen-based modern tree cover reconstructions and original AVHRR-based estimates match well in Siberia (Tarasov et al., 2007b), providing satisfactory es-
- ²⁵ timates of percent variance explained and RMSE for both total woody cover (r^2 =0.77, RMSE=11.69) and different woody-cover fractions, including broadleaved (r^2 =0.66, RMSE=3.31) and needleleaved (r^2 =0.79, RMSE=10.23) tree cover. However, the error bars are still relatively large to discuss such minor fluctuations in the vegetation cover.

CPD

5, 127-151, 2009

Vegetation and climate in southern Siberia since 15 kyr BP



The Younger Dryas (YD) Stadial is well recognized in the diatom and sedimentary records from Lake Baikal (e.g. Prokopenko et al., 1999; Morley et al., 2005). However, up to date the pollen-based studies from the region reported rather weak YD signal, suggesting that the YD cooling had little effect on regional vegetation and thus on the pollen assemblages (e.g. Bezrukova et al., 2005; Demske et al., 2005; Shichi et al., 2008). In the KTK1 pollen and vegetation records the YD event is well defined. Both the significant decrease in the woody cover percentages from ca. 25% during AL to below 10% around 12.4 kyr BP and the replacement of boreal woodland by shrubby tundra communities help to place this event within ca. 12.8–11.6 kyr BP interval. Our reconstruction results also suggest that deterioration of the regional climate during YD was less pronounced in the vegetation records than during the earlier late glacial interval.

The onset of the Holocene interglacial conditions is marked in the KTK1 record by the increase in arboreal pollen percentages, return to boreal woodland vegetation and increase in woody cover to above 25% at 11.5 kyr BP. Since that time boreal forest be-

- ¹⁵ increase in woody cover to above 25% at 11.5 kyr BP. Since that time boreal forest became a major feature of the landscape around Kotokel. However, it took about 1 kyr until forest coverage in the area reached maximum (above 50%) between 10.5 and 7.5 kyr BP. Our pollen record shows a sharp increase in *Pinus-sylvestris* type percentages (8–32%) between 7.4 and 6.7 kyr BP, reflecting the spread of Scots pine in the
- ²⁰ region after 7 kyr BP. This feature is in good agreement with other pollen records from the Lake Baikal region and from the broader areas of Siberia (e.g. MacDonald et al., 2000; Bezrukova et al., 2005; Demske et al., 2005; Andreev and Tarasov, 2007). The reconstructed 10%-decrease in woody cover percentages dated to ca. 7–6.5 kyr BP, indicating that the vegetation cover became similar to that observed today can be ex-
- ²⁵ plained by the partial degradation of the birch forest around Kotokel suggested by the KTK1 pollen record.

Our results suggest distinct vegetation and environmental changes around Lake Kotokel since ca. 15 kyr BP, which can be related to the regional and global climate dynamics (Fig. 4). The first major spread of the boreal forest vegetation in the region at



ca. 13.5 kyr BP was characterized by significant climate amelioration (e.g. increase in T_{w} , T_{c} and P_{ann}) associated with the BO/AL Interstadial (Fig. 4e). Similar changes in temperature and precipitation values were derived from the Buguldeika pollen record from southern Baikal (Tarasov et al., 2007a). However, already at ca. 12.5 kyr BP the BMA reconstruction suggests a decrease in T_{w} and T_{c} and in P_{ann} dawn to the pre-5 interstadial values (Fig. 4a-c) (Tarasov et al., 2007a; this study). This major climate deterioration is in line with the reconstructed shift in vegetation from boreal woodland to much more open tundra landscape. The cold and dry climate oscillation derived from the KTK1 pollen record is perfectly synchronous to the YD Interstadial seen in the oxygen isotope records from Greenland (e.g. Stuiver et al., 1995; Fig. 4e) and from

10

15

China (e.g. Yuan et al., 2004; Fig. 4d).

The onset of a relatively warm and wet climate occurred after ca. 11.6 kyr BP, suggesting the onset of the Holocene interglacial conditions in line with the isotope records, indicating warming in the North Atlantic region and strengthening of the East Asian monsoon (Fig. 4). The most favorable conditions for the taiga growing with $T_{w} \sim 17 - 18^{\circ}$ C, $T_{c} \sim -19^{\circ}$ C and $P_{ann} \sim 500 - 550$ mm occurred ca. 10.8–7.3 kyr BP. In the

- Buguldeika record west of Kotokel the onset of the optimal conditions (e.g. $T_{w} \sim 16^{\circ}$ C, $T_c \sim -21^{\circ}$ C, $P_{ann} \sim 480$ mm) is reconstructed for the 9–7 kyr BP interval, e.g. 1.8 kyr later (Tarasov et al., 2007a). Whether this delay in the onset of the reconstructed "climatic
- optimum" represents local phenomena or spatial variations in the regional climate or 20 the results are influenced by other (non-climatic) factors, such as problems with the dating of the Baikal bottom sediments (e.g. Colman et al., 1996) and under-representation of some widely distributed woody taxa (e.g. larch) more pronounced in the pollen assemblages of the sediment cores recovered from larger lake requires further investi-
- gation. However, results of the recent study are consistent with the results derived 25 from the Continent and Buguldeika pollen records (Fig. 1), suggesting that the most favourable climate conditions are typical for the first half of the Holocene, as well as the Eemian Interglacial.



About 7-6.5 kyr BP the climate reconstructions from the KTK1 and Buguldeika records demonstrate decrease in temperature and precipitation in both records, but more pronounced in the Buguldeika record (Tarasov et al., 2007a). The reconstructed drop in T_c , and P_{ann} values is particularly important for discussion of the regional changes in the vegetation cover (e.g. Demske et al., 2005; Shichi et al., 2008; Fig. 3 5 in this study). Comparison of the precipitation curve (Fig. 4a) derived from the KTK1 record with the δ^{18} O data from the Dongge cave (Fig. 4d) demonstrates similarity between the two records, suggesting that in the Lake Baikal region the Holocene precipitation and tree-cover dynamics may be linked to shifts in intensity of the Pacific monsoon. The Holocene changes in the thermal parameters derived from the KTK1 10 pollen assemblages are in agreement with the δ^{18} O data from Greenland. The late Holocene cooling trend, a characteristic feature of the Northern Hemisphere and particularly of the North Atlantic region (Wanner et al., 2008), is especially pronounced in the winter temperature curve.

Acknowledgements. This study was done when Elena Bezrukova was a guest scientist at FU Berlin, funded by the German Research Foundation (DFG) via the project TA 540/1 "Comparison of climate and carbon cycle dynamics during late Quaternary interglacials using a spectrum of climate system models, ice-core and terrestrial archives". We thank O. N. Shestakova and RFFI Grant 06-05-64 671; Hikaru Takahara and the Russian-Japanese research team members, who took part in the coring campaign at Lake Kotokel in 2004, supported by the grants RFBR 05-05-66 942 and JSPS 16 255 009. The paper is dedicated to the memory of Dominique Jolly, who was among the pioneers introducing quantitative vegetation reconstruction methods in northern Eurasia.



The publication of this article is financed by CNRS-INSU.

5, 127-151, 2009 Vegetation and climate in southern Siberia since 15 kyr BP P. E. Tarasov et al. **Title Page** Introduction Abstract Conclusions References Figures Tables Back Close Full Screen / Esc Printer-friendly Version Interactive Discussion

References

- Alpat'ev, A. M., Arkhangel'skii, A. M., Podoplelov, N. Y., and Stepanov, A. Y.: Fizicheskaya geografiya SSSR (Aziatskaya chast'), Vysshaya Shkola, Moscow, 359 pp., 1976 (in Russian).
 Andreev, A. A. and Tarasov, P. E.: Pollen records, postglacial: Northern Asia, in: Encyclopedia
- ⁵ of Quaternary Science, edited by: Elias, S. A., Elsevier, Amsterdam, Netherlands, vol. 4., 2721–2729, 2007.
 - Bartlein, P. J., Webb III, T., and Fleri, E. C.: Holocene climatic change in the northern Midwest: pollen-derived estimates, Quaternary Res., 22, 361–374, 1984.
 - Battarbee, R. W., Gasse, F., and Stickley, C. E. (Eds.): Past climate variability through Europe and Africa, Developments in Paleoenvironmental Research, 6, Springer, Dordrecht, 2004.
- BDP-Members: Preliminary results of the first drilling on Lake Baikal, Buguldeika site, southeastern Siberia, Quatern. Int., 37, 3–17, 1997.
 - BDP-Members: A continuous record of climate changes for the last five million years from the bottom sediment of Lake Baikal, Russian Geology and Geophysics, 39, 135–154, 1998 (in
- 15 Russian).

10

BDP-Members: A new Quaternary record a regional tectonic, sedimentation and paleoclimate changes from drill core BDP-99 at Posolskaya Bank, Lake Baikal, Quatern. Int., 136(1), 105–121, 2005.

Berglund, B. E. and Ralska-Jasiewiczowa, M.: Pollen analysis and pollen diagrams, In: Hand-

²⁰ book of Holocene Palaeoecology and Palaeohydrology, edited by: Berglund, B. E., Interscience, New-York, 455–484, 1986.

Bezrukova, E. V.: Paleogeografiya Pribaikal'ya v pozdnelednikov'e i golotsene, Nauka, Novosibirsk, 128 pp., 1999 (in Russian).

Bezrukova, E. V., Abzaeva, A. A., Letunova, P. P., Kulagina, N. V., Vershinin, K. E., Belov, A. V.,

- ²⁵ Orlova, L. A., Danko, L. V., and Krapivina, S. M.: Post-glacial history of Siberian spruce (*Picea obovata*) in the Lake Baikal area and the significance of this species as a paleoenvironmental indicator, Quatern. Int., 136, 47–57, 2005.
 - Bezrukova, E. V., Krivonogov, S. K., Takahara, H., Letunova, P. P., Shichi, K., Abzaeva, A. A., Kulagina, N. V., and Zabelina, Yu. S.: Ozero Kotokel opornyi razrez pozdnelednikovya i
- ³⁰ golotsena Vostochnoi Sibiri, Doklady Academii Nauk, Geografiya, 420(2), 1–6, 2008. Colman, S. M., Jones, G. A., Rubin, M., King, J. W., Peck, J. A., and Orem, W. H.: AMS radiocarbon analyses from Lake Baikal, Siberia. Challenges of dating sediments from a large,

5, 127-151, 2009

Vegetation and climate in southern Siberia since 15 kyr BP



oligotrophic lake, Quaternary Sci. Rev., 15, 669-684, 1996.

- Danzeglocke, U., Jöris, O., and Weninger, B.: CalPal-2007^{online}, available at: http://www.calpal-online.de, 2008.
- DeFries, R. S. Townshend, J. R. G., and Hansen, M. C.: Continuous fields of vegetation char-
- ⁵ acteristics at the global scale at 1-km resolution, J. Geophys. Res., 104, 16911–16923, 1999.
 - Demske, D., Heumann, G., Granoszewski, W., Nita, M. Mamakowa, K., Tarasov, P. E., and Oberhänsli, H.: Late glacial and Holocene vegetation and regional climate variability evidenced in high-resolution pollen records from Lake Baikal, Global Planet. Change, 46, 255– 279, 2005.
- 10

15

Frenzel, B., Pecsi, B., and Velichko, A. A. (Eds.): Atlas of Palaeoclimates and Palaeoenvironments of the Northern Hemisphere, Late Pleistocene – Holocene, Hungarian Academy of Sciences, Budapest, Gustav Fisher Verlag, Stuttgart, 146 pp., 1992.

Galaziy, G. I. (Ed.): Baikal Atlas. Federal Agency for Geodesy and Cartography of Russia, Moscow, 160 pp., 1993 (in Russian).

- Grachev, M. A., Vorobyova, S. S., Khlystov, O. M., Bezrukova, E. V., Weinberg, E. V., Goldberg, E. L., Granina, L. Z., Kornakova, E. G., Lazo, F. I., Levina, O. V., Letunova, P. P., Otinov, P. V., Pirog, V. V., Fedotov, A. P., Yaskevich, S. A., Bobrov, V. A., Sukhorukov, F. V., Rezchikov, V. I., Fedorin, M. A., Zolotarev, K. V., and Kravchinsky, V. A.: Signal of the pale-
- oclimates of Upper Pleistocene in the sediments of Lake Baikal, Russ. Geol. Geophys., 38, 957–980, 1997.
 - Grichuk, V. P.: An experiment in reconstructing some characteristics of climate in the Northern Hemisphere during the Atlantic Period of Holocene, in: Holocene, edited by: Neustadt, M. I., Nàuka, Moscow, 41–57, 1969 (in Russian).
- Grichuk, V. P.: Late Pleistocene vegetation history, in: Late Quaternary environments of the Soviet Union, edited by: Velichko, A. A., University of Minnesota Press, Minneapolis, USA, 155–178, 1984.

Grimm, E. C.: TILIA 2.0 Version b.4 (Computer Software), Illinois State Museum, Research and Collections Center, Springfield, 1993.

³⁰ Grimm, E. C.: CONISS: A FORTRAN 77 program for stratigraphically constrained cluster analysis by the methods of incremental sum of squares, Comput. Geosci., 13, 13–15, 1987. Grimm, E. C.: TGView, Illinois State Museum, Research and Collections Center, Springfield, 2004. CPD

5, 127-151, 2009

Vegetation and climate in southern Siberia since 15 kyr BP



- Guiot, J.: Methodology of the last climatic cycle reconstruction from pollen data, Palaeogeogr. Palaeocl., 80, 49–69, 1990.
- Guiot, J., Pons, A., de Beaulieu, J.-L., and Reille, M.: A 140 000-year climatic reconstruction from two European pollen records, Nature, 338, 309–313, 1989.
- ⁵ Horiuchi, K., Minoura, K., Hoshino, K., Oda, T., Nakamura, T., and Kawai, T.: Palaeoenvironmental history of Lake Baikal during the last 23 000 years, Palaeogeogr. Palaeocl., 157, 95–108, 2000.
 - Iversen, J.: Viscum, Hedera and Ilex as climatic indicators. A contribution to the study of pastglacial temperature climate, Geol. Foeren. Foerhandl., 66, 463–483, 1944.
- Kageyama, M., Peyron, O., Pinot, S., Tarasov, P., Guiot, J., Joussaume, S., Ramstein, G., and PMIP participating groups: The Last Glacial Maximum climate over Europe and western Siberia, a PMIP comparison between models and data, Clim. Dynam., 17, 23–43, 2001.
 - Khotinskii, N. A.: Holocene climatic changes, in: Late Quaternary Environments of the Soviet Union, edited by: Velichko, A. A., University of Minnesota Press, Minneapolis, 305–312, 1984.
- ¹⁵ 1984. Khursevich, G. K., Karabanov, E. B., Prokopenko, A. A., Williams, D. F., Kuzmin, M. I., Fedenya, S. A., and Gvozdkov, A. A.: Insolation regime in Siberia as a major factor controlling diatom production in Lake Baikal during the past 800 000 years, Quatern. Int., 90–91, 47–58, 2001.
- ²⁰ Korde, N. V.: Bottom-sediments biostratigraphy of the Kotokol' Lake, in: Mezozoiskie i kainozoiskie ozera Sibiri, Nauka, Moscow, 150–170, 1968 (in Russian).
 - MacDonald, G. M., Velichko, A. A., Kremenetski, C. V., Borisova, O. K., Goleva, A. A., Andreev, A. A., Cwynar, L. C., Riding, R. T., Forman, S. L., Edwards, T. W. D., Aravena, R., Hammarlund, D., Szeicz, J. M., and Gattaulin, V. N.: Holocene treeline history and climate change across northern Eurasia, Quaternary Res., 53, 302–311, 2000.
 - Molozhnikov, V. N.: Rastitel'nye soobshchestva Pribaikal'ya, Nauka, Novosibirsk, 271 pp., 1986 (in Russian).
 - Morley, D. W., Leng, M. J., Mackay, A. W., and Sloane, H. J.: Late glacial and Holocene environmental change in the Lake Baikal region documented by oxygen isotopes from diatom
- ³⁰ silica, Global Planet. Change, 46, 221–233, 2005.

25

Nakagawa, T., Tarasov, P., Kotoba, N., Gotanda, K., and Yasuda, Y.: Quantitative pollen-based climate reconstruction in Japan: application to surface and late Quaternary spectra, Quaternary Sci. Rev., 21, 2099–2113, 2002.

5, 127–151, 2009	
Vegetation and climate in southern Siberia since 15 kyr BP P. E. Tarasov et al.	
Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
I	۶I
•	
Back	Close
Full Screen / Esc	
Printer-friendly Version	

Title Page Introduction Abstract Conclusions References Figures Tables Back Close Full Screen / Esc Printer-friendly Version Interactive Discussion

- New, M., Lister, D., Hulme, M., and Makin, I.: A high-resolution data set of surface climate over global land areas, Clim. Res., 21, 1–25, 2002.
- Oberhänsli, H. and Mackay, A.: Introduction to "Progress towards reconstructing past climate in Central Eurasia, with special emphasis on Lake Baikal, Global Planet. Change, 46, 1–7, 2005.

5

15

Overpeck, J. T., Webb, III T., and Prentice, I. C.: Quantitative interpretation of fossil pollen spectra, dissimilarity coefficients and the method of modern analogs, Quaternary Res., 23, 87–108, 1985.

Prentice, C. I., Guiot, J., Huntley, B., Jolly, D., and Cheddadi, R.: Reconstructing biomes from palaeoecological data: a general method and its application to European pollen data at 0 and 6 ka, Clim. Dynam., 12, 185–194, 1996.

Prentice, I. C., Jolly, D., and BIOME 6000 participants: Mid-Holocene and glacial maximum vegetation geography of the northern continents and Africa, J. Biogeogr., 27, 507–519, 2000.

Prokopenko, A. A. and Williams, D. F.: Deglacial methane emission signals in the carbon isotopic record of Lake Baikal, Earth Planet. Sci. Lett., 218, 135–147, 2004.

Prokopenko, A. A., Williams, D. F., Karabanov E. B., and Khursevich, G. K.: Response of Lake Baikal ecosystem to climate forcing and *p*CO₂ change over the last glacial/interglacial transition, Earth Planet. Sci. Lett., 172, 239–253, 1999.

Seppä, H. and Birks, H. J. B.: July mean temperature and annual precipitation trends during

- the Holocene in the Fennoscandian tree-line area, pollen-based reconstructions, Holocene, 11, 527–539, 2001.
 - Seppä, H. and Birks, H. J. B.: Holocene climate reconstructions from the Fennoscandian treeline area based on pollen data from Toskaljavri, Quaternary Res., 57, 191–199, 2002.

Shichi, K., Takahara, H., Krivonogov, S. K., Bezrukova, E. V., Kashiwaya, K., and Takehara, A.:

- ²⁵ Vegetation and climate records for the last 50 kyr from Lake Kotokel, the middle Lake Baikal area, East Siberia, Quatern. Int., in press, 2008.
 - Stuiver, M., Grootes, P. M., and Braziunas, T. F.: The GISP2 d18O climate record of the past 16 500 years and the role of the sun, ocean and volcanoes, Quaternary Res., 44, 341–354, 1995.
- Takahara, H., Krivonogov, S. K., Bezrukova, E. V., Miyoshi, N., Morita, Y., Nakamura, T., Hase, Y., Shinomiya, Y., and Kawamuro, K.: Vegetation history of the southeastern and eastern coasts of Lake Baikal from bog sediments since the last interstade, in: Lake Baikal: a Mirror in Time and Space for Understanding Global Change Processes, edited by: Mi-

5, 127-151, 2009

Vegetation and climate in southern Siberia since 15 kyr BP

noura, K., Elsevier, Amsterdam, 108–118, 2000.

5

Tarasov, P. E., Harrison, S. P., Saarse, L., Pushenko, M. Ya, Andreev, A. A., Aleshinskaya, Z. V., Davydova, N. N., Dorofeyuk, N. I., Efremov, Yu. V., Khomutova, V. I., Sevastyanov, D. V., Tamosaitis, J., Uspenskaya, O. N., Yakushko, O. F., and Tarasova, I. V.: Lake status records from the former Soviet Union and Mongolia: data base documentation, NOAA Paleoclima-

tology Publications Series Report No. 2, Boulder, 274 pp., 1994.

- Tarasov, P. E., Webb III, T., Andreev, A. A., Afanas'eva, N. B., Berezina, N. A., Bezusko, L. G., Blyakharchuk, T. A., Bolikhovskaya, N. S., Cheddadi, R., Chernavskaya, M. M., Chernova, G. M., Dorofeyuk, N. I., Dirksen, V. G., Elina, G. A., Filimonova, L. V., Glebov, F. Z.,
- Guiot, J., Gunova, V. S., Harrison, S. P., Jolly, D., Khomutova, V. I., Kvavadze, E. V., Osivopa, I. M., Panova, N. K., Prentice, I. C., Saarse, L., Sevastyanov, D. V., Volkova, V. S., and Zernitskaya, V. P.: Present day and mid-Holocene biomes reconstructed from pollen and plant macrofossil data from the former Soviet Union and Mongolia, J. Biogeogr., 25, 1029–1054, 1998.
- ¹⁵ Tarasov, P. E., Guiot, J., Cheddadi, R., Andreev, A. A., Bezusko, L. G., Blyakharchuk, T. A., Dorofeyuk, N. I., Filimonova, L. V., Volkova, V. S., and Zernitskaya, V. P.: Climate in northern Eurasia 6000 years ago reconstructed from pollen data, Earth Planet. Sci. Lett., 171, 635– 645, 1999.

Tarasov, P. E., Volkova, V. S., Webb III, T., Guiot, J., Andreev, A. A., Bezusko, L. G.,

Bezusko, T. V., Bykova, G. V., Dorofeyuk, N. I., Kvavadze, E. V., Osipova, I. M., Panova, N. K., and Sevastyanov, D. V.: Last Glacial Maximum biomes reconstructed from pollen and plant macrofossil data from Northern Eurasia, J. Biogeogr., 27(3), 609–620, 2000.

Tarasov, P. E., Dorofeyuk, N. I., and Vipper, P. B.: The Holocene dynamics of vegetation in Buryatia, Stratigr. Geo. Correl.+, 10, 88–96, 2002.

- Tarasov, P. E., Granoszewski, W., Bezrukova, E. V., Brewer, S., Nita, M., Abzaeva, A. A., and Oberhänsli, H.: Quantitative reconstruction of the Last Interglacial vegetation and climate based on the pollen record from Lake Baikal, Russia, Clim. Dynam., 25, 625–637, 2005.
 - Tarasov, P., Bezrukova, E., Karabanov, E., Nakagawa, T., Wagner, M., Kulagina, N., Letunova, P., Abzaeva, A., Granoszewski, W., and Riedel, F.: Vegetation and climate dynamics during
- the Holocene and Eemian interglacials derived from Lake Baikal pollen records, Palaeogeogr. Palaeocl., 252, 440–457, 2007a.
 - Tarasov, P. E., Williams J. W., Andreev, A. A., Nakagawa, T., Bezrukova, E. V., Herzschuh, U., Igarashi, Y., Müller, S., Werner, K., and Zheng, Z.: Satellite- and pollen-based quantita-

5, 127–151, 2009

Vegetation and climate in southern Siberia since 15 kyr BP



147

tive woody cover reconstructions for northern Asia: Verification and application to late-Quaternary pollen data, Earth Planet. Sci. Lett., 264, 284–298, 2007b.

- Vipper, P. B. and Smirnov, N. N.: The studying of biogeocenozes history by the fresh water lakes bottom deposits, Obshchie metody izucheniya istorii sovremennykh ekosistem, Nauka,
- Moscow, 14–39, 1979 (in Russian). 5

15

- Wanner, H., Beer, J., Bütikofer, J., Crowley, T. J., Cubasch, U., Flückiger, J., Goosse, H., Grosjean, M., Joos, F., Kaplan, J. O., Küttel, M., Müller, S. A., Prentice, I. C., Solomina, O., Stocker, T. F., Tarasov, P., Wagner, M., and Widmann, M.: Mid- to late Holocene climate change - an overview, Quaternary Sci. Rev., 27(19-20), 1791-1828, 2008.
- Williams, D. F., Peck, J., Karabanov, E. B., Prokopenko, A. A., Kravchinsky, V., King, J., and 10 Kuzmin, M. I.: Lake Baikal record of continental climate response to orbital isolation during the past 5 million years. Science, 278, 1114-1117, 1997.
 - Williams, D. F., Kuzmin, M. I., Prokopenko, A. A., Karabanov, E. B., Khursevich, G. K., and Bezrukova, E. V.: The Lake Baikal drilling project in the context of a global lake drilling initiative, Quatern. Int., 80-81, 3-18, 2001.
 - Williams, J. W., Shuman, B. N., Webb III, T., Bartlein, P. J., and Leduc, P. L.: Late Quaternary vegetation dynamics in North America: Scaling from taxa to biomes, Ecol. Monogr., 74, 309-334, 2004.

Yuan, D. X., Cheng, H., Edwards, R. L., Dykoski, C. A., Kelly, M. J., Zhang, M. L., Qing, J. M.,

Lin, Y. S., Wang, Y. J., Wu, J. Y., Dorale, J. A., An, Z. S., and Cai, Y. J.: Timing, duration, and 20 transitions of the Last Interglacial Asian monsoon, Science, 304, 575–578, 2004.



5, 127-151, 2009

Vegetation and climate in southern Siberia since 15 kvr BP







CPD

5, 127–151, 2009

Vegetation and climate in southern Siberia since 15 kyr BP







CPD

5, 127–151, 2009





Fig. 3. Local pollen assemblages zones **(a)** and time series of individual vegetation types (biomes) dominating in the study area since 15 kyr BP **(b)**, along with the qualitative characteristic of vegetation **(c)**, and quantitative changes in woody cover percentages **(d)** derived from the KTK1 pollen record, plotted along the time axis.









