

*Climate of the Past Discussions* is the access reviewed discussion forum of *Climate of the Past*

# Potential analysis reveals changing number of climate states during the last 60 kyr

V. N. Livina<sup>1</sup>, F. Kwasniok<sup>2</sup>, and T. M. Lenton<sup>1</sup>

<sup>1</sup>School of Environmental Sciences, University of East Anglia, Norwich, UK

<sup>2</sup>School of Engineering, Computing and Mathematics, University of Exeter, Exeter, UK

Received: 28 August 2009 – Accepted: 9 September 2009 – Published: 30 September 2009

Correspondence to: V. N. Livina (v.livina@uea.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

CPD

5, 2223–2237, 2009

## Potential analysis

V. N. Livina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

We develop and apply a new statistical method of potential analysis for detecting the number of states of a geophysical system, from its recorded time series. Estimation of the degree of a polynomial potential allows us to derive the number of potential wells in a system. The method correctly detects changes in the number of wells in artificial data. In ice-core proxy records of Greenland paleotemperature, a reduction in the number of climate states from two to one is detected sometime prior to the last glacial maximum (LGM), 23–19 kyr BP. This bifurcation can be interpreted as loss of stability of the warm interstadial state of the Dansgaard-Oeschger events. In data spanning the last glacial termination, up to four climate states are detected, plausibly representing the LGM, Bolling-Allerod, Younger Dryas, and the Holocene. The proposed method can be applied to a wide range of geophysical time series exhibiting bifurcations.

## 1 Introduction

Knowing the number of states in a geophysical system is crucial for understanding its underlying dynamics and for modeling its behavior. Of particular interest are changes in the number of states which represent bifurcations of a system. Commonly the number of system states is inferred either by simply looking at time series data, or on the basis of theoretical models. Yet when there are two or more states present in a system the precise number is not always obvious from examining time series by eye (see e.g. Fig. 2b). In particular, what looks like a system with two states may actually contain more states. Consequently we have developed a more formal (though still necessarily approximate) method of estimating the number of system states from time series data.

In studies of paleoclimate changes, it is common to describe the glacial-interglacial cycles as shifts between different climate states. For example, simple models have assumed either two (glacial, interglacial) (Saltzman and Verbitsky, 1993) or three (mild glacial, full glacial, interglacial) (Paillard, 2001) states for Northern Hemisphere ice vol-

## Potential analysis

V. N. Livina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Potential analysis**

V. N. Livina et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ume. Within the last ice age, the Dansgaard-Oeschger (DO) events that are recorded in Greenland ice cores (Dansgaard et al., 1993), North Atlantic paleoclimate records (Bond et al., 1997), and are now recognized further afield (Wang et al., 2001), are commonly treated as a two-state system (cold stadial, warm interstadial). The DO events are widely explained in terms of a model (Stommel, 1961; Rahmstorf, 1995; Ganopolski and Rahmstorf, 2001) for bistability of the Atlantic ocean thermohaline circulation, coupled to changes in sea-ice cover and atmospheric circulation. Potentially, different combinations of Northern Hemisphere ice volume and ocean-atmosphere-sea-ice state could give rise to multiple climate states as recorded in Greenland. Ice volume clearly affects the stability regime of the ocean-atmosphere-sea-ice as the Holocene interglacial has not exhibited the extreme DO variability seen in the last ice age.

Here we apply our new method to examine how many climate states are present in high-resolution Greenland Ice-Core Project (GRIP) (Dansgaard et al., 1993) and North Greenland Ice-Core Project (NGRIP) (NGRIP project, 2004) records, as a function of time through the last 60 kyr (part of the last ice age and the Holocene). The precise mechanisms causing transitions between climate states do not concern us here. We make the simplest assumption that multiple climate states can be approximated by a non-oscillatory polynomial potential, albeit one that changes through time. Furthermore, we assume that transitions between climate states are triggered purely by stochastic noise. This is consistent with recent analysis of the DO events (especially in the GRIP and NGRIP records), which does not support any underlying periodicity (Ditlevsen et al., 2007).

## 2 Dynamical framework

We treat the climate system through this interval as a nonlinear dynamical system which can possess multiple states, with shifts between these different climate states induced by stochastic forcing. A one-dimensional conceptual model for this is given by

the Langevin equation

$$\dot{z} = -U'(z) + \sigma\eta, \quad (1)$$

where  $U(z)$  is a potential function,  $\sigma$  is the noise level and  $\eta$  is a Gaussian white noise with zero mean and unit variance. The state variable  $z$  represents a large-scale climate variable like Greenland paleotemperature and is here identified with ice-core proxy ( $\delta^{18}\text{O}$  stable water isotope) records. In (Ditlevsen, 1999), the Fokker-Plank equation was used to derive the potential of the GRIP calcium record, with white non-Gaussian noise and two noise terms in the Langevin equation.

The shape of the potential is given by a polynomial

$$U(z) = \sum_{i=1}^L a_i z^i, \quad (2)$$

where the order  $L$  is even and the leading coefficient  $a_L$  is positive for Eq. (1) to possess a stationary solution. The order of the polynomial controls the complexity of the potential. Increasing values of  $L$  allow more states to be accommodated; for example, a fourth-order polynomial can capture a system with two states (double-well potential) (Kwasniok and Lohmann, 2009a,b).

The number of system states is estimated by means of a polynomial fit of the probability density function of the data. Suppose the system is governed by Eq. (1). The corresponding Fokker-Planck equation for the probability density function  $p(z, t)$

$$\partial_t p(z, t) = \partial_z [U'(z)p(z, t)] + \frac{1}{2}\sigma^2 \partial_z^2 p(z, t) \quad (3)$$

has a stationary solution given by (Gardiner, 2004)

$$p(z) \sim \exp[-2U(z)/\sigma^2]. \quad (4)$$

Given this one-to-one correspondence between the potential and the stationary probability density of the system, the potential can be reconstructed from time series data

## Potential analysis

V. N. Livina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the system as

$$U = -\frac{\sigma^2}{2} \log p_d, \quad (5)$$

where  $p_d$  is the empirical probability density of the data. This is estimated using a standard Gaussian kernel estimator (Silverman, 1986). Then least-square fits of  $-\log p_d$  weighted with the probability density of the data (Kwasniok and Lohmann, 2009a) with polynomials of increasing even order  $L$  are calculated, starting with  $L=2$ , until a negative leading coefficient  $a_L$  is encountered. The polynomial of highest degree before first obtaining a negative leading coefficient is considered the most appropriate representation of the probability density of the time series, both locally and globally, avoiding overfitting of sampling fluctuations in the probability density.

The number of states  $S$  in the system is then determined as

$$S = 1 + \frac{l}{2}, \quad (6)$$

where  $l$  is the number of inflection points of the fitted polynomial potential of appropriate degree  $L$  as described above. This definition takes into account not only the degree of the polynomial but its actual shape. We only look at even-order potentials with positive leading coefficient. These have positive curvature both at minus and plus infinity. Thus, inflection points can only occur in pairs (if any). Any potential has at least one state (with no inflection points). Then we count one further state for each pair of inflection points. This can be either a real minimum (well) or just a flattening in the potential corresponding to degeneracies in the potential (Kwasniok and Lohmann, 2009b); definition (6) accommodates both possibilities. The number of inflection points is numerically given as the number of sign changes in the second derivative on a fine enough mesh.

Examples of the polynomial fits for simulated data are given in Fig. 1. Given double-well potential data ( $U(z)=z^4-2z^2$ ), the probability density of the time series has bimodal shape. The empirical potential given as  $U=-\frac{\sigma^2}{2} \log p_d$  can be fitted by polynomials of

## Potential analysis

V. N. Livina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Potential analysis**

V. N. Livina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



various orders, but only those of degree 2 and 4 have a positive leading coefficient. Higher-order polynomials have negative leading coefficients. Although fitting the data better and better locally, they are clearly inadequate globally. The fourth-order polynomial is the appropriate representation of the probability density and the method identifies the time series as double-well potential.

A simplified version of definition (6) would be to only count the relative minima (real wells) in the potential. We opt here for the more comprehensive definition of system states as degenerate potentials have been shown to occur in the context of ice-core records (Kwasniok and Lohmann, 2009a,b). Moreover, even if the system possesses real potential wells (for example, the artificial data shown in Fig. 2) changes in the number of states (bifurcations) tend to be picked up more quickly when using the inflection point definition as this is a more sensitive feature.

### 3 Data

To test the method for estimating the number of system states, we generated four sets of artificial data for four different potentials: one-well ( $U(z) = z^2$ ), double-well ( $U(z) = z^4 - 2z^2$ ), triple-well ( $U(z) = z^6 - 4.5z^4 + 5z^2$ ) and four-well potential ( $U(z) = z^8 - 6.5z^6 + 13z^4 - 8z^2$ ) (Fig. 2a). Simulated data of these systems were obtained by numerically integrating Eq. (1) using the Euler scheme. The noise level  $\sigma = 1.5$  was adopted. The sampling interval is 0.05 system time units. For each of the potentials, 150 time units worth of data were generated, resulting in 3000 data points. The subsets are combined to produce a record with 12 000 data points, corresponding to time running from 0 to 600 (Fig. 2b).

Having tested the methods on artificial data, we examined ice-core proxy records of paleotemperature at two different sites 325 km apart in Greenland; GRIP (Dansgaard et al., 1993) (Fig. 3a) and NGRIP (NGRIP project, 2004) (Fig. 3c), on the most recent GICC05 time scale (Svensson et al., 2008). The data are  $\delta^{18}\text{O}$  stable water isotope records, which are a proxy for past air temperature at the ice core sites. The records

can also be influenced by changing water source temperatures and snowfall seasonality. There are significant differences between the two records during the last ice age, with NGRIP systematically colder (depleted in  $^{18}\text{O}$ ), and more variable, when Northern Hemisphere ice sheets are more extensive (NGRIP project, 2004). This may be because NGRIP received a greater fraction of colder air coming over the northern side of the Laurentide ice sheet. Importantly, the difference between the two records does not show a component of millennial variability. Here we consider the last 60 kyr NGRIP series with resolution 20 yr and a part of the GRIP series (which is 112 kyr long) in the same interval (60–0 kyr BP) with the same temporal resolution of 20 yr. For the detection of changing numbers of states we use sliding windows of varying length through each dataset.

## 4 Results

We visualize the estimated number of system states as a colour contour plot, expressed as a function of the time at the end of the data window ( $x$ -axis, aligned with the data time scale) and the time length of the data window ( $y$ -axis). As the results are plotted at the ends of their corresponding time windows, each point in the contour plot should be compared with the segment of the time series data ending at that particular point, and having the length of the corresponding time window.

In the artificial data, for the smallest windows, there are sporadic misidentifications of the number of states, due to poor statistics (Fig. 2c). Generally time windows of order 400 data points (20 time units) are sufficient to get reliable results. There are delays in detection, which grow with the size of the sliding windows, and are typically 40–50% of the window size. For the 3- to 4-well transition the delay in detection is somewhat longer than for the 1- to 2-well or 2- to 3-well transitions. These limitations should be remembered in interpreting the results for the Greenland paleotemperature data.

The GRIP (Fig. 3a) and NGRIP (Fig. 3c) paleotemperature data are highly variable and non-stationary. Nevertheless, the algorithm detects a number of interesting com-

## Potential analysis

V. N. Livina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mon features in both records (Fig. 3b,d). Over 60–25 kyr BP, 2 climate states are most commonly detected, consistent with the conventional view of the Dansgaard-Oeschger events. Exceptions of 1-state detection appear to be caused when long intervals of steady cooling, e.g. circa 55–50 kyr BP, occupy a significant part of the analysis window.

The most pronounced feature, common to both records, is a change from 2–well to 1–well potential, generally detected in the interval 27–20 kyr BP (the timing dependent on window size), and inferred to have occurred prior to its detection. The transition is most sharply detected in the GRIP data as having occurred prior to 23 kyr BP for a range of intermediate window sizes. In the NGRIP data, the transition is less pronounced, consistent with the greater noise level in this dataset obscuring the detection of states. In both records, the shift to a 1–well potential is persistent, indicating a bifurcation in the climate system that occurred late in the last ice age but prior to the Last Glacial Maximum (LGM) 23–19 kyr BP (Yokoyama et al., 2000).

The return of a second state in the climate is only detected in both records, for most window sizes, around 12 kyr BP (i.e. at the end of the Younger Dryas), although it is sporadically detected earlier, from the time of the Bolling warming, especially in the NGRIP data. This reflects the growing influences of changes during the deglaciation on the algorithm. The number of states detected increases up to 4 as the data spanning the interval of the last glacial termination come to dominate the analysis window. These four climate states plausibly represent the LGM, Bolling-Allerod warm interval, Younger Dryas cold interval, and the Holocene.

As the end of the analysis window advances into the Holocene, the estimated number of system states declines. However, the number of climate states only reduces to 1 for both datasets when the analysis window comprises only the data since around 10 kyr BP, i.e. the start of the Holocene. For the GRIP data, there is some indication that the 8.2 kyr BP event can influence the algorithm sufficiently that 2 climate states are detected when using small windows that span the event. The 8.2 kyr BP event is also visible in the NGRIP data but has poorer signal-to-noise ratio, so does not exert

## Potential analysis

V. N. Livina et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the same influence on the algorithm.

Given that the ice-core time series are much shorter (in terms of the number of recorded transitions between states) than the simulated time series used above to verify the method the question arises how robust our results on the ice-core data are statistically. The mean recurrence time (Kramers waiting time) between DO events is 2.8 kyr which is about the same as in the artificial data in nondimensional system units. Hence the maximum window size of 20 kyr in Fig. 3b and d corresponds to 20 units on the vertical axis in Fig. 2c. This is just enough to enable a correct detection of the number of states with very high probability. Together with the fact that our results are robust for a range of window sizes between 10 and 20 kyr this gives us some confidence that our conclusions are reliable.

## 5 Conclusions

The results support the widely held view of the Dansgaard-Oeschger events that they represent switches between 2 states of the climate system, which were present from before the start of our analysis 60 kyr BP. Most interesting is the detection of a bifurcation in the climate system that reduced the number of states from 2 to 1, sometime prior to the last glacial maximum (LGM) 23–19 kyr BP. This can be interpreted as the loss of a stable warm interstadial state. This result is not inconsistent with the conventional labelling of DO warming event 2 occurring at 23.4 kyr BP timing according to (Rahmstorf, 2003) or the earlier DO events 3 at 27.8 kyr BP and 4 at 29.0 kyr BP. Rather we suggest that although transitions to a warm state were still triggered up to the LGM, the resulting state became not even marginally stable. In the case of DO event 2 at least (and possibly 3 and even 4), the system was always destined to revert back to a cold state at a rate determined by its internal dynamics and/or the removal of some forcing factor. Indeed shortening of the duration of the warm interstadial state can be seen in the timeseries (Fig. 3a, c) as time progresses toward the LGM.

Clearly a warm state reappears during the deglaciation, in the form of the Bolling

### Potential analysis

V. N. Livina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Potential analysis**

V. N. Livina et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

warming event (conventionally labelled DO event 1 at 14.6 kyrBP; Rahmstorf, 2003), but whether it is appropriate to liken this to the earlier interstadials or the Holocene is unclear from our analysis. Likewise whether the Younger Dryas cool event should be likened to the earlier cold stadial state is ambiguous. Clearly Northern Hemisphere ice volume had declined significantly prior to either of these events, and was undergoing a transition from glacial to interglacial states. For the last 10 kyr of the Holocene, under interglacial ice volume only 1 stable Atlantic ocean-atmosphere-sea-ice state is detected. We interpret the 8.2 kyrBP cooling event as a meltwater induced transient change in ocean-atmosphere-sea-ice from which the system was always destined to recover. Our tentative detection of up to 4 climate states could represent full glacial with only one stable ocean-atmosphere-sea-ice state, intermediate ice volume with 2 different ocean-atmosphere-sea-ice states and full interglacial with only one stable ocean-atmosphere-sea-ice state.

In this view the stability regime of the Atlantic ocean-atmosphere-sea-ice system is reshaped and sometimes bifurcated by longer timescale changes in ice volume, as suggested by a number of models, e.g. (Ganopolski and Rahmstorf, 2001; Colin de Verdiere et al., 2006). Bistability of the Atlantic ocean-atmosphere-sea-ice system is present for intermediate Northern Hemisphere ice volumes, whereas under glacial maximum conditions there is only one stable cold state for Greenland temperature, and under interglacial conditions there is only a single (different) stable warm state.

*Acknowledgements.* We would like to thank J. Kantelhardt and Yu. Sapronov for discussions and P. Ditlevsen for useful comments on the manuscript. The research was supported by NERC through project NE/F005474/1.

**References**

Bond, G. Showers, W., Cheseby, M., Lotti, R., Almasi, P., de Menocal, P., Priore, P., Cullen, H., Hajdas, I., and Bonani, G.: A Pervasive Millennial-Scale Cycle in North Atlantic Holocene and Glacial Climates, *Science*, 278, 1257–1266, 1997. 2225

- Colin de Verdiere, A., Ben Jelloul, M., and Sevellec, F.: Bifurcation structure of thermohaline millennial oscillations, *J. Climate*, 19, 5777–5795, 2006. 2232
- Dansgaard, W., Johnson, S. J., Clausen, H. B., Dahljensen, D., Gundestrup, N. S., Hammer, C. U., Hvidberg, C. S., Steffensen, J. P., Sveinbjornsdottir, A. E., Jouzel, J., and Bond, G.: Evidence for general instability of past climate from a 250 kyr ice-core record, *Nature*, 364, 218–220, 1993. 2225, 2228
- Ditlevsen, P.D.: Observation of  $\alpha$ -stable noise induced millennial climate changes from an ice-core record, *Geophys. Res. Lett.*, 26(10), 1441–1444, 1999. 2226
- Ditlevsen, P. D., Andersen, K. K., and Svensson, A.: The DO-climate events are probably noise induced: statistical investigation of the claimed 1470 years cycle, *Climate of the Past*, 3(1), 129–134, 2007. 2225
- Gardiner, C. W.: *Handbook of Stochastic Methods for Physics, Chemistry and the Natural Sciences* (Springer Series in Synergetics), Springer; 3rd edition, 415 pp., 2004. 2226
- Ganopolski, A. and Rahmstorf, S.: Rapid changes of glacial climate simulated in a coupled climate model, *Nature*, 409, 153–158, 2001. 2225, 2232
- Kwasniok, F. and Lohmann, G.: Deriving dynamical models from paleoclimatic records: application to glacial millennial-scale climate transitions, *Phys. Rev. E*, in press, 2009a. 2226, 2227, 2228
- Kwasniok, F. and Lohmann, G.: A stochastic nonlinear oscillator model for glacial millennial-scale climate variability derived from ice-core data. *Nonl. Proc. Geophys.*, in press, 2009b. 2226, 2227, 2228
- North Greenland Ice Core Project members High-resolution record of Northern Hemisphere climate extending into the last interglacial period, data are available at: <http://www.iceandclimate.nbi.ku.dk/data/> *Nature*, 431, 147–151, 2004. 2225, 2228, 2229
- Paillard, D.: Glacial cycles: toward a new paradigm, *Reviews of Geophys.*, 39(3), 325–346, 2001. 2224
- Rahmstorf, S.: Bifurcation of the Atlantic thermohaline circulation in response to changes in the hydrological cycle, *Nature*, 378, 145–149, 1995. 2225
- Rahmstorf, S.: Timing of abrupt climate change: a precise clock, *Geophys. Res. Lett.*, 30(10), p. 1510, 2003. 2231, 2232
- Saltzman, B. and Verbitsky, M. Y.: Multiple instabilities and modes of glacial rhythmicity in the Plio-Pleistocene: a general theory of late Cenozoic climatic change, *Climate Dynamics*, 9, 1–15, 1993. 2224

**Potential analysis**

V. N. Livina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Silverman, B. W.: Density estimation for statistics and data analysis, Chapman and Hall, 1986. 2227
- Stommel, H.: Thermohaline convection with two stable regimes of flow, *Tellus XIII*, 2, 224–230, 1961. 2225
- 5 Svensson, A., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Davies, S. M., Johnsen, S. J., Muscheler, R., Parrenin, F., Rasmussen, S. O., Roethlisberger, R., Seierstad, I., Steffensen, J. P., and Vinther, B. M.: A 60000 year Greenland stratigraphic ice core chronology, *Climate of the Past*, 4, 47–57, 2008. 2228
- 10 Wang, Y. J., Cheng, H., Edwards, R. L., An, Z. S., Wu, J.Y., Shen, C. C., and Dorale, J. A.: A High-Resolution Absolute-Dated Late Pleistocene Monsoon Record from Hulu Cave, China., *Science*, 294, 2345–2348, 2001. 2225
- Yokoyama Y., Lambeck, K., DeDekker, P., Johnston, P., and Fifield, L.K.:Timing of the Last Glacial Maximum from observed sea-level minima, *Nature*, 406, 713–716, 2000. 2230

CPD

5, 2223–2237, 2009

## Potential analysis

V. N. Livina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




## Potential analysis

V. N. Livina et al.



**Fig. 1.** Empirical potential (magenta curve) of double-well potential artificial data and its polynomial fits of 2nd, 4th and 6th orders. The parabolic fit (2nd order) is not as accurate as the 4th-order polynomial, whereas the fit of 6th order approximates the potential only locally, becoming negative for large  $|z|$ . The numerical algorithm chooses the highest order of polynomial that approximates the potential globally with positive leading coefficient; in this case it is the 4th-order polynomial. The method correctly classifies the data as double-well potential.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




**Fig. 2.** Detection of the number of states in artificial data: **(a)** four different potentials with increasing number of wells from one to four; **(b)** the resulting time series; **(c)** contour plot of the number of detected states versus time and size of sliding window (red – one well, green – two, cyan – three, purple – four). The number of detected wells is mapped at the end of the sliding time windows.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Potential analysis

V. N. Livina et al.



**Fig. 3.** Potential analysis of Greenland ice-core data: **(a)** GRIP time series; **(b)** contour plot of the number of detected states versus time and size of sliding window, mapped at the end of the sliding time windows (red – one well, green – two, cyan – three, purple – four). **(c)** NGRIP time series; **(d)** contour plot of the number of detected states versus time and size of sliding window, as in (b).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)