



## Abstract

A new deep ice core drilling, TALDICE, has been successfully handled by a European team at Talos Dome, in the Ross Sea sector of East Antarctica, down to 1620 m depth. Using stratigraphic markers and a new inverse method, we produce the first official chronology of the ice core, called TALDICE-1. We show that it notably improves an a priori chronology resulting from a one-dimensional ice flow model, and that it is in agreement with a posteriori controls of the resulting accumulation rate and thinning function along the core. An absolute uncertainty of only 300 yr is obtained in the course of the last deglaciation. This uncertainty remains lower than 600 yr over Marine Isotope Stage 3, back to 50 kyr BP. The phasing of the TALDICE ice core climate record with respect to the central East Antarctic plateau and Greenland records can thus be determined with a precision allowing for a discussion of the mechanisms at work at sub-millennial time scales.

## 1 Introduction

The sequence of the last deglaciation in Greenland is characterized by the succession of an abrupt warming (Bolling-Allerod, 14.7 kyr BP, Steffensen et al., 2008) followed by an abrupt cooling (Younger Dryas, 12.9 kyr BP, Stuiver et al., 1995) before a rapid return (11.7 kyr BP, Severinghaus et al., 1998) to interglacial conditions (Alley, 2000). Such millennial scale climate variability is similar to the one observed during the last glacial period, and has counterpart climatic features in Antarctica. The Antarctic signal consists of a slow warming which precedes the abrupt transition in Greenland by a few hundreds to thousand years. When Greenland temperature abruptly rises, a slow cooling starts in Antarctica (Jouzel et al., 1987). Previous investigations of central East Antarctic ice cores have revealed the same consistent North-South timing and phasing for the sequence of the last deglaciation (Blunier et al., 1998; EPICA Community Members, 2006).

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**TALDICE-1 age scale  
of the Talos Dome  
deep ice core, East  
Antarctica**

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



However studies of coastal ice cores in Antarctica have revealed a different pattern in two cases. The Taylor Dome site, situated near the Ross Sea, reveals a more Greenlandic-like evolution during the deglaciation (Steig et al., 1998), whereas the first deglacial warming recorded in the Law Dome ice core situated in the Indian Ocean sector ends before the abrupt warming of the Bølling/Allerød (BA) in Greenland (Morgan et al., 2002). Currently only few other ice cores from Antarctic coastal or near-coastal sites cover the last 25 kyr (for example Siple Dome (Brook et al., 2005), Byrd (Sowers and Bender, 1995; Blunier et al., 1998) and Epica Dronning Maud Land (EDML) (EPICA Community Members, 2006) ice cores) and the ice layers spanning the last deglaciation can be located close to bedrock where accurate dating is challenging. Moreover coastal cores often undergo high variability of snow accumulation due to wind scouring, which can also affect the dating quality. For example the Siple Dome ice core situated west of the Ross Sea has recorded a nearly negligible snow deposition rate around 15 kyr BP, leading to hiatus in the isotopic signal (Brook et al., 2005).

In order to document the regional variability of Antarctic climate, and its time relationship with climatic changes in other regions of the Earth (such as Greenland and the North Atlantic), a new deep ice core (1620 m) has been recovered at 5 km from the peripheral dome of Talos Dome (159°11' E, 72°49' S, 2315 m) on the eastern edge of the East Antarctic plateau during the field seasons 2004–2008. The international TALDICE (TALos Dome Ice CorE) project (<http://www.taldice.org>) conducted by a consortium of five European nations led by Italy, is a contribution to the international project IPICS (International Partnerships in Ice Core Science; <http://www.pages-igbp.org/ipics/index.html>) of the International Polar Year 2007/09. Talos Dome is an ice dome, roughly equally distant (~300 km) to the Southern Indian Ocean and the Ross Sea. Its near-coastal location leads to a current accumulation rate of 8.5 cm ice equivalent per year (cm ie/yr) (Frezzotti et al., 2007; Stenni et al., 2002) i.e. about three to four times higher than on the East Antarctic plateau, providing decadal time resolution during the last deglaciation period.

**TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica**

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In addition to its focus on the last deglaciation (Stenni et al., 2010), the TALDICE project also intends to investigate the millennial-scale variability of the last glacial period, down to at least 50 kyr BP, and to compare its shape and phasing with respect to other East-Antarctic ice core records. Studying the sequence of climatic events between different areas directly relies on the quality of the ice core chronologies. The main techniques currently available to build an ice core age scale can be grouped into four categories: (1) layer-counting, a technique which requires annual layer resolution in snow deposition and in trace element measurements, unfortunately not applicable in most Antarctic sites because the accumulation rates are not sufficient; (2) tuning of ice records to insolation changes which is only applicable for long (>100 kyr) records and usually results in large dating uncertainties (several kyr); (3) matching age markers to other well-dated time series; (4) glaciological models simulating the thinning and flow of snow/ice layers through the ice sheet.

Absolute ages are needed to analyse the links between climate and orbital forcing (e.g. (Kawamura et al., 2007), and to depict the sequences of events recorded in different archives. Air trapped in polar ice cores have the unique property of containing global tracers of the atmosphere such as CH<sub>4</sub> and  $\delta^{18}\text{O}_{\text{atm}}$  that globally display the same temporal variations on the different drilling sites. It is thus possible to build a relative dating of different polar ice cores that can provide insights on the timing of climatic events between Greenland and Antarctica and between different sites in Antarctica. A firn core was dated at Talos Dome by using seasonal variations in non-sea-salt sulphate concentrations coupled with the recognition of atomic bomb marker level (1965–1966) and sulphate spikes due to the most important volcanic events in the last 800 years (Stenni et al., 2002). Due to incorrect identification of missing seasonal signals and absence of absolute volcanic chronology before 1000 AD, it is impossible to build an absolute layer-counting age scale for ages older than the last millennium. As a consequence, we chose to build a relative glaciological age scale for Talos Dome, based on the absolute GICC05 age scale from Greenland down to 50 kyr (Svensson et al., 2008) and on the EDC3 age scale from Antarctica for older ages (Parrenin et al.,

2007a), using both records synchronizations and glaciological modelling.

A new dating method based on inverse techniques has been recently developed (Lemieux-Dudon et al., 2009). It first requires background scenarios for the accumulation rate, the thinning function and the Close-Off-Depth in Ice Equivalent (CODIE) and the difference of depth for a given age between the trapped gas and the ice matrix ( $\Delta$ depth), which must be provided by simulations (ice flow and snow densification modelling). It is then complemented with chronological data provided by age markers (data enabling to associate a time with a specific depth). The efficiency of this method has already been underlined by improving the consistency between EPICA/Dome C (EDC), EPICA Dronning Maud Land (EDML), and NorthGRIP age scales (Lemieux-Dudon et al., 2010) over the last 50 kyr.

We aim here at applying this inverse method to obtain the TALDICE-1 age scale. We provide the background scenario by carrying simulations with a 1-D ice flow model (Parrenin et al., 2004), coupled with a densification model (Arnaud et al., 2000). We define age markers through the use of  $\text{CH}_4$  and water isotopic profiles in the TALDICE and other ice cores. In order to match our new age scale to the absolute GICC05 age scale from Greenland (Svensson et al., 2008), most tie points are chosen through the synchronization of the TALDICE and Greenland methane records.

In this paper we first present the tie points (Sect. 2.4) and describe the background scenario (Sect. 3) and the use of the inverse method (Sect. 4). Results and uncertainties are then discussed, including a posteriori controls of the resulting TALDICE-1 age scale and of the glaciological entities reconstructed by the model.

## 2 Data and methods

The quality of the chronology strongly depends on the number, distribution and quality of tie points along the record. The central tool used here to define tie points relies on synchronizing the TALDICE and other ice core  $\text{CH}_4$  profiles.

### TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**TALDICE-1 age scale  
of the Talos Dome  
deep ice core, East  
Antarctica**

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CH<sub>4</sub> measurements have already been used many times for relative dating of ice cores (Blunier and Brook, 2001; Blunier et al., 1998, 2007; Chappellaz et al., 1997). During glacial abrupt events, CH<sub>4</sub> shows large and abrupt changes with time, usually associated with Greenland temperature changes (Huber et al., 2006). Its atmospheric lifetime of about 10 yr (IPCC, 2001) is larger than the interhemispheric mixing time (~1 yr), thus making its mixing ratio changes of global significance. Similar CH<sub>4</sub> records should thus be obtained in any ice core of both hemispheres. The limits of the CH<sub>4</sub> tie points are (1) the existence of an inter-polar gradient, not always well known, bringing error in tie point definition when CH<sub>4</sub> changes abruptly (Dällenbach et al., 2000); (2) the smoothing of atmospheric composition changes which can differ from one ice core to the other due to different gas trapping conditions (Spahni et al., 2003); (3) the presence of time intervals where CH<sub>4</sub> does not vary significantly, such as Marine Isotopic Stage 2 (MIS 2). Still, CH<sub>4</sub> is particularly useful to date new Antarctic ice cores by matching them with the Greenlandic ones, as the reference CH<sub>4</sub> signal is well-known back to ~110 kyr BP (Brook et al., 1996), and as the Greenland ice cores can be absolutely dated by annual layer counting (Andersen et al., 2006; Svensson et al., 2008).

Below 1428 m of depth in the TALDICE core, corresponding to a gas age of ≈141 kyr BP, the resolution of our CH<sub>4</sub> measurements is not good enough for synchronization purposes. We thus extend the chronology by matching the TALDICE δ<sup>18</sup>O water isotopic profile with the EDC one on the EDC3 age scale (Parrenin et al., 2007a), making the assumption that the main features of climatic variations occurred in phase at both sites. This part of the chronology could be refined later on when more gas measurements will be available.

Finally a posteriori control of the final gas age scale have been provided by measurements of the isotopic composition of molecular oxygen contained in the air enclosed in the ice (δ<sup>18</sup>O<sub>atm</sub>).

## 2.1 Methane measurements

CH<sub>4</sub> has been measured both at Grenoble (LGGE) and Bern laboratories, using two different methods. First, both LGGE and Bern used the same method as for previous ice cores, consisting of a wet extraction method combined with gas chromatographic analyses (Chappellaz et al., 1997; Spahni et al., 2005) The obtained LGGE values are increased by 6 ppbv in order to be in good agreement with Bern values (Spahni et al., 2005).

798 samples have been measured between 73 and 1620 m of depth, with a mean 1 $\sigma$  error of 10 ppbv. The mean depth resolution is 2 m, and increases to 0.5 m during sharp transitions. Second, part of the record (between 667 and 817 m and between 1132 m and 1487 m of depth) has been measured using a new on-line technique where the gas coming out of a continuous-flow analysis (CFA) system is separated from water with a membrane and then transferred to a gas chromatograph, thus providing ~20 cm depth resolution, but with a lower precision of the measurements (1 $\sigma$  of 15 to 20 ppbv; Schüpbach et al., 2009).

## 2.2 Water isotopic measurements

Measurements of the oxygen isotope composition ( $\delta^{18}\text{O}$ ) of water in the TALDICE samples were conducted on discrete samples averaging over a length of 1 m. The  $\delta^{18}\text{O}$  measurements were carried out in France (Laboratoire des Sciences du Climat et l'Environnement) and in Italy (University of Trieste and Parma) using the well-known CO<sub>2</sub> water equilibration technique (Epstein et al., 1953). Intercomparisons of reference waters among the involved laboratories were conducted over the analysis period and the same water standard was used in all laboratories. The precision of  $\delta^{18}\text{O}$  measurements is  $\pm 0.05\text{‰}$ . The  $\delta^{18}\text{O}$  data provided here are the results of at least duplicate measurements over 1620 samples.

## 2.3 Molecular oxygen isotopic measurements ( $\delta^{18}\text{O}_{\text{atm}}$ )

We measured the  $\delta^{18}\text{O}_{\text{atm}}$  on 145 TALDICE samples between 583 and 1402 m, using the analytical technique described in (Landais et al., 2003). The mean  $2\sigma$  precision is 0.05‰.

## 2.4 Synchronization and age marker assignments

Down to 1191 m of depth ( $\approx 50$  kyr BP) we matched the  $\text{CH}_4$  profile of TALDICE with the Greenland Composite methane curve published by Blunier et al. (2007) which is based on the absolute layer-counted GICC05 age scale. The GICC05 age scale is considered as the current reference for ice cores absolute dating down to about 50 kyr BP (Svensson et al., 2008), what allows to consider that the Taldice chronology should show a good coherency with absolute ages during this period.

Before 50 kyr BP, the most accurate and detailed  $\text{CH}_4$  profile available is the one measured on the EDC ice core (Loulergue et al., 2008). We thus extended the  $\text{CH}_4$  synchronization between TALDICE and EDC profiles on the EDC3 age scale between 50 and 141 kyr BP (back to the onset of the penultimate deglaciation), corresponding to the depth range 1228-1428 m in the TALDICE ice core. The shift of the TALDICE matching between Greenland and EDC  $\text{CH}_4$  records takes place at 53.5 kyr BP (tie point corresponding to the Dansgaard/Oeschger #14 warming). A similar method has already been applied at EDML (EPICA Community Members, 2006) by synchronizing EDML and Greenland Composite methane profiles in order to investigate the Greenland-Antarctic timing of events down to 55 kyr BP.

$\text{CH}_4$  synchronization is performed by visual wiggle-matching, tie points being chosen at mid-slope or maximum or minimum of  $\text{CH}_4$  changes, depending on the signal shape. For the synchronization exercise between TALDICE and EDC ice cores we consider that each tie point between TALDICE and EDC should bear the same  $\text{CH}_4$  mixing ratio (no influence of the interpolar gradient). With this method, forty-four age markers have been defined for the gas phase of the TALDICE core (Figures and Tables 1 and 2).

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**TALDICE-1 age scale  
of the Talos Dome  
deep ice core, East  
Antarctica**

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We also define fourteen tie points between TALDICE and EDC  $\delta^{18}\text{O}$  records, at mid-slope or maximum of the sharpest climatic transitions between 1443 and 1556 m in TALDICE, covering the period between 159.2 and 280 kyr BP on the EDC3 time scale (Table 3). This relies on the hypothesis of synchronous climate changes in Eastern Antarctica on multi-millennial time scales, as suggested by the coherency observed between various ice core records (Watanabe et al., 2003). This assumption is also corroborated by the comparison of TALDICE and EDC in the younger part of cores (Stenni et al., 2010). Current  $\text{CH}_4$  data resolution in the corresponding TALDICE depth range becomes too low to provide good markers with respect to the EDC  $\text{CH}_4$  profile. Contrary to the first forty-four  $\text{CH}_4$  tie points, the additional fourteen ones defined by the  $\delta^{18}\text{O}$  records are located in the ice phase.

The uncertainty on the absolute age of each tie point is obtained by adding the uncertainty of the visual matching between  $\text{CH}_4$  or  $\delta^{18}\text{O}$  records, and the inherent uncertainty of the GICC05 and EDC3 age scales at the corresponding ages. It also includes the error induced by the interhemispheric  $\text{CH}_4$  gradient that leads to synchronizations errors reaching up to 50 yr (Dällenbach et al., 2000). The visual matching uncertainty amounts to less than 50 yr during the sharp transitions corresponding to the BA and Younger Dryas (YD) events, taking also into account the analytical uncertainty. On the other hand, the uncertainty amounts to several millennia for the oldest TALDICE ice, the largest portion of it being related with the poor resolution of the available  $\delta^{18}\text{O}$  profile. Future  $\delta^{18}\text{O}$  measurements at 5-cm depth resolution will help us to improve this part of the TALDICE time scale.

In total, fifty-eight age markers are now available to constrain the dating scenarios provided by glaciological models, the highest concentration of tie points being during the deglaciation (11 to 18 kyr BP) and during MIS3 (29 to 55 kyr BP) periods.

### 3 A priori scenario

The inverse method (Lemieux-Dudon et al., 2010) used to build the TALDICE-1 age scale requires prior guess for the accumulation rate (expressed in ice equivalent (ie)), the thinning function, and the CODIE. The latter is critical as the synchronization of CH<sub>4</sub> records leads to age constraints on the gas chronology. In order to translate the CH<sub>4</sub> time markers into chronological constraints on the ice chronology, we need to estimate past TALDICE close-off depths (COD) where air bubbles are formed, and thus the age difference between the trapped gases and the surrounding ice ( $\Delta$ age). At first order, both parameters depend on the accumulation rate and on surface temperature, determining for any given site at which depth pores get enclosed at the bottom of the firn.

#### 3.1 Prior guess: 1-D ice flow model and snow densification model

In order to calculate a first a priori scenario for the three entities, we use the 1-D ice flow model developed by (Parrenin et al., 2007b) which combines ice flow modeling with a firn densification model (Arnaud et al., 2000). It calculates the age of the ice at a given depth with the following equation:

$$\text{age}(z) = \int_0^z \frac{1}{T(z)az} dz \quad (1)$$

where  $z$  is depth (m),  $T$  is the thinning ratio and  $a$  is the initial annual layer thickness (i.e. accumulation rate in ice equivalent), deduced from the isotopic composition of the ice. The required density profile has been measured in the field at Talos Dome.

The ice flow model is then coupled with a firn densification model (Arnaud et al., 2000) which uses the ice age scenario combined with past accumulation rate and temperature scenario deduced from the water isotopic data to provide past CODIE and  $\Delta$ age scenario, enabling to calculate a chronology associated with the gas phase of the ice core.

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Past accumulation rates and temperature changes are usually deduced from the deuterium content of the ice ( $\delta D$ ). Since the full TALDICE  $\delta D$  record is not available yet, the  $\delta^{18}\text{O}$  data have been multiplied by 8 (Merlivat, 1978) to build a synthetic  $\delta D$  record. This approximation is based on the assumption that the second order parameter deuterium excess ( $d = \delta D - 8 \delta^{18}\text{O}$ ) is negligible, but we must keep in mind that recent studies have shown variations of  $\sim 10\%$  of  $d$  in Antarctica during the last deglaciation period (unpublished data). We finally used the following equations:

$$T = T^0 + \alpha \Delta \delta D_{\text{corr}} \quad (2)$$

and

$$A = A^0 \exp(\beta \Delta \delta D_{\text{smo}}) \quad (3)$$

where  $T^0$  and  $A^0$  are the present-day surface temperature ( $^{\circ}\text{C}$ ) and accumulation rate (cm of ice equivalent per year) at the drilling site.  $\Delta \delta D_{\text{corr}}$  corresponds to the temporal deviation from the present-day deuterium content, corrected for past variations in the global ocean using the  $\delta^{18}\text{O}$  record derived from the benthic foraminifera stacks (Lisiecki and Raymo, 2005), itself corrected for deep-water temperature effects.  $\Delta \delta D_{\text{smo}}$  is a 50-yr average of  $\Delta \delta D_{\text{corr}}$  and  $\alpha$  represents the slope of the present-day spatial isotopic thermometer.  $\beta$  is a coefficient defining the glacial-interglacial amplitude of accumulation changes.

The model contains several poorly known parameters (namely  $\alpha$ ,  $\beta$ ,  $p$ ,  $\Delta H$ ) and also requires input information about current conditions prevailing at the site (see Table 4). Unknown parameters are determined here by carrying several simulations until the best overall fit with the independent age markers is obtained.

### 3.2 Elevation changes at Talos Dome

The thinning function at TALDICE could have been affected by past elevation changes at the dome. It is thus important to estimate and take these changes into account in

the ice flow simulations. Constraints on past elevation changes in the Talos Dome area are available since the last deglaciation; they are summarized in Table 5. Geomorphological studies carried through the Outback Nunataks range and in the Ross Sea sector revealed past glacier tracks, suggesting an increase of the dome altitude to a maximum of  $170\pm 20$  m above the current altitude, reached between 19 and 13 kyr BP. Elevation changes during the Last Glacial Maximum (LGM) have been triggered by the progression of the grounding line up to the continental margin in response to the glacial sea level lowering (Anderson et al., 2002). The maximum elevation change is consistent with analyses of moraine deposits in the upper part of Rennick Glacier (Meneghel et al., 1999), a few tens of km East of Talos Dome, which suggest an increase of the glacier thickness by about 200 m during the first part of the last termination.

The increased dome elevation during part of the deglaciation was favoured by the increase of accumulation rate. We suggest that the dome started to lower its elevation when the grounding line retreated from its LGM position between 14.3 and 12 kyr BP (Anderson et al., 2002). Mackintosh et al. (2007) measured exposure ages of mountains dipsticks across the Framnes Mountains in Mac Robertson Land in East Antarctica and concluded to a maximum thickness of the East Antarctic ice sheet around 13 kyr BP. Studies performed in other coastal Antarctic areas at the onset of the Holocene around 10 kyr BP, such as moraine dating in the Mary Byrd Land region (Ackert et al., 1999), suggest that the elevation was still  $100\pm 50$  m higher than today. We thus extend these conclusions to the case of Talos Dome as well. Radiochronological measurements on penguin guano, shells and seal skin (Baroni and Hall, 2004) suggest that the retreat of the East Antarctic ice sheet margin reached Terra Nova Bay shortly before 7 kyr BP. Such evolution of the dome elevation in the course of the last deglaciation and the early Holocene contrasts with its counterpart on the East Antarctic plateau, where the altitude was lower during the LGM, due to the dominant effect of reduced glacial accumulation rates (Ritz et al., 2001). Glacial geology observation and  $C^{14}$  dating point out that the Ross Ice Sheet started to grow at Mc Murdo Sound around 30 kyr BP (Baroni, C., personal communication, 2010). We thus make the assumption

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that the altitude of the dome was similar to today at 30 kyr BP and progressively increased toward its maximum at around 16 kyr BP.

Past elevation changes at TALDICE are incorporated as input information in the 1-D ice flow model.

### 5 3.3 Results obtained with the 1-D ice flow model

The resulting a priori chronology appears in good agreement with some age markers, while differences remain with others (Fig. 3). The use of a glaciological model tends to smooth the depth-age relationship, leading to disagreements of up to 30 000 yr with some of the tie points older than 140 kyr BP.

10 Actually because of the numerous poorly-known parameters, as well as a simple description of the ice flow, the 1-D model is not able to take into account the ice anisotropy, eventual past migrations of the dome summit, or any piece of information related to lateral limits, inducing errors in the thinning function scenario. Besides it also neglects the fact that relationships between accumulation rate, temperature and water isotopic  
15 data could have been different in the past (Masson-Delmotte et al., 2006; Stenni et al., 2001).

## 4 TALDICE-1 age scale based on an inverse method

### 4.1 Principles

20 The inverse method used here to improve the glaciological model output is a probabilistic inverse approach based on a Bayesian inference. It attempts to find the best compromise between the a priori ice flow model scenario and chronological information from tie points (Lemieux-Dudon et al., 2009). It optimizes the three key glaciological entities that vary along the core: the accumulation rate (expressed in ice equivalent (ie)), the total thinning function, and the CODIE.

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The probabilistic combination of uncertainties of the model and of the tie points also enables to estimate confidence intervals on the new dating.

## 4.2 Input parameters

In addition to the background scenario (based on ice flow and snow densification simulations) a background error covariance matrix, which accounts for the uncertainties on the background dating scenario must be assessed in order to run the inverse method. In this study, we made the following assumptions: (i) no error correlation between the accumulation rate and the thinning function, as argued by (Parrenin et al., 2007b) who show that the total thinning function is only weakly sensitive to accumulation changes; (ii) no error correlation between the accumulation rate and CODIE (further studies are necessary to test this latter assumption). The resulting error covariance matrix is therefore a diagonal block with three sub-matrices, respectively associated to the three glaciological entities (thinning, accumulation and CODIE). These sub-matrices are fully defined by setting a standard deviation vector and a correlation matrix.

The standard deviation and correlation profiles were chosen of the same order of magnitude as proposed in (Lemieux-Dudon et al., 2009, 2010). A constant standard deviation with depth (set to 40%) is assessed for the accumulation rate and the CODIE (more precisely to the logarithm of these quantities because errors are assumed to be lognormally distributed), while a growing function with depth is chosen for the thinning function (Lemieux-Dudon, 2009, 2010). The correlation profiles are normalized Gaussian functions with a correlation length parameter. This latter parameter is in time unit for the accumulation and CODIE and set to 4000 yr. It is in depth unit for the thinning function and set to 50 m. Further statistical analysis shall be performed to better constrain these parameters. A posteriori controls on the background settings can be conducted by testing the new chronology against independent paleoclimatic data (see Sect. 5.1.3).

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 5 Results and discussion

### 5.1 TALDICE ice and gas chronologies and confidence intervals

#### 5.1.1 The new gas and ice age scales

Figure 2 shows the age/depth relationship at TALDICE according to the a priori glaciological model and after using the inverse method. They are both compared to age markers, and the confidence interval on the ice chronology calculated by the inverse method is also provided.

The TALDICE-1 chronology provides a very good agreement with all age markers. As already observed by (Lemieux-Dudon et al., 2010) the relative chronological uncertainty decreases at the location of age markers and increases during periods where few or no age markers are available. In addition, the uncertainty increases with depth.

The TALDICE-1 age scale provides an uncertainty of  $\sim 300$  yr over the last deglaciation and is lower than 600 yr during the D/O events of MIS 3. The relative uncertainty reaches 2 kyr at the MIS5.5 onset. By comparison with the EDC3 and EDML1 time scales (Parrenin et al., 2007a; Ruth et al., 2007), their uncertainty over the last deglaciation, the MIS 3 period and the MIS 5.5 onset is estimated to be 0.4 kyr, 1.5 kyr and 3 kyr, respectively. The relatively small uncertainty in the TALDICE-1 chronology enables us to investigate in detail the timing of the millennial-scale climatic events throughout the last climatic cycle.

Before the last interglacial period, the errors associated with tie points becomes considerably larger than for younger TALDICE ice; the chronological uncertainty remains lower than 3 kyr down to 205 kyr BP, and then largely increase up to  $\sim 30$  kyr for the oldest part of the record.

CPD

6, 1733–1776, 2010

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 5.1.2 Corrections applied to the background scenario

Figure 3 shows the ratio between the accumulation rate, thinning rate and CODIE values deduced from the inverse method and those estimated by the background scenario, focusing on the last 50 kyr where age markers are more frequent.

Corrections applied to the accumulation rate usually remain lower than the  $\pm 20\%$  error margin usually considered for the accumulation rate deduced from isotopic data (Blunier et al., 2007; Huber et al., 2006). Only during the MIS 2 period the inverse method requires a decrease of 20% compared to the initial accumulation rate.

Corrections applied to the CODIE do not exceed 16%, and are consistent with typical uncertainties associated with modeled and measured close-off depth (Goujon et al., 2003; Landais et al., 2006).

The correction applied to the modeled thinning function remains below 5% for the last 16 kyr. It then increases to about 22% at 25 kyr BP, followed by a decrease and a constant level below 15% until around 40 kyr BP where small successive DO events occur. The tie point assignment becomes more uncertain during this time interval and leads to larger changes in the thinning function deduced by the inverse method.

## 5.1.3 A posteriori evaluation of the TALDICE-1 age scale

### A posteriori evaluation with a firn densification model

The CODIE determined with the inverse method can be evaluated a posteriori by transforming it into a  $\Delta$ age and by comparing it with the output of the firn densification model of Goujon et al. (2003) which calculates the CODIE and  $\Delta$ age using the temperature and accumulation rate scenarios from the inverse model. Indeed surface temperature and accumulation rate are known to be the two main parameters controlling firn densification (Schwander et al., 1997).

The  $\Delta$ age deduced from the CODIE of the inverse method and the one calculated by the firn densification model agree very well all along the core (Fig. 4). The inverse

CPD

6, 1733–1776, 2010

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





model-based  $\Delta$ age remains well within the range of  $\Delta$  ages provided by the densification model when accumulation rates vary in a  $\pm 20\%$  range around the accumulation rate calculated with the inversion. The mean difference between both gas chronologies amounts to only 16 yr. However, differences reach up to 450 yr over D/O events #9, 10 and 11. This reflects the difficulty to define accurate  $\text{CH}_4$  tie points for these small events.

### A posteriori evaluation with a dust event and volcanic sulfate peaks

Ice core dust records from the East Antarctic plateau reveal a large decrease from the LGM to the Holocene (Delmonte et al., 2002). This decrease is also observed at TALDICE, with a relatively sharp minimum at 764.9 m of depth, occurring almost in phase with the Antarctic Isotopic Maximum (AIM) 1. The EDC counterpart for this feature is observed at 421.3 m of depth (Stenni et al., 2001). As the dust source at both sites is similar (Patagonia) (Delmonte et al., 2010) the dust minimum should be synchronous. It has been dated at  $14.7 \pm 0.1$  kyr BP at EDC according to the new EDC age scale (Lemieux-Dudon et al., 2010). The TALDICE-1 age scale dates it at  $14.8 \pm 0.1$  kyr BP.

The validity of the dust control point is confirmed using three sulfate volcanic peaks observed in both EDC and TALDICE and expected to be synchronous (M. Severi, personal communication, 2010). For instance the deepest of the three sulfate peaks recorded at 766.09 m of depth at TALDICE and dated at  $14.9 \pm 0.1$  ky according to the TALDICE-1 age scale, corresponds to the volcanic peak observed at 422.07 m of depth at EDC and dated at  $14.7 \pm 0.1$  ky according to the new EDC chronology (Lemieux-Dudon et al., 2010). The dust and volcanic sulfate peaks thus strengthen our confidence in the TALDICE-1 chronology in the middle of the last deglaciation.

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## A posteriori evaluation using $\delta^{18}\text{O}$ of molecular oxygen

As in the case of atmospheric  $\text{CH}_4$ , the isotopic composition of molecular oxygen in the atmosphere ( $\delta^{18}\text{O}_{\text{atm}}$ ) is globally homogeneous and shows changes associated with climate variations (Bender et al., 1994; Sowers et al., 1991). It can thus complement  $\text{CH}_4$  tie points, especially during periods of time when the atmospheric  $\text{CH}_4$  mixing ratio does not vary much in the atmosphere. Down to 50 kyr BP we choose to compare TALDICE  $\delta^{18}\text{O}_{\text{atm}}$  to the EDML record on the new EDML age scale (Lemieux-Dudon et al., 2010), both chronologies being thus tied to the GICC05 chronology (Fig. 5). For older ages, where TALDICE methane is synchronized to the EDC one on the EDC3 chronology, we compare TALDICE and EDML  $\delta^{18}\text{O}_{\text{atm}}$  records on the EDML1 age scale which is tied to the EDC3 one (Ruth et al., 2007), and to the Vostok  $\delta^{18}\text{O}_{\text{atm}}$  record (Petit et al., 1999) which has been previously put on the EDC3 age scale (Loulergue et al., 2008). New EDML  $\delta^{18}\text{O}_{\text{atm}}$  measurements before 80 kyr BP were performed at LSCE with the same analytical method.

The three  $\delta^{18}\text{O}_{\text{atm}}$  records appear in very good agreement. We particularly focused our measurements on the deglaciation period, where we observe an extremely good fit between EDML and TALDICE records. Those observations strengthen again our confidence on the final TALDICE-1 gas age scale and on the choice of the preliminary age markers.

The few points measured between 60 and 85 kyr BP and after 100 kyr BP enable to catch the sharp  $\delta^{18}\text{O}_{\text{atm}}$  transition between 70 and 80 kyr BP, taking place at a time of reduced  $\text{CH}_4$  variations. It reasonably confirms the suitability of TALDICE-1 for this time interval.

CPD

6, 1733–1776, 2010

### TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 5.2 How physically realistic are the glaciological parameters resulting from the inverse model?

### 5.2.1 Accumulation rate reconstruction

#### Comparison with the water isotopic signal

Overall, the accumulation rate calculated by the inverse method over the full TALDICE time scale is very close to the one deduced from the water isotopes, as can be seen in Fig. 6, thus giving confidence that the standard relationship expected between the accumulation rate and isotopic data is also valid at Talos Dome on long time scales (Frezzotti et al., 2004, 2007; Magand et al., 2004). Still, a difference of 25% is observed around 23 kyr BP (Fig. 6). The amplitude of the last glacial-interglacial accumulation rate changes thus appears slightly more pronounced with the scenario from the inverse method or, conversely, underestimated with the classical approach. These differences could be partly explained by the fact that the 1-D model is not able to simulate whether relationships between accumulation rate/temperature and isotopic data have been different in the past (Masson-Delmotte et al., 2006; Stenni et al., 2001).

As previously explained, the link between the accumulation rate and the water isotopic signal is represented in the models by the parameter  $\beta$ , which consists in a constant value in the classical approach. However the inverse method requires a  $\beta$  value varying in time, around a mean value of 0.0198 associated with a correlation coefficient  $R^2=0.93$  between accumulation rate and deuterium data. We add to the Fig. 6 the accumulation rate calculated for a constant  $\beta=0.0198$ . It seems that the inverse accumulation rate mainly differs from this relationship in the period ranging between the AIM1 and the Early Holocene, showing higher values, whereas it differs from the direct model accumulation rate during LGM low values.

The large accumulation rate decrease during the LGM, and to a lesser extent during MIS 6, could reflect the sensitivity of this coastal site to changes of moisture source due to sea ice extent and presence of an ice sheet in the Ross sea embayment, both

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



possibly affecting snow precipitation linked to cyclonic activity and wind-scouring. Indeed, radar layer studies have shown that the LGM snow accumulation became nearly negligible at Taylor Dome (also in the Ross sea embayment), probably in response to changes in storm tracks (Morse et al., 1998, 2007). Talos Dome is influenced by moisture originating mainly from the Indian (around 50 %) and in a lesser extent from the Pacific/Ross Sea (30%) sectors of the Southern Ocean (Scarchilli et al., 2010). The snowfalls originating from the Pacific Ocean reach Talos Dome after crossing the Ross Sea, where extensive presence of sea ice occurs also during summer time. During the LGM and most of the deglaciation, the presence of the Ross Ice Sheet extending up to the continental margin could have contributed to dry the air masses coming from the Ross Sea area. The same phenomenon could explain the different ratio between accumulation and water isotopes between the LGM and the Holocene.

### A posteriori control with $^{10}\text{Be}$ data

The cosmogenic isotope Beryllium-10 ( $^{10}\text{Be}$ ) signal in ice cores is influenced by changes in its production, deposition processes and snow accumulation. During a deglaciation, the snow accumulation drastically changes and at first order,  $^{10}\text{Be}$  concentration can be used to constrain it. Previous studies have shown that  $^{10}\text{Be}$  is mostly dry deposited on the East Antarctic plateau, enabling to directly connect  $^{10}\text{Be}$  concentrations in the ice [ $^{10}\text{Be}$ ] with the snow accumulation rate (Delmas, 1992; Raisbeck and Yiou, 1985), adding a constant term for the wet deposition effect.

$^{10}\text{Be}$  measurements have been performed on TALDICE ice covering the last deglaciation on the new Accelerator Mass Spectrometer, ASTER, installed at CEREGE. They show a near-doubling of [ $^{10}\text{Be}$ ] in LGM ice compared with the Holocene one, similar to the Vostok Antarctic site (Raisbeck et al., 1992) or to the GISP2 and GRIP Greenland sites (Finkel and Nishiizumi, 1997; Raisbeck et al., 1992).  $^{10}\text{Be}$  concentration and snow accumulation rate are inversely correlated. Calculating the ratios [ $^{10}\text{Be}$ ]<sub>AIM1</sub>/[ $^{10}\text{Be}$ ]<sub>Holocene</sub> and [ $^{10}\text{Be}$ ]<sub>LGM</sub>/[ $^{10}\text{Be}$ ]<sub>Holocene</sub>, we obtain 1.23 and 1.82, respectively. The former is similar to the Holocene/AIM1 ratio of accumulation

rate deduced from the inverse method. On the other hand the inverse method provides a Holocene/LGM accumulation rate ratio of 2.20, whereas the ratio deduced from water isotopes is 1.80 (Fig. 6). This means that either the more reduced accumulation rate during the LGM deduced from the inverse method is incorrect (meaning that the method gives too much weight to accumulation rate changes relative to thinning function changes) or that other parameters than accumulation rate, such as cosmogenic isotope production, wet versus dry deposition changes and/or, dust and aerosol variations or atmospheric transport, also modulated the TALDICE LGM/Holocene [ $^{10}\text{Be}$ ] values.

## 5.2.2 Thinning function

Broadly speaking, the a priori ice-flow model calculations provide a thinning function in good agreement with the one deduced by the inverse method, especially once altitudinal changes have been taken into account over the deglaciation. However the dome may have slightly moved through time (Urbini et al., 2008), generating a different deformation regime than uniaxial compression over a certain depth range in the ice. Thus local differences between the ice-flow and inverse thinning function can be expected and are indeed observed.

There is no direct proxy to quantify the thinning function with depth. Some physical properties of ice, e.g. the *c*-axis orientation (fabric) and the crystal size, allow one to qualitatively assess changes in the deformation regime, which could reflect and/or generate changes in the local thinning, in association with changes in ice properties.

The *c*-axis distribution of crystals reflects their orientation and is known to be a consequence of the strain-rate history experienced by ice during transport from the ice sheet surface (Castelnau and Duval, 1994; Durand et al., 2006). *C*-axes rotate during the deformation towards the main direction of compression and away from the tensional direction (Van der Veen and Whillans, 1994). Consequently, the resulted anisotropy of the ice will affect the flow of ice (Gagliardini and Meyssonier, 2000; Mangeney et al., 1997). We can thus qualitatively compare the crystal orientation with the reconstructed

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



thinning function, to look for possibly common patterns.

The fabric evolution has been measured on the TALDICE ice using an automatic fabric analyser (Montagnat, in preparation). The *c*-axis orientation data are expressed as the orientation tensor  $a^{(2)}$  as defined in (Durand et al., 2006). The eigenvalues of this orientation tensor are used to express the evolution of the *c*-axis orientation with depth (Durand et al., 2006; Montagnat et al., 2010). The evolution of the eigenvalues  $a_1^{(2)}$ , expressing the progressive fabric clustering toward the vertical direction (a value of 1 means that all crystal *c*-axes are vertically oriented and 3 would correspond to an isotropic fabric), is plotted in Fig. 7 and compared with the modeled thinning functions.

Several interesting features show up in the common trends between crystal orientation evolution and thinning functions:

- Around 700–750 m (11.5–12.4 kyr BP) the fabric evolution shows an increasing slope at the time when the ice-flow and inverse thinning functions start to diverge in their main trend.
- Between 800 and 900 m (16.4–25.2 kyr BP) a clear increase in the rate of fabric clustering appears, in relation with the largest divergence between the inverse thinning function and the one resulting from the ice-flow model without altitudinal changes. In parallel a reduction in mean crystal size has been observed (Montagnat, n.d.), suggesting a pinning effect on grain growth associated with increasing dust concentration in the same depth range, an effect similar to observations on the EDC ice core (Durand et al., 2007). Smaller grains lead to a change in ice rheology and ice viscosity (Cuffey et al., 2000), providing a causal relationship with the increased thinning function deduced from the inversion.
- Both fabric and inverse thinning function evolutions become negligible, a coherent feature as the thinning function is directly related with the strain evolution.
- Around 1100–1150 m (42.8–46.6 kyr) both the fabric and the thinning function evolution record an increasing rate of change.

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Around 1200–1250 m (51.2–58.0 kyr) the fabric evolution gets very low again while the evolution of the thinning rate slows down. A maximum of fabric concentration could be reached at this point.

5 These observations point to the limits of the 1-D ice flow model estimate of the thinning function and confirm added value of an inverse approach. In order to come now to a quantitative and mechanistic link between crystal evolution and thinning function at the dome, an anisotropic 3-D ice flow model would be required (Durand et al., 2006; Gillet-Chaulet et al., 2006).

## 6 Conclusions

10 Thanks to the quality of CH<sub>4</sub> tie points, to the implementation of a new inverse method for dating ice cores, and to the availability of a posteriori information allowing us to evaluate the produced chronology, we have built the first official dating – called TALDICE-1- of the new coastal ice core drilled at Talos Dome in Antarctica. Although this chronology will evolve in the future, in particular for periods of time older than 50 kyr BP and  
15 during the Holocene, with the addition of new control points and with better constraints on accumulation rate changes and on the thinning of ice layers with depth, it already provides us a solid background dating giving confidence in the conclusions raised from this new deep ice core, in terms of phasing between Greenland, the East Antarctic plateau and Talos Dome during the last deglaciation (Stenni et al., 2010) and during  
20 MIS 3 (Buiron et al., 2010).

The TALDICE-1 age scale provides an age uncertainty relative to the GICC05 age scale of about 300 yr for the ice record covering the last deglaciation; this uncertainty remains lower than 600 yr back to 50 kyr BP during the millennial-scale events which occurred during the last glacial period. The resulting gas and ice chronologies are  
25 consistent with  $\Delta$ age estimated by a firn densification model (Goujon et al., 2003), using the input climatic conditions resulting from the inverse method. Such consistency

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is usually a real challenge for ice core dating (Loulergue et al., 2007) and makes a clear bonus to use the TALDICE-1 chronology for further climatic investigations.

The next steps to improve the TALDICE age scale should include higher resolution records notably over the deepest part of the core (below 1420 m), and the use of the  $^{10}\text{Be}$  peak associated with the Laschamp event at 41.5 kyr BP, which will provide a firm marker of TALDICE  $\Delta\text{age}$  under glacial conditions (Loulergue et al., 2007).

Eventually proxies currently used as a posteriori information, like the  $\delta^{18}\text{O}_{\text{atm}}$  profile or dust and sulphate contents, could be later directly used as input chronological information in the inverse method, when additional data connected to the ice core age will be available during periods of low methane variations.

*Acknowledgements.* Talos Dome Ice core Project (TALDICE), a joint European program led by Italy, is funded by national contributions from Italy, France, Germany, Switzerland and the United Kingdom. The main logistical support was provided by PNRA at Talos Dome. We thank the logistic and drilling TALDICE team for their efficiency over each drilling season. Funding support for the dating work was provided by the Institut National des Sciences de l'Univers (INSU) through its program LEFE. We thank Mélanie Baroni from CEREGE for providing data and interpretation of  $^{10}\text{Be}$ , Mirko Severi for providing sulfate data and Barbara Delmonte for providing dust data. Thanks also to Frédéric Prié and Bénédicte Minster for their help during isotopic measurements at LSCE, Saclay. We are deeply indebted to our late colleague Jean-Marc Barnola for his help concerning the use of the firn densification model. This is TALDICE publication no. xx.



The publication of this article is financed by CNRS-INSU.

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## References

- Ackert, R. P., Barclay, D., Borns, H., Calkin, P., Kurz, M., Fastook, J., and Steig, E.: Measurements of Past Ice Sheet Elevations in Interior West Antarctica, *Science*, 286(5438), 276–280, doi:10.1126/science.286.5438.276, 1999.
- 5 Alley, R. B.: The Younger Dryas cold interval as viewed from central Greenland, *Quaternary Sci. Rev.*, 19(1–5), 213–226, doi:10.1016/S0277-3791(99)00062-1, 2000.
- Andersen, K. K., Svensson, A., Johnsen, S. J., Rasmussen, S. O., Bigler, M., Röthlisberger, R., Ruth, U., Siggaard-Andersen, M. L., Peder Steffensen, J., Dahl-Jensen, D. and others: The Greenland Ice Core Chronology 2005, 15–42ka. Part 1: constructing the time scale, *Quaternary Sci. Rev.*, 25(23-24), 3246–3257, 2006.
- 10 Anderson, J. B., Shipp, S. S., Lowe, A. L., Wellner, J. S., and Mosola, A. B.: The Antarctic Ice Sheet during the Last Glacial Maximum and its subsequent retreat history: a review, *Quaternary Sci. Rev.*, 21(1–3), 49–70, 2002.
- Arnaud, L., Barnola, J., and Duval, P.: Physical modeling of the densification of snow/firn and ice in the upper part of polar ice sheets, *Physics of Ice Core Records*, 285–305, 2000.
- 15 Baroni, C. and Hall, B. L.: A new Holocene relative sea-level curve for Terra Nova Bay, Victoria Land, Antarctica, *J. Quaternary Sci.*, 19(4), 377–396, doi:10.1002/jqs.825, 2004.
- Bender, M., Sowers, T., and Labeyrie, L.: The Dole Effect and its variations during the last 130,000 years as measured in the Vostok Ice Core, *Global Biogeochem. Cycles*, 8(3), 363–376, doi:199410.1029/94GB00724, 1994.
- 20 Blunier, T. and Brook, E. J.: Timing of millennial-scale climate change in Antarctica and Greenland during the last glacial period, *Science*, 291(5501), 109–112, 2001.
- Blunier, T., Chappellaz, J., Schwander, J., Dallenbach, A., Stauffer, B., Stocker, T. F., Raynaud, D., Jouzel, J., Clausen, H. B., Hammer, C. U. and Johnsen, S. J.: Asynchrony of Antarctic and Greenland climate change during the last glacial period, *Nature*, 394(6695), 739–743, doi:10.1038/29447, 1998.
- 25 Blunier, T., Spahni, R., Barnola, J.-M., Chappellaz, J., Loulergue, L., and Schwander, J.: Synchronization of ice core records via atmospheric gases, *Clim. Past*, 3, 325–330, doi:10.5194/cp-3-325-2007, 2007.
- 30 Brook, E. J., White, J. W., Schilla, A. S., Bender, M. L., Barnett, B., Severinghaus, J. P., Taylor, K. C., Alley, R. B., and Steig, E. J.: Timing of millennial-scale climate change at Siple Dome, West Antarctica, during the last glacial period, *Quaternary Sci. Rev.*, 24(12–13), 1333–1343,

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2005.

Brook, E. J., Sowers, T., and Orchardo, J.: Rapid Variations in Atmospheric Methane Concentration During the Past 110,000 Years, *Science*, 273(5278), 1087–1091, doi:10.1126/science.273.5278.1087, 1996.

5 Buiron, D., Stenni, B., Frezzotti, M., Chappellaz, C., Masson-Delmotte, V. and The TALDICE Consortium: A 50,000-year climatic record from the new coastal TALDICE ice core: consequences on millennial-scale variability features through the Antarctic continent, in preparation, 2010.

10 Capron, E., Landais, A., Lemieux-Dudon, B., Schilt, A., Masson-Delmotte, V., Buiron, D., Chappellaz, J., Dahl-Jensen, D., Johnsen, S. and Leuenberger, M.: Synchronising EDML and NorthGRIP ice cores using  $\delta^{18}\text{O}$  of atmospheric oxygen ( $\delta^{18}\text{O}_{\text{atm}}$ ) and  $\text{CH}_4$  measurements over MIS5 (80–123 kyr), *Quaternary Sci. Rev.*, 29(1–2), 222–234, doi:10.1016/j.quascirev.2009.07.014, 2010.

15 Castelnau and Duval: Simulations of anisotropy and fabric development in polar ices, *Ann. Glaciol.*, 20, 277–282, 1994.

Chappellaz, J., Brook, E., Blunier, T., and Malaizé, B.:  $\text{CH}_4$  and  $\delta^{18}\text{O}$  of  $\text{O}_2$  records from Antarctic and Greenland ice: A clue for stratigraphic disturbance in the bottom part of the Greenland Ice Core, *J. Geophys. Res.*, 102(C12), 26,547–26,557, doi:199710.1029/97JC00164, 1997.

20 Cuffey, K. M., Thorsteinsson, T., and Waddington, E. D.: A renewed argument for crystal size control of ice sheet strain rates, *J. Geophys. Res.*, 105(B12), 27,889–27,894, doi:200010.1029/2000JB900270, 2000.

25 Dällenbach, A., Blunier, T., Flückiger, J., Stauffer, B., Chappellaz, J. and Raynaud, D.: Changes in the atmospheric  $\text{CH}_4$  gradient between Greenland and Antarctica during the Last Glacial and the transition to the Holocene, *Geophys. Res. Lett.*, 27(7), 1005–1008, doi:200010.1029/1999GL010873, 2000.

Delmas, R. J.: Free tropospheric reservoir of natural sulfate, *J. Atmos. Chem.*, 14(1), 261–271, doi:10.1007/BF00115238, 1992.

30 Delmonte, B., Petit, J. R., and Maggi, V.: LGM-Holocene changes and Holocene millennial-scale oscillations of dust particles in the EPICA Dome C ice core, East Antarctica, *Ann. Glaciol.*, 35, 306–312, 2002.

Delmonte, B., Andersson, P., Schöberg, H., Hansson, M., Petit, J., Delmas, R., Gaiero, D., Maggi, V., and Frezzotti, M.: Geographic provenance of aeolian dust in East Antarctica

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



during Pleistocene glaciations: preliminary results from Talos Dome and comparison with East Antarctic and new Andean ice core data, *Quaternary Sci. Rev.*, 29(1–2), 256–264, doi:10.1016/j.quascirev.2009.05.010, 2010.

Durand, G., Gillet-Chaulet, F., Svensson, A., Gagliardini, O., Kipfstuhl, S., Meyssonier, J., Parrenin, F., Duval, P., and Dahl-Jensen, D.: Change in ice rheology during climate variations - implications for ice flow modelling and dating of the EPICA Dome C core, *Clim. Past*, 3, 155–167, doi:10.5194/cp-3-155-2007, 2007.

Durand, G., Gagliardini, O., Thorsteinsson, T., Svensson, A., Kipfstuhl, S., and Dahl-Jensen, D.: Ice microstructure and fabric: an up-to-date approach for measuring textures, *J. Glaciol.*, 52, 619–630, doi:10.3189/172756506781828377, 2006.

EPICA Community Members: One-to-one coupling of glacial climate variability in Greenland and Antarctica, *Nature*, 444(7116), 195–198, doi:10.1038/nature05301, 2006.

EPICA Community Members, F.: Eight glacial cycles from an Antarctic ice core, *Nature*, 429(6992), 623–628, doi:10.1038/nature02599, 2004.

Epstein, S., Buchsbaum, R., Lowenstam, L., and Urey, H.: Carbonate-water isotopic temperature scale, *Geol. Soc. Am. Bull.*, 62, 417–425, 1953.

Finkel, R. C. and Nishiizumi, K.: Beryllium 10 concentrations in the Greenland Ice Sheet Project 2 ice core from 3–40 ka, *J. Geophys. Res.*, 102(C12), 26,699–26,706, doi:199710.1029/97JC01282, 1997.

Frezzotti, M., Bitelli, G., de Michelis, P., Deponti, A., Forieri, A., Gandolfi, S., Maggi, V., Mancini, F., Remy, F., Tabacco, I. E., et al.: Geophysical survey at Talos Dome, East Antarctica: the search for a new deep-drilling site, *Ann. Glaciol.*, 39(1), 423–432, 2004.

Frezzotti, M., Urbini, S., Proposito, M., Scarchilli, C., Gandolfi, S. and others: Spatial and temporal variability of surface mass balance near Talos Dome, East Antarctica, *J. Geophys. Res.*, 112, 1–5, 2007.

Gagliardini, O. and Meyssonier, J.: Simulation of anisotropic ice flow and fabric evolution along the GRIP-GISP2 flowline, central Greenland, *Ann. Glaciol.*, 30, 217–223, 2000.

Gillet-Chaulet, F., Gagliardini, O., Meyssonier, J., Zwinger, T., and Ruokolainen, J.: Flow-induced anisotropy in polar ice and related ice-sheet flow modelling\*, *J. Non.-Newton Fluid.*, 134(1–3), 33–43, doi:10.1016/j.jnnfm.2005.11.005, 2006.

Goujon, C., Barnola, J., and Ritz, C.: Modeling the densification of polar firn including heat diffusion: Application to close-off characteristics and gas isotopic fractionation for Antarctica and Greenland sites, *J. Geophys. Res.*, 108, 18 pp., doi:200310.1029/2002JD003319, 2003.

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Huber, C., Leuenberger, M., Spahni, R., Fluckiger, J., Schwander, J., Stocker, T., Johnsen, S., Landais, A., and Jouzel, J.: Isotope calibrated Greenland temperature record over Marine Isotope Stage 3 and its relation to CH<sub>4</sub>, *Earth Planet. Sc. Lett.*, 243(3–4), 504–519, doi:10.1016/j.epsl.2006.01.002, 2006.
- 5 IPCC: “Climate Change 2001 – The Scientific Basis”, Contribution of Working Group I to the 3rd assessment of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, 881 pp., 2001.
- Jouzel, J., Lorius, C., Petit, J. R., Genthon, C., Barkov, N. I., Kotlyakov, V. M., and Petrov, V. M.: Vostok ice core: a continuous isotope temperature record over the last climatic cycle (160,000 years), *Nature*, 329(6138), 403–408, doi:10.1038/329403a0, 1987.
- 10 Kawamura, K., Parrenin, F., Lisiecki, L., Uemura, R., Vimeux, F., Severinghaus, J. P., Hutterli, M. A., Nakazawa, T., Aoki, S., Jouzel, J., Raymo, M. E., et al.: Northern Hemisphere forcing of climatic cycles in Antarctica over the past 360,000 years, *Nature*, 448(7156), 912–916, doi:10.1038/nature06015, 2007.
- 15 Landais, A., Barnola, J. M., Kawamura, K., Caillon, N., Delmotte, M., Van Ommen, T., Dreyfus, G., Jouzel, J., Masson-Delmotte, V., Minster, B. and others: Firn-air delta<sup>15</sup>N in modern polar sites and glacial-interglacial ice: a model-data mismatch during glacial periods in Antarctica?, *Quaternary Sci. Rev.*, 25(1–2), 49–62, 2006.
- Landais, A., Chappellaz, J., Delmotte, M., Jouzel, J., Blunier, T., Bourq, C., Caillon, N., Cherrier, S., Malaizé, B., Masson-Delmotte, V., et al.: A tentative reconstruction of the last interglacial and glacial inception in Greenland based on new gas measurements in the Greenland Ice Core Project (GRIP) ice core, *J. Geophys. Res.*, 108, 4563–4572, 2003.
- Lemieux-Dudon, B., Parrenin, F., and Blayo, E.: A probabilistic method to construct a common and optimal chronology for an ice core, *Physics of Ice Core Records*, 2009.
- 25 Lemieux-Dudon, B., Blayo, E., Petit, J., Waelbroeck, C., Svensson, A., Ritz, C., Barnola, J., Narcisi, B. M., and Parrenin, F.: Consistent dating for Antarctic and Greenland ice cores, *Quaternary Sci. Rev.*, 29(1–2), 8–20, doi:10.1016/j.quascirev.2009.11.010, 2010.
- Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic delta O-18 records, *Paleoceanography*, 20(2), PA2007, doi:10.1029/2005PA001164, 2005.
- 30 Loulergue, L., Parrenin, F., Blunier, T., Barnola, J.-M., Spahni, R., Schilt, A., Raisbeck, G., and Chappellaz, J.: New constraints on the gas age-ice age difference along the EPICA ice cores, 0-50 kyr, *Clim. Past*, 3, 527–540, doi:10.5194/cp-3-527-2007, 2007.
- Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola,

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



J., Raynaud, D., Stocker, T. F., and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH<sub>4</sub> over the past 800,000 years, *Nature*, 453(7193), 383–386, doi:10.1038/nature06950, 2008.

Mackintosh, A., White, D., Fink, D., Gore, D. B., Pickard, J., and Fanning, P. C.: Exposure ages from mountain dipsticks in Mac. Robertson Land, East Antarctica, indicate little change in ice-sheet thickness since the Last Glacial Maximum, *Geology*, 35(6), 551–554, 2007.

Magand, O., Frezzotti, M., Pourchet, M., Stenni, B., Genoni, L., and Fily, M.: Climate variability along latitudinal and longitudinal transects in East Antarctica, *Ann. Glaciol.*, 39(1), 351–358, 2004.

Mangeney, A., Califano, F., and Hutter, K.: A numerical study of anisotropic, low Reynolds number, free surface flow for ice sheet modeling, *J. Geophys. Res.*, 102(B10), 22,749–22,764, doi:199710.1029/97JB01697, 1997.

Masson-Delmotte, V., Dreyfus, G., Braconnot, P., Johnsen, S., Jouzel, J., Kageyama, M., Landais, A., Loutre, M.-F., Nouet, J., Parrenin, F., Raynaud, D., Stenni, B., and Tuerter, E.: Past temperature reconstructions from deep ice cores: relevance for future climate change, *Clim. Past*, 2, 145–165, doi:10.5194/cp-2-145-2006, 2006.

Meneghel, M., Bondesan, A., Salvatore, M. C., and Orombelli, G.: A model of the glacial retreat of upper Rennick Glacier, Victoria Land, Antarctica, *Ann. Glaciol.*, 29, 225–230, 1999.

Merlivat, L.: Molecular diffusivities of H<sub>2</sub> 160, HD160, and H<sub>2</sub> 180 in gases, *J. Chem. Phys.*, 69(6), 2864, doi:10.1063/1.436884, 1978.

Montagnat: Texture evolution along Talos Dome ice core, Antarctica., in preparation, 2010.

Morgan, V., Delmotte, M., Van Ommen, T., and Jouzel, J.: Relative Timing of Deglacial Climate Events in Antarctica and Greenland, *Science*, 297(5588), 1862–1864, doi:10.1126/science.1074257, 2002.

Morse, D. L., Waddington, E. D., and Rasmussen, L. A.: Ice deformation in the vicinity of the ice-core site at Taylor Dome, Antarctica, and a derived accumulation rate history, *J. Glaciol.*, 53(182), 449–460, 2007.

Morse, D. L., Waddington, E. D., and Steig, E. J.: Ice Age storm trajectories inferred from radar stratigraphy at Taylor Dome, Antarctica, *Geophys. Res. Lett.*, 25(17), 3383–3386, 1998.

Parrenin, F., Barnola, J.-M., Beer, J., Blunier, T., Castellano, E., Chappellaz, J., Dreyfus, G., Fischer, H., Fujita, S., Jouzel, J., Kawamura, K., Lemieux-Dudon, B., Loulergue, L., Masson-Delmotte, V., Narcisi, B., Petit, J.-R., Raisbeck, G., Raynaud, D., Ruth, U., Schwander, J., Severi, M., Spahni, R., Steffensen, J. P., Svensson, A., Udisti, R., Waelbroeck, C., and

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wolff, E.: The EDC3 chronology for the EPICA Dome C ice core, *Clim. Past*, 3, 485–497, doi:10.5194/cp-3-485-2007, 2007.

Parrenin, F., Dreyfus, G., Durand, G., Fujita, S., Gagliardini, O., Gillet, F., Jouzel, J., Kawamura, K., Lhomme, N., Masson-Delmotte, V., Ritz, C., Schwander, J., Shoji, H., Uemura, R., Watanabe, O., and Yoshida, N.: 1-D-ice flow modelling at EPICA Dome C and Dome Fuji, East Antarctica, *Clim. Past*, 3, 243–259, doi:10.5194/cp-3-243-2007, 2007.

Parrenin, F., Remy, F., Ritz, C., Siegert, M. J., and Jouzel, J.: New modeling of the Vostok ice flow line and implication for the glaciological chronology of the Vostok ice core, *J. Geophys. Res.*, 109, 1–14, 2004.

Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J., Basile, I., Bender, M., Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., et al.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399(6735), 429–436, doi:10.1038/20859, 1999.

Raisbeck, G. M., Yiou, F., Jouzel, J., Petit, J. R., Barkov, M. I., and Bard, E.: 10 Be deposition at Vostok, Antarctica, during the last 50,000 years and its relationship to possible cosmogenic production variations during this period, Bard E & Broecker WS eds: *The Last Deglaciation: Absolute and Radiocarbon Chronologies*, NATO ASI series, I, 2, 127–140, 1992.

Raisbeck, G. M. and Yiou, F.: 10 Be in polar ice and atmospheres, *Ann. Glaciol.*, 7, 138–140, 1985.

Ritz, C., Rommelaere, V., and Dumas, C.: Modeling the evolution of Antarctic ice sheet over the last 420,000 years: Implications for altitude changes in the Vostok region, *J. Geophys. Res.*, 106(D23), 31,943–31,964, doi:200110.1029/2001JD900232, 2001.

Ruth, U., Barnola, J.-M., Beer, J., Bigler, M., Blunier, T., Castellano, E., Fischer, H., Fundel, F., Huybrechts, P., Kaufmann, P., Kipfstuhl, S., Lambrecht, A., Morganti, A., Oerter, H., Parrenin, F., Rybak, O., Severi, M., Udisti, R., Wilhelms, F., and Wolff, E.: “EDML1”: a chronology for the EPICA deep ice core from Dronning Maud Land, Antarctica, over the last 150 000 years, *Clim. Past*, 3, 475–484, doi:10.5194/cp-3-475-2007, 2007.

Scarchilli, C., Frezzotti, M., and Ruti, P.: Snow precipitation at the Talos Dome core site in East Antarctica: Provenance, seasonality and blocking factors, *Clim. Dynam.*, in review, 2010.

Schüpbach, S., Federer, U., Kaufmann, P. R., Hutterli, M. A., Buiron, D., Blunier, T., Fischer, H., and Stocker, T. F.: A New Method for High-Resolution Methane Measurements on Polar Ice Cores Using Continuous Flow Analysis, *Environ. Sci. Technol.*, 43(14), 5371–5376, doi:10.1021/es9003137, 2009.

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Schwander, J., Sowers, T., Barnola, J., Blunier, T., Fuchs, A., and Malaizé, B.: Age scale of the air in the summit ice: Implication for glacial-interglacial temperature change, *J. Geophys. Res.*, 102(D16), 19,483–19,493, doi:199710.1029/97JD01309, 1997.

Severinghaus, J. P., Sowers, T., Brook, E. J., Alley, R. B., and Bender, M. L.: Timing of abrupt climate change at the end of the Younger Dryas interval from thermally fractionated gases in polar ice, *Nature*, 391(6663), 141–146, 1998.

Sowers, T. and Bender, M.: Climate Records Covering the Last Deglaciation, *Science*, 269(5221), 210–214, doi:10.1126/science.269.5221.210, 1995.

Sowers, T., Bender, M., Raynaud, D., Korotkevich, Y. S., and Orchardo, J.: The  $\delta^{18}\text{O}$  of atmospheric  $\text{O}_2$  from air inclusions in the Vostok Ice Core: Timing of  $\text{CO}_2$  and ice volume changes during the penultimate deglaciation, *Paleoceanography*, 6(6), 679, doi:10.1029/91PA02023, 1991.

Spahni, R., Chappellaz, C., Stocker, T., Loulergue, L., Hausmann, G., Kawamura, K., Flückiger, J., Schwander, J., Raunaud, D., Masson-Delmotte, V., and Jouzel, J.: Atmospheric Methane and Nitrous Oxide of the Late Pleistocene from Antarctic Ice Cores, *Science*, 310(5752), 1317–1321, doi:10.1126/science.1120132, 2005.

Spahni, R., Schwander, J., Flückiger, J., Stauffer, B., Chappellaz, J., and Raynaud, D.: The attenuation of fast atmospheric  $\text{CH}_4$  variations recorded in polar ice cores, *Geophys. Res. Lett.*, 30(11), 1571–1574, 2003.

Steffensen, J. P., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Fischer, H., Goto-Azuma, K., Hansson, M., Johnsen, S. J., Jouzel, J., Masson-Delmotte, V., et al.: High-Resolution Greenland Ice Core Data Show Abrupt Climate Change Happens in Few Years, *Science*, 321(5889), 680–684, doi:10.1126/science.1157707, 2008.

Steig, E. J., Brook, E., White, J., and Sucher, C.: Synchronous Climate Changes in Antarctica and the North Atlantic, *Science*, 282(5386), 92–95, doi:10.1126/science.282.5386.92, 1998.

Stenni, B., Buiron, D., Frezzotti, M., and The Taldice Consortium: Unified Antarctic and Greenland climate seesaw during the last deglaciation, *Nature Geosciences*, submitted, 2010.

Stenni, B., Masson-Delmotte, V., Johnsen, S., Jouzel, J., Longinelli, A., Monnin, E., Rothlisberger, R., and Selmo, E.: An oceanic cold reversal during the last deglaciation, *Science*, 293(5537), 2074–2077, 2001.

Stenni, B., Proposito, M., Gragnani, R., Flora, O., Jouzel, J., Falourd, S. and Frezzotti, M.: Eight centuries of volcanic signal and climate change at Talos Dome (East Antarctica), *J. Geophys. Res.*, 107(D9), 4076, doi:10.1029/2000JD000317, 2002.

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Stuiver, M., Grootes, P. M., and Braziunas, T. F.: The GISP2 [ $\delta$ ]18O Climate Record of the Past 16,500 Years and the Role of the Sun, Ocean, and Volcanoes, *Quaternary Res.*, 44(3), 341–354, doi:10.1006/qres.1995.1079, 1995.

5 Svensson, A., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Davies, S. M., Johnsen, S. J., Muscheler, R., Parrenin, F., Rasmussen, S. O., Rthlisberger, R., Seierstad, I., Steffensen, J. P., and Vinther, B. M.: A 60 000 year Greenland stratigraphic ice core chronology, *Clim. Past*, 4, 47–57, doi:10.5194/cp-4-47-2008, 2008.

10 Urbini, S., Frezzotti, M., Gandolfi, S., Vincent, C., Scarchilli, C., Vittuari, L., and Fily, M.: Historical behaviour of Dome C and Talos Dome (East Antarctica) as investigated by snow accumulation and ice velocity measurements, *Global Planet. Change*, 60(3–4), 576–588, 2008.

Van der Veen, C. and Whillans, I.: Development of fabric in ice, *Cold Cold Reg. Sci. Technol.*, 22(2), 171–195, doi:10.1016/0165-232X(94)90027-2, 1994.

15 Watanabe, O., Jouzel, J., Johnsen, S., Parrenin, F., Shoji, H., and Yoshida, N.: Homogeneous climate variability across East Antarctica over the past three glacial cycles, *Nature*, 422(6931), 509–512, doi:10.1038/nature01525, 2003.



**Table 1.** Tie points defined by CH<sub>4</sub> record synchronization between TALDICE and Greenland ice cores on GICC05 age scale.

Greenland reference ice core	NorthGRIP depth (m)	TALDICE depth (m)	Gas age (yr BP)	Uncertainty (yr)
GRIP	846.6	420	5200	200
GRIP	1219.2	553	8250	150
GRIP	1470.3	684	11300	200
GRIP	1489.9	695	11 650	100
GRIP	1525.2	738	12 850	100
GRIP	1604.2	784	14 680	100
GRIP	1647.6	820	16 550	500
GRIP	1672.8	844	176 50	300
GRIP	1861.3	935	27 550	300
GRIP	1891.5	953	28 900	200
GRIP	1941.0	985	32 100	200
GRIP	1962.6	994	33 250	200
GRIP	1976.4	1006	33 850	200
GRIP	1991.8	1015	34 800	200
GRIP	2008.6	1025	35 450	200
GRIP	2029.0	1040	36 700	200
NGRIP	2072.2	1062	38 350	200
GRIP	2076.7	1070	38 650	200
NGRIP	2093.2	1082	39 800	200
NGRIP	2110.5	1095	40 850	200
NGRIP	2126.1	1099	41 600	200
NGRIP	2132.5	1111	42 100	200
NGRIP	2155.5	1123	43 300	200
NGRIP	2169.8	1135	44 400	200
GISP	2224.0	1170	47 200	200
GRIP	2252.4	1191	49 270	400

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 2.** Tie points defined by CH<sub>4</sub> record synchronization between TALDICE and EDC ice core on the EDC3 age scale.

EDC depth (m)	TALDICE depth (m)	Gas age – (yr BP)	Uncertainty (yr)
924.9	1228	53 500	200
944.5	1239	55 150	200
998	1264	59 800	1000
1089.1	1302	69 600	400
1106.6	1308	71 200	1000
1145.3	1316	75 000	1000
1197.8	1326	80 000	800
1239.0	1333	83 400	800
1302.7	1345	89 500	500
1369.3	1356	96 000	500
1439.6	1370	103 000	800
1457.2	1373	105 000	1500
1515.4	1380	112 000	1000
1544.8	1385	115 200	1500
1622.0	1394	121 700	2000
1717.3	1410	128 500	1000
1763.6	1420	132 000	2000
1817.4	1428	141 000	2000

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 3.** Tie points defined by water isotopic record synchronization between TALDICE and EDC ice core based on the EDC3 age scale.

EDC depth (m)	TALDICE depth (m)	Associated age (yr BP)	Uncertainty (yr)
1888.7	1443	159 200	2000
1903.1	1462	162 000	2000
1938.0	1471	169 000	2000
1953.0	1480	172 000	2000
1979.0	1484	177 500	2000
2022.2	1493	187 300	2000
2031.2	1495	189 000	2000
2072.0	1500	197 500	2000
2105.5	1512	203 000	3000
2123.8	1516	206 500	3000
2197.1	1526	222 000	3000
2299.8	1545	243 200	4000
2318.1	1548	247 000	4000
2410.4	1558	290 000	5000

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 4.** Input parameters used in the 1-D ice flow model (Parrenin et al., 2007b).

Parameter	Meaning	Value	References and comments
$\beta$	Glacial-interglacial amplitude of accumulation rate changes	0,0165	Tuned for best fit of age markers
$\alpha$	spatial slope of the present-day isotopic thermometer	5,04‰/°C	(Frezzotti et al., 2004; Magand et al., 2004)
$A_0$	Present-day accumulation rate	8 cm ie/yr	Tuned for best fit of age markers; 10% lower than (Stenni et al., 2002)
$T_0$	Present-day surface temperature	−41 °C	(Frezzotti et al., 2004)
$p$	parameter controlling the (non)-linearity of the vertical velocity profile of ice	3	(Parrenin et al., 2007b)
$\Delta H$	Past elevation changes	Table 4	See Sect. 3.2

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Table 5.** Estimated elevation changes (positive numbers indicate an increase) in the TALDICE area between 0 and 30 kyr BP. Error bars are estimated from bibliographic descriptions (Stenni et al., 2010).

Age (kyr BP)	Elevation change (m)
0	0
7	0±20
10	100±50
12.7	170±50
16	170±20
19	100±50
25	20±50
30	0±50

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.



**Fig. 1. (A):** synchronization of the TALDICE CH<sub>4</sub> record (purple curve) with the Greenland CH<sub>4</sub> composite on GICC05 age scale (Blunier et al., 2007) (blue curve) back to 53 kyr BP, and with the EDC CH<sub>4</sub> record (grey curve) on EDC3 age scale (Louergue et al., 2008) for ages older than 53 kyr BP. **(B):** synchronization between TALDICE (Stenni et al., 2010) (purple curve) and EDC (grey curve) ice cores on the EDC3 age scale (EPICA Community Members, 2004) using δ<sup>18</sup>O of ice. For both graphs, dashed lines indicate the location of each tie point.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version


Interactive Discussion

**TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica**

D. Buiron et al.



**Fig. 2.** TALDICE-1 age/depth relationship for the ice (blue lines) and for the gas (black lines), from the a priori glaciological model (dashed lines), and after applying the inverse method (continuous lines), with a zoom on the period between 8 and 50 kyr BP; Bottom: uncertainty associated with the chronology deduced from the inverse method and zoomed between 8 and 50 kyr BP.



**Fig. 3.** Ratio between a posteriori and a priori values of glaciological entities plotted against the TALDICE-1 age scale: from top to bottom: CODIE (black line with symbols) accumulation rate (grey line), and thinning function (black line).

**TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica**

D. Buiron et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





**TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica**

D. Buiron et al.




**Fig. 4.**  $\Delta$ age evolution on the TALDICE-1 age scale, calculated by the inverse model (light blue line) and by the firn densification model (dark blue line). The grey shade represents the error interval based on accumulation rate uncertainties of  $\pm 20\%$ . The maximum difference recorded between  $\Delta$ age estimates from the inverse method and from the firn densification model is pointed by the arrow: it occurs around 40 kyr BP and reaches 400 yr. The difference remains below  $\pm 200$  yr for other times.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**TALDICE-1 age scale  
of the Talos Dome  
deep ice core, East  
Antarctica**

D. Buiron et al.



**Fig. 5.**  $\delta^{18}\text{O}_{\text{atm}}$  measured along the TALDICE ice core (black line and symbols) on the TALDICE-1 age scale, compared with its counterparts from EDML (blue line with rounded symbols; Capron et al., 2010) and Vostok (purple line; Petit et al., 1999). Inside frame: zoom on the LGM and the last deglaciation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.



**Fig. 6.** Accumulation rate history deduced from the water isotopic profile (light blue line) and after adjustment by the inverse method (dark blue line). The grey area represents the error interval for an accumulation rate varying by  $\pm 20\%$  around the value deduced from the inverse method. Inside frame: zoom over the last deglaciation period. We add the  $^{10}\text{Be}$  constraints for Holocene and LGM periods (stars), as well as the accumulation rate calculated by the traditional isotopic equation for a value of  $\beta=0.0198$  (purple line).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica

D. Buiron et al.



**Fig. 7.** Comparison of thinning functions deduced from the a priori ice-flow model (dashed grey line), by the ice-flow model taking into account altitudinal changes (grey line) and by the inverse model (black line). They are shown together with the eigenvalues  $a_1$  of the second order orientation tensor measured along the TALDICE ice core (blue diamonds). Grey shades underline changes of slope in the fabric evolution and the blue shade highlights a depth range where reduced crystal sizes have been measured (not shown).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion