

***Interactive comment on* “Surface circulation patterns and the pathways of sea surface carbon dioxide (CO₂) off northern Chile (~27.5° S) between 30 and 10 kyr BP: global and/or local forcing?” by J. A. Placencia et al.**

Anonymous Referee #1

Received and published: 31 March 2010

This manuscript suffers from at least two major flaws and therefore should not be published.

The authors attempt to reconstruct the partial pressure of carbon dioxide (pCO₂) in surface waters near Chile by comparing the carbon isotopic composition (d¹³C) of alkenones against the d¹³C of planktonic foraminifera. In doing so, it is assumed that the d¹³C of the foraminifera records the d¹³C of dissolved inorganic carbon without any fractionation. It is further assumed that the d¹³C of alkenones is offset from the d¹³C of DIC by an amount that is related in a simple linear way to the pCO₂ of surface

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



water (actually, to the aqueous CO₂ concentration, but I use pCO₂ here because the authors convert aqueous CO₂ to pCO₂ in presenting their results). Hence, the authors imply that pCO₂ can be calculated by comparing d¹³C of alkenones against d¹³C of foraminifera.

The first failure of this approach is that the d¹³C of alkenones is not offset from the d¹³C of DIC by a simple linear relationship that is a function only of pCO₂. As has been pointed out by Bidigare et al. (1997) and by Popp et al. (1998), both cited in this paper, the carbon isotope fractionation between haptophytes (producers of alkenones) and DIC depends also on cell size and on cell growth rate. Growth rate, in turn, depends on many factors, including light, macronutrients (e.g., phosphate) and micronutrients (e.g., iron). The authors note that iron supply varied over the course of their record, and they also note that this may have influenced the growth rates of organisms. If so, then this would have also impacted the fractionation of carbon isotopes between alkenones and DIC. Without a mechanism to correct for changes in growth rate, it is impossible to derive meaningful estimates of pCO₂.

Second, the d¹³C of planktonic foraminifera is not representative of the d¹³C of DIC, nor is the d¹³C of planktonic foraminifera related to the d¹³C of DIC by a constant factor. In particular, the carbon isotope fractionation between foraminifera shells and DIC is strongly dependent of the concentration of carbonate ion. This sensitivity to carbonate ion concentration is described in many papers by Howie Spero, Jelle Bijma, and their coworkers (e.g., Spero et al., *Nature*, 390 (1997) 497-500; Russell and Spero, *Paleoceanography*, 15 (2000) 43-52). The carbonate ion of the surface ocean worldwide changed appreciably over the course of deglaciation associated with the rise in atmospheric CO₂. More important here, changes in upwelling and in the biological consumption of CO₂ in the region offshore of Chile would have imposed changes in the ambient carbonate ion concentration even larger than those experienced in less dynamic regimes. It is not possible to constrain the carbonate ion concentration from the available data. Consequently, it is not possible to constrain the changes through

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

time in the carbonate ion effect on fractionation of $d^{13}C$ between foraminifera and DIC. For these reasons, meaningful reconstruction of surface ocean pCO_2 by the methods employed here are not possible. Therefore, this paper should not be published.

Interactive comment on Clim. Past Discuss., 6, 347, 2010.

CPD

6, C59–C61, 2010

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C61

