

This discussion paper is/has been under review for the journal *Climate of the Past* (CP).  
Please refer to the corresponding final paper in CP if available.

# A multi-proxy perspective on millennium-long climate variability in the Southern Pyrenees

**M. Morellón**<sup>1,2,3</sup>, **A. Pérez-Sanz**<sup>3</sup>, **J. P. Corella**<sup>2,3</sup>, **U. Büntgen**<sup>4</sup>, **J. Catalán**<sup>5</sup>,  
**P. González-Sampériz**<sup>3</sup>, **J. J. González-Trueba**<sup>6</sup>, **J. A. López-Sáez**<sup>7</sup>, **A. Moreno**<sup>2,3</sup>,  
**S. Pla**<sup>5</sup>, **M. Á. Saz-Sánchez**<sup>8</sup>, **P. Scussolini**<sup>9,12</sup>, **E. Serrano**<sup>10</sup>, **F. Steinhilber**<sup>1</sup>,  
**V. Stefanova**<sup>11</sup>, **T. Vegas-Vilarrúbia**<sup>12</sup>, and **B. Valero-Garcés**<sup>2,3</sup>

<sup>1</sup>Department of Surface Waters, Eawag, Überlandstrasse 133, 8600 Dübendorf, Switzerland

<sup>2</sup>Laboratorio Internacional de Cambio Global (LINCGlobal), PUC-CSIC, Departamento de Ecología, Facultad de Ciencias Biológicas, PUC, Alameda 340, PC 6513677, Santiago, Chile

<sup>3</sup>Departamento de Procesos Geoambientales y Cambio Global, Instituto Pirenaico de Ecología (IPE) – CSIC, Campus de Aula Dei, Avda. Montañana 1005, 50059 Zaragoza, Spain

<sup>4</sup>Swiss Federal Research Institute for Forest, Snow and Landscape (WSL), ETH domain, Zürich, 8903 Birmensdorf, Switzerland

<sup>5</sup>Centre d'Estudis Avançats de Blanes (CEAB) – CSIC, C/Accés a la Cala St. Francesc, 14, Blanes 17300, Spain

<sup>6</sup>Departamento de Geografía, Prehistoria y Arqueología, Calle Tomás y Valiente s/n, 01006 Vitoria-Gasteiz, Spain

**A multi-proxy  
perspective on  
millennium-long  
climate variability**

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**A multi-proxy perspective on millennium-long climate variability**

M. Morellón et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

<sup>7</sup> Centro de Ciencias Humanas y Sociales, CCHS-CSIC, C/Albasanz, 26–28, 28037 Madrid, Spain

<sup>8</sup> Departamento de Geografía y Ordenación del Territorio, Facultad de Filosofía y Letras, Universidad de Zaragoza, Calle Pedro Cerbuna 12, 50009 Zaragoza, Spain

<sup>9</sup> Faculty of Earth and Life Sciences, Department of Marine Biogeology, Vrije Universiteit, De Boelelaan 1085, 1081HV Amsterdam, The Netherlands

<sup>10</sup> Subunidad de Geografía Física Departamento de Geografía, Facultad de Filosofía y Letras, Plaza del Campus s/n, 47011 Valladolid, Spain

<sup>11</sup> Limnological Research Center, University of Minnesota, 310 Pillsbury Drive SE, Minneapolis, MN 55455, USA

<sup>12</sup> Department of Ecology, Faculty of Biology, University of Barcelona, Av. Diagonal 645, 08028 Barcelona, Spain

Received: 23 September 2011 – Accepted: 24 September 2011 – Published: 4 October 2011

Correspondence to: M. Morellón (mario.morellon-marteles@eawag.ch)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Abstract

A review of selected, well-dated, multiproxy paleoclimatic records (lacustrine, dendrochronological, geomorphological) characterizes the main environmental changes occurred in the Southern Pyrenees during the last millennium. Warmer and relatively arid conditions prevailed during the Medieval Climate Anomaly (MCA, <1300 AD), with a significant development of xerophytes and Mediterranean vegetation and limited deciduous tree formations (mesophytes). The Little Ice Age (LIA, 1300–1800 AD) was generally colder and moister, with an expansion of deciduous taxa and cold-adapted mountainous conifers. Two major phases occurred within this period: (i) a transition MCA-LIA, characterized by fluctuating, moist conditions and relatively cold temperatures (ca. 1300 and 1600 AD); – (ii) a second period, characterized by coldest conditions and higher humidity, coinciding with maximum (recent) glacier advances (ca. 1600–1850 AD). After the LIA a warming and more arid phase started coinciding with glacier retreat, and interrupted by a short-living cooling episode during the late 19th to early 20th centuries. Some records suggest a response to solar activity with colder and slightly moister conditions during solar minima. Centennial-scale hydrological fluctuations are in phase with reconstructions of NAO variability, which appears to be the main forcing for humidity in the region during the last millennium.

## 1 Introduction

Climate reconstructions for the last millennium are crucial for our understanding of past natural variability and for assessing the influence of external forcing factors such as solar activity and volcanic eruptions. The perspective of a current global warming scenario has provoked an increasing interest in studies of climate fluctuations prior to the industrial era (Jansen et al., 2007). Furthermore, long-term palaeoclimatic perspectives offer an excellent basis to improve predictive models (Solomon et al., 2007). Past climate conditions have been reconstructed using various proxy data, such as

CPD

7, 3049–3089, 2011

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tree rings (Büntgen et al., 2011), lake sediments (Last and Smol, 2001), historical documents (Vicente-Serrano and Cuadrat, 2007), marine sediments (Abrantes et al., 2005), corals (Goodkin et al., 2008), speleothems (Proctor et al., 2002), and ideally a combination thereof (Jones et al., 2009).

5 In spite of an ample array of associated uncertainties (Frank et al., 2010), most climate reconstructions agree on the existence of two different stages with distinct climatic conditions during the last millennium: a period of relatively high temperatures from ~ 800–1300 AD, known as the Medieval Climate Anomaly (MCA); and a phase with relatively cooler temperatures from ~ 1350–1850 AD, the so-called Little Ice Age  
10 (LIA). Unfavourable climate conditions during the LIA had also a considerable impact in European human societies (Fagan, 2000), as they likely contributed to the decline of health and economic wealth of medieval societies triggered by the Black Death (Büntgen et al., 2011), for instance, and led to the abandonment of different areas such as Norse settlements in Greenland (Patterson et al., 2010; D’Andrea et al., 2011).  
15 Despite the precise chronology, the spatial extent and mechanisms behind these climatic phases still remain a subject of debate (Paasche and Bakke, 2010).

The impact of the LIA on Central European ecosystems and societies has been widely documented (Büntgen et al., 2011; Magny et al., 2011b, Denton and Broecker, 2008). However, its environmental response in the Mediterranean region is not yet well understood (Touchan et al., 2011; Esper et al., 2007; Büntgen et al., 2008, 2010b; Luterbacher et al., 2005). Several studies have suggested that the hydrological impact of climate changes will affect dramatically snow-fed watersheds, such as the headwaters of many rivers located in the Pyrenees (Beniston, 2003; García-Ruiz et al., 2011). The Southern Pyrenees constitute the main source of water resources nowadays, supplying more than 45 % of the annual runoff to the surrounding semi-arid Ebro Basin, a relatively densely populated area of NE Spain (López-Moreno and García-Ruiz, 2004). To understand future climate changes induced by global warming is necessary to assess the impact of climate change on the hydrological cycle and the range of climate variability and threshold values during the last millennia.

**A multi-proxy perspective on millennium-long climate variability**

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





as towards the Aquitania (France) and Ebro Depressions (Spain) (north and south, respectively) (Fig. 1b).

The regional climate is subject to Atlantic and Mediterranean influences and the effect of macro-relief on precipitation and temperature, which leads to a high climatic heterogeneity over short distances (López-Moreno et al., 2008). Annual precipitation exceeds 600 mm and sometimes even reaches 2000 mm at the highest altitudes. Most of the annual precipitation occurs during the cold season in the Atlantic areas but during spring and autumn in the Mediterranean regions, with relatively dry summers. The annual averaged 0°C isotherm is located at ~ 2700 m a.s.l. (Del Barrio et al., 1990). In higher altitudes vegetation is mainly composed of subalpine and alpine pastures (*Carex curvula* and *Festuca scoparia*) with conifers (*Pinus uncinata* and *Juniperus* sp.) like the only arboreal component. Small forested areas of *Fagus sylvatica*, *Fraxinus excelsior*, *Corylus avellana*, *Betula pubescens*, *Acer campestre*, *Sorbus aucuparia* and *Sorbus aria* can be found above 1800 m a.s.l. and mixed forest with *Pinus sylvestris*, *Fagus sylvatica* and *Abies alba* are also common (Peinado Lorca and Rivas-Martínez, 1987). Between 1800 and 1500 m, *Pinus sylvestris* and *Quercus faginea* forests are dominant, while at around 1000 m Mediterranean influence increases and the vegetation landscape is composed by *Quercus rotundifolia* and *Quercus pubescens*, depending on the annual precipitation. Foothills are mainly covered by Mediterranean pine and oak forests (*Quercus ilex-rotundifolia*, *Q. faginea*, *Q. pubescens*), with dense shrub land (*Rhamnus*, *Pistacia*, *Arbutus*, *Prunus*, *Cornus*, *Buxus*, *Genisteae*, *Cistaceae*, etc.). *Corylus avellana* and other phreatophytes such as *Salix*, *Fraxinus*, *Populus* or *Ulmus* are only located in the riverine soils (Blanco-Castro et al., 1997). There are two main types of lakes in the Pyrenean domain (Fig. 1a): (i) karstic lakes, located at the Southern foothills of the Pyrenees, in the so-called External Ranges, between 600 and 1000 m a.s.l., in the transition between Atlantic and Mediterranean climate types and vegetation domains (García and Gómez, 2008) and subjected to a considerable human influence during the last millennia (Arreo, Estanya, Montcortès); (ii) high-mountain, glacial lakes located in the Axial Pyrenees, above 1800 m a.s.l., characterized by colder

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and more humid conditions, near to the timberline (e.g., Basa de la Mora, Redon) where human impact is limited. Currently in the Pyrenees there are 21 glaciers with a total extension of about 495 ha. On the Spanish side there are 29 ice bodies, from which only 10 can be considered as glaciers (260 ha) (González Trueba et al., 2008), and thus, appear the most relevant for this study. Nevertheless, LIA remains such as moraines and proglacial materials do exist in more than 100 deglaciated cirques in the Pyrenean high mountains. These deposits are located in the current glaciated and non-glaciated massifs (Fig. 1b) (Serrano and Martínez de Pisón, 1994; Grove and Gelatly, 1995; Chueca et al., 1998a,b). The Pyrenean glaciers are reduced in thickness and sheltered by the highest summits of the range. They underwent the last LIA glacier advance, although they display current morphological features that imply a significant loss in relation to the external moraines, bevelled shapes, burials, and even some disappearance of ice bodies. From 1880 to 1980 AD at least 94 glaciers became extinct, while 17 glaciers have disappeared on the Spanish side since the 1980s.

### 3 Paleoclimate archives

For this study, we selected paleoclimate sequences from published studies that fulfill the following requirements: (i) reconstructions are based on natural archives, (ii) these records have independent chronologies based either on absolute dating (tree ring or lacustrine varves counting, calibrated radiometric dating) or documentary sources and (iii) climate variability is considered as the main forcing mechanism of environmental changes reconstructed at these sites. Paleoclimatic archives analyzed here comprise: (i) dendroclimatic reconstructions (Fig. 1a), (ii) lake and marine-littoral sequences (Fig. 1a), and (iii) geomorphologic evidences of glacier advances or retreats (Fig. 1b). Reconstructed environmental variables vary from climatic parameters (temperature, precipitation, altitude anomaly) to evidences of environmental changes directly related to these parameters (changes in vegetation cover, runoff and effective moisture fluctuations reconstructed by lake level changes and/or salinity fluctuations).

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Although anthropogenic activities often imply similar modifications in the landscape to those produced by climatic variability, comparing different type of data and employing a multidisciplinary methodology in palaeoenvironmental sequences, allows to discern between both climate and human forcings (Moreno et al., 2008; Morellón et al., 2011; Rull et al., 2011).

### 3.1 Lake and marine-littoral sequences

Unravelling the complex interplay between climate changes and human impact in Pyrenean lakes and littoral marine sequences is complex and needs the use of multiple proxy-data and robust chronologies based on either radiometric techniques ( $^{137}\text{Cs}$ ,  $^{210}\text{Pb}$ ,  $^{14}\text{C}$ ) or varve counting or a combination of these two methods. Furthermore, the correlation with historical events detected in most of the sequences provides further chronological support (e.g., onset and/or intensification of farming, introduction of particular cultivars, deforestation events, Rull et al., 2011). Most of the lacustrine sequences used in this review have been analyzed following a multidisciplinary strategy, comprising a variety of proxies: (i) sedimentology (sedimentary facies, microfacies and varve sub-layering), (ii) elemental and isotopic geochemistry, and (iii) biological indicators (pollen, diatoms, chironomids and chrysophytes). Other lake and peatbog records (i.e., Tramacastilla and Bubal, Montserrat-Martí, 1992; Bosc dels Estanyons, Miras et al., 2007; Riu dels Orris fen, Ejarque et al., 2010; and Lake Burg, Bal et al., 2011, among others), have not been considered for this paper because of the lack of a multiproxy analysis or their exclusive focus on human activities.

#### 3.1.1 Lake Arreo

Lake Arreo (42°46' N, 2°59' W, 655 m a.s.l.) is a karstic lake located in the Westernmost part of the foothills of the Pyrenees. The chronology of the sequence is based on the combination of varve counting,  $^{137}\text{Cs}/^{210}\text{Pb}$ , and  $^{14}\text{C}$  dating and spans the last 2600 yr (Corella Aznar, 2011). The main paleoindicators analyzed include sedimentary facies

CPD

7, 3049–3089, 2011

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and microfacies, elemental geochemistry, pollen and diatoms. The sensitivity of some of these proxies has been demonstrated by its successful correlation with instrumental climate data series and limnological monitoring data at shorter timescales (Corella et al., 2011).

The onset of the LIA in the Arreo sequence (1300 AD) is characterized by a better development and preservation of biogenic varves, decreased content in aragonite and gypsum, and corresponding lower values of strontium, and the increase in the centric diatom *Cyclotella distinguenda* and arboreal pollen. These proxies are indicative of the development of higher lake levels and associated meromixis, and a forest recovery likely caused by moister conditions (Table 1). Additionally, an increase in *Typha* indicates the flooding of the littoral platform previously emerged during the MCA. The decrease in deciduous *Quercus* since AD 1870 points towards more arid conditions since the end of the LIA. An arid phase was detected between 1730–1790 AD, characterized by a deposition of clay-rich facies (Fig. 2) and high strontium content in the sediment (Corella Aznar, 2011) (Table 1, Fig. 3). Nevertheless, the presence of biogenic varves, and the dominant presence of *Cyclotella distinguenda* and the low Sr values indicates relatively high lake levels during the mid-19th–mid-20th century (Corella et al., 2011). The increase in gypsum and strontium (Fig. 3) in the sediment suggests that lake waters were more concentrated than during warmer mid-20th century.

### 3.1.2 Lake Estanya

The Balsas de Estanya karstic system (42°02' N, 0°32' E, 670 m a.s.l.) is located at the foothills of the External Ranges, in the Central-Eastern Pre-Pyrenees (Fig. 1a). The sedimentary sequence has been dated by  $^{137}\text{Cs}/^{210}\text{Pb}$  and radiocarbon and spans the last 21 ka.

The multi-proxy analysis of sediment cores recovered in the distal areas of the main lake, including sedimentology, elemental and isotopic geochemistry and biological indicators (pollen, diatoms and chironomids) reveals arid conditions during the MCA followed by an increase in humidity during the LIA, starting at 1300 AD and lasting to

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A multi-proxy perspective on millennium-long climate variability**

M. Morellón et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

1850 AD. Consistently, pollen reflects warmer and drier conditions for the MCA, with a landscape dominated by junipers and Mediterranean elements, a relatively low abundance of mesophytes and a poorly developed aquatic component. The increase in humidity during the LIA is characterized by increase of riparian and hygrophytes (5 *Salix*, *Ulmus*, *Tamarix*, *Potamogeton*, *Myriophyllum*, *Nuphar*, *Lemna*), and a recovery of mesophytes (*Betula*, *Corylus*, *Alnus*, deciduous *Quercus*, *Abies*) (Fig. 2, Table 1). Likewise diatom assemblages suggests low water levels and saline conditions during the MCA (Morellón et al., 2011).

A reconstruction of salinity, based on the results from Principal Component Analyses applied to the high-resolution geochemical dataset (see details in Morellón et al., 10 2011) identifies large hydrological fluctuations. Other biological proxies (diatoms, chironomids) and sedimentological indicators show a coherent evolution (Table 1, Fig. 3). The LIA in Estanya is characterized by complex internal fluctuations with increased water balance during the solar minima.

### 15 3.1.3 Lake Montcortès

Lake Montcortès (42°19' N, 0°59' E, 1027 m a.s.l.) is located 50 km north-east of Estanya (Fig. 1a). The composite sequence of Montcortès spans the last 5.3 ka and constitutes the longest varved, annually resolved sequence recovered to date in the IP (Corella Aznar, 2011). The age model is particularly robust because of the combination of varve counting, <sup>210</sup>Pb and radiocarbon dating. 20

The analysis of microfacies and sublayering of laminae revealed colder conditions during the LIA (1350–1850 AD) (Corella Aznar, 2011). During this period, most of the calcite laminae show a fine-coarse sub-layering interpreted as a result of prolonged winter conditions with a delayed warming in spring, which may cause supersaturation leading to rapid nucleation of small calcite crystals. Higher Fe/Mn ratios also suggest more frequent and longer anoxic conditions at the hypolimnion (Corella Aznar, 25 2011), and increased abundance of planktic diatom *Cyclotella cyclopuncta* points to higher lake levels (Scussolini et al., 2011). The lack of diatom valves during the MCA

is explained through the synergy of arid conditions and higher anthropogenic pressure on the lake. The presence of the low Mediterranean scrub community reflects warmer conditions during the MCA, and its disappearance during the LIA in the 15th century, together with the increase in forest, are interpreted as colder and more humid conditions during the LIA (Rull et al., 2011). Towards the end of the LIA (ca. 1700–1900 AD) intermittent abundance of diatoms and occasional predominance of pennate, littoral taxa point to fluctuating lake levels; in more recent times, the presence of oligotrophic species marks the shift to poorer nutrient levels (Fig. 3).

### 3.1.4 Lake Basa de la Mora

Lake Basa de la Mora (42°32' N, 0°19' E, 1914 m a.s.l.) is a shallow, glacial-karstic lake located in the Cotiella Massif, in the Central-Eastern area of the Internal Ranges of the Pyrenees (Fig. 1a).

This radiocarbon-dated sedimentary sequence displays the highest resolution for the Holocene period in the region (117 cm kyr<sup>-1</sup>). The multiproxy analyses of sediment cores recovered at the deepest part of the lake revealed more arid conditions in medieval times (900–1300 AD), as indicated by: a rise in *Juniperus*, a decrease in mesophytes and aquatic plants and the highest percentages of heliophyte herbs (including *Artemisia*, although this taxum could also indicate the existence of anthropogenic activities in the area) (Table 1, Fig. 2), and a decreased in clastic input, represented by lower Si content (Fig. 3). The MCA is also characterized by high organic and carbonate productivity related to a larger development of the palustrine area and, possibly, the lowest lake level of the last 2000 yr. During the most of the LIA (1300–1750 AD), an opposite pattern occurs, leading to higher values of elements associated to detrital input due to an increase in runoff (Moreno et al., 2011b; Pérez-Sanz, 2009). Sedimentological and geochemical signatures for the last 200 yr are coherent again with lower lake levels and drier climate conditions, although some relatively humid intervals (e.g., 1900–1970 AD) occur.

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.1.5 Lake Redon

Lake Redon (42°38' N, 0°46' E; 2240 m a.s.l.) is located in the high altitudes in the Maladeta batholith, an area characterized by low human impact (summer occasional sheep grazing) (Fig. 1a). The age model of the sequence is based on <sup>210</sup>Pb dating and 12 radiocarbon dates, covering the entire Holocene (Camarero et al., 1998). Chrysophytes cysts are used as proxies of winter/spring temperatures (Pla and Catalán, 2005) and diatoms for water alkalinity (Catalan et al., 2009), which eventually can be related to air temperature of the snow-free period (summer and autumn). For both proxies, transfer functions were developed, based on calibration sets of more than one hundred lakes. A recent study has confirmed the relationship between chrysophyte assemblages and spring temperatures, as well as the mechanism behind it (Pla-Rabes and Catalan, 2011).

Interestingly, the two records did not show a parallel behaviour (Fig. 3). A decrease of snow period temperatures (winter/spring) would have started earlier than a similar decrease in snow-free period temperatures (summer-autumn). While summer temperature was still high during 11th and 12th centuries, winter temperatures decrease sharply. According to Redon record summer cooling occurred abruptly by 1200 AD and reached the lowest temperature values by 1700 AD. The snow period temperatures improved progressively from the 12th century but showed cold phases in the 14th, 18th and early 20th centuries.

### 3.1.6 Portlligat Bay

Portlligat Bay (42°17' N; 3°17' E, 0 m a.s.l.) is a shallow (maximum depth, 7.1 m) and small inlet located in the NE Mediterranean coast of the IP, connected to the sea (Fig. 1a). Pollen associations (López-Sáez et al., 2009) (Fig. 2) indicate relatively dry conditions and high temperatures during the MCA, following the drop of *Pinus sylvestris* formations and the development of Mediterranean taxa such as Cistaceae and *Quercus suber*. The sequence shows a different pattern of forest evolution during the LIA

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with higher percentages of *Pinus sylvestris* and lower Mediterranean formations. These could be related to colder temperatures. Concentration of fungal ascospores Type 18 has been used for the reconstruction of humidity (Van Geel, 1978) and fluctuating but more humid conditions were recorded from 1250 AD to the early 1800s (Table 1). Maximum humidity was reached from 1450 to 1550 AD, whereas the most arid conditions took place during the 19th and 20th centuries and, secondarily, during the MCA (1000–1150 AD) (López-Sáez et al., 2009) (Fig. 2).

## 3.2 Dendroclimatological reconstructions

### 3.2.1 Gerber-Sobreestivo (GER-SOB) chronology

Two timberline sites (GER and SOB) of similar ecological and climatic conditions were selected for this reconstruction (see Büntgen et al., 2008 for details). Samples include living and dry-dead pine (*Pinus uncinata* Mill.) trees of all age-classes. The Gerber site (GER) (42°38' N, 1°60' E, 2200–2350 m a.s.l.) is located within the central-eastern part of the Pyrenees, whereas the Sobreestivo site (42°41' N, 0°06' E, 2300–2400 m a.s.l.), is located 70 km west of the GER site (Fig. 1a). Temperature records from the Pic du Midi de Bigorre mountain observatory (43°04' N, 0°09' E, 2862 m a.s.l.) were used for proxy calibration trials. Methodological details can be found in Büntgen et al. (2008). This annually resolved record of June–September temperature variations robustly spans the 1260–2005 AD period, and thus constitutes the longest annually-resolved warm season temperature record for the Pyrenees.

Relatively high summer temperatures were recorded during most of the 1300–1500 AD period and the 20th century, separated by a rather prolonged cooling from around 1450–1850 AD, corresponding to the LIA (Table 1, Fig. 3). The six warmest decades occurred during the 20th century with the following four warmest being reconstructed for the period 1360–1440 AD (Büntgen et al., 2008).

CPD

7, 3049–3089, 2011

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.2.2 Capdella record

Five chronologies from several sites in the Spanish-Pyrenees were used for this reconstruction: Aigües Tortes and Larra, obtained from *Pinus uncinata* R. samples, and Pinobajo, Trapa and Ibónciecho, from *Pinus sylvestris* (Saz Sánchez, 2003). Standard techniques were applied for the sampling, construction of chronologies and reconstruction of climatic variables (Fritts, 1990). Among 20 and 25 samples were extracted from living trees to construct each chronology. Tree-rings widths were measured with a resolution of 0.01 mm using an ANIOL measuring table (Aniol, 1983). Absolute dating of tree-rings was checked by program COFECHA (Holmes, 1986). ARSTAN was used to standardize ring-width measures in order to eliminate low frequency non climatic signal and to combine standardized indices into site chronologies (Cook and Kairiukstis, 1990). Rainfall records from Capdella weather station (42°28' N, 0°59' E, 1407 m a.s.l.) were used for proxy calibration. Finally, using PRECON (Fritts and Shashkin, 1995) chronologies and climatic data are calibrated in order to obtain a response function. Methodological details can be found in (Saz Sánchez, 2003) and (Chueca Cía et al., 2005) (Table 1, Fig. 3). The reconstructed precipitation shows relatively higher values than in recent times for the last 500 yr, with several marked decade – long humid periods around AD 1500, 1600, 1700 and the early 20th century and with relative rainfall minima centred at ca. 1800 AD, 1900 AD and an abrupt decreasing trend during the late 20th century.

### 3.3 Evidences of glacier fluctuations

Extensive research carried out by (González Trueba et al., 2008) demonstrated that glaciers occurred in more than 20 different massifs of the Pyrenees during the LIA, with a wide range of mean (Equilibrium line altitudes) ELAs above between 2650 m a.s.l. in the oceanic-influenced massifs (Balaitous and Infierno massifs among others) and 2900 m a.s.l. in the continental massifs (Maladeta and Posets massifs among others). Ten massifs presenting glaciers were considered relevant for this study (Fig. 1b).

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The most common aspect was the development of glacier in the cirques with reduced or absent tongues. This study is based on geomorphologic and morphostratigraphical surveys and historical archives and demonstrates the rapid response of these ice bodies to climatic variations.

5 The historical maximum extent of glaciers in the Pyrenees took place between 1600 and 1750 AD, followed by a retreat during the next 50 yr, alternating with minor re-advances responsible of the development of numerous small moraines (Fig. 3). A secondary advance, characterized by a fast but short growth of ice bodies was recorded during the 19th century, lasting until the 1920s. The temperatures estimated from the lowering of the ELAs are 0.7–0.9°C colder than present-day conditions in this area. A generalized phase of glaciers retreat occurred since the 1870s, with minor re-advances during the last decades of the 19th century and the 1920s. Since then, glaciers underwent a continuous retreat, becoming ice patches in most of the cirques affected by LIA glaciations, with minor expansions in 1945 and 1964 AD in the French slopes. Since the 1970s–1980s, a drastic retreat started, continuing till the present (Fig. 3) (González Trueba et al., 2008).

#### 4 Millennium-long environmental changes in the Southern Pyrenees

Selected records from the Southern Pyrenees, including lacustrine and marine-littoral sequences, dendroclimatological records, and reconstructions of glacier fluctuations, provide evidences of a broadly consistent pattern of climatic fluctuations during the last millennium. The overall good agreement of a wide range of sequences with different locations and altitudes, likely subjected to variable human activity, suggests that they respond to a common climatic forcing. Comparison within these sequences and correlation with other regional and global records enable us to discuss the timing, intensity and chronology of the main environmental changes occurred prior, during and after the LIA and their relationship with climatic forcing mechanisms (Fig. 3).

CPD

7, 3049–3089, 2011

### A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cattle raising, resulting in a change to mesotrophic conditions (López-Sáez et al., 2009) (Table 1, Fig. 2).

The increase in flooding activity in the alluvial systems of the nearby Central Ebro Basin, whose headwaters are located in the Pyrenees, also indicate increased humidity during this period (Sancho et al., 2007; Constante et al., 2011). The occurrence of predominantly humid conditions during the LIA has been widely detected elsewhere in the IP (Moreno et al., 2011a): (i) higher lake levels were reconstructed in the north (e.g., La Cruz Lake, Iberian Range, Julià et al., 1998) and in the south of Spain (Martín-Puertas et al., 2008, 2010), (ii) increased continental runoff reconstructed from marine sequences (e.g., Nieto-Moreno et al., 2011), (iii) increased Westerlies in the W Mediterranean (Moreno et al., 2011b), (iv) increased storm activity in the nearby French Mediterranean Coast (Dezileau et al., 2011) and (v) more frequent floods in the Tagus headwaters (Moreno et al., 2008; Benito et al., 2003). Accordingly, increased water availability has been reconstructed in other areas of the W Mediterranean, such as Morocco (Détriché et al., 2009; Esper et al., 2007) and in the Alps (Magny et al., 2008, 2010).

A change in calcite sublayering in Lake Montcortès to dominant Fine-Coarse textures indicative of predominantly lower temperatures was recorded after ca. 1350 AD (Corella Aznar, 2011), coinciding with colder conditions in Lake Redon, both in winter and summer temperatures (Pla and Catalán, 2005; Catalan et al., 2009). Other evidences of widespread colder conditions in the region come from the neighbouring regions of Landes and Aquitaine (SW France) where loess deposits occurred from 0.8 to 0.2 ka BP (Bertran et al., 2011). The lack of temperature data from GER-SOB dendroclimatic reconstruction prior to 1300 AD makes it difficult to attribute an exact date to the onset of cold conditions associated to the LIA. In fact, Büntgen et al. (2008) reconstructed the four warmest decades of pre-industrial times during the period 1360–1440 AD, which indicates the predominance of relatively mild thermal conditions during the first part of the LIA. However, a drop in temperatures occurred during the mid to late 15th century, consistent with a decrease in winter temperatures in Redon.

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A multi-proxy perspective on millennium-long climate variability**

M. Morellón et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Other indirect evidences of temperature changes also indicate that the coldest conditions of the LIA in NE Iberian Peninsula did not take place until the 15th–16th centuries. As an example, the nearby Ebro River (Fig. 1a), fed by many tributaries from the Southern Pyrenees, froze near its mouth in the town of Tortosa (Tarragona), 17 times since ca. 1400 AD. According to historical reports (Puente, 2007), the maximum frequency (13 out of 17) of these frost episodes and intensity occurred between the 16th and 18th centuries. Historical reports also document the presence of an extensive network of ice stores that were built and maintained between the 16th and 19th centuries along the Mediterranean Coast of the IP (Font-Tullot, 1986), which evidences the occurrence of long-lasting low temperatures during this period. Thus, and in spite of the limitations of the datasets exposed above, moister conditions associated with the LIA started earlier (1300–1350 AD) than the main decrease in temperatures (1400–1500 AD). Consistently, global climate reconstructions indicate a decrease in temperatures along the 15th century (Mann and Jones, 2003), associated to a decline in solar irradiance (Steinilber et al., 2009; Bard et al., 2000).

Maximum (recent) glacier advances in the Southern Pyrenees took place from 1600 to 1750 AD, leading to a slower development during the late 18th and early 19th century. The chronology of this phase coincides with reconstructions for the European Alps carried out by (Schaefer et al., 2009). In spite of the occurrence of cold and humid conditions along the LIA, glacier advances required a combination of high winter precipitation and low summer temperatures (Paasche and Bakke, 2010). According to this dataset, these two conditions were probably not fulfilled during the early phases of the LIA in the Pyrenees, between 1300 and 1600 AD. Coldest summer temperatures, accompanied by relatively cold winter conditions in Redon occurred from 1500 to 1700 AD. Consistently, the most humid conditions, indicated by minimum salinity and maximum lake levels and/or clastic input, were not reached until the period 1500–1600 AD (Fig. 3) and maximum precipitation in Capdella was recorded by tree rings ca. 1600 AD (Saz Sánchez, 2003), coinciding with a phase of dominant negative NAO conditions (Fig. 3).

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



It is noteworthy that stable predominantly negative NAO conditions did not occur until this period 1550 AD (Fig. 3), emphasizing the importance of winter precipitations needed for glacier growth in the Pyrenees. In fact, although Alpine glaciers started their LIA-related re-advance during the 13th century, they did not reach their maximum and most continuous temporal extension until the period 1600 AD (Denton and Broecker, 2008). However, Calderone Glacier in Gran Sasso Massif (Italy), one of the Southernmost glacier in Mediterranean Europe, experienced a minor advance in the period 1313–1373 AD, prior to a maximum advance later on during the LIA, coinciding with higher lake levels in lakes Acesa and Fucino centered around 500 cal yrs BP (see (Giraudi et al., 2011) and references therein).

We can broadly identify two major phases of environmental change occurred during the LIA in the Southern Pyrenees: (i) a transitional phase MCA-LIA, characterized by fluctuating, moist conditions and relatively cold temperatures, between ca. 1300 and 1600 AD; and (ii) a second period, extending from ca. 1600 to 1850 AD, characterized by lower temperatures and higher humidity, leading to maximum glacier advances. In spite of the predominance of cold and humid conditions along the LIA, a lower amplitude hydrological and thermal internal variability was recorded in most of the sites (Fig. 3), as discussed below.

### 4.2 LIA to recent warming transition

An aridification trend started synchronously in lakes Arreo, Estanya, Montcortès and Basa de la Mora after ca. 1800 AD. Higher water concentration and/or more frequent oxic conditions in these settings indicate lower lake levels and decreased water balance. Pollen records show a new increase of human pressure and no aridity trends can be clearly extracted. Nevertheless, humidity reconstruction based on non-pollen palynomorphs in Portlligat indicates an abrupt decrease starting ca. 1750 AD, coinciding with an abrupt reduction of *Castanea* and *Vitis* culture while cereal cultivation is completely interrupted; the percentage of several arboreal elements (*Fagus*, *Abies*, *Corylus*, *Quercus suber* type) are also reduced or even disappear.

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Decreased humidity has been also recorded in lakes in the North of the IP (e.g., Taravilla, Moreno et al., 2008; Lake Enol, López-Merino et al., 2011) and S Spain (e.g., Doñana, Sousa and García-Murillo, 2003; Lake Zoñar, Martín-Puertas et al., 2008). Marine cores around the IP also documented reduced river input associated with lower humidity (e.g., Nieto-Moreno et al., 2011). A similar hydrological response was detected in other areas of the W Mediterranean, like the French Jura (Magny et al., 2008) or Morocco (Détriché et al., 2009).

However, Capdella tree ring sequence display a more complex variability, characterized by relative rainfall minima centred at ca. 1800 AD, 1900 AD and an abrupt decreasing trend during the late 20th century (Fig. 3).

Temperatures started to rise at the same time, between 1800 and 1850 AD, according to GEB-SOB record, reaching their maximum values during the 20th century. In fact, dendroclimatological reconstructions carried out in Morocco also show an increase in aridity after the mid-20th century (Esper et al., 2007). A change in calcite sub-layering also occurred in Montcortès at this time (Fig. 3). Consistently, glaciers started to retreat continuously after 1850 AD, with a brief readvance between the last decades of the late 19th century and the 1920s (González Trueba et al., 2008), coinciding with lower winter temperatures at Lake Redon (Pla and Catalán, 2005) (Fig. 3). Glacier readvances at this time have also been documented in other Mediterranean areas, like the Apennines (Giraudi, 2005), the Balkans (Hughes, 2010), Sierra Nevada (Gómez Ortiz et al., 2002, 2009), or the Cantabrian Mountains (González-Trueba, 2006, 2007; Serrano et al., 2011). Interestingly, this episode correlates with an increase in rainfall documented in Capdella and a decrease in NAO values during the early 20th century (Trouet et al., 2009), coinciding with a relative maximum in precipitation recorded in Capdella and a relative temperature minima in the GER-SOB records (Fig. 3).

The onset of more arid conditions reconstructed for lake sediments in Arreo occurred right afterwards, since 1870 AD. Finally a further decrease in temperatures occurred in the mid-20th century, as documented by GER-SOB and Redon records. Minor glacier stabilisation phases described by González-Trueba et al. (2008) might also be related

with this short-living climate deterioration phase, in the frame of a generalized warming trend.

## 5 Regional context and paleoclimatic implications of the LIA in the Pyrenees

Temperature fluctuations in the Southern Pyrenees display a similar pattern compared to other reconstructions from Central Europe (Büntgen et al., 2010a) and with other Northern Hemisphere palaeoclimate records at centennial timescales (e.g., Jones et al., 1998; Mann and Bradley, 1999) showing consistency with changes in solar activity (Steinilber et al., 2009; Bard et al., 2000; Vaquero et al., 2002), global temperature fluctuations (Mann et al., 2008), and NAO index (Trouet et al., 2009). Another natural forcing factor of the climate besides solar activity are volcanic eruptions (Gao et al., 2008). Comparison of climate records and volcanic activity does not show a clear pattern in the Pyrenees. On the one hand, some climate records show the expected cooling after volcanic eruptions, for example after the eruption of Tambora in 1815 AD. On the other hand, the largest eruption in the past 1500 yr (about 2.5 times larger stratospheric sulphate injection than Tambora) of an unknown volcano in the 1258 AD does not show up in any of the analysed climate records. Both, low solar activity and volcanic eruptions, lead to a cooling at ground compared to higher solar activity and no volcanic eruptions. In the past 1000 yr volcanic eruptions coincided with times of low solar activity as e.g. during the LIA and only a few eruptions occurred during the MCA when solar activity was relatively high. Thus, climate changes found in the climate reconstructions and interpreted as results due to low solar activity might partly be due to volcanic eruptions.

### 5.1 The footprint of solar activity

In spite of the limited chronological resolution of most of the records, derived from radiocarbon dating, several oscillations in the hydrological and thermal dynamics of

CPD

7, 3049–3089, 2011

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A multi-proxy perspective on millennium-long climate variability**

M. Morellón et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the reviewed sequences during the LIA suggest that colder temperatures and at least, more positive water balances and/or increased runoff dominated during periods of diminished solar activity, the so-called grand solar minima such as: the Oort (1010–1050 AD), Wolf (1282–1342 AD, onset of the LIA), Spörer (1460–1550 AD), Maunder (1645–1715 AD) and Dalton (1790–1830 AD) minima, in agreement with recent research carried out in Central Europe (e.g., Magny et al., 2008, 2011a).

Contrasting patterns have been recorded for the Oort Minimum: although lower winter and summer temperatures were recorded in Lake Redon and relatively more humid conditions were reconstructed in Arreo, variable conditions occur in the rest of the lakes (Estanya, Montcortès and Basa de la Mora). However, during the Wolf Minimum (1282–1342 AD) all lakes experienced more humid conditions, evidenced by increased lake levels, lower water concentration and/or increased sediment delivery. Although Portlligat Bay has recorded more humid conditions in the Eastern Pyrenees during the LIA, in contrast to drier environments during the MCA, no significant hydrological response occurs for this solar minimum. In agreement with the expected pattern, an abrupt decrease in temperature was recorded in tree-ring based reconstructions of GER-SOB and higher altitude (colder) conditions occurred in Lake Redon (Fig. 3).

Colder conditions also occurred in the Pyrenees during the Spörer Minimum, as indicated by lower temperatures in GER-SOB dendroclimatic reconstruction, calcite sublayering in Montcortès and lower winter temperature in Redon. Interestingly, within this phase, lasting for almost 100 yr (1460–1550 AD), generally recorded as a relatively humid period, all the lacustrine sequences reviewed in this paper show a short living reversed pattern (more negative water balance) centred around 1500 AD. Higher mean annual rainfall was also recorded in Capdella tree-ring sequence, with an abrupt decrease in precipitations at the end of this phase.

During the Maunder Minimum reduced summer temperatures were recorded both in GER-SOB and Redon, coinciding with a major phase of glacier advances. Most of the lake sequences clearly evidence higher effective moisture and/or increased runoff. However, the easternmost records of the region reviewed in this paper (Montcortès,

Portlligat) show a less clear response, probably due to particular, local conditions. Mean annual rainfall reconstructed from Capdella also shows a relative maximum, less evident than the one corresponding to the Spörer Minimum (Fig. 3).

In contrast, a clear temperature drop becomes evident in GER-SOB towards the end of the Dalton Minimum, coinciding with a series of volcanic eruptions at the beginning of the 19th century, and contemporaneous to a short phase of glacier advances. Although temporal resolution of the Lake Redon record is too coarse compared to the duration of Dalton Minimum, slightly colder summer, and warmer winter temperatures occurred at this site. Colder but variable thermal conditions were recorded in Montcortès. Generally humid conditions occurred in most of the lakes. However, Capdella only recorded relatively more humid conditions at the beginning of this phase. The short duration of the Dalton Minimum event (ca. 40 yr) and its relatively lower amplitude compared to the Spörer Minimum might be responsible for these differences. The last significant 'major' phase of glacier advance also took place at the early 19th century, slightly delayed if compared with the other records, likely due to the longer response time of glaciers to abrupt climate fluctuations.

Although most of the reviewed sequences display colder and more humid conditions during the grand solar minima, the timing and intensity of the environmental response is highly variable. The influence of volcanic eruptions, other regional and local climate mechanisms and/or human influence might be responsible for this. Additionally, the chronological resolution of these records in some cases (e.g., lacustrine records, glacier fluctuations) or the limited sensitivity of proxies may obscure the impact of this high-frequency climate variability.

## 5.2 Hydrological variability and the North Atlantic Oscillation

Hydrological fluctuations in the Southern Pyrenees appear to be in phase with the NAO reconstruction for the last millennium (Trouet et al., 2009). In agreement with most of the IP (Moreno et al., 2011a), arid conditions prevailed during the MCA, when reconstructed NAO values were predominantly positive (Trouet et al., 2009) (Fig. 3). During

### A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A multi-proxy perspective on millennium-long climate variability**

M. Morellón et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the instrumental period, there is clear evidence of interannual precipitation variations in the W Mediterranean area, with the highest positive rainfall anomaly located over N Iberia at the mean time as high pressure is dominant in N Europe (Cullen and deMenocal, 2000). Negative rainfall anomalies during the MCA and vice-versa during the LIA for the W Mediterranean region have been also reproduced by gridded high-resolution precipitation reconstructions (Pauling et al., 2006) and ECHO-G model simulations (Liu et al., 2009). Widespread proxy evidences for this centennial-scale pattern also exist (Graham et al., 2011). Furthermore, correlation maps between NAO and winter precipitation over the IP for the period 1950–1990 reproduced by models display particularly high negative values for the Southern Pyrenees within the IP (Gómez-Navarro et al., 2011) reinforcing this NAO-climatic forcing for the last centuries.

Accordingly, high frequency mean annual rainfall variability captured in tree-ring record of Capdella (Saz Sánchez, 2003) displays a good correlation with NAO reconstruction and also humidity patterns reconstructed from lake sediments agree on multi-centennial timescales, reproducing this transition from arid conditions during the MCA into higher humidity during the LIA (Fig. 3). However, lake records show an increase in humidity starting at ca. 1300 AD, whereas, although NAO values show a decreasing trend after the MCA, most negative values are not reached until 1600 AD. Higher humidity recorded in lakes during the period 1300–1600 AD might have been amplified by increased aquifer recharge resulting from reduced evaporation and thus, more positive water balance driven by lower temperatures during the LIA (e.g., Cacho et al., 2010 and references therein). Glacier maximum development coincides with the coldest and most humid period of the LIA in the Southern Pyrenees (1600–1850 AD) and displays a similar pattern compared to other Mediterranean mountains, like the Apennines (Giraudi, 2005) and the Balkans (Hughes, 2010), among others (González Trueba et al., 2008). The decoupling between temperature reconstructions and solar activity during recent times points towards another influence, which is probably due to human activity, in agreement with many other studies from all over the world (Solomon et al., 2007).

Main phases of environmental change during the last millennium are synchronous at a regional scale in the Iberian Peninsula (Moreno et al., 2011a) and also can be recognized across the Mediterranean area. However, there is an antiphase behaviour in the hydrological signal, with dry MCA and wet LIA in the western Mediterranean records and the opposite in some Eastern Mediterranean lakes (Roberts et al., 2011). This Mediterranean antiphase pattern is mostly caused by the higher influence of the NAO in westernmost areas. Thus, the NAO appears to be the main climatic mechanism controlling rainfall patterns at least at centennial to multidecadal timescales prior to the instrumental period.

## 6 Conclusions

Paleoclimatic records from the Southern Pyrenees, including lacustrine and marine-littoral sequences, tree rings, and records of glacier fluctuations, evidence a broadly consistent pattern at centennial to multi-decadal timescales for hydrological and thermal fluctuations during the last millennium, likely due to a common external solar forcing and NAO variability.

Warmer and arid conditions prevailed during the MCA (up to 1300 AD) and colder and humid conditions, evidenced by higher lake levels, increased runoff and a change in dominant vegetation during the LIA (1300–1800 AD). Although the LIA is characterized by an internal complex structure, two major phases of environmental change can be identified in all the records within this period: (i) a transitional phase MCA-LIA, characterized by fluctuating, moist conditions and relatively cold temperatures (ca. 1300 and 1600 AD); and (ii) a second period, characterized by lower temperatures and higher humidity, coinciding with maximum (recent) glacier advances in the second half of the LIA (ca. 1600–1850 AD)

A synchronous increase in temperature and aridity, coinciding with glacier retreat is found in all the records after 1800 AD, coinciding with the end of the LIA. A short-living

CPD

7, 3049–3089, 2011

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

return towards colder conditions and glacier stabilisation occurred during the late 19th–early 20th century, leading to warmer and more arid conditions afterwards.

Climatic oscillations reconstructed for the last millennium are consistent with other reconstructions from the Western Mediterranean area and glacier fluctuations are also in phase with other Western European (i.e., Alps) and Mediterranean mountains (e.g., Apennines and Balkans). A broadly consistent response to phases of low solar activity has also been found, generally characterized by colder and more humid conditions. We did not find a clear pattern regarding the climatic response on volcanic eruptions. While some climate records show the expected cooling due to the volcanic eruption of Tambora in 1815 AD, the same records do not show a significant change due to the strongest volcanic eruption in the past 1500 yr in 1258 AD from an unknown volcano. Furthermore, hydrological changes reconstructed in this region indicate a strong relationship with reconstructed NAO variability that appears to be the main climatic mechanism controlling high- to mid-frequency variations in precipitation totals during the last millennium.

*Acknowledgement.* This research has been funded by the Spanish InterMinistry of Science and Technology (CICYT), through the projects LIMNOCAL (CGL2006-13327-C04-01), GLOB-ALKARST (REN2003-09130-C02-02), GRACCIE-CONSOLIDER (CSD2007-00067), DINAMO (CGL2009-645-07992) and CRYOMONT (CGL2010-19724). Additional support was provided by the Spanish National Parks Agency through the project HORDA (083/2009). M. Morellón is supported by a Postdoctoral fellowship funded by the Spanish Ministry of Education and Science through FECYT (Spanish Foundation for Science and Technology). A. Pérez-Sanz is supported by a pre-doctoral fellowship funded by the Regional Aragonese Government and A. Moreno acknowledges the “Ramón y Cajal” postdoctoral program for funding.

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## References

- Abrantes, F., Lebreiro, S., Rodrigues, T., Gil, I., Bartels-Jonsdottir, H., Oliveira, P., Kissel, C., and Grimalt, J. O.: Shallow-marine sediment cores record climate variability and earthquake activity off Lisbon (Portugal) for the last 2000 years, *Quaternary Sci. Rev.*, 24, 2477–2494, 2005.
- Aniol, R. W.: Tree-ring analysis using catras, *Dendrochronologia*, 1, 45–53, 1983.
- Bal, M.-C., Pelachs, A., Perez-Obiol, R., Julia, R., and Cunill, R.: Fire history and human activities during the last 3300 cal yr bp in Spain's Central Pyrenees: the case of the Estany de Burg, *Palaeogeogr. Palaeoclimatol.*, 300, 179–190, 2011.
- Bard, E., Raisbeck, G., Yiou, F., and Jouzel, J.: Solar irradiance during the last 1200 years based on cosmogenic nuclides, *Tellus B*, 52, 985–992, 2000.
- Beniston, M.: Climatic change in mountain regions: a review of possible impacts, *Clim. Change*, 59, 5–31, 2003.
- Benito, G., Díez-Herrero, A., and Fernández de Villalta, M.: Magnitude and frequency of flooding in the Tagus basin (Central Spain) over the last millennium, *Clim. Change*, 58, 171–192, 2003.
- Bertran, P., Bateman, M. D., Hernandez, M., Mercier, N., Millet, D., Sitzia, L., and Tastet, J.-P.: Inland aeolian deposits of south-west france: Facies, stratigraphy and chronology, *J. Quaternary Sci.*, 26, 374–378, 2011.
- Blanco-Castro, E., Casado, M., Costa, M., Escribano, R., Garcí a Antón, M., Génova, M., Gómez, A., Moreno, J., Morla, C., Regato, P., and Sainz Ollero, H.: Los bosques ibéricos. Una interpretación geobotánica, *Planeta, Barcelona (Spain)*, 572 pp., 1997.
- Büntgen, U., Frank, D., Grudd, H., and Esper, J.: Long-term summer temperature variations in the Pyrenees, *Clim. Dynam.*, 31, 615–631, 2008.
- Büntgen, U., Franke, J., Frank, D., Wilson, R., González-Rouco, F., and Esper, J.: Assessing the spatial signature of European climate reconstructions, *Clim. Res.*, 41, 125–130, doi:10.3354/cr00848, 2010a.
- Büntgen, U., Trouet, V., Frank, D., Leuschner, H. H., Friedrichs, D., Luterbacher, J., and Esper, J.: Tree-ring indicators of German summer drought over the last millennium, *Quaternary Sci. Rev.*, 29, 1005–1016, doi:10.1016/j.quascirev.2010.01.003, 2010b.
- Büntgen, U., Tegel, W., Nicolussi, K., McCormick, M., Frank, D., Trouet, V., Kaplan, J. O., Herzig, F., Heussner, K.-U., Wanner, H., Luterbacher, J. R., and Esper, J.: 2500

### A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



years of European climate variability and human susceptibility, *Science*, 331, 578–582, doi:10.1126/science.1197175, 2011.

Cacho, I., Valero-Garcés, B., and González-Sampériz, P.: Revisión de las reconstrucciones paleoclimáticas en la península ibérica desde el último periodo glacial, in: *Clima en España: Pasado Presente y Futuro. Informe de Evaluación del Cambio Climático Regional*, edited by: Pérez, F. F. and Boscolo, R., Red temática Clivar-España, Madrid (Spain), 2010.

Camarero, L., Masqué, P., Devos, W., Ani-Ragolta, I., Catalan, J., Moor, H. C., Pla, S., and Sanchez-Cabeza, J. A.: Historical variations in lead fluxes in the Pyrenees (Northeast Spain) from a dated lake sediment core, *Water Air Soil Pollut.*, 105, 439–449, 1998.

Catalan, J., Pla, S., García, J., and Camarero, L.: Climate and CO<sub>2</sub> saturation in an alpine lake throughout the Holocene, *Limnol. Oceanogr.*, 54, 2542–2552, 2009.

Chueca Cía, J. C., Julián Andrés, A., Saz Sánchez, M. A., Creus Novau, J., and López Moreno, J. I.: Responses to climatic changes since the Little Ice Age on Maladeta Glacier (Central Pyrenees), *Geomorphology*, 68, 167–182, 2005.

Chueca, J., Peña, J. L., Lampre, F., García Ruiz, J. M., and Martí, C. E.: Los glaciares del pirineo aragonés: estudio de su evolución y extensión actual, edited by: Departamento de Geografía, Urbanismo y Ordenación del Territorio, Universidad de Zaragoza, Zaragoza (Spain), 1998a.

Chueca, J., Peña, J. L., Lampre, F., and Julián, A.: La pequeña edad del hielo en el pirineo central y meridional. Inferencias paleoambientales a partir de datos geomorfológicos, in: *Las Huellas Glaciares de las Montañas Españolas*, edited by: Gómez Ortiz, A., and Pérez Alberti, A., Universidad de Santiago de Compostela, Santiago de Compostela (Spain), 307–328, 1998b.

Constante, A., Peña, J. L., Muñoz, A., and Picazo, J.: Climate and anthropogenic factors affecting alluvial fan development during the late Holocene in the Central Ebro valley, Northeast Spain, *Holocene*, 21, 275–286, doi:10.1177/0959683610378873, 2011.

Cook, E. R. and Kairiukstis, L. A.: *Methods of Dendrochronology*, Kluwer Academic Publishers, London (UK), 394 pp., 1990.

Corella Aznar, J. P.: *Climate and Human Impact in Northern Spain Since Mid-Holocene: the High Resolution Records of Lakes Arreo and Montcortès*, Unpublished Phd Thesis, Department of Earth Sciences, University of Zaragoza, Zaragoza (Spain), 223 pp., 2011.

Corella, J., Amrani, A., Sigró, J., Morellón, M., Rico, E., and Valero-Garcés, B.: Recent evolution of Lake Arreo, Northern Spain: influences of land use change and climate, *J. Paleolimnol.*,

**A multi-proxy perspective on millennium-long climate variability**

M. Morellón et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

doi:10.1007/s10933-010-9492-7, in press, 2011.

Cullen, H. M. and de Menocal, P. B.: North Atlantic influence on Tigris–Euphrates streamflow, *Int. J. Climatol.*, 20, 853–863, 2000.

D'Andrea, W. J., Huang, Y., Fritz, S. C., and Anderson, N. J.: Abrupt Holocene climate change as an important factor for human migration in West Greenland, *P. Natl. Acad. Sci. USA*, doi:10.1073/pnas.1101708108, 2011.

Del Barrio, G., Creus, J., and Puigdefabregas, J.: Thermal seasonality of the high mountain belts of the Pyrenees, *Mountain Res. Develop.*, 10, 227–233, 1990.

Denton, G. H. and Broecker, W. S.: Wobbly ocean conveyor circulation during the Holocene?, *Quaternary Sci. Rev.*, 27, 1939–1950, 2008.

Détriché, S., Bréhéret, J. G., Soulié-Märsche, I., Karrat, L., and Macaire, J. J.: Late Holocene water level fluctuations of Lake Afourgagh (Middle-Atlas Mountains, Morocco) inferred from charophyte remains, *Palaeogeogr. Palaeocl.*, 283, 134–147, 2009.

Dezileau, L., Sabatier, P., Blanchemanche, P., Joly, B., Swingedouw, D., Cassou, C., Castaings, J., Martinez, P., and Von Grafenstein, U.: Intense storm activity during the Little Ice Age on the French Mediterranean coast, *Palaeogeogr. Palaeocl.*, 299, 289–297, 2011.

Ejarque, A., Miras, Y., Riera, S., Palet, J. M., and Orengo, H. A.: Testing micro-regional variability in the Holocene shaping of high mountain cultural landscapes: a palaeoenvironmental case-study in the Eastern Pyrenees, *J. Archaeol. Sci.*, 37, 1468–1479, doi:10.1016/j.jas.2010.01.007, 2010.

Esper, J., Frank, D., Buntgen, U., Verstege, A., Luterbacher, J., and Xoplaki, E.: Long-term drought severity variations in Morocco, *Geophys. Res. Lett.*, 34, L17702, 2007.

Fagan, B. M.: *The Little Ice Age: how climate made history, 1300–1850*, Basic Books, New York (USA), 2000.

Font-Tullot, I.: Cambios climáticos en la Península Ibérica durante el último milenio, con especial referencia a la “pequeña edad glacial”, in: *Quaternary Climate in Western Mediterranean*, Proceedings of the Symposium on Climatic Fluctuations During the Quaternary in the Western Mediterranean Regions, edited by: López-Vera, F., Universidad Autónoma de Madrid, Madrid (Spain), 1986.

Frank, D. C., Esper, J., Raible, C. C., Buntgen, U., Trouet, V., Stocker, B., and Joos, F.: Ensemble reconstruction constraints on the global carbon cycle sensitivity to climate, *Nature*, 463, 527–530, 2010.

Fritts, H. C. and Shashkin, A. V.: Modelling tree-ring structure as related to temperature,

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- precipitation and day length, in: *Tree Rings as Indicators of Ecosystem Health*, edited by: Lewis, T. E., CRC Press, Boca Raton, FL, 17–57, 1995.
- Gao, C., Robock, A., and Ammann, C.: Volcanic forcing of climate over the past 1500 years: an improved ice core-based index for climate models, *J. Geophys. Res.*, 113, D23111, 2008.
- 5 García, M. B. and Gómez, D.: Flora del pirineo aragonés. Patrones espaciales de biodiversidad y su relevancia para la conservación, *Pirineos*, 162, 71–88, 2008.
- García-Ruiz, J. M., López-Moreno, J. I., Vicente-Serrano, S. M., Lasanta-Martínez, T., and Beguería, S.: Mediterranean water resources in a global change scenario, *Earth-Sci. Rev.*, 105, 121–139, 2011.
- 10 Giraudi, C.: Middle to late Holocene glacial variations, periglacial processes and alluvial sedimentation on the higher Apennine massifs (Italy), *Quaternary Res.*, 64, 176–184, 2005.
- Giraudi, C., Magny, M., Zanchetta, G., and Drysdale, R. N.: The Holocene climatic evolution of mediterranean italy: a review of the continental geological data, *Holocene*, 21, 105–115, doi:10.1177/0959683610377529, 2011.
- 15 Gómez-Navarro, J. J., Montávez, J. P., Jerez, S., Jiménez-Guerrero, P., Lorente-Plazas, R., González-Rouco, J. F., and Zorita, E.: A regional climate simulation over the Iberian Peninsula for the last millennium, *Clim. Past*, 7, 451–472, doi:10.5194/cp-7-451-2011, 2011.
- Gómez-Ortiz, A., Schulte, L., Salvador, F., Sánchez, S., and Simón, M.: Geomorphological map of Sierra Nevada. Glacial and periglacial geomorphology, *Consejería de Medio Ambiente. Junta de Andalucía, Granada (Spain)*, 2002.
- 20 Gómez-Ortiz, A., Palacios, D., Schulte, L., Salvador-Franch, F., and Plana-Castellví, J. A.: Evidences from historical documents of landscape evolution after Little Ice Age of a mediterranean high mountain area, Sierra Nevada, Spain (eighteenth to twentieth centuries), *Geograf. Ann. Ser. A Phys. Geogr.*, 91, 279–289, doi:10.1111/j.1468-0459.2009.00370.x, 2009.
- 25 González-Trueba, J. J.: Topoclimatical factors and very small glaciers in Atlantic Mountain of SW Europe: the Little Ice Age glacier advance in Picos de Europa (NW Spain), *Z. Gletscherkunde Glazialgeol.*, 39, 115–125, 2006.
- González-Trueba, J. J.: *La Pequeña Edad del Hielo en los Picos de Europa*, Universidad de Cantabria, Santander (Spain), 2007.
- 30 González-Trueba, J. J., Moreno, R. M., Martínez de Pison, E., and Serrano, E.: “Little Ice Age” glaciation and current glaciers in the Iberian Peninsula, *Holocene*, 18, 551–568, doi:10.1177/0959683608089209, 2008.
- Goodkin, N. F., Hughen, K. A., Doney, S. C., and Curry, W. B.: Increased multidecadal variability

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the North Atlantic oscillation since 1781, *Nature Geosci.*, 1, 844–848, 2008.

Graham, N., Ammann, C., Fleitmann, D., Cobb, K., and Luterbacher, J.: Support for global climate reorganization during the “medieval climate anomaly”, *Clim. Dynam.*, 37, 1217–1245, doi:10.1007/s00382-010-0914-z, 2011.

5 Grove, J. M. and Gellatly, A. F.: Little Ice Age fluctuations in the Pyrenees, *Z. Gletscherkunde Glacialgeol.*, 31, 199–206, 1995.

Holmes, R.: Dendrochronology program library. Guide for computer program cofecha, Laboratory of Tree-Ring Research, Arizona, USA, 1986.

10 Hughes, P. D.: Little Ice Age glaciers in the Balkans: Low altitude glaciation enabled by cooler temperatures and local topoclimatic controls, *Earth Surf. Process. Landforms*, 35, 229–241, 2010.

15 Jansen, E., Overpeck, J., Briffa, K. R., Duplessy, J.-C., Joos, F., Masson-Delmotte, V., Olago, D., Otto-Bliesner, B., Peltier, W. R., Rahmstorf, S., Ramesh, R., Raynaud, D., Rind, D., Solomina, O., Villalba, R., and Zhang, D.: Palaeoclimate, in: *Climate Change 2007: The Physical Science Basis*, Intergovernmental Panel on Climate Change, Cambridge, United Kingdom and New York, NY, USA, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, 2007.

20 Jones, P. D., Briffa, K. R., Barnett, T. P., and Tett, S. F. B.: High-resolution palaeoclimatic records for the last millennium: interpretation, integration and comparison with general circulation model control-run temperatures, *Holocene*, 8, 455–471, doi:10.1191/095968398667194956, 1998.

25 Jones, P. D., Briffa, K. R., Osborn, T. J., Lough, J. M., van Ommen, T. D., Vinther, B. M., Luterbacher, J., Wahl, E. R., Zwiers, F. W., Mann, M. E., Schmidt, G. A., Ammann, C. M., Buckley, B. M., Cobb, K. M., Esper, J., Goosse, H., Graham, N., Jansen, E., Kiefer, T., Kull, C., Kuttel, M., Mosley-Thompson, E., Overpeck, J. T., Riedwyl, N., Schulz, M., Tudhope, A. W., Villalba, R., Wanner, H., Wolff, E., and Xoplaki, E.: High-resolution palaeoclimatology of the last millennium: a review of current status and future prospects, *Holocene*, 19, 3–49, doi:10.1177/0959683608098952, 2009.

30 Julià, R., Burjachs, F., Dasí, M. J., Mezquita, F., Miracle, M. R., Roca, J. R., Seret, G., and Vicente, E.: Meromixis origin and recent trophic evolution in the Spanish mountain Lake la Cruz, *Aq. Sci. Res. Across Bound.*, 60, 279–299, 1998.

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Last, W. M. and Smol, J. P.: Tracking environmental change using lake sediments, Developments in paleoenvironmental research series, Kluwer Academic Publishers, Norwell, MA, USA, 2001.

Liu, J., Wang, B., Ding, Q., Kuang, X., Soon, W., and Zorita, E.: Centennial variations of the global monsoon precipitation in the last millennium: results from echo-g model, *J. Climate*, 22, 2356–2371, doi:10.1175/2008jcli2353.1, 2009.

López-Merino, L., Moreno, A., Leira, M., Sigró, J., González-Sampériz, P., Valero-Garcés, B. L., López-Sáez, J. A., Brunet, M., and Aguilar, E.: Two hundred years of environmental change in Picos de Europa national park inferred from sediments of Lago Enol, Northern Iberia, *J. Paleolimnology*, doi:10.1007/s10933-011-9546-5, in press, 2011.

López-Moreno, J. I. and García-Ruiz, J. M.: Influence of snow accumulation and snowmelt on streamflow in the Central Spanish Pyrenees, *Hydrol. Sci. J.*, 49, 1–802, 2004.

López-Moreno, J. I., Goyette, S., and Beniston, M.: Climate change prediction over complex areas: spatial variability of uncertainties and predictions over the Pyrenees from a set of regional climate models, *Int. J. Climatol.*, 28, 1535–1550, 2008.

López-Sáez, J. A., López-Merino, L., Mateo, M. Á., Serrano, Ó., Pérez-Díaz, S., and Serrano, L.: Palaeoecological potential of the marine organic deposits of *posidonia oceanica*: a case study in the NE Iberian peninsula, *Palaeogeogr., Palaeoclimatol., 271*, 215–224, 2009.

Magny, M., Gauthier, E., Vanniere, B., and Peyron, O.: Palaeohydrological changes and human-impact history over the last millennium recorded at Lake Joux in the Jura Mountains, Switzerland, *Holocene*, 18, 255–265, doi:10.1177/0959683607086763, 2008.

Magny, M., Arnaud, F., Holzhauser, H., Chapron, E., Debret, M., Desmet, M., Leroux, A., Millet, L., Revel, M., and Vannière, B.: Solar and proxy-sensitivity imprints on paleohydrological records for the last millennium in West-Central Europe, *Quaternary Res.*, 73, 173–179, 2010.

Magny, M., Bossuet, G., Ruffaldi, P., Leroux, A., and Mouthon, J.: Orbital imprint on Holocene palaeohydrological variations in West-Central Europe as reflected by lake-level changes at cerin (Jura Mountains, Eastern France), *J. Quaternary Sci.*, 26, 171–177, 2011a.

Magny, M., Peyron, O., Gauthier, E., Vannière, B., Millet, L., and Vermot-Desroches, B.: Quantitative estimates of temperature and precipitation changes over the last millennium from pollen and lake-level data at Lake Joux, Swiss Jura Mountains, *Quaternary Res.*, 75, 45–54, 2011b.

Mann, M. E. and Bradley, R. S.: Northern Hemisphere temperatures during the past millennium:

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



inferences, uncertainties, and limitations, *Geophys. Res. Lett.*, 26, 759–762, 1999.

Mann, M. E. and Jones, P. D.: Global surface temperatures over the past two millennia, *Geophys. Res. Lett.*, 30, doi:10.1029/2003GL017814, 2003.

Mann, M. E., Zhang, Z., Hughes, M. K., Bradley, R. S., Miller, S. K., Rutherford, S., and Ni, F.: Proxy-based reconstructions of hemispheric and global surface temperature variations over the past two millennia, *P. Natl. Acad. Sci. USA*, 105, 13252–13257, doi:10.1073/pnas.0805721105, 2008.

Martín-Puertas, C., Valero-Garcés, B. L., Mata, M. P., González-Sampériz, P., Bao, R., Moreno, A., and Stefanova, V.: Arid and humid phases in Southern Spain during the last 4000 years: the Zonar Lake record, Cordoba, Holocene, 18, 907–921, doi:10.1177/0959683608093533, 2008.

Martín-Puertas, C., Jiménez-Espejo, F., Martínez-Ruiz, F., Nieto-Moreno, V., Rodrigo, M., Mata, M. P., and Valero-Garcés, B. L.: Late Holocene climate variability in the Southwestern Mediterranean region: an integrated marine and terrestrial geochemical approach, *Clim. Past*, 6, 807–816, doi:10.5194/cp-6-807-2010, 2010.

Miras, Y., Ejarque, A., Riera, S., Palet, J. M., Orenge, H., and Euba, I.: Dynamique holocène de la végétation et occupation des Pyrénées andorranes depuis le néolithique ancien, d'après l'analyse pollinique de la tourbière de bosc dels estanyons (2180 m, Vall del Madriu, Andorre), *Compt. Rend. Palevol.*, 6, 291–300, 2007.

Montserrat-Martí, J.: Evolución glacial y postglacial del clima y la vegetación en la vertiente sur del pirineo: Estudio palinológico., *Monografías del instituto pirenaico de ecología-csic*, Zaragoza, 147 pp., 1992.

Morellón, M., Valero-Garcés, B., González-Sampériz, P., Vegas-Vilarrúbia, T., Rubio, E., Rieradevall, M., Delgado-Huertas, A., Mata, P., Romero, Ó., Engstrom, D. R., López-Vicente, M., Navas, A., and Soto, J.: Climate changes and human activities recorded in the sediments of Lake Estanya (NE Spain) during the medieval warm period and Little Ice Age, *J. Paleolimnol.*, doi:10.1007/s10933-009-9346-3, in press, 2011.

Moreno, A., Valero-Garcés, B. L., González-Sampériz, P., and Rico, M.: Flood response to rainfall variability during the last 2000 years inferred from the Taravilla Lake record (Central Iberian Range, Spain), *J. Paleolimnol.*, 40, 943–961, 2008.

Moreno, A., Morellón, M., Martín-Puertas, C., Frigola, J., Canals, M., Cacho, I., Corella, J. P., Pérez, A., Belmonte, Á., Vegas-Vilarrúbia, T., González-Sampériz, P., and Valero-Garcés, B. L.: Was there a common hydrological pattern in the Iberian Peninsula region

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



during the medieval climate anomaly?, PAGES News, 19, 16–18, 2011a.

Moreno, A., Pérez, A., Frigola, J., Nieto-Moreno, V., Rodrigo-Gámiz, M., González-Sampériz, P., Morellón, M., Martín-Puertas, C., Corella, J. P., Belmonte, Á., Sancho, C., Cacho, I., Herrera, G., Canals, M., Jiménez-Espejo, F., Martínez-Ruiz, F., Vegas, T., and Valero-Garcés, B. L.: The medieval climate anomaly in the Iberian Peninsula reconstructed from marine and lake records, *Quaternary Sci. Rev.*, in review, 2011b.

Nieto-Moreno, V., Martínez-Ruiz, F., Giral, S., Jiménez-Espejo, F., Gallego-Torres, D., Rodrigo-Gámiz, M., García-Orellana, J., Ortega-Huertas, M., and de Lange, G. J.: Tracking climate variability in the western Mediterranean during the Late Holocene: a multiproxy approach, *Clim. Past Discuss.*, 7, 635–675, doi:10.5194/cpd-7-635-2011, 2011.

Paasche, Ø. and Bakke, J.: Defining the Little Ice Age, *Clim. Past Discuss.*, 6, 2159–2175, doi:10.5194/cpd-6-2159-2010, 2010.

Patterson, W. P., Dietrich, K. A., Holmden, C., and Andrews, J. T.: Two millennia of North Atlantic seasonality and implications for Norse colonies, *P. Natl. Acad. Sci. USA*, 107, 5306–5310, doi:10.1073/pnas.0902522107, 2010.

Pauling, A., Luterbacher, J., Casty, C., and Wanner, H.: Five hundred years of gridded high-resolution precipitation reconstructions over Europe and the connection to large-scale circulation, *Clim. Dynam.*, 26, 387–405, 2006.

Peinado Lorca, M. and Rivas-Martínez, S.: La vegetación de España, Colección aula abierta, 544 pp., 1987.

Pérez-Sanz, A.: Reconstrucción paleoambiental del ibón de la basa de la mora (pireneo central): Primeros resultados del análisis palinológico. Trabajo de fin de máster, Master Ciencias de la Tierra, Universidad de Zaragoza, Zaragoza (Spain), 2009.

Pla-Rabes, S. and Catalan, J.: Deciphering chrysophyte responses to climate seasonality, *J. Paleolimnol.*, 46, 139–150, 2011.

Pla, S. and Catalán, J.: Chrysophyte cysts from lake sediments reveal the submillennial winter/spring climate variability in the Northwestern Mediterranean region throughout the Holocene, *Clim. Dynam.*, 24, 263–278, 2005.

Proctor, C., Baker, A., and Barnes, W.: A three thousand year record of North Atlantic climate, *Clim. Dynam.*, 19, 449–454, 2002.

Puente, J. M.: Cuando el río se helaba. Las heladas históricas del ebro a su paso por tortosa, *Rev. Aficion. Meteorol.*, 1, 2007.

Roberts, N., Moreno, A., Valero-Garcés, B. L., Corella, J. P., Jones, M., Allcock, S.,

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Woodbridge, J., Morellón, M., and Luterbacher, J.: Palaeolimnological evidence for a bipolar climate see-saw in the Mediterranean since 900 AD, *Global Planet. Change*, in press, 2011.
- Rull, V., González-Sampériz, P., Corella, J., Morellón, M., and Giral, S.: Vegetation changes in the Southern Pyrenean flank during the last millennium in relation to climate and human activities: the Montcortès lacustrine record, *J. Paleolimnol.*, 1–18, doi:10.1007/s10933-010-9444-2, in press, 2011.
- Sancho, C., Peña, J. L., Muñoz, A., Benito, G., McDonald, E., Rhodes, E. J., and Longares, L. A.: Holocene alluvial morphopedosedimentary record and environmental changes in the Bardenas Reales Natural Park (NE Spain), *Catena*, doi:10.1016/j.catena.2007.09.011, 2007.
- Saz Sánchez, M. A.: Temperaturas y precipitaciones en la mitad Norte de España Desde el Siglo XV. Estudio dendroclimático, Serie investigación, Publicaciones del Consejo de la Naturaleza de Aragón, Zaragoza, Spain, 2003.
- Schaefer, J. M., Denton, G. H., Kaplan, M., Putnam, A., Finkel, R. C., Barrell, D. J. A., Andersen, B. G., Schwartz, R., Mackintosh, A., Chinn, T., and Schluchter, C.: High-frequency Holocene glacier fluctuations in New Zealand differ from the northern signature, *Science*, 324, 622–625, doi:10.1126/science.1169312, 2009.
- Scussolini, P., Vegas-Vilarrúbia, T., Rull, V., Corella, J., Valero-Garcés, B., and Gomà, J.: Middle and late Holocene climate change and human impact inferred from diatoms, algae and aquatic macrophyte pollen in sediments from Lake Montcortès (NE Iberian Peninsula), *J. Paleolimnol.*, doi:10.1007/s10933-011-9524-y, in press, 2011.
- Serrano, E. and Martínez de Pisón, E.: Geomorfología y evolución glaciar en el pirineo aragonés oriental, in: *El Glaciario Surpirenaico. Nuevas Aportaciones*, edited by: Martí, C. E. and García, J. M., Geoforma Ediciones, 33–64, 1994.
- Serrano, E., González-Trueba, J. J., Sanjosé, J. J., and Del Río, L. M.: Ice patch origin, evolution and dynamics in a temperate high mountain environment: the Jou Negro, Picos de Europa (NW Spain), *Geograf. Ann. Ser. A Phys. Geogr.*, 93, 57–70, doi:10.1111/j.1468-0459.2011.00006.x, 2011.
- Solomon, S., Qin, D., Manning, M., Marquis, M., Averyt, K., Tignor, M. M. B., Miller, H., and Chen, Z.: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, 2007.
- Sousa, A. and García-Murillo, P.: Changes in the wetlands of Andalusia (Doñana natural park,

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- SW Spain) at the end of the Little Ice Age, *Clim. Change*, 58, 193–217, 2003.
- Steinhilber, F., Beer, J., and Fröhlich, C.: Total solar irradiance during the Holocene, *Geophys. Res. Lett.*, 36, L19704, 2009.
- Touchan, R., Anchukaitis, K., Meko, D., Sabir, M., Attalah, S., and Aloui, A.: Spatiotemporal drought variability in Northwestern Africa over the last nine centuries, *Clim. Dynam.*, 37, 237–252, doi:10.1007/s00382-010-0804-4, 2011.
- 5 Trouet, V. R., Esper, J., Graham, N. E., Baker, A., Scourse, J. D., and Frank, D. C.: Persistent positive North Atlantic oscillation mode dominated the medieval climate anomaly, *Science*, 324, 78–80, doi:10.1126/science.1166349, 2009.
- 10 Valero-Garcés, B. and Moreno, A.: Iberian lacustrine records: responses to past and recent global changes in the Mediterranean region, *J. Paleolimnol.*, in press, 2011.
- Van Geel, B.: A palaeoecological study of Holocene peat bog sections in Germany and The Netherlands, based on the analysis of pollen, spores and macro- and microscopic remains of fungi, algae, cormophytes and animals, *Rev. Palaeobot. Palynol.*, 25, 1–120, 1978.
- 15 Vaquero, J. M., Gallego, M. C., and García, J. A.: A 250-year cycle in naked-eye observations of sunspots, *Geophys. Res. Lett.*, 29, doi:10.1029/2002GL014782, 2002.
- Vicente-Serrano, S. and Cuadrat, J.: North Atlantic oscillation control of droughts in North-East Spain: evaluation since 1600 AD, *Climatic Change*, 85, 357–379, 2007.

**Table 1.** Paleoclimatic sequences from the Southern Pyrenees reviewed in this paper, including climate/hydrology and vegetation conditions recorded for the different chronological stages defined in this paper: Late MCA (1000–1300 AD), LIA (1300–1800 AD) and Post-LIA (after 1800 AD).

Site and location	Type	Environmental conditions					
		Late MCA (1000–1300 AD) Hydrology/ climate Vegetation	LIA (1300–1800 AD) Hydrology/ climate Vegetation	Post-LIA (after 1800 AD) Hydrology/ climate Vegetation			
Arreo 42°46' N 2°59' W 655 m a.s.l.	Lake	Low lake levels, high salinity, holomictic conditions	Fall in mesophytes, mainly deciduous <i>Quercus</i> and development of ruderals, meadows and pastures	High lake levels, lower salinity, meromictic conditions	Deciduous forest recovery (mainly <i>Quercus</i> sp. and <i>Fagus</i> ). Decrease in human activities (meadows and pastures)	Low lake levels, higher salinity, holomictic conditions	Deforestation of oak and beech forest and development of <i>Pinus</i> formations (including recent <i>Pinus radiata</i> cultivation)
Estanya 42°02' N 0°32' E 670 m a.s.l.	Lake	Low lake levels, high salinity, variable runoff	Increase of <i>Juniperus</i> and Mediterranean taxa. Reduced proportions and variety of aquatic plants	Higher lake levels, lower salinity, variable runoff	Development of Mesophytes, peak of <i>Pinus nigra</i> type, and increase of aquatic taxa	High lake levels, reduced runoff	Recovery of open mixed Mediterranean/deciduous formations with Pine. Maximum expansion of cultivated areas
Montcoriès 42°19' N 0°59' E 1027 m a.s.l.	Lake	Warm conditions, reduced anoxia, higher runoff	Establishment of Mediterranean shrubland with increase in evergreen <i>Quercus</i> and decrease of pines. Development of cultivated taxa	Cold conditions, higher anoxia, lower runoff	Increase in AP formations, specially conifers and deciduous taxa. Decrease in shrubland and heliophytes	Warm conditions, lower anoxia, higher runoff	Expansion of conifer formations and anthropogenic influenced areas (mainly pastures with ruderals)
Basa de la mora 42°32' N 0°19' E 1904 m a.s.l.	Lake	High carbonate precipitation, decreased runoff	Development of <i>Juniperus</i> and heliophytes and decrease in all deciduous taxa	Lower carbonate precipitation, higher clastic input	Fall in <i>Juniperus</i> and heliophytes and increase in Mesophytes and Pteridophytes	High carbonate precipitation, decreased runoff	Recovery of mixed conifer ( <i>Pinus</i> and <i>Juniperus</i> )/deciduous taxa formations
Redon 42°38' N 0°46' E 2240 m a.s.l.	Lake	Higher summer-autumn temperatures, relatively low winter-spring conditions	–	Lower summer-autumn temperatures, relatively low winter-spring temperatures	–	Relatively high summer-autumn temperatures, high winter-spring temperatures	–
Portlligat 42°17' N 3°17' E 0 m a.s.l.	Shallow marine	–	Increase in <i>Quercus suber</i> , deciduous <i>Quercus</i> forests and ranean shrublands ( <i>Cistus ladanifer</i> )	–	Expansion of <i>Pinus sylvestris</i> and reduction in <i>Quercus suber</i> and Mediterranean shrublands	–	Recovery of <i>Quercus suber</i> and deciduous <i>Quercus</i> forests in mixed formations with conifers
Gerber 42°38' N 1°60' E 2200–2350 m a.s.l.	Dendro-climatic	–	–	Lower temperatures	–	Higher temperatures	–
Sobreestivo 42°41' N 0°06' E 2300–2400 m a.s.l.	Dendro-climatic	–	–	High, variable rainfall	–	Relatively high, stable rainfall	–

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.



**Fig. 1.** (A) Location of the different records reviewed in this paper within the Pyrenees, including lacustrine, dendroclimatic and marine-littoral reconstructions (see legend). (B) Detailed map of the different massifs with the presence of glaciers during the LIA (modified from González-Trueba et al., 2008).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 2.** Selected pollen taxa and group diagrams from lakes Arreo, Basa de la Mora, Estanya and Portlligat and correlation between them. In Arreo sequence Mesophytes curve is composed by deciduous *Quercus*, *Fagus* and *Corylus*; Heliophytes group by *Artemisia*, *Chenopodiaceae*, *Asteraceae*; and Hydro-hygrophytes includes *Cyperaceae* and *Typha*. In Basa de la Mora, Heliophytes group is formed by *Ephedra distachya*, *E. fragilis*, *Artemisia*, *Cichorioideae*, *Asteroidae*, *Carduaceae*, *Chenopodiaceae*, *Plantago*, and *Helianthemum*; whereas Mesophytes comprise: deciduous *Quercus*, *Alnus*, *Carpinus*, *Salix*, *Ulmus*, *Populus*, *Fraxinus* and *Juglans*. Estanya groups of Mesophytes and Hydro-hygrophytes are composed by *Betula*, *Corylus*, *Alnus*, *Tilia*, *Ulmus*, *Populus*, *Salix*, *Fagus*, and deciduous *Quercus*; and *Cyperaceae*, *Typha*, *Potamogeton*, *Myriophyllum*, *Nuphar* and *Lemna*, respectively. Finally, Portlligat hydro-hygrophytes curve includes: *Cyperaceae*, *Myriophyllum*, *Asplenium*, *Pteridophyta monolete* and *Pteridophyta trilete*.

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. (Caption on next page.)

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

## A multi-proxy perspective on millennium-long climate variability

M. Morellón et al.

**Fig. 3. (A)** Selected records from the Southern Pyrenees reviewed in this paper, from top to bottom: Capdella tree-ring based mean annual rainfall (30× moving average) (Saz Sánchez, 2003), GER-SOB summer temperature tree-ring based reconstruction (original data and 20× moving average) (Büntgen et al., 2008), diatom alkalinity-based summer-autumn temperature and chrysophyte-based winter-spring temperatures reconstruction in Lake Redon (Pla and Catalán, 2005; Catalan et al., 2009), phases of advance and retreat of the Pyrenean glaciers (González Trueba et al., 2008), calcite sublayering (negative values correspond to fine-coarse sequences and vice-versa for positive values) (Corella Aznar, 2011) and diatom C : P ratio (Scussolini et al., 2011) in Lake Montcortès, Si (cps) content in Lake Basa de la Mora (original data and 30× moving average) (Moreno et al., 2011), Sr (cps) content in Lake Arreo sequence (original data and 30× moving average) (Corella Aznar, 2011) and XRF-based salinity reconstruction and humid/arid phases in Lake Estanya (Morellón et al., 2011). **(B)** Supplementary regional and global records, from top to bottom: NH temperature reconstruction (Mann and Jones, 2003), solar irradiance (Steinhilber et al., 2009), and NAOms reconstruction (Trouet et al., 2009). Vertical yellow bars represent the chronology of the grand sunspot minima and temporal divisions Medieval Climate Anomaly (MCA), Little Ice Age (LIA) and Industrial Era (IND. ERA) are also indicated at the uppermost part of the figure.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion