

Interactive comment on “Glacial CO₂ cycle as a succession of key physical and biogeochemical processes” by V. Brovkin et al.

Anonymous Referee #1

Received and published: 27 June 2011

The manuscript entitled “Glacial CO₂ cycle as a succession of key physical and biogeochemical processes” by Brovkin et al. Simulates changes in the carbon cycle over the last glacial inception and deglaciation with a model of intermediate complexity CLIMBER 2. The model is forced with changes in orbital parameters and radiative forcing due to CO₂, CH₄ and N₂O. The model is coupled to an ice-sheet model which simulates about 110m lower sea level at the LGM. The processes thus taken into account in the study are temperature, salinity, circulation changes as well as responses of the ocean-sediment system, iron fertilization and land carbon uptake/release.

It is an interesting study, worth publishing in Climate of the Past, as transient simulations of the carbon system since the last interglacial have not been really performed before and can provide additional information on the behavior of the marine carbon

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



cycle.

However I think that some elements are missing for a good understanding of the carbon system in these transient simulations. Mainly, land carbon changes and the weathering variations imposed in these simulations really have to be described in more details if not shown for the paper to be publishable (see below).

1) CLIMBER-2 comprises a simple vegetation model and the carbon changes in this terrestrial model are taken into account in experiment PCBL. However it is just mentioned in the text that the land carbon stock at pre-industrial times is lower (-200 GtC) than at the last interglacial. The total glacial/interglacial change in carbon stock is never mentioned. As land carbon changes have a significant impact on atmospheric CO₂ and the marine carbon system (d¹³C_{dic}), the authors should give more information about the land carbon changes and eventually show the evolution on a plot. In addition a difference of 200GtC between 125ka B.P. And 0 ka B.P. seems quite large, the authors could comment on that.

2)The model is apparently forced by changes in weathering. These changes in weathering are scaled by the changes in runoff. Weathering changes can have a non negligible effect on the marine carbon system. It is therefore also important to at least state in the text if not show the exact weathering changes applied: the direction, the amplitude... also did silicate and carbonate weathering changes equally? I know that G/I/G weathering changes are very uncertain and that estimates of carbonate weathering changes go in all direction. I do not contest how it is implemented in the model but I think that the reader really needs to have more details on this.

3)You briefly mention changes in export production in the text. Again it might be nice to have a % change at the LGM if not a timeseries of XP changes for simulation PCBL. You could also mention the difference in export production simulated with and without iron fertilization.

4)I appreciate the separation of factors and the fact that the 4 simulations (P, PC, PCB

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



& PCBL) are shown in figure 2. However for clarity, it would be nice if the different contributions of temperature, salinity, circulation, CaCO₃ compensation, iron fertilization to the glacial CO₂ drawdown were clearly stated in the text. In addition, the authors could comment on the impact of sediment processes on atmospheric CO₂.

5) On fig. 4. the [CO₃] changes obtained are quite high. The authors go quite fast at discussing their results compared to paleoproxies. In the Equatorial Pacific, Yu core GGC48 displays in agreement with proxies about 15 μmol/L change. However core GGC15 displays no change. How are the changes at high latitude? Higher or smaller than the ones showed for 30S:30N? I would expect the Southern Ocean changes to be even higher. Rickaby 2010 find a ~45 μmol/L changes in the Weddell Sea, which is usually described as a “upper limit” change (Zeebe and Marchitto, 2010, Nat. Geosciences). In addition p1777, L4-7 do not seem correct. As seen on fig 4. [CO₃] in both the Atlantic and Pacific decrease the deglaciation. In both the Atlantic and Pacific there is high [CO₃] content at 10 ka B.P.

6) About millennial-scale changes in pCO₂ You mention L27 that the rise in pCO₂ (10–20 ppmv) during millennial-scale AMOC shut down is mainly due to DIC decrease between 1 and 3 km in the Indo-Pacific region. Is it over the whole region or centered into a more specific area? You suggest this is due to the weakening of the reverse cell of the Indo-Pacific overturning circulation by 2 Sv. First I am a little surprised that just a 2 Sv change leads to a 10 ppm CO₂ rise. Then could you please precise which water mass you are talking about. Are you saying that the AAIW weakens by 2 Sv? L. 6, p1782, you could also cite Obata et al 2007 (j. Clim, 20)

Minor point: fig1 a is not so useful, changes in land carbon and weathering would be much more informative.

Interactive comment on Clim. Past Discuss., 7, 1767, 2011.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)