

Moisture-source
temperature
estimated from ice
core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Ranges of moisture-source temperatures estimated from Antarctic ice core stable isotope records over the glacial-interglacial cycles

R. Uemura^{1,*}, V. Masson-Delmotte¹, J. Jouzel¹, A. Landais¹, H. Motoyama², and B. Stenni³

¹IPSL-Laboratoire des Sciences du Climat et l'Environnement (LSCE/IPSL), UMR8212, CEA-CNRS-UVSQ, 91191, France

²National Institute of Polar Research, Research Organization of Information and Systems, Tachikawa, Tokyo 190-8518, Japan

³Department of Mathematics and Geosciences, University of Trieste, via E. Weiss 2, 34128, Trieste, Italy

*now at: Department Chemistry, Biology and Marine Science, Faculty of Science, University of the Ryukyus, 1 Senbaru, Nishihara, Okinawa, 903-0213, Japan

Received: 27 December 2011 – Accepted: 17 January 2012 – Published: 30 January 2012

Correspondence to: R. Uemura (ruemura@sci.u-ryukyu.ac.jp)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

A single isotope ratio (δD or $\delta^{18}\text{O}$) of water is widely used as an air-temperature proxy in Antarctic ice cores. These isotope ratios, however, do not solely depend on air-temperature but also on the extent of distillation of heavy isotopes out of atmospheric water vapor from an oceanic moisture source to a precipitation site. The temperature changes at the oceanic moisture source (ΔT_{source}) and at the precipitation site (ΔT_{site}) can be retrieved by using deuterium-excess (d) data. A new d record from Dome Fuji, Antarctica is produced spanning the past 360 000 yr and compared with records from Vostok and EPICA Dome C ice cores. To retrieve ΔT_{source} and ΔT_{site} information, different linear regression equations have been proposed using theoretical isotope distillation models. A major source of uncertainty lies in the coefficient of regression, β_{site} which is related to the sensitivity of d to ΔT_{site} . We show that different ranges of temperature and selections of isotopic model outputs may increase the value of β_{site} by a factor of two. To explore the impacts of this coefficient on the reconstructed temperatures, we apply for the first time the exact same methodology to the isotope records from the three Antarctica ice cores. We show that uncertainties in the β_{site} coefficient strongly affect (i) the glacial-interglacial magnitude of ΔT_{source} ; (ii) the imprint of obliquity in ΔT_{source} and in the site-source temperature gradient. By contrast, we highlight the robustness of ΔT_{site} reconstruction using water isotopes records.

1 Introduction

Climate records preserved in Antarctic ice cores provide an important clue for the climate history over the several hundred thousand years. Among many deep ice cores from Antarctica, Vostok (Petit et al., 1999), EPICA Dome C (hereafter EDC) (EPICA-CommunityMembers, 2004; Jouzel et al., 2007) and Dome Fuji cores (hereafter DF) (Watanabe et al., 2003) have provided climate and environmental information covering multiple glacial-interglacial cycles. δD and $\delta^{18}\text{O}$ of water have been widely used

CPD

8, 391–434, 2012

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

as proxies for air-temperature, albeit with uncertainties in temperature reconstructions arising from possible changes in precipitation intermittency (Steig et al., 1994; Sime et al., 2009; Laepple et al., 2011), ice sheet elevation (Masson-Delmotte et al., 2011), or evaporation conditions (Boyle, 1997).

5 The latter source of uncertainty can be explored using a second-order isotopic information available through deuterium-excess ($d = \delta D - 8 \times \delta^{18}O$) (Dansgaard, 1964). The d parameter gives mainly access to processes affected by kinetic fractionation and has been used as a tracer of evaporation conditions such as ocean surface temperature and relative humidity in moisture source ocean (Merlivat and Jouzel, 1979).
10 The d value of modern atmospheric vapor above the Southern Ocean shows a clear dependency on sea surface temperature and relative humidity at the moisture source (Uemura et al., 2008). The initial information is then modified by subsequent meteorological processes (e.g. precipitation, snow formation, and cloud microphysics). The Antarctic precipitation d value is therefore also affected by kinetic fractionation during the snow formation, whose importance is expressed by a supersaturation function (Jouzel and Merlivat, 1984; Risi et al., 2010; Winkler et al., 2012), and by equilibrium distillation at very cold temperatures.

At first order, past variations of δD and d in ice cores can be assumed to mostly depend on polar temperature (ΔT_{site}), moisture source temperature (ΔT_{source}), and ocean isotopic composition ($\delta^{18}O_{\text{sw}}$). Source and site climatic information can therefore be extracted from ice core δD and d records, using a linear regression model based on Rayleigh distillation isotopic model and/or more sophisticated models (Vimeux et al., 2001; Cuffey and Vimeux, 2001; Stenni et al., 2001, 2010). This dual isotope approach allows us to correct site temperature estimate for moisture source effects. Such site and source temperature inversions have been conducted with different methodologies (different adjustments of the linear regression coefficients) for Vostok (Cuffey and Vimeux, 2001; Vimeux et al., 2002), EDC (Stenni et al., 2001, 2003, 2010), and EPICA Dronning Maud Land (EDML) (Stenni et al., 2010) ice cores. These studies have revealed the importance of obliquity in modulating Vostok source temperature

Moisture-source temperature estimated from ice core

R. Uemura et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(Vimeux et al., 2002). It has also been shown that source correction enhances the correlation between Vostok ΔT_{site} and atmospheric CO_2 during glacial inception (Cuffey and Vimeux, 2001), and that glacial millennial Antarctic Isotopic Maxima largely reflect changes in site temperature (Stenni et al., 2003). However, there is no well established methodology to reconstruct site and source temperatures.

Indeed, it is not possible to assess the full range of unknown source parameters (source temperature, relative humidity, wind speed) with only two isotope measurements (δD and $\delta^{18}\text{O}$) (Stenni et al., 2010). To overcome this problem, different approaches have been deployed to account for the ocean isotope changes and relative humidity effect (Vimeux et al., 2001; Cuffey and Vimeux, 2001; Stenni et al., 2001, 2003). The largest difference between these studies lies in the coefficient describing the sensitivity of d to ΔT_{site} , hereafter β_{site} ($= -\partial d / \partial T_{\text{site}}$), for which different studies have used values ranging from ~ 0.5 (Cuffey and Vimeux, 2001; Stenni et al., 2001) to 1.2 (Salamatin et al., 2004). In this manuscript, we will review the causes for these differences and their impacts on ΔT_{source} and ΔT_{site} reconstructions for the three central East Antarctic ice cores using published isotope profiles from Vostok and EDC, and a new deuterium excess record from DF.

This manuscript is organized as follows: ice-core data and isotope modeling are described in Sect. 2 while causes for the β_{site} differences and their impacts on ΔT_{source} and ΔT_{site} are discussed in Sect. 3 and site and source reconstructions in Sect. 4.

2 Isotope data and modeling

2.1 Ice core isotopic data

We used the three deep ice cores drilled in East Antarctica, DF, Vostok and EDC. The DF data were extended and revised in this paper. Two ice cores were drilled at the DF station ($77^{\circ}19' \text{S}$, $39^{\circ}42' \text{S}$, elevation 3810 m). The first 2503 m-depth core (DF1) was drilled in 1996, and revealed the $\delta^{18}\text{O}$ and δD records back to ~ 340 kyr

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



BP (thousands of years before present) (Watanabe et al., 2003; Uemura et al., 2004). Then, the other core (Dome Fuji 2nd, DF2) reached near bedrock in 2007 at the depth of 3035.2 m (Motoyama, 2007), and its $\delta^{18}\text{O}$ record from a ~ 130 m-length (2399.5–2531.5 m depth) core was published (Uemura et al., 2010). Here, we show δD and d records from this ~ 130 m-length part of the DF2 core at 10 cm-length sampling from every 50 cm ($n = 265$). The δD data from the overlapping part (2400–2500 m) shows remarkable similarity between DF1 and DF2 (Fig. 1).

The DF isotope ratios were measured using an equilibration method (Uemura et al., 2007) at National Institute of Polar Research, Japan. Routine measurements (2006 to 2008) of a reference water sample show that reproducibilities (1σ) of this water are 0.3‰ for δD , 0.04‰ for $\delta^{18}\text{O}$, and 0.4‰ for d . Duplicate analyses for 119 ice core samples confirm these estimates. In order to obtain the revised d record for the entire DF core, a new data set was made by 1) subtracting artificial isotope offsets from DF1 record (see below) and 2) replacing the DF1 data by DF2 data for overlapping part (2400–2503 m). First, there are offsets between DF1 and DF2 ($\delta_{\text{DF1}} - \delta_{\text{DF2}} = 0.8\text{‰}$ for δD , and 0.24‰ for $\delta^{18}\text{O}$). Re-measurement of DF1 δ values, based on ~ 60 samples re-cut from the archived DF1 core, were consistent with that of DF2 data, but larger than that of old DF1 data. Therefore, the gap could be caused by problems in the storage of the DF1 samples preserved in glass vials as liquid. The offsets of δ values were subtracted from the published data (Watanabe et al., 2003; Uemura et al., 2004). Second, an enlarged view of the records clearly shows that the d data are less noisy for DF2 than for DF1, suggesting that the precision of the new data is improved by replicate measurements of many ($\sim 30\%$) samples (Fig. 1). Thus, we replaced DF1 record by DF2 for the overlapping period.

For Vostok (78°30' S, 106°54' E, elevation 3488 m), the d data are available back to ~ 420 kyr BP (Vimeux et al., 2001). We restricted the ice core comparison over the past 250 kyr because the deeper part of the data is likely affected by changes in ice origin due to ice flow (Vimeux et al., 2001). For the EDC core (75°06' S, 123°23' E, elevation 3233 m), we used the recently published record spanning over 140 kyr BP (Stenni et al.,

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2010). Another d record from EDML (75°00' S, 00°09' E, elevation 2892 m) is also available for the same period but was not considered here because of uncertainties in elevation/ice origin impact (Stenni et al., 2010).

Precision of $\delta^{18}\text{O}$ and δD measurements of both the Vostok and EDC cores are 0.05‰ and 0.5‰ respectively (Vimeux et al., 2001; Stenni et al., 2010) with a final precision of 0.6‰ on d . Therefore, the isotopic precisions of the three ice cores are similar. The temporal-resolution of the data sets, however, differs one from the other. The average temporal-resolution of the DF data is ~ 270 yr, except for the ~ 70 yr resolution during the Holocene. For Vostok, the average temporal resolution is smaller than ~ 220 yr for our period of interest (Vimeux et al., 2001). The EDC core shows the highest temporal resolution; ~ 18 yr for the Holocene, ~ 50 yr for MIS2 and ~ 42 yr for MIS5.5 (Stenni et al., 2010).

The comparison of the ice core isotopic records first requires to place them on a common age scale. As shown in Fig. 2, significant differences (up to ~ 5 kyr) exist between the glaciological age scales established for Vostok (GT4 age-scale by Petit et al., 1999) and for EDC (EDC3 age-scale by Parrenin et al., 2007) and the O_2/N_2 orbital age scale for DF (Kawamura et al., 2007). Here, we use the DF age scale as a target and synchronized the Vostok, EDC and our new stacked DF record using visual δD matching (Fig. 2). We used 21 tie points for DF-Vostok fitting. The difference of the age-scales is ~ 1 kyr before 80 kyr BP, and increased up to ~ 5 kyr around 120 kyr BP. We used nine tie points for DF-EDC fitting. The difference of the age-scales is ~ 1 kyr before 80 kyr BP, and increased up to ~ 4 kyr for the older part. For the calculation of ΔT_{site} and ΔT_{source} , all the data were re-sampled with 200-yr resolution, which is under-sampling for EDC data, and slightly over-sampling for Vostok and DF data.

2.2 Raw d -excess and δD data

The raw d records from the DF, Vostok, and EDC ice cores show common low frequency variability (~ 40 kyr) (Fig. 2). This obliquity imprint is significantly smaller in DF than in the Vostok and EDC cores. In particular, minima of DF d record at about 20, 40,

Moisture-source temperature estimated from ice core

R. Uemura et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and 90 kyr BP are less distinct. Such difference, however, cannot be found in the older section (200–250 kyr BP) of DF and Vostok. Another difference can be found during the glacial terminations. From 20 to 10 kyr BP (Termination I), the onset of d increases in DF and Vostok appears to lag that of EDC. From 140 kyr to 130 kyr BP (Termination II), a similar lag is observed. It is difficult to interpret the raw d record because it is an empirically defined second-order parameter. In fact, a different definition of d parameter can produce different glacial interglacial variations (see Sect. 3 and Appendix A). Climatic interpretation of d will be discussed after inferring ΔT_{source} from isotopic data (Sect. 4).

2.3 Isotope inversion method

In order to retrieve the climatic information from the isotope ratios, we need to evaluate several coefficients using an isotope model. Sensitivities of δD and d to ΔT_{site} , ΔT_{source} , relative humidity (RH), and ocean isotopic composition ($\delta^{18}\text{O}_{\text{sw}}$) were estimated using a mixed cloud isotopic model (MCIM) developed by Ciais and Jouzel (1994). We used a linear inversion method which is very similar procedure performed for Vostok (Cuffey and Vimeux, 2001) and EDC records (Stenni et al., 2001, 2003, 2010; Masson-Delmotte et al., 2004).

There are several differences among the inversion methods used for various ice cores. First, the coefficients for the correction for $\delta^{18}\text{O}_{\text{sw}}$ were used as fixed parameters in several studies (Vimeux et al., 2001, 2002; Cuffey and Vimeux, 2001; Stenni et al., 2001). This coefficient, however, can be exactly calculated as a function of the isotopic composition of the ice (Vimeux et al., 2001; Jouzel et al., 2003; Kavanaugh and Cuffey, 2003), as follows:

$$\delta^{18}\text{O}_{\text{corr}} = \delta^{18}\text{O}_{\text{ice}} - \delta^{18}\text{O}_{\text{sw}} \frac{(1 + \delta^{18}\text{O}_{\text{ice}})}{(1 + \delta^{18}\text{O}_{\text{sw}})} \quad (1)$$

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$\delta D_{\text{corr}} = \delta D_{\text{ice}} - \delta D_{\text{SW}} \frac{(1 + \delta D_{\text{ice}})}{(1 + \delta D_{\text{SW}})} \quad (2)$$

Here, $\delta^{18}\text{O}_{\text{corr}}$ and δD_{corr} are the isotope ratios of the ice ($\delta^{18}\text{O}_{\text{ice}}$ and δD_{ice}) corrected for ocean isotope changes ($\delta^{18}\text{O}_{\text{sw}}$ and δD_{sw}). An estimate of the global ice volume component of sea-water $\delta^{18}\text{O}$ (Bintanja and Wal, 2008) was used for the $\delta^{18}\text{O}_{\text{sw}}$ record. The δD_{sw} can be calculated on the assumption that $\delta D_{\text{sw}} = 8 \times \delta^{18}\text{O}_{\text{sw}}$. Thus, the d value corrected for the ocean isotope effect (d_{corr}) can be obtained by: $d_{\text{corr}} = \delta D_{\text{corr}} - 8\delta^{18}\text{O}_{\text{corr}}$. Following recent studies (e.g. Stenni et al., 2010), we apply this correction before conducting the multiple linear regression.

Second, δD and d in polar precipitation preserve source effects and therefore depends on climatic parameters affecting evaporation kinetics, including ΔT_{source} but also relative humidity (RH) and wind-speed regimes (Merlivat and Jouzel, 1979; Petit et al., 1991). For Vostok, the RH effect was explicitly included in the ΔT_{source} by using a linear relation found in outputs of Atmospheric General Circulation Models ($\Delta h \sim -0.4\Delta T_{\text{source}}$) (Vimeux et al., 2001; Cuffey and Vimeux, 2001). For EDC, the RH effect was not included, restricting the analyses of source effects to ΔT_{source} only (Stenni et al., 2001). Here, we adopted the latter approach for the sake of simplicity. Thus, the results of multiple linear regressions can be expressed as:

$$\Delta \delta D_{\text{corr}} = \gamma_{\text{site}} \Delta T_{\text{site}} - \gamma_{\text{source}} \Delta T_{\text{source}} \quad (3)$$

$$\delta d_{\text{corr}} = -\beta_{\text{site}} \Delta T_{\text{site}} + \beta_{\text{source}} \Delta T_{\text{source}} \quad (4)$$

All the γ and β coefficients are positive numbers expressed in ‰K^{-1} . The Δ represents deviations from the present reference values (in this study “present” is defined as the averaged value of the past 2 kyr). The ΔT_{site} , mean annual temperature at ground level, is calculated from the mean annual temperature of condensation (ΔT_{c}) by assuming a constant relation ($\Delta T_{\text{c}} = 0.67\Delta T_{\text{site}}$) (Jouzel and Merlivat, 1984). This is of course an approximation as Antarctic inversion strength varies through places, time and weather conditions. A detailed analysis of condensation temperature conducted

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



using half-century-long meteorological observations from Vostok (Ekaykin, 2003) recently confirmed the validity of this approach for this site.

Finally, a kinetic isotope fractionation occurs during ice crystal growth because of super-saturation over ice, parameterized as a linear function of temperature and tuned to reproduce the spatial d vs. δ distribution in Antarctica (Petit et al., 1991). We used a linear function ($S_i = 1.020 - 0.0037T$) which best reproduces the present-day surface snow d vs. δD distribution from various parts of Antarctica (Masson-Delmotte et al., 2008), and from the Japanese-Swedish IPY 2007/2008 traverse (Fujita et al., 2011) covering the Syowa-Dome F route (Fig. 3). In this study, all the simulations were conducted using this fixed super super-saturation function.

2.4 Present-day isotope tuning

The first step for estimating the sensitivity coefficients is to fit the model output to the present-day snow isotopic distribution (Fig. 3). An important problem of this tuning is the difficulty to simulate present-day snow isotope ratios by using the MCIM. At the DF site, the present-day (past 2 kyr average $\pm 1\sigma$) isotope ratios are $\delta^{18}\text{O} = -55.09 \pm 0.16\text{‰}$, $\delta D = -426.3 \pm 1.4\text{‰}$, and $d = 14.4 \pm 0.3\text{‰}$. With our MCIM tuning, these isotope ratios can only be reached with a very cold site temperature, -61.0°C , which is significantly lower than the 10 m-depth temperature of -57.7°C (Motoyama et al., 2005) and 1-yr observations at 4 m-height of -54.8°C (Fujita and Abe, 2006). The difficulties to simulate present-day δ values might also reflect the difference between annual mean temperature and precipitation weighted temperature (e.g. Masson-Delmotte et al., 2011), and/or differences in prescribed surface air-pressures. In fact, the air pressure of DF is the lowest among the coring sites since its elevation is the highest (3810 m a.s.l.) compared with the VK (3488 m a.s.l.) and EDC (3233 m a.s.l.). Lower air-pressure results in smaller distillation in the model, and thus produces higher simulated δ values.

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MCIM to simulate present-day snow isotope ratios (“Isotope fitted”), as described in Sect. 2.4. Two additional simulations were conducted to reproduce the present-day temperature (-55°C). The first case used $T_{\text{source}} = 18^{\circ}\text{C}$ (“Temperature fitted 1”) and the second case used T_{source} as 11°C (“Temperature fitted 2”). The latter T_{source} value was estimated by GCM (Delaygue et al., 2000) and adopted for Vostok reconstruction (Vimeux et al., 2001). With these hypotheses on moisture source, the simulated isotope ratios were systematically higher than the values observed in the DF ice core. This prevents us from selecting the model results compatible with the DF ice core data (see Sect. 3.2). Instead, we performed sensitivity tests by varying ΔT_{site} and ΔT_{source} by 2°C , and performed multiple regressions on all the data.

Table 2 clearly shows that β_{site} decreases as the simulated δD increases. Changes in β_{site} are the largest ($\sim 130\%$) compared to the smaller changes in the other coefficients (from -20% to 30%). This can be explained by the increasing d - δD slope simulated in response to δD depletion. Figures 3 and 4 illustrate the non-stability of the linear slope between d and δD .

Note that the β_{site} estimation (0.6) of “Temperature fitted 2” is comparable to the values (0.5) used in previous ice core reconstructions. Thus, we conclude that most of the discrepancy found in previous studies can be explained by the choices of different methodologies: whether the model is tuned to reproduce the observed ice core isotope ratios (Salamatin et al., 2004), or is driven by realistic air-temperatures (Cuffey and Vimeux, 2001; Stenni et al., 2001). Our approach accounting for large slopes at cold temperatures allows us to explore this source of uncertainty on reconstructing site and source temperatures.

3.2 Data selection for regressions

Another methodology that affects the β_{site} value is the data selection criteria used for establishing the multiple linear regressions coefficients. In previous studies, the regression analyses were performed on the simulated δ values in response to a wide range of randomly changed climate conditions (ΔT_{site} , ΔT_{source}) (e.g. Masson-Delmotte et al.,

2004). The large range of results obtained for DF with this approach is illustrated in Fig. 4 as open circles.

In contrast, we propose now to restrict the range of site and source conditions for which the modeled isotopic results are consistent with the glacial-interglacial ice core data. Note that our model results account for sea water isotopic composition corrections; this is a marginal source of uncertainty in the model-data comparison. We therefore selected a subset of the modelled isotope values (and corresponding ΔT_{site} and ΔT_{source}) displayed as black circles in Fig. 4, and performed the multiple linear regression analysis on this dataset. We illustrate in Fig. 4 the importance of this data selection for the d - δD slopes, which increase with isotopic depletion and temperature ranges. Generally, excluding higher δD data results in larger d - δD slope and thus larger β_{site} coefficient. Our approach, thus, will result in larger β_{site} coefficient, compared to the previous approach.

These results suggest that our methodology, using the MCIM model with deliberately cold conditions and selected data, should be seen as an upper end approach for the β_{site} coefficient, compared to previous approaches which did not sufficiently account for the large slope at cold temperatures.

3.3 Different sensitivity coefficients

The isotope inversion was performed to retrieve the ΔT_{site} and ΔT_{source} information using the method described above ($S_i = 1.020 - 0.003T$, and present T_{site} and simulated δ values are shown as *isotope fitted* in Table 1). The linear relations obtained for the DF ice core are:

$$\Delta \delta D_{\text{corr}} = 7.7 \Delta T_{\text{site}} - 3.2 \Delta T_{\text{source}} (R^2 = 0.999, n = 14, p < 0.001), \quad (5)$$

$$\delta d_{\text{corr}} = -1.3 \Delta T_{\text{site}} + 1.6 \Delta T_{\text{source}} (R^2 = 0.989, n = 14, p < 0.001). \quad (6)$$

The linear correlations are strong, thus the statistical uncertainty of each coefficient due to multiple regression analysis is very small (0.03 to 0.05). The uncertainty caused by the data selection for the regression analysis (Sect. 3.2) is also small. In fact, if we

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



remove several (1 to 5) points located around the “corner” of data subset (Fig. 4), the uncertainties of the linear coefficients are 0.01 to 0.04. Such statistical regression error analysis, however, does not represent the total uncertainties on the estimated ΔT_{site} and ΔT_{source} . It does not account for uncertainties on MCIM parameterization tunings (e.g. supersaturation), and on climate parameters not taken into account (precipitation intermittency, inversion strength, changes in wind speed and/or relative humidity. For example, Winkler et al. (2011) show that the β_{site} depends on the tuning of supersaturation, and an error range was estimated as 1.29 to 2.04.

We have also applied the exact same methodology for EDC and Vostok, using the same tuning of MCIM and performing the linear analyses on the range of model outputs consistent with each ice core isotopic dataset. Table 1 compares our results with previous regressions. Among four coefficients in the equations, only β_{site} is significantly different from previous estimate, which is now 2.5 times larger than previous one (~ 0.5). As discussed in Sects. 3.1 and 3.2, differences found in β_{site} between previous studies can be explained by the different methodologies: whether the model is tuned to reproduce the isotope ratios or air-temperature.

The value of β_{site} , which seems high for East Antarctica sites, is actually very dependent on the d definitions and on the range of temperatures studied. To understand the influence of the historical linear definition of d on the β_{site} , we tested a logarithmic definition of d (see Appendix A for detail). The results confirm that the value of β_{site} significantly depends on the definition of d . A logarithmic definition can remove the influence of ΔT_{site} on d spatial variations, and thus significantly reduce the value of β_{site} . But this new definition does not change anything to the temperature reconstructions, because β_{site} act as a correction factor for the empirically defined d parameter.

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Reconstructed ΔT source and ΔT site

4.1 ΔT_{source} and ΔT_{site} of Dome F

The new coefficients obtained in this study (Eqs. 5 and 6) were applied to the DF ice core data to estimate past variations in ΔT_{source} and ΔT_{site} (Fig. 5). The new ΔT_{source} vary between -5.9°C to 2.9°C over the past 360 kyr. The record shows maxima of ΔT_{source} occurring during Marine Isotope Stage (MIS) 5e and MIS 9e with values respectively $2.6 \pm 0.3^{\circ}\text{C}$ and $2.9 \pm 0.3^{\circ}\text{C}$ higher than present. A glacial minimum at 19 kyr BP is $-5.9 \pm 0.3^{\circ}\text{C}$. Here, temperatures are based on the average and $\pm 1\sigma$ of 1 kyr duration around the maximum or minimum point. This ΔT_{source} reconstruction remains high when ΔT_{site} and δD drop during glacial inceptions, linked with high d values during the onset of glaciations. The DF ΔT_{site} derived from the new coefficients ranges between $-7.8 \pm 0.2^{\circ}\text{C}$ to $4.8 \pm 0.5^{\circ}\text{C}$ over the past 360 kyr. Maximum ΔT_{site} occur during MIS 9e, with temperature $4.8 \pm 0.5^{\circ}\text{C}$ warmer than that of present-day average.

In order to show the importance of the value of β_{site} , other ΔT_{source} and ΔT_{site} records which were obtained using the smaller values of β_{site} (0.5) and other coefficients used in Cuffey and Vimeux (2001) (Table 1) are shown in Fig. 5a. For clarity, we labeled this ΔT_{source} and ΔT_{site} records based on the smaller β_{site} coefficient as $\Delta T_{\text{source}}^{\text{low}}$ and $\Delta T_{\text{site}}^{\text{low}}$, respectively. Two major differences are identified for the source temperature records. First, the glacial-interglacial amplitude of the new ΔT_{source} is approximately twice larger than that of $\Delta T_{\text{source}}^{\text{low}}$. Second, the timing of the interglacial-glacial inception maxima is strongly affected: small β_{site} coefficients induce a maximum $\Delta T_{\text{source}}^{\text{low}}$ at the onset of glaciations (MIS 5d), while large β_{site} coefficients place ΔT_{source} maxima in phase with Antarctic warmth (MIS 5e). With the exception of glacial inceptions, the new ΔT_{source} reconstruction is also much more similar to ΔT_{site} .

In contrast to the sensitivity of ΔT_{source} to the uncertainties in regression coefficients, the ΔT_{site} shows much less sensitivity (Fig. 5c). Indeed, the difference between ΔT_{site} and $\Delta T_{\text{site}}^{\text{low}}$ is $\sim 0.6^{\circ}\text{C}$, which is within typical error (-10 – 30%) of temperature

CPD

8, 391–434, 2012

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



estimations (Jouzel et al., 2003). The previously published DF ΔT_{site} (which corresponds to $\Delta T_{\text{site}}^{\text{low}}$) estimated MIS 9e to be warmer than the present by $4.0 \pm 0.3^\circ\text{C}$ (Kawamura et al., 2007) (Fig. 5c). Our revised estimate lies above this previous value, as it produces a MIS 9e ΔT_{site} estimate of $4.8 \pm 0.5^\circ\text{C}$. For MIS 5e, previous and our DF ΔT_{site} estimates are $3.5 \pm 0.1^\circ\text{C}$ and $3.7 \pm 0.4^\circ\text{C}$, respectively. The ΔT_{site} records from six Antarctic ice cores also show similar degree of warming during interglacial periods (Masson-Delmotte et al., 2011). A GCM study, based on increased CO_2 projections, however suggested that MIS 5e isotope temperature could be 6 to 10°C higher than that of present day (Sime et al., 2009). This estimation is still significantly larger than our reconstruction. Thus, these results suggest that moisture source effects cannot explain the model-data discrepancy.

4.2 Comparison with the other cores

We applied the same procedure as DF for Vostok and EDC data, and obtained the ΔT_{site} and ΔT_{source} records using the coefficients obtained in this study (Table 1). Figure 6 shows the ΔT_{source} , ΔT_{site} , and the temperature gradient records.

The glacial-interglacial changes in ΔT_{source} are again significantly larger than the estimates using previous MCIM tuning. For Vostok, the new ΔT_{source} decreases to about $-6.0 \pm 0.1^\circ\text{C}$ around 19 kyr BP (temperatures are based on the 1 kyr average $\pm 1\sigma$), which is lower than the previous estimation of $\sim -4^\circ\text{C}$ (Vimeux et al., 2002). For the EDC, the ΔT_{source} decrease to about $-8.1 \pm 0.3^\circ\text{C}$ at 19 kyr BP, which is much lower than the previous estimation of -3 to -4°C (Stenni et al., 2010). This large EDC ΔT_{source} variation results from a larger β_{site} value, which means that a larger proportion of δD is added to the raw d record.

The glacial-interglacial changes in ΔT_{site} are close to the previous estimations. The Vostok ΔT_{site} estimations based on this study decreased to about -7.1°C at 19 kyr BP and reached the minimum at -7.8°C at 21.7 kyr BP. These values are almost the same as the previous estimation (Vimeux et al., 2002). For the EDC, our ΔT_{site} estimate decreases to about $-9.9 \pm 0.2^\circ\text{C}$ at 19 kyr BP, slightly larger than the previous estimate

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of -8 to -9 °C at 20 kyr BP without accounting for elevation corrections (Stenni et al., 2003, 2010). The larger ΔT_{site} results from combination of relatively larger variation in δD and larger γ_{site} values. In summary, as for DF case, the new ΔT_{site} estimations of Vostok and EDC cores are consistent with the previous estimations within typical error (-10 – 30 %) (Jouzel et al., 2003). Interestingly, at glacial inceptions, all the ΔT_{site} records show warmer value than classical δD -temperature estimation, and thus the co-variation of ΔT_{site} and CO_2 found in Vostok core (Cuffey and Vimeux, 2001) is generally supported.

The temperature gradients ($\Delta T_{\text{source}} - \Delta T_{\text{site}}$) of DF, EDC and Vostok show a strong 40-kyr cycle (Fig. 6). Such marked 40-kyr periodicity has not been found in previous reconstructions of temperature gradient at Vostok (Vimeux et al., 2002) and EDC (Stenni et al., 2010) as the gradient was strongly correlated with ΔT_{site} ($r^2 = 0.9$) (Stenni et al., 2010). The glacial-interglacial variations of the gradient of this study ranges from -2 °C to 4 °C. These variations are smaller than the previous ones by ~ 2 to 3 °C (~ 40 % smaller). The amplitude of temperature gradient ranged from -3 °C to 6 °C at Vostok (Vimeux et al., 2002), and -2 °C to 6 °C at EDC (Stenni et al., 2010). The gradient record of Vostok shows a gradual decreasing trend from past to present (i.e. the value of minima decreases), suggesting that the long-term trend might be affected by ice flow.

4.3 Comparison with SST records from ocean sediments

Which (low versus high coefficient) approach is most consistent with independent information? Here, we take advantage of sea surface temperature (SST) estimates produced thanks to biological and geochemical analyses of deep sea sediments, and compare our reconstructed ΔT_{source} with SST estimates. We are aware that different marine proxies can lead to significant uncertainties in reconstructions, which can also have seasonal biases (e.g. MARGO Project Members, 2009). The recent synthesis of LGM SST data depicts a latitudinal temperature gradient in the Southern Hemisphere oceans. Backward trajectory models suggests that the present-day main

Moisture-source temperature estimated from ice core

R. Uemura et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

moisture source for DF is located around 40–70° S ocean (Reijmer et al., 2002; Sode-
mann and Stohl, 2009). In the 42.5° S–52.5° S latitudinal band, the LGM minus present-
day anomaly is -3.8°C (MARGO Project Members, 2009). Figure 5b shows a stacked
Southern Ocean mid-latitude SST record (Barrows et al., 2007) that was made by com-
bining the ocean sediment records from the MD97-2120 (45°32' S, 174°56' E) from east
of New Zealand (Pahnke and Sachs, 2006) and MD88-770 (46°01' S, 96°27' E) from the
Indian Ocean (Labeyrie et al., 1996). The stacked SST record shows a relatively large
SST shift ($\sim 5^{\circ}\text{C}$) during Termination I. Another sediment (MD07-3076) retrieved in the
Atlantic sector of the Southern Ocean (44°09' S, 14°13' W) shows a glacial-interglacial
temperature change of $\sim 7^{\circ}\text{C}$ both during Termination I (Skinner et al., 2010) and Ter-
mination V (Vázquez Riveiros et al., 2010).

Our ΔT_{source} variation based on isotope-fitting is therefore only compatible with the
largest SST amplitude estimates. However, the moisture source likely moves between
glacial and interglacial periods, as suggested by water tagging in atmospheric models
(Delaygue et al., 2000) or abrupt shifts of d (Masson-Delmotte et al., 2010a). Dur-
ing glacial periods, northward shifts of the mean evaporation locations are expected
to compensate the impact of SST cooling into ΔT_{source} . While associated with un-
certainties on the actual moisture sources and the proxy-based SST reconstructions,
this comparison suggests that our large ΔT_{source} reconstruction is at the upper limit of
compatibility with SST data. Note that there are several limitations for the ΔT_{source} re-
construction caused by uncertainties on supersaturation and precipitation intermittency
in the past different climate conditions.

4.4 Obliquity and deuterium-excess

A strong 41-kyr Earth's obliquity periodicity is detected in many paleoclimatic SST
records, in both hemispheres (Cortijo et al., 1999; Sachs et al., 2001) including ice
cores (Jouzel et al., 2007). Based on these paleoclimatic evidence, obliquity was sug-
gested to be a key driver of glacial-interglacial dynamics (Vimeux et al., 1999, 2001;

CPD

8, 391–434, 2012

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Khodri et al., 2001; Raymo and Nisancioglu, 2003; Loutre et al., 2004; Huybers and Wunsch, 2005; Lee and Poulsen, 2008; Davis and Brewer, 2009).

The raw d record from Antarctic ice cores also shows a strong 41-kyr obliquity periodicity (Stenni et al., 2001, 2003, 2010; Vimeux et al., 1999; Uemura et al., 2004).

Previous studies suggested that the obliquity component of the Vostok d record could be related to the contrasting impact of obliquity on annual mean insolation and temperature at low and high latitudes (affecting SST), amplified by changes in the moisture transport induced by the annual insolation gradient between low and high latitudes (Vimeux et al., 1999; Loutre et al., 2004). In previous reconstructions, $\Delta T_{\text{source}}^{\text{low}}$ was very similar to the raw deuterium excess signals and showed a large obliquity imprint.

Interestingly, δD and d are anti-correlated in the 40-kyr band (Fig. 7). As a result, small differences in β_{site} have large impacts on the obliquity signal in ΔT_{source} . Spectral analysis shows that our new ΔT_{source} reconstruction indeed has weak obliquity signal (Fig. 8). It cannot therefore be ruled out that the strong obliquity signal found in raw d record of DF and Vostok cores is an artifact of site temperature. This does not apply to EDC where the obliquity imprint in the raw d is less clear.

As a result of our new reconstruction, spectral properties of our temperature gradient ($\Delta T_{\text{source}} - \Delta T_{\text{site}}$) from DF, Vostok and EDC cores appears mainly modulated by obliquity (Fig. 8) and without 100 kyr cycles. A recent modeling study (Antico et al., 2010) supports a large control of obliquity on latitudinal moisture transport.

Compared with δD , the strong imprint of obliquity is reduced in our ΔT_{site} reconstruction (Fig. 8), suggesting that it can be partly affected by moisture origin effects. The obliquity signal, however, remains in ΔT_{site} . Thus, the moisture source effect cannot fully explain the fact that climate models display a negligible impact of obliquity on the simulated annual mean Antarctic temperature, suggesting that it may result from model biases, such as a lack of austral sea ice in control simulations (Masson-Delmotte et al., 2010b).

Furthermore, our synchronized age scale is based on the imprint of precession on DF O_2/N_2 and is independent from orbital properties of the stable isotope records. In

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



our high β_{site} case, the site-source temperature gradient mainly exhibits obliquity periodicities, which is a requested characteristic for orbital tuning. Cross spectrum analysis shows that the mean lag of this reconstructed gradient behind the obliquity is 2.9 kyr. The gradients of EDC and VK also lag the obliquity by 2.9 and 3.0 kyr, respectively.

This differs from the observed 5 kyr lag between EDC δD and obliquity (Jouzel et al., 2007). The processes responsible for the observed lags remain elusive, and likely involved the interplay between ice sheets and ocean circulation. Further investigations of the mechanisms linking obliquity and site-source temperature gradient should be conducted, in order to use such information for orbital dating of ice cores. It will be important independent evidence to test the length of MIS 11, in which the accuracy of the EDC3 age scale (Parrenin et al., 2007) was questioned (Masson-Delmotte et al., 2010b; Pol et al., 2011).

5 Conclusions and perspective

We obtained a new d record from DF ice core spanning the past 360 000 yr and compared with records from Vostok and EDC ice cores. To interpret the d and δD records as ΔT_{source} and ΔT_{site} information, we used isotopic inversion based on the MCIM model as previously performed by Vimeux et al. (2002); Stenni et al. (2001) on the Vostok and EDC ice cores, respectively. A major source of uncertainty lies in the coefficient of regression, β_{site} which is related to the sensitivity of d to ΔT_{site} . We showed that different ranges of temperature and selections of isotopic model outputs may increase the value of β_{site} by a factor of two. The results suggests that the causes for the differences of β_{site} between the previous studies lie in the isotopic model tuning for the present-day, and the δD ranges selected for regression analyses because the β_{site} depends on the range of temperature or isotopic depletion considered.

For the first time, we applied the exact same temperature inversion methodology to the three Antarctica ice cores data set (DF, Vostok, and EDC). All ΔT_{source} records were very sensitive to the inversion method, with our new approach (isotope-fitting) leading

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

to amplitudes of ΔT_{source} variation $\sim 2^\circ\text{C}$ to 4°C larger compared to the previous estimations. For DF, the ΔT_{source} estimates yields values between -5.9°C to 2.9°C over the past 360 kyr, Comparisons with proxy-based SST records suggest that our large ΔT_{source} reconstruction is at the upper limit of compatibility with SST data. In contrast to the significant modification in ΔT_{source} , the ΔT_{site} records were not modified significantly and therefore appear robust. The differences between the current and previous studies are within published uncertainty (-10 – 30%) of ΔT_{site} estimations. The new ΔT_{source} estimates showed only weak 41-kyr obliquity periodicity, but the obliquity signal was found in the temperature gradient ($\Delta T_{\text{source}} - \Delta T_{\text{site}}$). The temperature gradient records of the three cores show a systematic anti-correlation with obliquity, providing new evidence for the importance of the Earth's obliquity as a driver of water cycle and climate dynamics. Such properties offer the perspective of using this obliquity imprint as a synchronization tool for dating ice cores.

The isotope inversion method involves simplification, and the procedure is challenging, in particular in Antarctica where isotope ratios are extremely depleted. Analyzing d at an event basis in modern condensing vapor and precipitation could be very useful, combined with moisture tracking analysis (Sodemann and Stohl, 2009) and isotopic modeling, to identify what are the correct or incorrect hypotheses and parameterizations in distillation models regarding the controls of d . Triple isotope ratios of oxygen (i.e. ^{17}O -excess) in water molecules (Landais et al., 2008; Luz and Barkan, 2010; Winkler et al., 2012) would also provide independent evidence for moisture source imprint on isotopically depleted snow. Finally, extensive field observation of d and ^{17}O -excess in clouds over Antarctica may provide better constraints on supersaturation function, and will allow to improve our understanding of the mechanism underlying the changes in d and ^{17}O -excess.

Logarithmic definition of d and meaning of β_{site}

The interplay of artifacts caused by the empirical d definition and by physical kinetic processes (evaporation and snow formation) restricts precise physical understanding of meaning to the value of two coefficients (β_{site} and β_{source}) in Eq. (4). The empirical definition (i.e. the constant slope of 8) of d itself induces changes in d because any process that does not follow the $\delta\text{D}/\delta^{18}\text{O}$ slope of 8 alters the d value. Indeed the $\delta\text{D}/\delta^{18}\text{O}$ slope can deviate from the mean value of 8 because of the combination of the nonlinearity of the ratio of equilibrium-fractionation factors and the crude approximation of $\ln(1 + \delta) \sim \delta$. In contrast to the historical linear definition of d , recent studies of triple oxygen isotope study of water have used the logarithmic framework for defining the ^{17}O -excess parameter (Barkan and Luz, 2007). Here, we examine how the historical definition of d affects the ice core d record and β_{site} by introducing a logarithmic based definition of deuterium excess.

In the normal δ plot, the slope of Global Meteoric Water Line (GMWL) apparently keeps a constant value “8” along with isotope distillation (Craig, 1961). However, it is difficult to understand the physical meaning of this constant slope because the ratio of equilibrium fractionation-factors of hydrogen and oxygen ($(\alpha_{\text{D}} - 1)/(\alpha_{18} - 1)$), which corresponds to the $\delta\text{D}/\delta^{18}\text{O}$ slope, increases from 8.7 to 9.6 when the temperature decreases from 20°C to 0°C (e.g. Jouzel, 1986). In principle, the increasing slope should be observed only in the $\log(1 + \delta)$ plot. This point was clearly described in the historic GMWL paper (Craig, 1961) as; “it is actually $\log(1 + \delta)$ which should be plotted for such a process ... The linear relation observed in Fig. 1 simply reflects a coincidence”. Recently, the limitation of normal δ diagrams is discussed in Miller (2002). In Antarctica, -50‰ and -400‰ in δ notation (either δD or $\delta^{18}\text{O}$) corresponds to -51.3‰ and -511‰ in $\log(1 + \delta)$ notation, respectively. Thus, $\ln(1 + \delta)$ cannot be approximated as δ .

**Moisture-source
temperature
estimated from ice
core**

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In order to show the differences of normal and logarithmic plot, global isotope data from Global Network of Isotopes in Precipitation (GNIP) database (e.g. Rozanski et al., 1993) and a recent compilation of Antarctic surface snow data (Masson-Delmotte et al., 2008) are shown in Fig. A1. Although, the GMWL appears to represent average isotopic composition, the slope is, in reality, non-linear. The d data shows parabolic curve against $\delta^{18}\text{O}$ with a minimum (\sim negative values in d) at $\sim -30\text{‰}$ in $\delta^{18}\text{O}$, and maxima ($\sim 15\text{--}20\text{‰}$ in d) at $\sim 0\text{‰}$ (equatorial) and $\sim -60\text{‰}$ in $\delta^{18}\text{O}$ (Antarctica) (Fig. A1a). Figure A1b shows the same data but on natural logarithmic plot. The data clearly show non-linear relationship between $\ln(1 + \delta\text{D})$ vs. $\ln(1 + \delta^{18}\text{O})$. A simple quadric fitting gives sufficiently good representation as:

$$\ln(1 + \delta\text{D}) = -2.85 \times 10^{-2} \times (\ln(1 + \delta^{18}\text{O}))^2 + 8.47 \times \ln(1 + \delta^{18}\text{O}) + 13.3 (R^2 = 0.999). \quad (\text{A1})$$

Unlike a constant slope of GMWL, the slope of this Global Meteoric Water Curve (GMWC) increases from ~ 8.5 to 12 as $\ln(1 + \delta^{18}\text{O})$ decreases. Figure A2 shows the slope and $\ln(1 + \delta^{18}\text{O})$ with corresponding annual-mean surface air-temperature. This slope generally agrees with the one expected from $(\alpha_{\text{D}} - 1)/(\alpha_{18} - 1)$. There is a clear gap between the transition of equilibrium fractionation factors for liquid-vapor and ice-vapor for ^{18}O (Majoube, 1971a, b) and for D (Merlivat and Nief, 1967). The gap was narrowed after incorporating the super saturation effect (Jouzel and Merlivat, 1984) (here, we used the S_i function described in Sect. 2.2 was used).

Based on the GMWC, a deuterium-excess parameter based on logarithmic δ , hereafter denoted as d_{ln} , is defined as:

$$d_{\text{ln}} = \ln(1 + \delta\text{D}) - (-2.85 \times 10^{-2} \times (\ln(1 + \delta^{18}\text{O}))^2 + 8.47 \times \ln(1 + \delta^{18}\text{O})). \quad (\text{A2})$$

In contrast to the d , which shows parabolic curve against $\delta^{18}\text{O}$, d_{ln} does not depend on $\ln(1 + \delta^{18}\text{O})$ (Fig. A1b). Note that d_{ln} is defined based on phenomenological data set. Thus, not only the equilibrium effect but also kinetic fractionation (like snow formation effect) is also eliminated.

Figure A3 shows the d_{ln} records of Antarctic ice cores. The d_{ln} ranges from about -15‰ to 5‰ . The glacial-interglacial amplitude is, thus, about two times larger than

the normal d variation. Unlike raw d records, the d_{In} records does not shows distinct ~ 40 kyr cycle nor the lag found in the glacial terminations. These results show that glacial-interglacial variations of d record are sensitive to its definition. Thus, great care should be taken to interpret such second-order parameter.

5 With the same methodology used for the d , the d_{In} data can be converting to the ΔT_{site} and ΔT_{source} . For the correction for ocean isotope changes, $\ln(1 + \delta)$ gives simpler formula than the case for normal δ .

$$\ln(1 + \delta^{18}\text{O})_{\text{corr}} = \ln(1 + \delta^{18}\text{O}_{\text{ice}}) - \ln(1 + \delta^{18}\text{O}_{\text{SW}}) \quad (\text{A3})$$

$$\ln(1 + \delta\text{D})_{\text{corr}} = \ln(1 + \delta\text{D}_{\text{ice}}) - \ln(1 + \delta\text{D}_{\text{SW}}) \quad (\text{A4})$$

10 Here, $\ln(1 + \delta^{18}\text{O})_{\text{corr}}$ and $\ln(1 + \delta\text{D})_{\text{corr}}$ are the isotope ratios of the ice corrected for ocean isotope changes. The $\delta\text{D}_{\text{SW}}$ can be calculated on the assumption that $\ln(\delta\text{D}_{\text{SW}} + 1) = -2.85 \times 10^{-2} \times (\ln(1 + \delta^{18}\text{O}_{\text{SW}}))^2 + 8.47 \times \ln(1 + \delta^{18}\text{O}_{\text{SW}})$. Thus, the d_{In} value corrected for the ocean isotope effect ($d_{\text{In,corr}}$) can be obtained by substituting $(1 + \delta^{18}\text{O})_{\text{corr}}$ and $(1 + \delta\text{D})_{\text{corr}}$ to the Eq. (A2). Then, the isotope inversion was performed to retrieve the ΔT_{site} and ΔT_{source} information. The relations obtained for the DF ice core are:

$$\Delta \ln(1 + \delta\text{D})_{\text{corr}} = 13.5\Delta T_{\text{site}} - 5.5\Delta T_{\text{source}} (R^2 = 0.999, n = 14, p < 0.001), \quad (\text{A5})$$

$$\delta d_{\text{In,corr}} = -0.4\Delta T_{\text{site}} + 1.8\Delta T_{\text{source}} (R^2 = 0.996, n = 14, p < 0.001). \quad (\text{A6})$$

We should note that we only modified the notation of δ and the definition of d . The two primary data sets (δD and $\delta^{18}\text{O}$) and inversion methodology are the same. Therefore, the ΔT_{site} and ΔT_{source} obtained based on Eqs. (A5) and (A6) are identical with those of obtained based on Eqs. (5) and (6). The values of coefficients of Eqs. (A5) and (A6) differ from those of Eqs. (5) and (6). The coefficient β_{site} (0.4) is significantly smaller than our estimation (1.3). This result clearly shows the fact that the value of β_{site} significantly depends on the definition of deuterium excess. In other word, β_{site} act as a correction factor for the empirically defined d parameter. Detailed and comprehensive analysis of d_{In} (e.g. other Antarctic sites, Greenland ice cores, and

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



global precipitation) is beyond the scope of this manuscript. At least for the case of DF, the smaller β_{site} value indicates that the d_{in} is better representation of source effect. In general, d_{in} would be suitable for comparison with similarly defined parameter, ^{17}O -excess.

- 5 *Acknowledgement.* We thank DF ice core group, Maki Nakada, Hironobu Yamada and Satoru Kikuchi for supporting isotopic measurement. We thank Kotaro Fukui for accessing the Antarctic snow isotope data. We also thank Alexey Ekaykin and Françoise Vimeux for comments. This research was supported by JSPS Postdoctoral Fellowships for Research Abroad and by a Grant-in-Aid for scientific research (S) (21221002) from the Ministry of Education, Science,
10 Sports and Culture, Japan and the French ANR Dome A project.



The publication of this article is financed by CNRS-INSU.

References

- 15 Antico, A., Marchal, O., Mysak, L. A., and Vimeux, F.: Milankovitch forcing and meridional moisture flux in the atmosphere: insight from a zonally averaged ocean-atmosphere model, *J. Climate*, 23, 4841–4855 doi:10.1175/2010JCLI3273.1, 2010.
- Barkan, E. and Luz, B.: Diffusivity fractionations of $\text{H}_2^{16}\text{O}/\text{H}_2^{17}\text{O}$ and $\text{H}_2^{16}\text{O}/\text{H}_2^{18}\text{O}$ in air and their implications for isotope hydrology, *Rapid Commun. Mass Spectrom.*, 21, 2999–3005, doi:10.1002/rcm.3180, 2007.
- 20 Barrows, T. T., Juggins, S., Deckker, P. D., Calvo, E., and Pelejero, C.: Long-term sea surface temperature and climate change in the Australian-New Zealand region, *Paleoceanography*, 22, PA2215, doi:10.1029/2006PA001328, 2007.

**Moisture-source
temperature
estimated from ice
core**

R. Uemura et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Bintanja, R. and Wal, R. S. W. V. D.: North American ice-sheet dynamics and the onset of 100 000-year glacial cycles, *Nature*, 454, 869–872, doi:10.1038/nature07158, 2008.
- Boyle, E. A.: Cool tropical temperatures shift the global $\delta^{18}\text{O}$ -T relationship: an explanation for the ice core $\delta^{18}\text{O}$ -borehole thermometry conflict?, *Geophys. Res. Lett.*, 24, 273–276, 1997.
- 5 Ciais, P. and Jouzel, J.: Deuterium and oxygen 18 in precipitation: Isotopic model, including mixed cloud processes, *J. Geophys. Res.*, 99, 16793–16803, 1994.
- Cortijo, E., Lehman, S., Keigwin, L., Chapman, M., Paillard, D., and Labeyrie, L.: Changes in meridional temperature and salinity gradients in the North Atlantic Ocean (30°–72° N) during the Last Interglacial Period, *Paleoceanography*, 14, 23–33, doi:10.1029/1998PA900004, 10 1999.
- Craig, H.: Isotopic variations in meteoric waters, *Science*, 133, 1702–1703, 1961.
- Cuffey, K. M. and Vimeux, F.: Covariation of carbon dioxide and temperature from the Vosotok ice core after deuterium-excess correction, *Nature*, 412, 523–527, 2001.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16, 436–468, 1964.
- 15 Davis, B. A. S. and Brewer, S.: Orbital forcing and role of the latitudinal insolation/temperature gradient, *Clim. Dynam.*, 32, 143–165, doi:10.1007/s00382-008-0480-9, 2009.
- Delaygue, G., Masson, V., Jouzel, J., and Koster, R. D.: The origin of Antarctic precipitation: a modelling approach, *Tellus B*, 52, 19–36, 2000.
- Ekeykin, A. A.: Meteorological regime of Central Antarctic and its role in the formation of isotope composition of snow thickness, Ph.D., University Joseph Fourier, Grenoble, 2003.
- 20 EPICA Community Members: Eight glacial cycles from an Antarctic ice core, *Nature*, 429, 623–628, doi:10.1038/nature02599, 2004.
- Fujita, K. and Abe, O.: Stable isotopes in daily precipitation at Dome Fuji, East Antarctica, *Geophys. Res. Lett.*, 33, L18503, doi:10.1029/2006GL026936, 2006.
- 25 Fujita, S., Holmlund, P., Andersson, I., Brown, I., Enomoto, H., Fujii, Y., Fujita, K., Fukui, K., Furukawa, T., Hansson, M., Hara, K., Hoshina, Y., Igarashi, M., Iizuka, Y., Imura, S., Ingvander, S., Karlin, T., Motoyama, H., Nakazawa, F., Oerter, H., Sjöberg, L. E., Sugiyama, S., Surdyk, S., Ström, J., Uemura, R., and Wilhelms, F.: Spatial and temporal variability of snow accumulation in Dronning Maud Land, East Antarctica, including two deep ice coring sites at Dome Fuji and EPICA DML, *The Cryosphere*, 5, 1057–1081, doi:10.5194/tc-5-1057-2011, 30 2011.
- Huybers, P. and Wunsch, C.: Obliquity pacing of the late Pleistocene glacial terminations, *Nature*, 434, 491–494, doi:10.1038/nature03401, 2005.

**Moisture-source
temperature
estimated from ice
core**

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jouzel, J.: Isotopes in cloud physics: multistep and multistage processes, *The terrestrial Environment B*, in: *Handbook of Environmental Isotope Geochemistry*, Elsevier, New York, 61–112, 1986.

Jouzel, J. and Merlivat, L.: Deuterium and oxygen 18 in precipitation: modeling of the isotopic effects during snow formation, *J. Geophys. Res.*, 89, 11749–11757, 1984.

Jouzel, J., Vimeux, F., Caillon, N., Delaygue, G., Hoffmann, G., Masson-Delmotte, V., and Parrenin, F.: Magnitude of isotope/temperature scaling for interpretation of Central Antarctic ice cores, *J. Geophys. Res.*, 108, 4361, doi:10.1029/2002JD002677, 2003.

Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B., Nouet, J., Barnola, J. M., Chappellaz, J., Fischer, H., Gallet, J. C., Johnsen, S., Leuenberger, M., Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J. P., Stenni, B., Stocker, T. F., Tison, J. L., Werner, M., and Wolff, E. W.: Orbital and millennial Antarctic climate variability over the past 800 000 years, *Science*, 317, 793–796, doi:10.1126/science.1141038, 2007.

Kavanaugh, J. L. and Cuffey, K. M.: Space and time variation of $\delta^{18}\text{O}$ and δD in Antarctic precipitation revisited, *Global Biogeochem. Cy.*, 17, 1017, doi:10.1029/2002GB001910, 2003.

Kawamura, K., Parrenin, F., Lisiecki, L., Uemura, R., Vimeux, F., Severinghaus, J. P., Hutterli, M., Nakazawa, T., Aoki, S., Jouzel, J., Raymo, M. E., Matsumoto, K., Nakata, H., Motoyama, H., Fujita, S., Goto-Azuma, K., Fujii, Y., and Watanabe, O.: Northern Hemisphere forcing of climatic cycles in Antarctica over the past 360 000 years, *Nature*, 448, 912–916, doi:10.1038/nature06015, 2007.

Khodri, M., Leclainche, Y., Ramstein, G., Braconnot, P., Marti, O., and Cortijo, E.: Simulating the amplification of orbital forcing by ocean feedbacks in the last glaciation, *Nature*, 410, 570–574, 2001.

Labeyrie, L., Labracherie, M., Gorfti, N., Pichon, J. J., Vautravers, M., Arnold, M., Duplessy, J.-C., Paterne, M., Michel, E., Duprat, J., Caralp, M., and Turon, J.-L.: Hydrographic changes of the Southern Ocean (Southeast Indian sector) over the last 230 kyr, *Paleoceanography*, 11, 57–76, 1996.

Laepple, T., Werner, M., and Lohmann, G.: Synchronicity of Antarctic temperatures and local solar insolation on orbital timescales, *Nature*, 471, 91–94, doi:10.1038/nature09825, 2011.

Landais, A., Barkan, E., and Luz, B.: Record of $\delta^{18}\text{O}$ and ^{17}O -excess in ice from Vostok Antarctica during the last 150 000 years, *Geophys. Res. Lett.*, 35, L02709,

doi:10.1029/2007GL032096, 2008.

Landais, A., Barkan, E., Vimeux, F., Masson-Delmotte, V., and Luz, B.: Combined analysis of water stable isotope ($H_2^{16}O$, $H_2^{17}O$, $H_2^{18}O$, $HD^{16}O$) in ice cores, in: Physics of Ice-Core Records II, edited by: Hondoh, T., Institute of Low Temperature Science, Hokkaido University Press, Sapporo, 315–327, 2009.

Lee, S.-Y. and Poulsen, C. J.: Amplification of obliquity forcing through mean annual and seasonal atmospheric feedbacks, *Clim. Past*, 4, 205–213, doi:10.5194/cp-4-205-2008, 2008.

Loutre, M.-F., Paillard, D., Vimeux, F., and Cortijo, E.: Does mean annual insolation have the potential to change the climate?, *Earth Planet. Sci. Lett.*, 221, 1–14, doi:10.1016/S0012-821X(04)00108-6, 2004.

Luz, B. and Barkan, E.: Variations of $^{17}O/^{16}O$ and $^{18}O/^{16}O$ in meteoric waters, *Geochim. Cosmochim. Acta*, 74, 6276–6286, doi:10.1016/j.gca.2010.08.016, 2010.

Majoube, M.: Fractionnement en oxygène-18 et en deutérium entre l'eau et sa vapeur, *J. Chim. Phys.*, 68, 1423–1436, 1971a.

Majoube, M.: Fractionnement en oxygène-18 entre la glace et la vapeur d'eau, *J. Chim. Phys.*, 68, 625–636, 1971b.

MARGO Project Members: Constraints on the magnitude and patterns of ocean cooling at the Last Glacial Maximum, *Nat. Geosci.*, 2, 127–132, doi:10.1038/NGEO411, 2009.

Masson-Delmotte, V., Stenni, B., and Jouzel, J.: Common millennial-scale variability of Antarctic and Southern Ocean temperatures during the past 5000 years reconstructed from the EPICA Dome C ice core, *The Holocene*, 14, 145–151, 2004.

Masson-Delmotte, V., Hou, S., Ekaykin, A., Jouzel, J., Aristarain, A., Bernardo, R. T., Bromwich, D., Cattani, O., Delmotte, M., Falourd, S., Frezzotti, M., Gallée, H., Genoni, L., Isaksson, E., Landais, A., Helsen, M. M., Hoffmann, G., Lopez, J., Morgan, V., Motoyama, H., Noone, D., Oerter, H., Petit, J. R., Royer, A., Uemura, R., Schmidt, G. A., Schlosser, E., Simões, J. C., Steig, E. J., Stenni, B., Stievenard, M., Broeke, M. R. V. D., Wal, R. S. W. V. D., Berg, W. J. V. D., Vimeux, F., White, J. W. C., and Vimeux, F.: A review of Antarctic surface snow isotopic composition: observations, atmospheric circulation, and isotopic modeling, *J. Climate*, 21, 3359–3387, doi:10.1175/2007JCLI2139.1, 2008.

Masson-Delmotte, V., Stenni, B., Blunier, T., Cattani, O., Chappellaz, J., Cheng, H., Dreyfus, G., Edwards, R. L., Falourd, S., Govin, A., Kawamura, K., Johnsen, S. J., Jouzel, J., Landais, A., Lemieux-Dudon, B., Laurantou, A., Marshall, G., Minster, B., Mudelsee, M., Pol, K., Rothlisberger, R., Selmo, E., and Waelbroeck, C.: Abrupt change of Antarctic

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



moisture origin at the end of Termination II, *P. Natl. Acad. Sci. USA*, 107, 12091–12094, doi:10.1073/pnas.0914536107, 2010a.

Masson-Delmotte, V., Stenni, B., Pol, K., Braconnot, P., Cattani, O., Falourd, S., Kageyama, M., Jouzel, J., Landais, A., Minster, B., Barnola, J. M., Chappellaz, J., Krinner, G., Johnsen, S., Röthlisberger, R., Hansen, J., Mikolajewicz, U., and Otto-Bliesner, B.: EPICA Dome C record of glacial and interglacial intensities, *Quaternary Sci. Rev.*, 29, 113–128, doi:10.1016/j.quascirev.2009.09.030, 2010b.

Masson-Delmotte, V., Buiron, D., Ekaykin, A., Frezzotti, M., Gallée, H., Jouzel, J., Krinner, G., Landais, A., Motoyama, H., Oerter, H., Pol, K., Pollard, D., Ritz, C., Schlosser, E., Sime, L. C., Sodemann, H., Stenni, B., Uemura, R., and Vimeux, F.: A comparison of the present and last interglacial periods in six Antarctic ice cores, *Clim. Past*, 7, 397–423, doi:10.5194/cp-7-397-2011, 2011.

Merlivat, L. and Jouzel, J.: Global climatic interpretation of the deuterium-oxygen 18 relationship for precipitation, *J. Geophys. Res.*, 84, 5029–5033, 1979.

Merlivat, L. and Nief, G.: Fractionnement isotopique lors des changements d'état solide-vapeur et liquide-vapeur de l'eau à des températures inférieures à 0°C, *Tellus*, 19, 122–127, 1967.

Miller, M. F.: Isotopic fractionation and the quantification of ¹⁷O anomalies in the oxygen three-isotope system: an appraisal and geochemical significance, *Geochim. Cosmochim. Acta*, 66, 1881–1889, 2002.

Motoyama, H.: The second deep ice coring project at Dome Fuji, Antarctica, *Sci. Drill.*, 5, 41–43, doi:10.2204/iodp.sd.5.05.2007, 2007.

Motoyama, H., Hirasawa, N., Satow, K., and Watanabe, O.: Seasonal variations in oxygen isotope ratios of daily collected precipitation and wind drift samples and in the final snow cover at Dome Fuji Station, Antarctica, *J. Geophys. Res.*, 110, D11106, doi:10.1029/2004JD004953, 2005.

Pahnke, K. and Sachs, J. P.: Sea surface temperatures of southern midlatitudes 0–160 kyr B.P., *Paleoceanography*, 21, PA2003, doi:10.1029/2005PA001191, 2006.

Paillard, D., Labeyrie, L., and Yiou, P.: Macintosh Program performs time-series analysis, *Eos Trans. AGU*, 77, 379, doi:10.1029/96EO00259, 1996.

Parrenin, F., Barnola, J.-M., Beer, J., Blunier, T., Castellano, E., Chappellaz, J., Dreyfus, G., Fischer, H., Fujita, S., Jouzel, J., Kawamura, K., Lemieux-Dudon, B., Loulergue, L., Masson-Delmotte, V., Narcisi, B., Petit, J.-R., Raisbeck, G., Raynaud, D., Ruth, U., Schwander, J., Severi, M., Spahni, R., Steffensen, J. P., Svensson, A., Udisti, R., Waelbroeck, C., and

Moisture-source temperature estimated from ice core

R. Uemura et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Wolff, E.: The EDC3 chronology for the EPICA Dome C ice core, *Clim. Past*, 3, 485–497, doi:10.5194/cp-3-485-2007, 2007.

Petit, J. R., White, J. W. C., Young, N. W., Jouzel, J., and Korotkevich, Y. S.: Deuterium excess in recent Antarctic snow, *J. Geophys. Res.*, 96, 5113–5122, 1991.

5 Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J.-M., Basile, I., Bender, M., Chappel-
laz, J., Davisk, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M., Lipenkov, V. Y.,
Lorius, C., Peâpin, L., Ritz, C., Saltzman, E., and Stievenard, M.: Climate and atmospheric
history of the past 420 000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429–436,
1999.

10 Pol, K., Debret, M., Masson-Delmotte, V., Capron, E., Cattani, O., Dreyfus, G., Falourd, S.,
Johnsen, S., Jouzel, J., Landais, A., Minster, B., and Stenni, B.: Links between MIS 11
millennial to sub-millennial climate variability and long term trends as revealed by new high
resolution EPICA Dome C deuterium data – A comparison with the Holocene, *Clim. Past*, 7,
437–450, doi:10.5194/cp-7-437-2011, 2011.

15 Raymo, M. E. and Nisancioglu, K.: The 41 kyr world: Milankovitch's other unsolved mystery,
Paleoceanography, 18, 1011, doi:10.1029/2002PA000791, 2003.

Reijmer, C. H., Broeke, M. R. V. D., and Scheele, M. P.: Air parcel trajectories and snowfall
related to five deep drilling locations in Antarctica based on the ERA-15 dataset, *J. Climate*,
15, 1957–1968, 2002.

20 Risi, C., Landais, A., Bony, S., Jouzel, J., Masson-Delmotte, V., and Vimeux, F.: Understanding
the ¹⁷O excess glacial-interglacial variations in Vostok precipitation, *J. Geophys. Res.*, 115,
D10112, doi:10.1029/2008JD011535, 2010.

Rozanski, K., Araguás-Araguás, L., and Gonfiantini, R.: Isotopic patterns in modern global
precipitation, in: *Climate Changes in Continental Isotopic Records*, edited by: Swart, P. K.,
25 Lohmann, K. C., McKenzie, J., and Savin, S., Geophysical Monograph, 78, American Geo-
physical Union, Washington DC, 1–36, 1993.

Sachs, J. P., Anderson, R. F., and Lehman, S. J.: Glacial surface temperature of the Southeast
Atlantic Ocean, *Science*, 293, 2077–2079, 2001.

Salamatin, A. N., Ekaykin, A. A., and Lipenkov, V. Y.: Modelling isotopic composition in precipi-
tation in Central Antarctica, *Mater. Glyatsiol. Issled.*, 97, 24–34, 2004.

30 Sime, L. C., Wolff, E. W., Oliver, K. I. C., and Tindall, J. C.: Evidence for warmer interglacials in
East Antarctic ice cores, *Nature*, 462, 342–345, doi:10.1038/nature08564, 2009.

Skinner, L. C., Fallon, S., Waelbroeck, C., Michel, E., and Barker, S.: Ventilation

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- of the deep Southern Ocean and deglacial CO₂ rise, *Science*, 328, 1147–1151, doi:10.1126/science.1183627, 2010.
- Sodemann, H. and Stohl, A.: Asymmetries in the moisture origin of Antarctic precipitation, *Geophys. Res. Lett.*, 36, L22803, doi:10.1029/2009GL040242, 2009.
- 5 Steig, E. J., Grootes, P. M., and Stuiver, M. S.: *Science*, 266, 1885–1886, doi:10.1126/science.266.5192.1885, 1994.
- Stenni, B., Masson-Delmotte, V., Johnsen, S., Jouzel, J., Longinelli, A., Monnin, E., Röthlisberger, R., and Selmo, E.: An oceanic cold reversal during the last deglaciation, *Science*, 293, 2074–2077, 2001.
- 10 Stenni, B., Jouzel, J., Masson-Delmotte, V., Röthlisberger, R., Castellano, E., Cattani, O., Falourd, S., Johnsen, S. J., Longinelli, A., Sachs, J. P., Selmo, E., Souchez, R., Stefensen, J. P., and Udisti, R.: A late-glacial high-resolution site and source temperature record derived from the EPICA Dome C isotope records (East Antarctica), *Earth Planet. Sci. Lett.*, 217, 183–195, doi:10.1016/S0012-821X(03)00574-0, 2003.
- 15 Stenni, B., Masson-Delmotte, V., Selmo, E., Oerter, H., Meyer, H., Röthlisberger, R., Jouzel, J., Cattani, O., Falourd, S., and Fischer, H.: The deuterium excess records of EPICA Dome C and Dronning Maud Land ice cores (East Antarctica), *Quaternary Sci. Rev.*, 29, 146–159, doi:10.1016/j.quascirev.2009.10.009, 2010.
- Tzedakis, P. C., Roucoux, K. H., De Abreu, L., and Shackleton, N. J.: The duration of forest stages in Southern Europe and interglacial climate variability, *Science*, 306, 2231–2235, doi:10.1126/science.1102398, 2004.
- 20 Uemura, R., Yoshida, N., Kurita, N., Nakawo, M., and Watanabe, O.: An observation-based method for reconstructing ocean surface changes using a 340 000-year deuterium excess record from the Dome Fuji ice core, Antarctica, *Geophys. Res. Lett.*, 31, L13216, doi:10.1029/2004GL019954, 2004.
- 25 Uemura, R., Matsui, Y., Motoyama, H., and Yoshida, N.: Deuterium and oxygen-18 determination of microliter quantities of a water sample using an automated equilibrator, *Rapid Commun. Mass Spectr.*, 21, 1783–1790, doi:10.1002/rcm.3022, 2007.
- Uemura, R., Matsui, Y., Yoshimura, K., Motoyama, H., and Yoshida, N.: Evidence of deuterium excess in water vapor as an indicator of ocean surface conditions, *J. Geophys. Res.*, 113, D19114, doi:10.1029/2008JD010209, 2008.
- 30 Uemura, R., Abe, O., and Motoyama, H.: Determining the ¹⁷O/¹⁶O ratio of water using a water-CO₂ equilibration method: application to glacial–interglacial changes in ¹⁷O-excess

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from the Dome Fuji ice core, Antarctica, *Geochim. Cosmochim. Acta*, 74, 4919–4936, doi:10.1016/j.gca.2010.05.007, 2010.

Vázquez Riveiros, N., Waelbroeck, C., Skinner, L., Roche, D. M., Duplessy, J.-C., and Michel, E.: Response of South Atlantic deep waters to deglacial warming during Terminations V and I, *Earth Planet. Sci. Lett.*, 298, 323–333, doi:10.1016/j.epsl.2010.08.003, 2010.

Vimeux, F., Masson, V., Jouzel, J., Stievenard, M., and Petit, J. R.: Glacial-interglacial changes in ocean surface conditions in the Southern Hemisphere, *Nature*, 398, 410–413, 1999.

Vimeux, F., Masson, V., Delaygue, G., Jouzel, J., Petit, J. R., and Stievenard, M.: A 420 000 year deuterium excess record from East Antarctica: information on past changes in the origin of precipitation at Vostok, *J. Geophys. Res.*, 106, 31863–31873, 2001.

Vimeux, F., Cuffey, K. M., and Jouzel, J.: New insights into Southern Hemisphere temperature changes from Vostok ice cores using deuterium excess correction, *Earth Planet. Sci. Lett.*, 203, 829–843, 2002.

Watanabe, O., Jouzel, J., Johnsen, S., Parrenin, F., Shoji, H., and Yoshida, N.: Homogeneous climate variability across East Antarctica over the past three glacial cycles, *Nature*, 509–512, 2003.

Winkler, R., Landais, A., Sodemann, H., Dümbgen, L., Prié, F., Masson-Delmotte, V., Stenni, B., and Jouzel, J.: Deglaciation records of ^{17}O -excess in East Antarctica: reliable reconstruction of oceanic normalized relative humidity from coastal sites, *Clim. Past*, 8, 1–16, doi:10.5194/cp-8-1-2012, 2012.

Moisture-source temperature estimated from ice core

R. Uemura et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Comparison of the temperature inversion coefficients.

Site	β_{site}	β_{source}	β_{sw}	γ_{site}	γ_{source}	γ_{sw}	Reference
DF	1.3	1.6		7.7	3.2	–	This study
Vostok ¹	0.5	1.3	2.8	7.1	3.7	4.8	Cuffey and Vimeux (2001); Vimeux et al. (2002)
Vostok ^{1,2}	0.9, 1.2	1.1, 1.2	3.0, 2.9	6.8, 6.9	2.8, 3.6	4.5, 4.7	Salamantin (2004)
Vostok ^{1,3}	1.0	1.4	3	–	–	–	Landais et al. (2009)
Vostok ³	1.1	1.5	–	–	–	–	Risi et al. (2010)
Vostok	1.4	1.6		7.7	3.1	–	This study
EDC ¹	0.5	1.3	2.6	7.6	3.5	5.0	Senni et al. (2001); Masson (2004)
EDC	0.5	1.3	–	7.6	3.6	–	Stenni et al. (2003)
EDC	1.2	1.5	–	7.8	3.4	–	This study
East Antarctica ^{1,3}	1.29–2.04	1.31–1.5	3	–	–	–	Winkler et al. (2011)

¹ Corrections for ocean isotope changes were estimated using β_{sw} and γ_{sw} : $\delta d = -\beta_{\text{site}}\Delta T_{\text{site}} + \beta_{\text{source}}\Delta T_{\text{source}} - \beta_{\text{sw}}\Delta\delta^{18}\text{O}_{\text{sw}}$, and $\Delta\delta D = \gamma_{\text{site}}\Delta T_{\text{site}} - \gamma_{\text{source}}\Delta T_{\text{source}} + \gamma_{\text{sw}}\Delta\delta^{18}\text{O}_{\text{sw}}$. See Sect. 2.3 for detail.

² Condensation temperature at precipitation deposition site (ΔT_{c}) used in original paper was converted to temperature at ground level (ΔT_{site}) using a linear relation $\Delta T_{\text{c}} = 0.67\Delta T_{\text{site}}$ (Jouzel and Merlivat, 1984). Thus, the d/T_{c} slopes (1.7 and 1.4 for 2 different trajectories) correspond to the values of 1.2 and 0.9 for β_{site} .

³ Variations of relative humidity was estimated as an independent parameter.

Moisture-source temperature estimated from ice core

R. Uemura et al.

Table 2. Test of β_{site} sensitivity to the source and site temperatures.

Tuning strategy	Simulated DF isotope ratio			Temperatures (°C)		Sensitivity coefficient			
	$\delta\text{D}(\text{‰})$	$\delta^{18}\text{O}(\text{‰})$	$d(\text{‰})$	DF site	Source	β_{site}	β_{source}	Y_{site}	Y_{source}
Isotope fitted	−422.7	−54.6	14.5	−61	18	1.3	1.6	7.7	3.2
Temperature fitted 1	−377.1	−46.2	6.5	−55	18	0.8	1.3	7.6	3.4
Temperature fitted 2	−351.1	−43.9	−0.2	−55	11	0.6	1.2	7.9	4.1

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.




Fig. 1. δ D and deuterium-excess records from the DF ice cores. **(a)** δ D and d records from DF1 (black lines) and DF2 (blue lines) cores. **(b)** Enlarged view of the d records during the overlapping period. Here, $\delta = R_{\text{sample}}/R_{\text{VSMOW}} - 1$, and R_{sample} and R_{VSMOW} are the isotope ratios (D/H and $^{18}\text{O}/^{16}\text{O}$) of sample and VSMOW, respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.




Fig. 2. Deuterium-excess and δD records of Antarctic ice cores. **(a)** Deuterium-excess (d) records from DF (red line), Vostok (blue line), and EDC (green line) cores plotted on the O_2/N_2 orbital age scale for DF. **(b)** δD records of the three cores plotted on the O_2/N_2 orbital age scale for DF. **(c)** δD records of the three cores plotted on the original age-scales (see text). Numbers indicates corresponding Marine Isotope Stage (MIS) (Tzedakis et al., 2004). All the isotope ratios are expressed as the deviations (Δ) from the present (past 2 kyr average) value.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.




Fig. 3. Isotope ratios of Antarctic surface snow and ice cores. Regression curves of surface snow data from various parts of Antarctica (black dashed-dotted line) (Masson-Delmotte et al., 2008), and from Japanese-Swedish IPY 2007/2008 traverse covering Syowa-Dome F route (black dotted line). A simulated curve is shown as blue solid line. Dots represent ice-core data of DF over the past 360 kyr (red), Vostok over the past 250 kyr (blue), and EDC over the past 140 kyr (green). Here, δD and d are corrected for $\delta^{18}O_{sw}$, see text for detail.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.




Fig. 4. Simulated δD and d values in Antarctica. Trajectories of present-day (black dotted line) and glacial conditions (red dotted line) are shown. Each point (solid or open circles) represents the simulated snow isotope values at DF site under a specific site and source temperature setting, shown as $(\Delta T_{\text{site}}, \Delta T_{\text{source}})$ in $^{\circ}\text{C}$. The DF ice core data (blue dot), and data used for regression analysis were shown as black solid circle. Linear regression $d/\delta D$ slopes of a constant $\Delta T_{\text{source}} (= -6.0^{\circ}\text{C})$ obtained for the filled circles and all circles are shown as green solid and chain lines, respectively. Here, δD and d are corrected for $\delta^{18}\text{O}_{\text{sw}}$, see text for detail.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.




Fig. 5. Different ΔT_{source} and ΔT_{site} of DF and a stacked SST records. **(a)** Black line represents the ΔT_{source} record of DF core based on the coefficients ($\beta_{\text{site}} = 1.3$) obtained by isotope-fitting (this study). Blue line represents the $\Delta T_{\text{source}}^{\text{low}}$ record of DF core based on the coefficients ($\beta_{\text{site}} = 0.5$) used for Vostok reconstruction (temperature-fitting) (Cuffey and Vimeux, 2001; Vimeux et al., 2002). Numbers indicate Marine Isotope Stages. **(b)** Black line is the ΔT_{source} of DF core, same as **(a)**, but with a stacked southern SST record of ocean sediments (Barrows et al., 2007) (red line). **(c)** The ΔT_{site} record based on the coefficients of this study (black line), $\Delta T_{\text{site}}^{\text{low}}$ (blue line), and previous ΔT_{site} reconstruction of DF core (temperature-fitting) (Kawamura et al., 2007).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.




Fig. 6. The ΔT_{site} , ΔT_{source} , and the temperature gradient derived from the three cores. Temperature records reconstructed from the DF (red line), Vostok (blue line), and EDC (green line) cores using the isotope-fitting method. **(a)** ΔT_{source} , **(b)** ΔT_{site} , **(c)** Temperature gradient ($= \Delta T_{\text{source}} - \Delta T_{\text{site}}$) and annual mean insolation gradient between 50 and 77° S (W m^{-2} , black dotted line).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.




Fig. 7. Obliquity components of the DF isotope records. **(a)** Earth's obliquity. **(b)** The raw d record of DF (black dotted line) and its filtered signal (red solid line). The filtered obliquity component was obtained by Gaussian filters with the Analyseries program (Paillard et al., 1996). A band-pass filter in the obliquity band ($f = 0.025 \pm 0.005$) was calculated (red solid line). **(c)** Filtered components in the DF d (red solid line) and δD (black solid line). **(d)** The δD (black dotted line) record of DF and its filtered signal (black solid line).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core core

R. Uemura et al.




Fig. 8. Power spectra of DF core isotope records. Power spectra of DF (red), Vostok (blue), and EDC (green); **(a)** δD , **(b)** d , **(c)** Ocean isotope change ($\delta^{18}O_{SW}$) over the past 360 kyr (Bintanja and Wal, 2008), **(d)** ΔT_{site} , **(e)** ΔT_{source} , and **(f)** temperature gradient ($\Delta T_{source} - \Delta T_{site}$). Spectral analysis is performed using Blackman–Tukey analysis with 50 % of series lag (for DF, VK, and $\delta^{18}O_{SW}$) and 70 % (for EDC) with the Analyseries software (Paillard et al., 1996).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.




Figure A1. Normal versus log plots for global meteoric water. **(a)** GMWL data from GNIP and Antarctic surface snow. GNIP data is based on 308 stations whose data contains the d data longer than 2 yr. The GNIP data weighted by amount of precipitation give a recession line ($\delta D = 8.18 \times \delta^{18}O + 11.2 (R^2=0.991)$). **(b)** Same as **(a)** but plot on the log plot. All the data give a regression curve (GMWC): $\delta D = -2.85 \times 10^{-2} \times (\delta^{18}O)^2 + 8.47 \times \delta^{18}O + 13.3 (R^2=0.999)$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Moisture-source temperature estimated from ice core

R. Uemura et al.




Figure A2. Slopes based on fractionation factors and GMWC. The $\delta D/\delta^{18}O$ slope is calculated using a differential of Eq. (7) ($= -5.70 \times 10^{-2} \times \ln(1 + \delta^{18}O) + 8.47$). Kinetic fractionation during snow formation was considered using a super-saturation function (see text).

Moisture-source temperature estimated from ice core

R. Uemura et al.




Figure A3. Logarithmic defined deuterium-excess and δD records of Antarctic ice cores. **(a)** Deuterium-excess based on logarithmic expression (d_{in}) records from DF (red line), Vostok (blue line), and EDC (green line) cores plotted on the O_2/N_2 orbital age scale for DF. **(b)** $\ln(\delta D + 1)$ records of the three cores plotted on the O_2/N_2 orbital age scale for DF.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)