

**Greenland
 $\delta^{18}\text{O}$ -accumulation
relationships**

S. L. Buchardt et al.

This discussion paper is/has been under review for the journal *Climate of the Past* (CP).
Please refer to the corresponding final paper in CP if available.

Investigating the past and recent $\delta^{18}\text{O}$ -accumulation relationship seen in Greenland ice cores

S. L. Buchardt, H. B. Clausen, B. M. Vinther, and D. Dahl-Jensen

Centre for Ice and Climate, NBI, University of Copenhagen, Juliane Maries Vej 30,
2100 Copenhagen OE, Denmark

Received: 30 July 2012 – Accepted: 14 August 2012 – Published: 27 August 2012

Correspondence to: S. L. Buchardt (lilja@gfy.ku.dk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Decadal means of $\delta^{18}\text{O}$ and accumulation rates from 52 ice core sites in Greenland are presented. The accumulation rates are derived from annual layers determined in the $\delta^{18}\text{O}$ curve. Investigation of the $\delta^{18}\text{O}$ -accumulation relationship across the ice divide reveals a significant Foehn effect with anticorrelation of $\delta^{18}\text{O}$ and accumulation on the lee side of the divide in Southern Greenland, while no effect is seen in Central Greenland. Furthermore, the sensitivity of accumulation rate to changes in temperature is found to be smaller in Northern Greenland than in the central and southern parts. Four sites in the data set contain sufficient recent data that the period of observed temperature rise from the 1990's and onwards can be investigated. All four sites are located close to the ice divide in Northern Greenland and while three sites show increased temperatures, none show evidence of increased accumulation.

1 Introduction

During the last decades, global temperatures have increased and are affecting global sea level which is currently rising at a rate of $3.11 \pm 0.6 \text{ mm yr}^{-1}$ (1993–2008) (Ablain et al., 2009). Continental ice is a significant contributor to this rise in sea level, but the uncertainty on the contribution is not well constrained, as pointed out by the IPCC panel in the AR4 report from 2007 (IPCC, 2007). The Greenland ice sheet constitutes the second largest continental ice mass and holds enough ice to increase global sea level by 7 m if melted completely. Observations have shown that coastal temperatures in Greenland have increased over the last decades (Box, 2002), the area of surface melting is growing (Abdalati and Steffen, 2001) and the ice loss has been estimated to $205 \pm 50 \text{ Gt yr}^{-1}$ (2005–2006) (AMAP, 2011). Few measurements exist from the interior part of the ice sheet, but GC-Net automated weather stations have observed rising temperatures (Box et al., 2009). RCM runs indicate that the rise in temperature should be followed by an increase in accumulation rate over the ice sheet, because the warmer

Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



air can contain more moisture (Box et al., 2006). Furthermore, several authors have determined empirical relationships between temperature and accumulation rates that also indicate that accumulation should increase with rising temperatures (Dahl-Jensen et al., 1993; Dansgaard et al., 1993; Johnsen et al., 1995). Dahl-Jensen et al. (1993) found the following relation between accumulation rates and $\delta^{18}\text{O}$ in the GRIP ice core.

$$A = 0.23 \exp(0.117(\delta^{18}\text{O} + 34.83)) \quad (1)$$

where A is the ice equivalent accumulation rate in m yr^{-1} and the $\delta^{18}\text{O}$ values are given in ‰. The factor 0.23 was chosen because the present day mean ice equivalent accumulation rate at GRIP is 0.23 m yr^{-1} .

In the following we will first investigate the relationship between temperature and accumulation rate by using ice core data from Greenland ice cores, and then we will use this as a foundation to look for recent changes in temperature and accumulation rate. Furthermore, we will investigate the geographical differences in the relationship between $\delta^{18}\text{O}$ and accumulation rate.

2 Data

Accumulation rates and temperatures for 52 sites in Greenland are inferred from ice core $\delta^{18}\text{O}$ data. The 52 sites and the time period covered at each site are listed in Table 1, and their locations are shown in Fig. 1. Most of the ice core data available are from cores drilled in the 1970's and 1980's and thus do not contain information on the recent warming period. Therefore, a number of firn cores were drilled and measured for $\delta^{18}\text{O}$ under the Ice2Sea Programme to either extend existing records to the present or to obtain data from new sites, thus increasing the amount of available data for this period. Where several cores exist from the same site, a composite $\delta^{18}\text{O}$ record has been created.

3 Method

It has been established that the annual mean $\delta^{18}\text{O}$ value at a given site on an ice sheet is closely related to the mean annual temperature at that site (Dansgaard, 1953, 1964). Johnsen et al. (1989) found the following empirical relation for sites located on the Greenland ice sheet

$$\delta^{18}\text{O} = 0.67 \times T - 13.7, \quad (2)$$

where T is the temperature in $^{\circ}\text{C}$ and the $\delta^{18}\text{O}$ is given in ‰. As this relationship is linear, we choose to compare the accumulation rates to $\delta^{18}\text{O}$ directly instead of to the temperatures calculated from $\delta^{18}\text{O}$.

All timescales used in this study are based on annual layer counting in the $\delta^{18}\text{O}$ curve. For low accumulation sites it is necessary to correct for the effect of smoothing by diffusion (Johnsen et al., 2000) before the records can be used for dating. The thickness of each annual layer as determined in the isotope curve is corrected for changes in density with depth in the firn using a Herron–Langway densification model (Herron and Langway Jr., 1980) and the deeper layers are corrected for flow-induced thinning using a 1-dimensional Dansgaard–Johnsen model (Dansgaard and Johnsen, 1969). Thus the $\delta^{18}\text{O}$ data are used to infer accumulation rates as well as temperatures. All accumulation rates discussed in this work are given in ice equivalent.

Our data set contains six cores with data reaching more than a few tens or hundreds of years back in time (cf. Table 1). These six long records are used to investigate the relation between $\delta^{18}\text{O}$ and accumulation on somewhat longer time scales of up to a few thousand years. These data are shown in Fig. 2.

In order to investigate the $\delta^{18}\text{O}$ -accumulation relationship in different parts of the Greenland ice sheet, the sites are divided into six groups based on their locations: Northwest (NW), Northeast (NE), Central west (CW), Central east (CE), Southwest (SW), and Southeast (SE). The boundary between the east and west groups is defined by the ice divide. Sites located on the ice divide have been sorted into the west group

in the northern and central groups but into the east group in the south, because the dominating moisture transport is from the southeast in Southern Greenland and from the west in the central and northern parts of the ice sheet. In Table 1 it is stated which group each site has been sorted into. The data from Renland have been omitted in this analysis, because Renland is a separate ice dome and thus not part of the main ice sheet.

4 Results and discussion

From Fig. 2 it is seen that the annual data show a quite noisy relationship between $\delta^{18}\text{O}$ and accumulation rate. By using averages over longer time periods, the relationship becomes clearer. As the changes we are interested in have happened over the last 1–2 decades we choose to use 10-yr mean values in the following to keep a balance between noise-level and time resolution. A data set consisting of average values of temperature and ice equivalent accumulation rate for each decade from 1900 to 2009 (or the part of this period covered by the core) is provided in the Supplement.

It is noteworthy from Fig. 2 that the scatter in $\delta^{18}\text{O}$ is comparable for the low accumulation sites (left) and the sites with higher accumulation (right), whereas the spread in accumulation seems smaller for the low accumulation sites. In the case of Dye-3, layers older than ~ 350 yr are affected by complicated upstream depositional conditions (Vinther et al., 2006). This probably explains at least part of the large spread in the calculated accumulation rates as we have not corrected for upstream effects.

Figure 3 shows 10 yr averages of accumulation rate and $\delta^{18}\text{O}$ for the 51 coring sites located on the main ice sheet. It is seen that no common Greenland relationship between $\delta^{18}\text{O}$ and accumulation exists. This is not surprising as the shape of the ice sheet is complex and moisture reaches different parts of the ice sheet via different routes. Assuming that the relationship between $\delta^{18}\text{O}$ and accumulation rate is exponential (Johnsen et al., 1989), least squares fits to the data in the six groups give the

Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



following relationships that we have chosen to state with a factor of 0.23 in front of the exponential to ease comparison with Eq. (1)

$$A = 0.23 \exp \left(0.099 \left(\delta^{18}\text{O} + 33.64 \right) \right) \quad \text{NW} \quad (3)$$

$$A = 0.23 \exp \left(0.023 \left(\delta^{18}\text{O} + 16.75 \right) \right) \quad \text{NE} \quad (4)$$

$$A = 0.23 \exp \left(0.140 \left(\delta^{18}\text{O} + 35.40 \right) \right) \quad \text{CW} \quad (5)$$

$$A = 0.23 \exp \left(0.137 \left(\delta^{18}\text{O} + 35.21 \right) \right) \quad \text{CE} \quad (6)$$

$$A = 0.23 \exp \left(-0.060 \left(\delta^{18}\text{O} + 20.18 \right) \right) \quad \text{SW} \quad (7)$$

$$A = 0.23 \exp \left(0.124 \left(\delta^{18}\text{O} + 34.25 \right) \right) \quad \text{SE} \quad (8)$$

Dye-3 data older than 300 yr were omitted when calculating the least square fit to the data from the SE area, because upstream effects cause artifacts in the layer thicknesses for older data.

In the southern part of the ice sheet, a positive correlation between $\delta^{18}\text{O}$ and accumulation is found east of the ice divide, while a negative correlation is seen to the west (Fig. 3: magenta and cyan, respectively). The dominating moisture transport across the ice divide is from the east, and the negative correlation west of the divide can be explained by the Foehn effect (warm, dry air). In the northern part of the ice sheet, all sites are located either on the divide or just east of it (black and green in Fig. 3). The sites east of the divide have lower accumulation than the sites that are located on the divide, but since there are only few data points in the NE group the relationship in Eq. (4) is not well defined. In Central Greenland no trend across the ice divide is seen (red and blue in Fig. 3). However, all sites in the CE group are located very close to the ice divide, so a Foehn effect east of the divide in Central Greenland can not be ruled out. However, the Foehn effect in the southern part of the ice sheet is seen for sites quite close to the divide, and the few data points from the northeast indicate

Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a quick drop in accumulation rate on the lee side of the divide. Perhaps the lack of an effect across the divide in Central Greenland can be explained by the fact that the area around the divide is broader and flatter and thus the orographic barrier is not as well defined as in South Greenland.

5 Table 2 shows the sensitivities of accumulation rate to changes in $\delta^{18}\text{O}$ and to changes in temperature assuming the relationship from Eq. (2). Two previously published values for GRIP are also shown for comparison. The values found in this study for the central part of the ice sheet agree with the value given by Dansgaard et al. (1993). Furthermore, the sensitivity seems to be lower in the north compared to the
10 central and southern parts of the ice sheet.

Out of the 52 sites in the data set, only five records contain information on the recent warming over the last couple of decades. The $\delta^{18}\text{O}$ record from ACT10A only goes back to 1994 AD, and since we have no information on previous conditions at this site, the record can not be used to draw conclusions about recent changes. Figure 4 shows
15 the mean values of accumulation and $\delta^{18}\text{O}$ for each decade since 1900 AD for the remaining four sites. The data show no significant increase in accumulation rate in recent years at any of the sites. However, the $\delta^{18}\text{O}$ values for the most recent decade at the NGRIP, B26 and NEEM sites are increasing indicating increasing temperatures. For NEEM and B26 the decade after 2000 AD has the “warmest” mean $\delta^{18}\text{O}$ value of
20 the whole data periods of 270 yr and 110 yr, respectively, while the NGRIP record of 1820 yr shows only four decades with higher mean $\delta^{18}\text{O}$ value than the 2000–2009 decade.

5 Conclusions

25 A data set consisting of decadal means of $\delta^{18}\text{O}$ and accumulation rates derived from annual layers determined in the $\delta^{18}\text{O}$ curves from 52 ice core sites in Greenland was presented. Out of these records, six span longer time periods than a few hundred years, and four can be used to study the period of recent warming from the 1990's and

CPD

8, 4105–4120, 2012

Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



onwards. From the long records it is clear that annual accumulation and $\delta^{18}\text{O}$ data are quite noisy. However, as the averaging period is increased, the relationship between accumulation and $\delta^{18}\text{O}$ becomes clearer.

Using 10 yr averages of $\delta^{18}\text{O}$ and accumulation rates for the 51 sites located on the main ice sheet, we found that no common $\delta^{18}\text{O}$ -accumulation relation exists for Greenland. Considering the complicated shape of the ice sheet and local differences in moisture transport this is not surprising. However, the data fall into six groups according to location. In Southern Greenland a significant Foehn effect is seen across the ice divide resulting in a negative correlation between $\delta^{18}\text{O}$ and accumulation on the lee (west) side of the ice divide. In Northern Greenland we only have few data on the lee side (east) of the ice divide, but the data that do exist show a sharp drop in accumulation rate across the divide and also indicate a significant change in the $\delta^{18}\text{O}$ -accumulation relationship. In Central Greenland, however, the data do not indicate a significant effect of the ice divide. This may be explained by the divide in Southern Greenland being more pronounced with steeper slopes on either side thus constituting a sharper barrier. The sensitivity of accumulation rate to changes in temperature seems to be smaller in Northern Greenland than in the central and southern areas.

The four records that contain recent data show no sign of increased accumulation rate after 2000 AD, but three of the records indicate very warm temperatures for that period. For NEEM and B26 the decade 2000–2009 has the warmest $\delta^{18}\text{O}$ value of the records of 270 yr, and 110 yr, respectively. In the 1820 yr long record from NGRIP only four decades have warmer $\delta^{18}\text{O}$ values than 2000–2009. At Camp Century no increase in accumulation rate or $\delta^{18}\text{O}$ can be seen, and the record from ACT10A is too short to draw any conclusions from. Thus, the ice core data indicate an increase in temperature after 2000 AD, but no increase in accumulation rate is observed. However, data is sparse and concentrated along the ice divide in Northern Greenland. New data series spanning at least the last 100 yr are needed from more sites in order to expand the above mentioned conclusions to wider areas of the ice sheet.

Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Supplementary material related to this article is available online at:
<http://www.clim-past-discuss.net/8/4105/2012/cpd-8-4105-2012-supplement.zip>.

Acknowledgements. We acknowledge the ice2sea project, funded by the European Commission's 7th Framework Programme through grant number 226375, ice2sea manuscript number 103.

References

- Abdalati, W. and Steffen, K.: Greenland ice sheet melt extent: 1979–1999, *J. Geophys. Res.*, 106, 33983–33988, 2001. 4106
- Ablain, M., Cazenave, A., Valladeau, G., and Guinehut, S.: A new assessment of the error budget of global mean sea level rate estimated by satellite altimetry over 1993–2008, *Ocean Sci.*, 5, 193–201, doi:10.5194/os-5-193-2009, 2009. 4106
- AMAP, 2011: Snow, Water, Ice and Permafrost in the Arctic (SWIPA): climate change and the cryosphere, Arctic monitoring and assessment programme (AMAP), Oslo, Norway, 538 pp., 2011. 4106
- Box, J. E.: Survey of Greenland instrumental temperature records: 1873–2001, *Int. J. Climatol.*, 22, 1829–1847, doi:10.1002/joc.852, 2002. 4106
- Box, J. E., Bromwich, D. H., Veenhuis, B. A., Bai, L.-S., Stroeve, J. C., Rogers, J. C., Steffen, K., Haran, T., and Wang, S.-H.: Greenland ice sheet surface mass balance variability (1988–2004) from calibrated Polar MM5 output, *J. Climate*, 19, 2783–2800, 2006. 4107
- Box, J. E., Yang, L., Bromwich, D. H., and Bai, L.-S.: Greenland ice sheet surface air temperature variability: 1840–2007, *J. Climate*, 22, 4029–4049, 2009. 4106
- Dahl-Jensen, D., Johnsen, S. J., Hammer, C. U., and Jouzel, J.: Past accumulation rates derived from observed annual layers in the GRIP ice core from Summit, Central Greenland, in: NATO ASI Series, Vol. 1, 12, Ice in the Climate System, edited by: Peltier, W. R., Springer, Berlin, 517–532, 1993. 4107, 4116, 4118, 4119
- Dansgaard, W.: The abundance of O^{18} in atmospheric water and water vapour, *Tellus* 5, 461–469, 1953. 4108
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus* 16, 436–468, 1964. 4108

Greenland $\delta^{18}O$ -accumulation relationships

S. L. Buchardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dansgaard, W. and Johnsen, S. J.: A flow model and a time scale for the ice core from Camp Century, Greenland, *J. Glaciol.*, 8, 215–223, 1969. 4108

Dansgaard, W., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N. S., Hammer, C. U., Hvidberg, C. S., Steffensen, J. P., Sveinbjörnsdottir, A. E., Jouzel, J., and Bond, G.: Evidence for general instability of past climate from a 250-kyr ice-core record, *Nature*, 364, 218–220, 1993. 4107, 4111, 4116

Herron, M. M. and Langway Jr., C. C.: Firn densification: an imperial model, *J. Glaciol.*, 25, 373–385, 1980. 4108

IPCC, Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007. 4106

Johnsen, S. J., Dansgaard, W. and White, J. W. C.: The origin of arctic precipitation under present and glacial conditions, *Tellus B*, 41, 452–468, 1989. 4108, 4109

Johnsen, S. J., Dahl-Jensen, D., Dansgaard, W., and Gundestrup, N.: Greenland palaeotemperatures derived from GRIP bore hole temperature and ice core isotope profiles, *Tellus B*, 47, 624–629, 1995. 4107

Johnsen, S. J., Clausen, H. B., Cuffey, K. M., Hoffmann, G., Schwander, J., and Creyts, T.: Diffusion of stable isotopes in polar firn and ice: The isotope effect in firn diffusion, in: *Physics of Ice Core Records*, edited by: Hondoh, T., Hokkaido University Press, Sapporo, 121–140, 2000. 4108

Vinther, B. M., Clausen, H. B., Johnsen, S. J., Rasmussen, S. O., Andersen, K. K., Buchardt, S. L., Dahl-Jensen, D., Seierstad, I. K., Siggaard-Andersen, M.-L., Steffensen, J. P., Svensson, A. M., Olsen, J., and Heinemeier, J.: A synchronized dating of three Greenland ice cores throughout the Holocene, *J. Geophys. Res.*, 111, D13102, doi:10.1029/2005JD006921, 2006. 4109

Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 1. Greenland ice cores dated by annual layer counting in the $\delta^{18}\text{O}$ record.

Ice core	Latitude	Longitude	Period covered by data	Group
ACT10A	65.69	-41.48	1994–2009	SE
B26*	77.25	-49.22	1900–2009	NW
Camp Century*	77.17	-61.10	1762–2009	NW
Crete*	71.12	-37.32	554–1973	CW
Dye-2*	66.48	-46.33	1743–1973	SW
Dye-3*	65.18	-43.83	-3816–1979	SE
Dye-3 1975 (D-2)	65.16	-43.94	1965–1974	SE
Dye-3 1975 (D-3)	65.16	-43.92	1965–1974	SE
Dye-3 1975 (D-4)	65.17	-43.90	1966–1974	SE
Dye-3 1975 (D-5)	65.17	-43.88	1965–1974	SE
Dye-3 1975 (D-6)	65.18	-43.87	1966–1974	SE
Dye-3 1980 Swiss core	65.20	-43.81	1834–1979	SE
Dye-3 1981 13B	65.09	-44.24	1965–1980	SE
Dye-3 1981 SDV	65.02	-44.65	1964–1980	SE
Dye-3 1982 7B	65.14	-44.02	1967–1979	SE
Dye-3 1983 4B	65.17	-43.93	1692–1981	SE
Dye-3 1984 5B	65.16	-43.88	1956–1982	SE
Dye-3 1984 9B	65.12	-44.09	1955–1983	SE
Dye-3 1984 11B	65.11	-44.16	1955–1983	SE
Dye-3 1984 15B	65.08	-44.30	1954–1983	SE
Dye-3 1984 16C	65.05	-44.33	1955–1983	SE
Dye-3 1984 18C	65.03	-44.39	1777–1983	SE
GRIP*	72.57	-37.62	-1846–1989	CW
Milcent 1973	70.30	-45.00	1177–1966	CW
NEEM*	77.45	-51.08	1737–2009	NW
NGRIP*	75.10	-42.32	187–2007	NW
North Central*	74.62	-39.60	1910–1976	NE
North Site 1972	75.77	-42.45	1925–1971	NE
Reniand*	71.31	-26.72	931–1986	-
Saddle North 1975 (SN)	66.17	-43.78	1967–1974	SE
Saddle North 1975 Satellite 1 (SNS-1)	66.47	-44.83	1961–1974	SW
Saddle South 1975 (SAS)	65.68	-44.33	1966–1974	SE
Saddle South 1975 satellite 1 (SDS-1)	65.70	-44.76	1964–1974	SW
Saddle South 1975 satellite 2 (SDS-2)	65.53	-44.12	1966–1974	SE
Saddle South 1975 satellite 3 (SDS-3)	65.83	-44.12	1966–1974	SE
Site A (Crete region) 1985	70.63	-35.82	1623–1984	CE
Site A-1 1975 (Dye-3 region)	67.50	-42.20	1965–1974	SE
Site A-1 1975 Satellite 1 (A-1-S-1)	67.03	-41.85	1967–1974	SE
Site A-1 1975 Satellite 2 (A-1-S-2)	67.85	-43.12	1961–1974	SW
Site B (Crete region) 1984	70.65	-37.48	1717–1983	CW
Site BDS 1975 (Dye-3 region)	64.50	-44.33	1965–1974	SE
Site C (Crete region) 1984	70.68	-38.78	1943–1983	CW
Site D (Crete region) 1984	70.63	-39.62	1767–1982	CW
Site E (Crete region) 1985	71.77	-35.85	1722–1984	CE
Site F (Crete region) 1985	71.50	-35.88	1925–1982	CE
Site G (Crete region) 1985	71.15	-35.83	1778–1984	CE
Site H (Crete region) 1985	70.87	-35.83	1933–1983	CE
South Dome 1975	63.53	-44.58	1869–1974	SE
South Dome 1975 Satellite 1 (DS-1)	63.60	-44.25	1968–1974	SE
South Dome 1975 Satellite 2 (DS-2)	63.55	-44.93	1962–1974	SW
South Dome 1975 Satellite 3 (DS-3)	63.70	-44.58	1964–1974	SW
SUMMIT 1974	72.28	-37.98	1904–1971	CW

* Record is a composite of several cores from the same site.

Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

Table 2. The sensitivity of accumulation rate A to changes in $\delta^{18}\text{O}$ and temperature T .

Group	$\frac{1}{A} \frac{\partial A}{\partial \delta}$ (‰ ⁻¹)	$\frac{1}{A} \frac{\partial A}{\partial T}$ (%K ⁻¹)
NW	9.9	6.6
NE	2.3	1.5
CW	14.0	9.4
CE	13.7	9.2
SW	-6.6	-4.2
SE	12.4	8.3
GRIP (from Dahl-Jensen et al., 1993)	11.7	7.8
GRIP (from Dansgaard et al., 1993)	14.4	9.6

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 1. The locations of the 52 ice core sites listed in Table 1. The five sites from which we have data for the most recent decade are indicated with labeled arrows.

Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. Mean values of accumulation rate and $\delta^{18}\text{O}$ for the six longest records in our data set. The averaging periods are 1 yr (green), 10 yr (yellow), 100 yr (cyan), and 1000 yr (black). The dotted grey line shows the relation from Dahl-Jensen et al. (1993) (Eq. 1). See Table 1 for the time periods covered by each data series.




Fig. 3. 10 yr averages of ice equivalent accumulation rate and $\delta^{18}\text{O}$ from 51 sites in Greenland. The sites have been sorted into six groups according to location: Northwest (black), Northeast (green), Central west (red), Central east (blue), Southwest (cyan), and Southeast (magenta). The least squares exponential fits to the data for each of the groups are shown in the corresponding colour and the relation from Dahl-Jensen et al. (1993) is shown for comparison (dotted grey curve). The insert shows the locations of the 51 core sites with colours indicating the grouping of the sites. For Dye-3, data is only shown for the last 300 yr as the thickness of layers older than this are affected by upstream conditions.

Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Greenland $\delta^{18}\text{O}$ -accumulation relationships

S. L. Buchardt et al.




Fig. 4. Average values of accumulation rates (blue) and $\delta^{18}\text{O}$ values (red) for each decade since 1900 AD. The blue and red dashed lines indicate the mean value $\pm 2\sigma$ of accumulation rate and $\delta^{18}\text{O}$, respectively, for the whole data period for each site. The periods of data for the four sites are: Camp Century: 1762–2009 AD, NEEM: 1737–2009 AD, B26: 1900–2009 AD, and NGRIP: 187–2007 AD.