

Abstract

High-resolution pollen analysis of core MD01-2430 from the Sea of Marmara (40°47.81' N, 27°43.51' E) allows us to reconstruct the vegetation response to climatic changes during the past 23 cal ka in the Northeastern Mediterranean. Variation in mesic/temperate forest cover indicates major climatic shifts connected to Heinrich Stadial 1, Bölling-Allerød, Younger Dryas and to the onset of the Holocene. Pollen-anthropogenic indicator approach was used to recognize human-induced landscape changes in the Sea of Marmara. The pollen-inferred onset of the Holocene occurs at ca. 11.5 cal ka, indicating that the Northeastern Mediterranean region represents a transitional zone where higher moisture availability supported an earlier forest expansion than the borderlands of the Aegean Sea and Black Sea. Three major forest retreats occurred during the Holocene at ca. 9.2, 5.5, and 2.1 cal ka while a weaker forest setback was detected at 7.9 cal ka. The Holocene forest setbacks are in phase with previously published alkenone-inferred sea-surface temperature decreases in the Sea of Marmara (core MD01-2430) reconstructed from the same archive. Our new pollen record testifies the sensitivity of Mediterranean forests to changes in moisture availability which is driven by changes in high-latitude climate drivers (North Atlantic Oscillations and/or Siberian High).

1 Introduction

Palynological studies carried out in the Northeastern Mediterranean region have shown that the vegetation of this region was sensitive to the climatic variations of the North Atlantic high latitudes (i.e. Heinrich events and Dangaars-Oeschger cycles) (Geraga et al., 2005; Fletcher et al., 2010a) as well as to short climatic shifts during the Holocene (Wijmstra, 1969; Roberts et al., 2001; Lawson et al., 2005; Tzedakis, 2007; Kotthoff et al., 2008a, 2011; Pross et al., 2009). Repeated forest contractions have been associated with regional sea surface temperature cooling events probably due to the

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



enhancement of the Siberian High (SH), (Rohling et al., 2002; Marino et al., 2009; Pross et al., 2009; Kotthoff et al., 2011), which is responsible of cold and dry spells in winter (Saaroni et al., 1996). However, despite the extensive palynological investigations carried out in Northeastern Mediterranean, the nature and timing of vegetation dynamics at the onset of the Holocene is not yet fully clarified (Tzedakis, 2007).

The Holocene reforestation in the northern border of the Aegean Sea occurs earlier (at 10.2 cal) (Kotthoff et al., 2008a), than in the Black Sea, where the establishment of forests is dated to anything between 9.5 and 8 cal ka (Atanassova, 2005; Mudie et al., 2007; Shumilovskikh et al., 2012). The Sea of Marmara lies in a transitional zone between the Aegean Sea and the Black Sea and may, therefore, be a key area to clarify the timing of early Holocene reforestation in the Northeastern Mediterranean region. Moreover, no high-resolution palaeoclimatic record exists for this region displaying a direct correlation between changes in sea surface temperatures and vegetation changes at times of rapid climatic shifts during the last deglaciation (such as Heinrich event 1 (HE 1), the Bölling-Allerød interstadial (B/A), and the Younger Dryas (YD)). The first overview of the vegetation dynamics around the Sea of Marmara for the past 33 ka was based on low-resolution pollen records that suggested colder and drier pre- and post-Late Glacial Maximum (LGM) conditions, a warm and humid early Holocene, and sustained anthropogenic impact on vegetation from 3.5 ¹⁴C kyr BP onwards (Mudie et al., 2002, 2007). Nonetheless unfavourable climatic conditions (e.g. dry/cold episodes) in the Eastern Mediterranean were also impacting prehistoric societies causing their collapse (Weiss et al., 1993; Vermoere et al., 2002).

In this paper we present a new pollen record from the topmost 7.5 m of a sediment core (MD01 2430) collected in the Sea of Marmara, covering the past 23 cal ka. The reconstructed SST from this sedimentary sequence revealed a series of cold spells during the last 15 000 yr (Vidal et al., 2010) and a surprisingly cold B/A interstadial which is in agreement with the dinocyst record (Londeix et al., 2009). Our main goals were (i) to characterize the Northeastern Mediterranean vegetation response, tempo, and amplitude to strong and weak North Atlantic abrupt climatic changes of the last

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



23 000 yr, (ii) to verify the unexpected low temperature recorded during the B/A as suggested by dinocysts analysis and alkenone-inferred SST done on the same core (see for details, Londeix et al., 2009; Vidal et al., 2010), (iii) to identify vegetation dynamics and climatic conditions (i.e. precipitation regime) prevailing during the early Holocene, and (iv) to propose the most likely atmospheric situation that can explain the climatic record of the Marmara region. To address these questions, we focused pollen analysis to obtain a high-temporal resolution record of vegetation dynamics for the LGM, HS 1 (~ 19–14.6 ka, Sanchez Goñi and Harrison, 2010; Stanford et al., 2011), B/A, YD and the Holocene.

Understanding past climatic changes and their impact on vegetation is an urgent need considering that from climate change projections over the Mediterranean emerges a pronounced decrease in precipitation and increase in temperature especially during the summer season (Giorgi and Lionello, 2008). It is likely that such changes in the precipitation regime will have an impact on water resources and consequently biodiversity, increase in forest fires, and reduction in crop production (Alcamo et al., 2007).

2 Regional setting

The Sea of Marmara is 275 km long and 80 km wide and connects the Aegean Sea with the brackish Black Sea via the 70 m deep Dardanelles Strait to the South and the 40 m deep Bosphorus Strait to the North (Fig. 1). The present oceanography of the Sea of Marmara is characterized by the outflow of brackish water from the Black Sea, whereas saline bottom water flows in from the Mediterranean Sea, which results in a surface-water salinity of 20.1–24.6 psu (Ünlüata et al., 1990). Major rivers flow into the Sea of Marmara only from the south. The Kocasu Rivers is the biggest modern river entering the southern shelf of the Sea of Marmar and it delivers about 90 % of the total suspended riverine sediment (Ünlüata et al., 1990; Çagatay et al., 2000; Kazancı et al., 2004). The Sea of Marmara is under a dry-summer subtropical climate (Köppen,

1923) with a mean winter, summer, and annual temperature of 3.8°C, 25.7°C, and 13.7°C respectively. Annual precipitation is ca. 660 mm yr⁻¹ (Unal et al., 2003; Kazancı et al., 2004) with most of the precipitation occurring during winter in association with the westerly depressions, while in summer the subtropical high-pressure area brings warm and dry conditions (Trewartha, 1981). In the Marmara region the potential vegetation is the eu-Mediterranean woodland near the coast of the Sea of Marmara and euxinian-type forest inland (Zohary, 1973; Roberts and Wright, 1993). The Euxinian forest is dominated by summergreen vegetation (*Quercus* species, *Fagus orientalis*, *F. sylvatica*, *Carpinus betulus*, *C. Orientalis*, and *Castanea*), *Pinus sylvestris*, and *Abies* (Zohary, 1973). The eu-Mediterranean woodland is a drought-resistant forest that extends from the coasts up to ca. 400 m elevation. The dominant species are evergreen oaks (*Quercus ilex*, *Q. Coccifera*, and *Q. infectoria*), *Pinus halepensis*, *Pistacia lentiscus*, *Olea europea*, *Ceratonia siliqua*, *Poaceae*, and herbs. At present, the landscape is very open and intensively farmed.

3 Materials and methods

The core MD01-2430 (40° 47.81' N, 27° 43.51' E) was retrieved during the MD123 cruise (R/V *Marion Dufresne*) in 2001 with a Calypso piston corer on the Western high separating the Central basin and the Tekirda basin at 580 m water depth and is 29 m long (Vidal et al., 2010). The base of the core section consists of homogeneous silty mud, gray with dark spots and thin pyrite-rich layers. A greyish-brown ash layer is present at 698–691 cm. At 350 cm a major lithologic transition corresponds to the transition from lacustrine to marine deposits. Between 320 and 230 cm a dark layer was interpreted as sapropel. A homogenous greenish silty mud characterizes the top of the sequence (Londeix et al., 2009; Vidal et al., 2010).

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1 Chronology and depth-age model

The chronology is based on six accelerator mass spectrometry radiocarbon dates on mollusc samples (Table 1, Vidal et al., 2010). Although the water reservoir age is still unknown for the Sea of Marmara, we decided to use the 400-yr correction as calculated for the Eastern Mediterranean Sea and Black Sea on modern mollusc shells (Siani et al., 2000). Indeed the first marine incursion in the Sea of Marmara has been estimated at ca. 14.7 cal ka (Vidal et al., 2010) and after that reservoir age in the bottom water of the Sea of Marmara are assumed to have been similar to the estimated reservoir age of the Mediterranean.

Geochemistry of the ash layer deposited at 698–691 cm in core MD01-2430 confirms that it corresponds to the marine tephra Y-2. This tephra represents the distal facies of the Cape Riva eruption of Santorini, and it had been reported in other marine sediment cores from the Sea of Marmara (Çagatay et al., 2000; Wulf et al., 2002). The Cape Riva eruption has been dated on deposits on land, in which four dates have a mean value and standard deviation of $18\,310 \pm 380$ $^{14}\text{C yr}$ (see for details, Pichler and Friedrich, 1976; Eriksen et al., 1990; Kwiecien et al., 2008).

In order to prepare the depth-age model, the radiocarbon dates were converted to calendar years BP with the CALIB rev 5 program (Stuiver et al., 1998). We used the Marine04 calibration curve (Hughen et al., 2004) for the dates based on molluscs and the IntCal04 dataset (Reimer et al., 2004) for the Y-2 tephra age. The depth-age model was based on non-parametric weighted regression within the framework of generalised additive models (Fig. 2). The resulting model and associated age estimates and 95 % confidence intervals are based on the combined uncertainty of the calibrated dates and the regression line (Birks and Heegaard, 2003; Heegaard et al., 2005). The mean estimated ages used in this paper are slightly different from those published in Vidal et al. (2010). However the depth-age model of Vidal et al. (2010) is within the confidence interval of our model (Fig. 2). The mean sedimentation rate during the glacial period ranges between about 70 and 30 cm kyr^{-1} (19 ± 4 yr cm^{-1}). Then during

the late-glacial and early Holocene, the sedimentation rate reaches the mean lowest value of $\sim 22 \text{ cm kyr}^{-1}$ ($46 \pm 6 \text{ yr cm}^{-1}$). Finally, above 20 cm depth the sedimentation rate increases to $31 \pm 15 \text{ yr cm}^{-1}$.

3.2 Pollen analysis

5 Samples of 1 cm^3 were used for pollen analysis, with *Lycopodium* tablets added for estimation of pollen concentration (Stockmarr, 1971). The samples were prepared as follows: coarse-sieving at $150 \mu\text{m}$, successive treatments with cold HCl at increasing strength (10%, 25%, and 50%), successive treatments with cold HF (HF 25% for 2 1/2 h, HF 70% 1 1/2 days), and micro-sieving ($10 \mu\text{m}$ mesh). Pollen grains and
10 spores were counted at a magnification of $\times 400$ and $\times 1000$ and identified using keys, pollen atlases (Moore et al., 1991; Reille, 1992–1998), and the reference collection of the UMR EPOC, University Bordeaux 1. A minimum of 20 taxa, 100 terrestrial pollen grains (excluding *Pinus*, aquatic plants, spores, and unknown), and a minimum of 200 objects (including *Lycopodium* and terrestrial pollen grains) were counted in each sample
15 (Finsinger and Tinner, 2005). The average terrestrial pollen sum is 160 ± 60 grains excluding *Pinus*, or 260 grains including *Pinus*. Pollen grains of *Pinus* are generally over-represented in marine pollen records and are therefore excluded from the pollen sum (Heusser and Balsam, 1977; Turon, 1984; Rossignol-Strick and Planchais, 1989). Maher (1981) showed how pollen percentages stabilize as the pollen sum increases, using the confidence intervals of the multinomial distribution. Rull (1987) indicated that
20 a pollen sum of 200 grains is statistically sufficient to produce reliable estimates. Indeed, after this value there are not significant variations in the widths of confidence intervals for percentage, while for pollen counts of 100 grains the variation is of ca. 1%.

25 The pollen types are grouped as follows: steppic (*Chenopodiaceae*, *Artemisia*, *Ephedra distachya*, and *E. fragilis*), mediterranean (*Quercus* evergreen type, *Olea*, *Phillyrea*, *Cistus*, and *Pistacia*), temperate (all trees and shrubs excluding *Pinus*, *Hippophaë*, *Quercus* evergreen type, *Olea*, *Phillyrea*, *Cistus* and *Pistacia*, *Ephedra*

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



distachya, *E. fragilis*), and upland herbs. The arboreal pollen group (AP) includes mediterranean and temperate pollen types. The program TILIA was used for plotting the pollen diagrams (Grimm, 1993). The identification of Holocene forest setbacks was based on shifts in AP percentage of at least 20% between adjacent. The sample resolution of the pollen record of 3–10 cm allowed for an average time resolution of 115 ± 100 yr (1 standard deviation-SD) during the glacial section, 420 ± 100 yr for the late glacial and 170 ± 90 yr for the Holocene. The pollen record was zoned using the visual zonation of the pollen curves on the basis of coincident changes occurring in multiple pollen taxa.

4 Results

Vegetation dynamics inferred from the pollen analysis

The pollen zone MD30-1 (22.5–21 cal ka) was dominated by herbs (*Cichorioideae*, *Aster* type, and *Centaurea nigra* type), steppic plants (*Artemisia* and *Chenopodiaceae*), *Hippophaë* and high pollen percentage of *Pinus* (not included in the pollen sum) (Fig. 3a). Pollen of temperate trees (mainly *Quercus* deciduous type and *Fagus*) were also present reaching two minima at 21.7 cal ka and 21.1 cal ka. These minima correspond to an increase in *Artemisia* and *Chenopodiaceae*. Occurrence of reworked pollen (mainly *Carya*) most probably testifies erosion of the slope around the basin of Marmara. The landscape was dominated by open land vegetation with *Hippophaë* and *Pinus* growing on sandy soils. This type of vegetation indicates a cold and rather dry climate during a period that falls in the Last Glacial Maximum (LGM, Mix et al., 2001) as also suggested by other records from the Sea of Marmara (Mudie et al., 2007).

At the beginning of zone MD30-2 (21–19.5 cal ka), pollen percentages of temperate trees (*Quercus deciduous* type, *Fagus* and *Carpinus*) increased at the cost of upland herbs, *Hippophaë* and *Pinus* (Fig. 3a). However, the pollen influx diagram indicates that *Hippophaë* was still an important component of the landscape (Fig. 3b) and the

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of *Sanguisorba minor* since 11.5 cal ka, followed by the appearance of *Pistacia* pollen. This feature is mirrored by the pollen influx values; although a sharp increase of oak pollen influx is only visible at ca. 10.5 cal ka. In this pollen zone an overall increase of the total pollen influx from two to six times corresponds to the deposition of Sapropel 1 (Fig. 3b). The occurrence of *Sanguisorba minor* and *Pistacia* can be considered as a biostratigraphic marker for the onset of the Holocene in the Eastern Mediterranean region (Rossignol-Strick, 1995; Turner and Sánchez Goñi, 1997; Kotthoff et al., 2008a). Mesic/temperate forest is the dominant vegetation during the early-middle Holocene, characterized by *Quercus deciduous* type, *Fagus*, *Abies*, *Ulmus*, *Carpinus*, and *Corylus* pollen. This type of vegetation indicates an increase of temperature and precipitation. Decrease of forest pollen percentages higher than 20% were detected at ca. 9.3–9.1 and a weaker forest setback occurred at 7.9 cal ka. These forest retreats correspond to an increase of steppic plants cutting off the long-term trend in increase of forest cover.

During the middle-late Holocene (zones MD30-8 and MD30-9) an overall slight reduction of the mesic forest is suggested by the pollen spectra, though this type of forest was still dominant. A marked forest setback is detected at ca. 5.5 cal ka, mainly due to a reduction of pollen percentages of *Quercus deciduous* type and increase in *Poaceae* and *Chenopodiaceae*. Then short-term minima of temperate and mediterranean tree pollen occur at ca 2.1 cal ka to which correspond increases of *Poaceae*, *Caryophyllaceae*, *Sanguisorba minor*, *Asteraceae*, *Cichorioideae*, and *Artemisia*. Increases in pollen percentage of herbs were coupled with the occurrence of anthropogenic-pollen indicators (Behre, 1990; Bottema and Woldring, 1990) such as *Cerealia* type, *Juglans* and *Olea* that occurred since 3.1, 2.4 and 2.3 cal ka respectively (Fig. 5). Probably human exploitation of the mesic/temperate forest became stronger. Similarly, in other sedimentary cores from the Sea of Marmara the disappearance of *Pinus* and *Quercus* forests and the spread of *Artemisia*, *Cichorioideae*, and *Chenopodiaceae* were detected after ca. 4000 yr BP (ca. 4.3–4.8 cal ka) and attributed to the human occupation of the area around the Sea of Marmara (Mudie et al., 2002, 2007).

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Discussion

5.1 Direct correlation between vegetation and sea surface changes in the Sea of Marmara region and integration with other records from the Northeastern Mediterranean region

5 The beginning of the studied sequence of core MD01-2430 falls in the Last Glacial Maximum (LGM, Mix et al., 2001) and according with the dinocysts flora (i.e. presence of *Spiniferites cruciformis*, *Pediastrum*, and fresh-water ostracods) the Sea of Marmara was an isolated mildly brackish (caspi-brackish). Dominance of *Spiniferites cruciformis* indicates a large seasonal contrast in the sea-surface temperatures (SST) (Londeix et al., 2009) (Fig. 4). At that time the vegetation was dominated by open land and *Hippophaë* (Fig. 3a) and some peaks in *Artemisia* and *Chenopodiaceae* pollen percentages were detected, indicating short-term dry periods. An increase of thermophilic dinocysts including subtropical/tropical coastal species (Londeix et al., 2009) occurred between 20.6 and 18.5 cal ka at about the time of an increase in temperate pollen types (*Quercus deciduous* type, *Fagus* and *Carpinus*) dated to 21–19.5 cal ka. Altogether these proxies indicate a change towards climatic conditions less severe than before (Fig. 4). Our data support recent model simulations for the LGM (21 ± 2 ka) that indicate the presence of cold-tolerant species (*Carpinus*, *Corylus*, *Fagus*, *Tilia*, *Fraxinus excelsior*, *Alnus*, *Quercus robur*, and *Ulmus*) on the western shore of the Sea of Marmara (Leroy and Arpe, 2007; Arpe et al., 2011).

AP percentages gradually decreased starting from about 19.5 cal ka at the benefit of steppic plants (Fig. 3a). Then a minimum in AP pollen percentages was reached between ca. 17–15 cal ka, pollen of steppic plants increased (up to 50 %) and pollen of semi-desert species (*Ephedra* sp.) were present. Similarly during this period (19.5–15 cal ka) thermophilic dinocysts almost disappeared: a first decrease is visible at ca. 18.4 cal ka and a second one at 16.4 cal ka. Cool dinocysts were instead present with stable values until 16.4 cal ka, when *Spiniferites cruciformis* became the dominant dinocyst (Fig. 4). Increase in steppic plants and cold dinocysts were also found in other

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cores from the Sea of Marmara (Mudie et al., 2007) and in the Aegean Sea (Geraga et al., 2010; Kotthoff et al., 2011) suggesting that at that time climatic conditions were more critical, e.g. colder and drier than the conditions during the LGM. Moreover this phase can be subdivided in a first half (up to 17 cal ka), which was probably wetter than the second half (17–15 cal ka). A similar dry/cold period with a two-phase pattern was also described for the Western Mediterranean (Naughton et al., 2007; Fletcher and Sanchez Goni, 2008; Fletcher et al., 2010a) and it was related to the HS 1, ~ 19–14.6 ka (Fig. 5). Based on lake-levels changes of Lake Lisan (paleo-Dead Sea), the link between Eastern Mediterranean drought phases and HSs events was present during the past 55 ka (Bartov et al., 2003). The proposed mechanism to explain this teleconnection accounts for an intensification of the Siberian High (SH) which is responsible for severe cold and dry winter conditions and polar/continental air outbreaks over the Aegean Sea (Rohling et al., 2002; Marino et al., 2009; Pross et al., 2009; Kotthoff et al., 2011). On the other hand, the collapse of the North Atlantic Deep Water circulation likely caused an input of cold water in the Mediterranean and a consequent reduction in the evaporation. In turn, this would have resulted in a reduction in the strength and the frequency of storms and therefore a decrease in precipitation over the Eastern Mediterranean region (Bartov et al., 2003).

Starting from 14.7 cal ka pollen percentages of steppic plants decreased and a sudden increase of *Quercus deciduous* type and AP is detected in the pollen record indicating an increase in temperatures and moisture. *Spiniferites cruciformis* is still dominant (Londeix et al., 2009), indicating probable cold winters and hot summers. Alkenone-derived SST indicates low temperature most probably during spring which is the period when coccoliths preferentially produce this compound (Fig. 4) (Vidal et al., 2010). A possible scenario that might explain the apparent contradiction between the climate evidences during the B/A is that the cold SST in Sea of Marmara is the result of the arrival of freshwater from melting Northern European ice during deglaciation, while our pollen data records the terrestrial vegetation. Another apparent contradiction is present when comparing the pollen and ostracods-derived stable-oxygen isotopes

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



record from core MD01-2430. The isotopes show a trend towards higher values since 14.5 cal ka which might be interpreted as an increase of evaporation. However, increase in moisture is evidenced by increase in mesic/temperate forest based on pollen data of core MD01-2430 and pollen records in the Aegean Sea (Kotthoff et al., 2008a, 2011) and in Turkey (Jones et al., 2007). This support the idea of Vidal et al. (2010) that attribute the increase in stable-oxygen isotopes values at the increase in precipitation-water $\delta^{18}\text{O}$ (3‰–4‰) and not to a drought phase during the B/A.

In addition in our record maximum of AP is reached towards the Allerød (Fig. 5) similarly to what was already described in Western Mediterranean (Combourieu Nebout et al., 2009; Fletcher et al., 2010b) suggesting a warming trend from the Bölling towards the Allerød. Our finding then support the idea of a climatic contrast during B/A with Greenland and North Europe expiring a cooling trend towards the end of the B/A while climate stability or warming was evidenced in Southern Europe (Genty et al., 2006; Combourieu Nebout et al., 2009).

In core MD01-2430 an optimum in dinoflagellates indicating cold conditions is again recorded between 13 and 12.15 ± 0.25 cal ka where *Bitectatodinium tepikiense* and *Spiniferites elongatus* are found with their highest values (Fig. 4). These species are winter cool-water indicators (Turon, 1984; Marino et al., 2009), and similarly alkenone-derived SST shows low values (ca. 10 °C, Vidal et al., 2010). This is in agreement with the results of pollen analysis that indicate a withdrawal of the mesic/temperate forest and increase of steppic plants, indicating cold and dry conditions during the YD. The YD has been already described in the Northeastern Mediterranean as an abrupt event characterized by a change towards more drought-tolerant type of vegetation (Wijmstra, 1969; Rossignol-Strick, 1995; Atanassova, 2005; Lawson et al., 2005; Mudie et al., 2007; Kotthoff et al., 2008a; Shumilovskikh et al., 2012). Pollen-based climatic reconstructions for the Aegean region have also shown a decline in annual temperature of ca. 6 °C and a precipitation decrease of ca. 300 mm (Dormoy et al., 2009; Kotthoff et al., 2011) during the YD.

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The onset of the Holocene in core MD01-2430 is marked by the establishment of a mesic/temperate forest with some element of the mediterranean forest which suggests the increase in temperature and water availability. The dinocysts record shows the association of cool and thermophilic species which might suggests a seasonal contrast in SST (Londeix et al., 2009) while alkenone SST show an increasing (Fig. 4) (Vidal et al., 2010). The late-glacial/Holocene transition as marked by the AP record in our record matches very well with changes in $\delta^{13}\text{C}$ values of a stalagmite located in Sofular Cave (Northwestern Anatolia) (Fleitmann et al., 2009), indicating a forested phase during the early-middle Holocene (Fig. 5). During the Holocene forest setbacks at ca. 9.2 and 5.5 are directly associated with decrease in SST in the Sea of Marmara (Fig. 4) highlighting a connection between the SST, the resulting moisture availability, and changes in the forest cover. A major forest retreat at 5.6 cal ka was also found in northern borderlands of the Aegean Sea (Kotthoff et al., 2008a), similarly a decrease in SST was found in the Aegean Sea (Marino et al., 2009). At ca. 5.5 cal ka decrease in winter-spring rain as determined by clay layer frequency are found in the Black Sea (Lamy et al., 2006) and a dry event is visible in stable-oxygen isotopes record of Soreq Cave (Fig. 5) (Bar-Matthews et al., 1997; Bar-Matthews and Ayalon, 2011). This drought phase coincides also with a major archaeological transition (Bar-Matthews and Ayalon, 2011; Roberts et al., 2011). Strikingly, our record shows only a weak forest retreat during the most pronounced shift in temperature at 7.9–8 cal ka as recorded by SST. Similarly $\delta^{13}\text{C}$ of Solfular Cave don't show any marked shifts in the values around 8 cal ka (Figs. 4 and 5) (Fleitmann et al., 2007). A pronounced vegetation shift is instead detected on the northern borderland of the Aegean Sea (Kotthoff et al., 2008b) and in Tenaghi Philippon pollen records (Pross et al., 2009) and it was associated with the 8.2 cal ka climate event. On the contrary a spreading of temperate forest was recently describe for the Black Sea region at ca. 8.3 cal ka and it was related to higher moisture availability (Shumilovskikh et al., 2012).

Further decreases in AP percentages from average values between adjacent samples of 10% might be identified at 10.4, 9.9, 6.1, 4.3, 2.9, 2.1, 1.2 and 0.3 cal ka

and interpreted as forest set-backs to which correspond an increase of steppic plants (Fig. 5). The minor oscillations prior the human impact (ca. 2.5 cal ka) can tentatively be correlated with increases in North-Atlantic ice-rafting – Bond’s events (Fig. 5) (Bond et al., 1997, 2001).

5.2 Early Holocene moisture availability in the Sea of Marmara region

The early Holocene in our sequence is characterized by a sudden expansion of temperate trees at ca. 11.5 ± 0.3 cal ka (Fig. 3a) indicating an increased availability of moisture. A warm and wet early Holocene (9.5 ^{14}C kyr BP; ca. 11 cal ka) was also invoked by Mudie et al. (2007) on the basis of pollen profiles from the Sea of Marmara. Holocene reforestation occurs at only at 12 cal ka in Nisi Fen (northern Greece) (Lawson et al., 2005), only at 10.2 cal ka on the northern border of the Aegean Sea according to marine pollen sequences (core SL 152, Kotthoff et al., 2008a), at ca. 8.5 ^{14}C kyr BP (9.5 cal ka) in the Southwestern Black Sea (Mudie et al., 2007), at ca. 8.5 cal ka in Southeastern Black Sea (Shumilovskikh et al., 2012), and at ca. 7.1–7.5 ^{14}C kyr BP (8 cal ka) in the Western Black Sea (Atanassova, 2005). However in core SL 152 (Aegean region) a decrease in the steppic plants (*Artemisia*, *Chenopodiaceae* and *Ephedra*) and expansions of herbs (Centaureaceae and *Cichorioideae*) occurred at ca. 11.6 cal ka indicating an increase of the water availability prior the forest expansion (Kotthoff et al., 2008a). A similar pattern can be seen in the Tenaghi Philippon sequence, where the reforestation occurs ca. 1000 yr after the decrease of the *Artemisia* and *Chenopodiaceae* (Pross et al., 2009).

The differing source areas for the pollen assemblages in the Sea of Marmara, Black Sea, and Aegean Sea might also explain the different timing of the reforestation. The Aegean sequence reflects the vegetation in the northern borderlands of the Aegean Sea (Kotthoff et al., 2008b), while the geographical position of the Sea of Marmara resulted in a pollen sequence that should reflect the vegetational changes on Thrace and Northwestern Anatolia. The fluvial component in the Sea of Marmara is represented only by rivers that flow from the Anatolia peninsula. However aerial pollen account

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for the main input in the Sea of Marmara sediments (van Zeist and Bottema, 1991; Mudie et al., 2002; Leroy et al., 2009). We can conclude that the pollen record of core MD01-2430 mainly represents Northwestern Anatolia. Stable-carbon isotopes of Sofular Cave suggest a presence of forested ecosystem since 10.5 cal ka in Northwestern Anatolia indicating that during the early Holocene moisture was available and that the vegetation response to favourable climatic conditions was fast (Fig. 5) (Fleitmann et al., 2009; Göktürk et al., 2011). Conversely, a dry early Holocene is detected by various proxies in the drier parts of the Near East, in sites located in southernmost Europe, and in continental zones. In these areas the reforestation occurred only after 8 ka due to persistent arid conditions that kept moisture levels below the tolerance threshold for tree growth (van Zeist and Bottema, 1991; Stevens et al., 2001; Bottema and Sarpaki, 2003; Wick et al., 2003; Tzedakis, 2007). We can conclude that the northern borderland of Southern Europe can be considered a transitional zone where the moisture level determined an earlier (ca. 11.5 cal ka) woodland expansion (e.g. Northern Anatolia as shown by our data) or a later expansion in ecological stressed areas (Tzedakis et al., 2004; Tzedakis, 2007).

The importance of the water availability (especially during winter) for the establishment of forest was highlighted by pollen-based climatic reconstructions (Kotthoff et al., 2008b; Dormoy et al., 2009). Recently climate simulations have shown that across the early Holocene stronger winter precipitation occurred over the Northeastern Mediterranean (near Turkey) while there is little evidence for summer precipitation in the Eastern Mediterranean (Brayshaw et al., 2011). The mechanism associated with winter precipitation in Eastern Mediterranean is not yet fully understood (Giorgi and Lionello, 2008). Evidence from paleo-data suggests that negative rainfall anomalies in the Black Sea region occurred during phases of high North Atlantic Oscillations (NAO) causing intervals of lower clay-layer frequencies (Lamy et al., 2006). A role of the Siberian High (SH) which is responsible for severe cold and dry winter conditions and polar/continental air outbreaks over the Aegean Sea (Rohling et al., 2002) is instead invoked to explain the evidence of 8.2 event in Tenaghi Philippon record (Pross

et al., 2009) and centennial-scale SST cooling events in the Aegean Sea (Marino et al., 2009). Finally comparison of palaeolimnological records from Eastern and Western Mediterranean has highlighted that NAO forcing alone cannot explain the hydrology pattern in Eastern Mediterranean since 900 AD (Roberts et al., 2012). In order to precisely establish the correlation between these two climate systems (SH/NAO) during the late-glacial and Holocene, it is necessary to make a comparison of our data with high-resolution pollen sequences from areas not affected by the SH (i.e. western part of the Mediterranean).

5.3 Evidences for human occupation during the Holocene

Holocene vegetation changes might be caused by variation in either climate or human activities or both. Human activities on the landscape can be detected by the increase of weeds and ruderal herbs (e.g. *Chenopodiaceae*, *Sanguisorba minor* type and *Cichorioideae*), domesticated species (e.g. *Cerealia* type, *Olea*, *Vitis* and *Juglans*) and an overall increase in non-arboreal pollen (NAP). However, all these pollen types can also be present in the natural vegetation, therefore care must be used in the interpretation of pollen diagrams of the Eastern Mediterranean (Behre, 1990; Bottema and Woldring, 1990).

Several Neolithic settlements (namely Illıpinar, Mentese, Barcın Höyük, Pendik, Fikirtepe, Yarımburkaz, and Toptepe) were established near the northern and southern shore of Sea of Marmara from ca. 8.4–7.4 cal ka (Roodenberg and Alpaslan-Roodenberg, 1999; Bottema et al., 2001; Rosenstock, 2005; Özdoğan, 2012) possibly causing a change in the landscape (Fig. 1). Palaeoenvironmental investigations connected with archaeological excavations were carried out in the area around Illıpinar (Fig. 1). The pollen spectra of Yenisehir core (the only sedimentary core dated) show a drop in tree pollen during the Late Neolithic/Early Chalcolithic (ca. 8–7.4 cal ka) and Late Chalcolithic/Early Bronze Age (ca. 5.7–4.6 cal ka), indeed during these periods the area was densely populated (Bottema et al., 2001).

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



On the basis of the pollen-anthropogenic approach (Behre, 1981) we can attest that our pollen diagram does not show clear traces of human activities during the early Holocene. A similar feature is clear in other pollen profiles located near Early Neolithic settlements, where a definite sign of human presence is not found, meaning that the human activities did not strongly impact the vegetation (van Zeist and Bottema, 1991; Willis, 1995; Bottema et al., 2001). On the other hand, in the pollen record of MD01-2430 core a deforestation phase was clearly detected at ca. 5.5 cal ka while a weaker forest setback occurred at 7.9 cal ka (Fig. 6). Considering chronological uncertainty, the forest setback at 5.5 cal ka might be related to the occupation phase around the nearby Illıpinar site during Late Chalcolithic/Early Bronze Age. Simultaneously, in the city of Kumtepe (ca. 5.5 cal ka) and later on in Troy (ca. 4.5 cal ka) there is clear evidence in increasing of openness (Korfmann, 2006; Riehl and Marinova, 2008). However the deforestation phase at 5.5 cal ka is also linked with a period of climatic deterioration (see previous section).

The presence in our pollen diagram of *Juglans*, *Olea*, *Phillyrea*, *Ericaceae*, *Cerealia* type, and *Centaurea nigra* type, as well as a steady increase of *Artemisia*, *Cichorioideae*, and trilete spores (possibly *Pteridium*) since ca. 2.5 cal ka indicates a marked change in the landscape (Fig. 6). This pollen assemblage suggests an opening of the forests in correspondence to an increase of human activities (e.g. cultivation of cereals and possibly arboriculture). The ancient city of Daskyleion lied on the shore of Lake Manyas nearby the south-western shore of Sea of Marmara (Fig. 1) from the Bronze Age to the Roman Period (ca. 2.6 cal ka–0.3 cal ka, Leroy et al., 2002). A pollen sequence from Lake Manyas indicates an increase of pollen percentage of *Juglans*, *Castanea*, *Vitis*, *Olea*, *Fraxinus ornus*, and *Quercus* (Leroy et al., 2002) which suggested a phase of intense human occupation characterised by arboriculture (the so-called Beyşehir Occupation Phase – BOP) (van Zeist and Bottema, 1991; Leroy et al., 2002). The BOP is reflected in several pollen diagram of Southwestern Turkey and it is dated from 3.5–1.4 cal ka (Eastwood et al., 1998). Although Lake Manyas lies few kilometres away from the Sea of Marmara, in our pollen diagram a clear BOP cannot be detected.

6 Conclusions

The marine pollen record from the Sea of Marmara, core MD01-2430, confirms the high vulnerability of the vegetation in Eastern Mediterranean (Northern Anatolia) to changes in the winter-spring precipitation regime and temperature. Steppic plants (such as *Artemisia* and *Chenopodiaceae*) dominated vegetation during periods with cold and dry climatic regime (e.g. HS1, YD), whereas *Cichorioideae* and *Asteraceae* and *Hippophaë* dominated during the LGM. We have shown a wet to dry trend within the HS1 in the Sea of Marmara region and drier and colder conditions than during the LGM, just as was found in the Western Mediterranean (Naughton et al., 2007; Fletcher et al., 2010a), and confirming that western storm tracks penetrated during the LGM as far as the Eastern Mediterranean (Kuhlemann et al., 2008; Laîné et al., 2009). Surprisingly our record does not show a time lag during the Holocene reforestation (ca. 11.5 cal ka), pointing to the fact that Northern Anatolia lies on a transitional area compared to the Near East or continental areas (see Black Sea records), where drought was a limiting factor up to 9–8 cal ka. In our record, as soon as moisture was sufficient forest vegetation expanded, probably also indicating presence of refugia of deciduous oak near the Sea of Marmara as predicted by model simulations (Arpe et al., 2011). The anthropogenic-pollen indicators approach shows that disentangling the climate from human signals in pollen sequences in the Eastern Mediterranean is still an open issue. Contemporaneous with a major regional human occupation, the major forest-withdrawal phase at 5.5 cal ka directly associated with SST lowering episodes can be linked to periods of low temperature and to winter precipitation according to pollen-independent palaeoclimatic proxies. Consistently with several paleoproxies in the Eastern Mediterranean the winter rainfall regime can be related to the SH or/and NAO. In the perspective of the future climate changes towards increased dry spells a better understanding of this climatic systems might be possible by the comparison of high-resolution paleoproxies from Eastern and Western Mediterranean.

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. We thank Miryam Bar-Matthews, Dominik Fleitmann and Laurence Vidal for respectively providing data on oxygen isotopes of Soreq Cave, carbon isotopes of Sofular Cave and sea-surface temperatures of core MD01-2430 that are present in Figs. 4 and 5; Fakke Gerritsen of the NIT (The Netherlands Institute in Turkey) in Istanbul for kindly providing assistance regarding data on prehistoric settlements in Anatolia. Guillemette Ménot and Laurence Vidal are acknowledged for constructive comments. We also thank Olga Kwiecien for discussion about the reservoir age of the Black Sea. Linguistic improvements of the manuscript by H.E. Wright Jr. are gratefully acknowledged as well as technical assistance given by Marie-Helen Castera. This is Bordeaux 1 University, UMR-CNRS5805 EPOC Contribution no. xx and Institut des Sciences de l'Evolution de Montpellier, University Montpellier 2 Contribution no (ISE-M 20xx-xxxx).



The publication of this article is financed by CNRS-INSU.

References

- Alcamo, J., Moreno, J. M., Novaky, B., Bindi, M., Corobov, R., Devoy, R. J. N., Giannakopoulos, C., Martin, E., Olesen, J. E., and Shvidenko, A.: Europe, in: *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Parry, M. P., Canziani, O. F., Palutikof, J. P., van der Linden, P. J., and Hanson, C. E., Cambridge University Press, Cambridge, 541–580, 2007.
- Almogi-Labin, A., Bar-Matthews, M., Shriki, D., Kolosovsky, E., Paterne, M., Schilman, B., Ayalon, A., Aizenshtat, Z., and Matthews, A.: Climatic variability during the last similar to 90 ka of the Southern and Northern Levantine Basin as evident from marine records and speleothems, *Quaternary Sci. Rev.*, 28, 2882–2896, 2009.

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Vegetation dynamics
in the Northeastern
Mediterranean region**

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Andersen, K. K., Svensson, A., Johnsen, S. J., Rasmussen, S. O., Bigler, M., Rothlisberger, R., Ruth, U., Siggaard-Andersen, M. L., Steffensen, J. P., Dahl-Jensen, D., Vinther, B. M., and Clausen, H. B.: The Greenland Ice Core Chronology 2005, 15–42 ka. Part 1: Constructing the time scale, *Quaternary Sci. Rev.*, 25, 3246–3257, 2006.
- 5 Arpe, K., Leroy, S. A. G., and Mikolajewicz, U.: A comparison of climate simulations for the last glacial maximum with three different versions of the ECHAM model and implications for summer-green tree refugia, *Clim. Past*, 7, 91–114, doi:10.5194/cp-7-91-2011, 2011.
- Atanassova, J.: Palaeoecological setting of the Western Black Sea area during the last 15 000 years, *Holocene*, 15, 576–584, 2005.
- 10 Bar-Matthews, M. and Ayalon, A.: Mid-Holocene climate variations revealed by high-resolution speleothem records from Soreq Cave, Israel and their correlation with cultural changes, *Holocene*, 21, 163–171, 2011.
- Bar-Matthews, M., Ayalon, A., and Kaufman, A.: Late quaternary paleoclimate in Eastern Mediterranean region from stable isotope analysis of speleothems at Soreq Cave, Israel, *Quaternary Res.*, 47, 155–168, 1997.
- 15 Bartov, Y., Goldstein, S. L., Stein, M., and Enzel, Y.: Catastrophic arid episodes in the Eastern Mediterranean linked with the North Atlantic Heinrich events, *Geology*, 31, 439–442, 2003.
- Behre, K.-E.: The interpretation of anthropogenic indicators in pollen diagrams, *Pollen et Spores*, 23, 225–245, 1981.
- 20 Behre, K.-E.: Some reflections on anthropogenic indicators and the record of prehistoric occupation phases in pollen diagrams from the Near East, in: *Man's Role in the Shaping of the Eastern Mediterranean Landscape*, edited by: Bottema, S., Entjes-Nieborg, G., van Zeist, W., Balkema, Rotterdam, 219–230, 1990.
- Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years, *Quaternary Sci. Rev.*, 10, 297–317, 1991.
- 25 Birks, H. J. B. and Heegaard, E.: Developments in age-depth modelling of Holocene stratigraphical sequences, *Pages News*, 11, 7–8, 2003.
- Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., deMenocal, P., Priore, P., Cullen, H., Hajdas, I., and Bonani, G.: A pervasive millennial-scale cycle in North Atlantic Holocene and Glacial Climates, *Science*, 278, 1257–1266, 1997.
- 30 Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M. N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I., and Bonani, G.: Persistent solar influence on North Atlantic climate during the Holocene, *Science*, 294, 2130–2136, 2001.

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bottema, S. and Sarpaki, A.: Environmental change in Crete: a 9000-yr record of Holocene vegetation history and the effect of the Santorini eruption, *Holocene*, 13, 733–749, 2003.
- Bottema, S. and Woldring, H.: Anthropogenic indicators in the pollen record of the Eastern Mediterranean, in: *Man's Role in the Shaping of the Eastern Mediterranean Landscape*, edited by: Bottema, S., Entjes-Nieborg, G., van Zeist, W., Balkema, Rotterdam, 231–264, 1990.
- Bottema, S., Woldring, H., and Kayan, I.: The late Quaternary vegetation history of Western Turkey, in: *The Ilipinar Excavations II*, edited by: Roodenberg, J. J., Thissen, L. C., Netherlands Instituut voor het Nabije Oosten, Leiden, 327–354, 2001.
- Boucher, K. (Ed.): *Global Climate*, The English Universities Press Ltd., London, 1974.
- Brayshaw, D. J., Rambeau, C. M. C., and Smith, S. J.: Changes in Mediterranean climate during the Holocene: insights from global and regional climate modelling, *Holocene*, 21, 15–31, 2011.
- Çagatay, M. N., Gorur, N., Algan, O., Eastoe, C., Tchapylyga, A., Ongan, D., Kuhn, T., and Kuscu, I.: Late Glacial-Holocene palaeoceanography of the Sea of Marmara: timing of connections with the Mediterranean and the Black Seas, *Mar. Geol.*, 167, 191–206, 2000.
- Combourieu Nebout, N., Peyron, O., Dormoy, I., Desprat, S., Beaudouin, C., Kotthoff, U., and Marret, F.: Rapid climatic variability in the west Mediterranean during the last 25 000 years from high resolution pollen data, *Clim. Past*, 5, 503–521, doi:10.5194/cp-5-503-2009, 2009.
- Dansgaard, W., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N. S., Hammer, C. U., Hvidbjerg, C. S., Steffensen, J. P., Sveinbjörnsdottir, A. E., Jouzel, J., and Bond, G.: Evidence for general instability of past climate from a 250-kyr ice-core record, *Nature*, 364, 218–220, 1993.
- Dormoy, I., Peyron, O., Combourieu Nebout, N., Goring, S., Kotthoff, U., Magny, M., and Pross, J.: Terrestrial climate variability and seasonality changes in the Mediterranean region between 15 000 and 4000 years BP deduced from marine pollen records, *Clim. Past*, 5, 615–632, doi:10.5194/cp-5-615-2009, 2009.
- Eastwood, W. J., Roberts, C. N., and Lamb, H. F.: Palaeoecological and archaeological evidence for human occupation in southwest Turkey: the Beysehir Occupation Phase, *Anatolian Studies*, 48, 69–86, 1998.
- Eriksen, U., Friedrich, W. L., Buchardt, B., Tauber, H., and Thomsen, M. S.: The Stronghyle caldera: geological, palaeontological and stable isotope evidence from radiocarbon-dated stromatolites from Santorini, in: *Thera and the Aegean World III. Volume Two: Earth Sci-*

**Vegetation dynamics
in the Northeastern
Mediterranean region**

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ences, edited by: Hardy, D. A., Keller, J., Galanopoulos, V. P., Flemming, N. C., and
Druitt, T. H., The Thera Foundation, London, 139–150, 1990.

Finsinger, W. and Tinner, W.: Minimum count sums for charcoal-concentration estimates in
pollen slides: reliability and potential errors, *Holocene*, 15, 293–297, 2005.

5 Fleitmann, D., Burns, S. J., Mangini, A., Mudelsee, M., Kramers, J., Villa, I., Neff, U., Al-
Subbary, A. A., Buettner, A., Hippler, D., and Matter, A.: Holocene ITCZ and Indian monsoon
dynamics recorded in stalagmites from Oman and Yemen (Socotra), *Quaternary Sci. Rev.*,
26, 170–188, 2007.

10 Fleitmann, D., Cheng, H., Badertscher, S., Edwards, R. L., Mudelsee, M., Göktürk, O. M.,
Fankhauser, A., Pickering, R., Raible, C. C., Matter, A., Kramers, J., and Tüysüz, O.: Timing
and climatic impact of Greenland interstadials recorded in stalagmites from Northern Turkey,
Geophys. Res. Lett., 36, L19707, doi:10.1029/2009GL040050, 2009.

Fletcher, W. J. and Sanchez Goni, M. F.: Orbital- and sub-orbital-scale climate impacts on
vegetation of the Western Mediterranean basin over the last 48 000 yr, *Quaternary Res.*, 70,
15 451–464, 2008.

Fletcher, W. J., Goni, M. F. S., Allen, J. R. M., Cheddadi, R., Combourieu-Nebout, N., Hunt-
ley, B., Lawson, I., Londeix, L., Magri, D., Margari, V., Muller, U. C., Naughton, F., Novenko, E.,
Roucoux, K., and Tzedakis, P. C.: Millennial-scale variability during the last glacial in vegeta-
tion records from Europe, *Quaternary Sci. Rev.*, 29, 2839–2864, 2010a.

20 Fletcher, W. J., Sanchez Goñi, M. F., Peyron, O., and Dormoy, I.: Abrupt climate changes of the
last deglaciation detected in a Western Mediterranean forest record, *Clim. Past*, 6, 245–264,
doi:10.5194/cp-6-245-2010, 2010b.

Genty, D., Blamart, D., Ghaleb, B., Plagnes, V., Causse, C., Bakalowicz, M., Zouari, K.,
Chkir, N., Hellstrom, J., Wainer, K., and Bourges, F.: Timing and dynamics of the last
25 deglaciation from European and North African $\delta^{13}\text{C}$ stalagmite profiles – comparison with
Chinese and South Hemisphere stalagmites, *Quaternary Sci. Rev.*, 25, 2118–2142, 2006.

Geraga, M., Tsaila-Monopolis, S., Ioakim, C., Papatheodorou, G., and Ferentinos, G.: Short-
term climate changes in the Southern Aegean Sea over the last 48 000 years, *Palaeogeogr.*
Palaeoclimatol., 220, 311–332, 2005.

30 Geraga, M., Ioakim, C., Lykousis, V., Tsaila-Monopolis, S., and Mylona, G.: The high-resolution
palaeoclimatic and palaeoceanographic history of the last 24 000 years in the Central
Aegean Sea, Greece, *Palaeogeogr. Palaeoclimatol.*, 287, 101–115, 2010.

- Giorgi, F. and Lionello, P.: Climate change projections for the Mediterranean region, *Global Planet. Change*, 63, 90–104, 2008.
- Göktürk, O. M., Fleitmann, D., Badertscher, S., Cheng, H., Edwards, R. L., Leuenberger, M., Fankhauser, A., Tüysüz, O., and Kramers, J.: Climate on the Southern Black Sea coast during the Holocene: implications from the Sofular Cave record, *Quaternary Sci. Rev.*, 30, 2433–2445, 2011.
- Grimm, E. C.: *Tilia* version 2.0 and *Tilia*Graph* 1.25, 1993.
- Heegaard, E., Birks, H. J. B., and Telford, R. J.: Relationships between calibrated ages and depth in stratigraphical sequences: an estimation procedure by mixed-effect regression, *Holocene*, 15, 612–618, 2005.
- Heusser, L. and Balsam, W. L.: Pollen Distribution in Northeast Pacific Ocean, *Quaternary Res.*, 7, 45–62, 1977.
- Hughen, K. A., Baillie, M. G. L., Bard, E., Warren Beck, J., Bertrand, C. J. H., Blackwell, P. G., Buck, C. E., Burr, G. S., Cutler, K. B., Damon, P. E., Edwards, R. L., Fairbanks, R. G., Friedrich, M., Guilderson, T. P., Kromer, B., McCormac, G., Manning, S., Ramsey, C. B., Reimer, P. J., Reimer, R. W., Remmele, S., Southon, J. R., Stuiver, M., Talamo, S., Taylor, F. W., van der Plicht, J., and Weyhenmeyer, C. E.: Marine04 marine radiocarbon age calibration, 0–26 cal kyr BP, *Radiocarbon*, 46, 1059–1086, 2004.
- Jones, M. D., Roberts, C. N., and Leng, M. J.: Quantifying climatic change through the last glacial–interglacial transition based on lake isotope palaeohydrology from Central Turkey, *Quaternary Res.*, 67, 463–473, 2007.
- Kazancı, N., Leroy, S., Ileri, Ö., Emre, Ö., Kibar, M., Öncel, S.: Late Holocene erosion in NW Anatolia from sediments of Lake Manyas, Lake Ulubat and the southern shelf of the Marmara Sea, *Turkey, Catena*, 57, 277–308, 2004.
- Köppen, W. (Ed.): *Die Klimate der Erde, Grundriss der Klimakunde*, De Gruyter, Berlin, 1923.
- Korfmann, M. (Ed.): *Troia – Archäologie eines Siedlungshügels und seiner Landschaft*, P. von Zabern, Mainz am Rhein, 2006.
- Kotthoff, U., Muller, U. C., Pross, J., Schmiedl, G., Lawson, I. T., van de Schootbrugge, B., and Schulz, H.: Lateglacial and Holocene vegetation dynamics in the Aegean region: an integrated view based on pollen data from marine and terrestrial archives, *Holocene*, 18, 1019–1032, 2008a.

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Vegetation dynamics
in the Northeastern
Mediterranean region**

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kotthoff, U., Pross, J., Muller, U. C., Peyron, O., Schmiedl, G., Schulz, H., and Bordon, A.: Climate dynamics in the borderlands of the Aegean Sea during formation of sapropel S1 deduced from a marine pollen record, *Quaternary Sci. Rev.*, 27, 832–845, 2008b.
- Kotthoff, U., Koutsodendris, A., Pross, J., Schmiedl, G., Bornemann, A., Kaul, C., Marino, G., Peyron, O., and Schiebel, R.: Impact of Lateglacial cold events on the Northern Aegean region reconstructed from marine and terrestrial proxy data, *J. Quaternary Sci.*, 26, 86–96, 2011.
- Kuhlemann, J., Rohling, E. J., Krumrei, I., Kubik, P., Ivy-Ochs, S., and Kucera, M.: Regional synthesis of Mediterranean atmospheric circulation during the last glacial maximum, *Science*, 321, 1338–1340, 2008.
- Kwiecien, O., Arz, H. W., Lamy, F., Wulf, S., Bahr, A., Rohl, U., and Haug, G. H.: Estimated reservoir ages of the Black Sea since the last glacial, *Radiocarbon*, 50, 99–118, 2008.
- Lainé, A., Kageyama, M., Salas-Méllia, D., Voltaire, A., Rivière, G., Ramstein, G., Planton, S., Tyteca, S., and Peterschmitt, J.: Northern Hemisphere storm tracks during the last glacial maximum in the PMIP2 ocean-atmosphere coupled models: energetic study, seasonal cycle, precipitation, *Clim. Dynam.*, 32, 593–614, 2009.
- Lamy, F., Arz, H. W., Bond, G. C., Bahr, A., and Patzold, J.: Multicentennial-scale hydrological changes in the Black Sea and Northern Red Sea during the Holocene and the Arctic/North Atlantic oscillation, *Paleoceanography*, 21, PA1008, doi:10.1029/2005pa001184, 2006.
- Lawson, I. T., Al-Omari, S., Tzedakis, P. C., Bryant, C. L., and Christanis, K.: Lateglacial and Holocene vegetation history at Nisi Fen and the Boras mountains, Northern Greece, *Holocene*, 15, 873–887, 2005.
- Leroy, S. A. G. and Arpe, K.: Glacial refugia for summer-green trees in Europe and South-West Asia as proposed by ECHAM3 time-slice atmospheric model simulations, *J. Biogeogr.*, 34, 2115–2128, 2007.
- Leroy, S., Kazanci, N., Ileri, O., Kibar, M., Emre, O., McGee, E., and Griffiths, H. I.: Abrupt environmental changes within a late Holocene lacustrine sequence south of the Marmara Sea (Lake Manyas, N-W Turkey): possible links with seismic events, *Mar. Geol.*, 190, 531–552, 2002.
- Leroy, S. A. G., Boyraz, S., and Gürbüz, A.: High-resolution palynological analysis in Lake Sapanca as a tool to detect recent earthquakes on the North Anatolian Fault, *Quaternary Sci. Rev.*, 28, 2616–2632, 2009.

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Londeix, L., Herreyre, Y., Turon, J.-L., and Fletcher, W.: Last Glacial to Holocene hydrology of the Marmara Sea inferred from a dinoflagellate cyst record, *Rev. Palaeobot. Palyno.*, 158, 52–71, 2009.
- 5 Maher, L. J.: Nomograms for computing 0.95 confidence limits of pollen data, *Rev. Palaeobot. Palyno.*, 32, 153–191, 1981.
- Marino, G., Rohling, E. J., Sangiorgi, F., Hayes, A., Casford, J. L., Lotter, A. F., Kucera, M., and Brinkhuis, H.: Early and middle Holocene in the Aegean Sea: interplay between high and low latitude climate variability, *Quaternary Sci. Rev.*, 28, 3246–3262, 2009.
- 10 Mayewski, P. A., Meeker, L. D., Twickler, M. S., Whitlow, S., Yang, Q. Z., Lyons, W. B., and Prentice, M.: Major features and forcing of high-latitude Northern Hemisphere atmospheric circulation using a 110 000-year-long glaciochemical series, *J. Geophys. Res.-Oceans*, 102, 26345–26366, 1997.
- Mix, A. C., Bard, E., and Schneider, R.: Environmental processes of the ice age: land, oceans, glaciers (EPILOG), *Quaternary Sci. Rev.*, 20, 627–657, 2001.
- 15 Moore, P. D., Webb, J. A., and Collinson, M. E. (Eds.): *Pollen Analysis*, Blackwell Scientific Publications, London, 1991.
- Mudie, P. J., Rochon, A., Aksu, A. E., and Gillespie, H.: Dinoflagellate cysts, freshwater algae and fungal spores as salinity indicators in Late Quaternary cores from Marmara and Black seas, *Mar. Geol.*, 190, 203–231, 2002.
- 20 Mudie, P. J., Marret, F., Aksu, A. E., Hiscott, R. N., and Gillespie, H.: Palynological evidence for climatic change, anthropogenic activity and outflow of Black Sea water during late Pleistocene and Holocene: centennial- to decadal-scale records from the Black and Marmara Seas, *Quaternary Int.*, 167, 73–90, 2007.
- Naughton, F., Goni, M. F. S., Desprat, S., Turon, J. L., Duprat, J., Malaize, B., Joli, C., Cortijo, E., Drago, T., and Freitas, M. C.: Present-day and past (last 25 000 years) marine pollen signal off Western Iberia, *Mar. Micropaleontol.*, 62, 91–114, 2007.
- 25 Özdogan, M.: Eastern Thrace the contact zone between Anatolia and the Balkans, in: *The Oxford Handbook of Anatolian Studies (10000-323BCE)*, edited by: Steadman, S. and McMahon, G., Oxford University Press, Oxford, UK, 898–923, 2011.
- 30 Pichler, H. and Friedrich, W.: Radiocarbon dates of santorini volcanics, *Nature*, 262, 373–374, 1976.

**Vegetation dynamics
in the Northeastern
Mediterranean region**

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pross, J., Kotthoff, U., Muller, U. C., Peyron, O., Dormoy, I., Schmiedl, G., Kalaitzidis, S., and Smith, A. M.: Massive perturbation in terrestrial ecosystems of the Eastern Mediterranean region associated with the 8.2 kyr BP climatic event, *Geology*, 37, 887–890, 2009.
- Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard-Andersen, M.-L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., and Ruth, U.: A new Greenland ice core chronology for the last glacial termination, *J. Geophys. Res.-Atmos.*, 111, D06102, doi:10.1029/2005JD006079, 2006.
- Reille, M. (Ed.): *Pollen et spores d'Europe et d'Afrique du nord*, Supplement 1 and 2, Laboratoire de Botanique historique et Palynologie, Marseille, 1992–1998.
- Reimer, P. J., Baillie, M. G. L., Bard, E., Bayliss, A., Warren Beck, J., Bertrand, C. J. H., Blackwell, P. G., Buck, C. E., Burr, G. S., Cutler, K. B., Damon, P. E., Lawrence Edwards, R., Fairbanks, R. G., Friedrich, M., Guilderson, T. P., Hogg, A. G., Hughen, K. A., Kromer, B., McCormac, G., Manning, S., Ramsey, C. B., Reimer, R. W., Remmele, S., Southon, J. R., Stuiver, M., Talamo, S., Taylor, F. W., van der Plicht, J., and Weyhenmeyer, C. E.: IntCal04 terrestrial radiocarbon age calibration, 0–26 calkyr BP, *Radiocarbon*, 46, 1029–1058, 2004.
- Riehl, S. and Marinova, E.: Mid-Holocene vegetation change in the Troad (W Anatolia): man-made or natural?, *Veg. Hist. Archaeobot.*, 17, 297–312, 2008.
- Roberts, N. and Wright Jr., H. E.: Vegetational, lake-level, and climatic history of the Near East and Southwest Asia, in: *Global Climates Since the past Last Glacial Maximum*, edited by: Wright, H. E. J., Kutzbach, J. E., Webb III, T., Ruddiman, W. F., Street-Perrott, F. A., and Bardein, P. J., University of Minneapolis Press, Minneapolis, 194–220, 1993.
- Roberts, N., Reed, J. M., Leng, M. J., Kuzucuoglu, C., Fontugne, M., Bertaux, J., Woldring, H., Bottema, S., Black, S., Hunt, E., and Karabiyyikoglu, M.: The tempo of Holocene climatic change in the Eastern Mediterranean region: new high-resolution crater-lake sediment data from Central Turkey, *Holocene*, 11, 721–736, 2001.
- Roberts, N., Eastwood, W. J., Kuzucuoğlu, C., Fiorentino, G., and Caracuta, V.: Climatic, vegetation and cultural change in the Eastern Mediterranean during the mid-Holocene environmental transition, *Holocene*, 21, 147–162, 2011.
- Roberts, N., Moreno, A., Valero-Garcés, B. L., Corella, J. P., Jones, M., Allcock, S., Woodbridge, J., Morellón, M., Luterbacher, J., Xoplaki, E., and Türkeş, M.: Palaeolimnological evidence for an east–west climate see-saw in the Mediterranean since AD 900, *Global Planet. Change*, 84–85, 23–34, 2012.

**Vegetation dynamics
in the Northeastern
Mediterranean region**

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Rohling, E. J., Mayewski, P. A., Abu-Zied, R. H., Casford, J. S. L., and Hayes, A.: Holocene atmosphere-ocean interactions: records from Greenland and the Aegean Sea, *Clim. Dynam.*, 18, 587–593, 2002.
- Roodenberg, J. J. and Alpaslan-Roodenberg, S.: The Neolithic in the Eastern Marmara: two examples of settlement, in: *Neolithic in Turkey*, edited by: Ozdogan, M. and Basgelen, N., Arkeoloji ve Sanat Yayinlari, Istanbul, 1–7, 1999.
- Rosenstock, E.: Höyük, Toumba and Mogila: a settlement form in Anatolia and the Balkans and its ecological determination 6500–5500 BC, in: *How did Farming Reach Europe?*, edited by: Lichter, C., Bizas, 2, 221–237, 2005.
- Rosignol-Strick, M.: Sea-land correlation of pollen records in the Eastern Mediterranean for the glacial-interglacial transition: biostratigraphy versus radiometric time-scale, *Quaternary Sci. Rev.*, 14, 893–915, 1995.
- Rosignol-Strick, M. and Planchais, N.: Climate patterns revealed by pollen and oxygen isotope records of a Tyrrhenian sea core, *Nature*, 342, 413–416, 1989.
- Rull, V.: A note on pollen counting in palaeoecology, *Pollen et Spores*, 29, 471–480, 1987.
- Saaroni, H., Bitan, A., Alpert, P., and Ziv, B.: Continental Polar outbreaks into the Levant and Eastern Mediterranean, *Int. J. Climatol.*, 16, 1175–1191, 1996.
- Sanchez Goñi, M. F. and Harrison, S. P.: Millennial-scale climate variability and vegetation changes during the Last Glacial: concepts and terminology, *Quaternary Sci. Rev.*, 29, 2823–2827, 2010.
- Shumilovskikh, L. S., Tarasov, P., Arz, H. W., Fleitmann, D., Marret, F., Nowaczyk, N., Plessen, B., Schlütz, F., and Behling, H.: Vegetation and environmental dynamics in the Southern Black Sea region since 18 kyr BP derived from the marine core 22-GC3, *Palaeogeogr. Palaeoclimatol.*, 337–338, 177–193, 2012.
- Siani, G., Paterne, M., Arnold, M., Bard, E., Métiévier, B., Tisnerat, N., and Bassinot, F.: Radiocarbon reservoir ages in the Mediterranean Sea and Black Sea, *Radiocarbon*, 42, 271–280, 2000.
- Stanford, J. D., Rohling, E. J., Bacon, S., Roberts, A. P., Grousset, F. E., and Bolshaw, M.: A new concept for the paleoceanographic evolution of Heinrich event 1 in the North Atlantic, *Quaternary Sci. Rev.*, 30, 1047–1066, 2011.
- Steffensen, J. P., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Fischer, H., Goto-Azuma, K., Hansson, M., Johnsen, S. J., Jouzel, J., Masson-Delmotte, V., Popp, T., Rasmussen, S. O., Röthlisberger, R., Ruth, U., Stauffer, B., Siggaard-Andersen, M.-L.,

**Vegetation dynamics
in the Northeastern
Mediterranean region**

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sveinbjörnsdóttir, À., Svensson, A., and White, J. W. C.: High-resolution Greenland ice core data show abrupt climate change happens in few years, *Science*, 321, 680–684, 2008.

Stevens, L. R., Wright Jr., H. E., and Ito, E.: Proposed changes in seasonality of climate during the Lateglacial and Holocene at Lake Zeribar, Iran, *Holocene*, 11, 747–755, 2001.

5 Stockmarr, J.: Tablets with spores used in absolute pollen analysis, *Pollen et Spores*, 13, 614–621, 1971.

Stuiver, M., Reimer, P. J., Bard, E., Beck, J. W., Burr, G. S., Hughen, K. A., Kromer, B., McCormac, F. G., van der Plicht, J., and Spurk, M.: INTCAL98 Radiocarbon Age Calibration, 24 000–0 cal BP, *Radiocarbon*, 40, 1041–1083, 1998.

10 Trewartha, G. T. (Ed.): *The Earth's Problem Climates*, The University of Wisconsin Press, Madison, 1981.

Turner, C. and Sánchez Goñi, M.-F.: Late glacial landscape and vegetation in Epirus, in: *Klithi: Palaeolithic Settlement and Quaternary Landscapes in Northwest Greece*, edited by: Bailey, G., McDonald Institute for Archaeological Research, Cambridge, 559–585, 1997.

15 Turon, J. L.: Direct land sea correlations in the Last Interglacial Complex, *Nature*, 309, 673–676, 1984.

Tzedakis, P. C.: Seven ambiguities in the Mediterranean palaeoenvironmental narrative, *Quaternary Sci. Rev.*, 26, 2042–2066, 2007.

20 Tzedakis, P. C., Frogley, M. R., Lawson, I. T., Preece, R. C., Cacho, I., and de Abreu, L.: Ecological thresholds and patterns of millennial-scale climate variability: The response of vegetation in Greece during the last glacial period, *Geology*, 32, 109–112, 2004.

Unal, Y., Kindap, T., and Karaka, M.: Redefining the climate zones of Turkey using cluster analysis, *Int. J. Climatol.*, 23, 1045–1055, 2003.

25 Ünlüata, Ü., Öguz, T., Latif, M., and Ozsoy, E.: On the physical oceanography of the Turkish Straits, in: *The Physical Oceanography of Sea Straits*, NATO ASI Ser., Ser. C, Vol. 318, edited by: Pratt, L. J., Kluwer Accademy, Dordrecht, 1990.

Vermoere, M., Bottema, S., Vanhecke, L., Waelkens, M., Paulissen, E., and Smets, E.: Palynological evidence for late-Holocene human occupation recorded in two wetlands in SW Turkey, *Holocene*, 12, 569–584, 2002.

30 Vidal, L., Ménot, G., Joly, C., Bruneton, H., Rostek, F., Çagatay, M. N., Major, C., and Bard, E.: Hydrology in the Sea of Marmara during the last 23 ka: implications for timing of Black Sea connections and sapropel deposition, *Paleoceanography*, 25, PA1205, doi:10.1029/2009PA001735, 2010.

- Vinther, B. M., Clausen, H. B., Johnsen, S. J., Rasmussen, S. O., Andersen, K. K., Buchardt, S. L., Dahl-Jensen, D., Seierstad, I. K., Siggaard-Andersen, M.-L., Stefansen, J. P., Svensson, A., Olsen, J., and Heinemeier, J.: A synchronized dating of three Greenland ice cores throughout the Holocene, *J. Geophys. Res.-Atmos.*, 111, doi:10.1029/2005JD006921, 2006.
- Weiss, H., Courty, M.-A., Wetterstrom, W., Guichard, F., Senior, L., Meadow, R., and Curnow, A.: The genesis and collapse of Third Millennium North Mesopotamian civilization, *Science*, 261, 995–1004, 1993.
- Wick, L., Lemcke, G., and Sturm, G.: Evidence of Lateglacial and Holocene climatic change and human impact in Eastern Anatolia: high-resolution pollen, charcoal, isotopic and geochemical records from the laminated sediments of Lake Van, Turkey, *Holocene*, 13, 665–675, 2003.
- Wijmstra, T. A.: Palynology of the first 30 meters of a 120 m deep section in Northern Greece, *Acta Bot. Neerl.*, 18, 511–527, 1969.
- Willis, K. J.: The pollen-sedimentological evidence for the beginning of agriculture in Southeastern Europe and Anatolia, *Porocilo o raziskovanju paleolitika, neolitika in eneolitika v Sloveniji*, 22, 9–24, 1995.
- Wulf, S., Kraml, M., Kuhn, T., Schwarz, M., Inthorn, M., Keller, J., Kuscu, I., and Halbach, P.: Marine tephra from the Cape Riva eruption (22 ka) of Santorini in the Sea of Marmara, *Mar. Geol.*, 183, 131–141, 2002.
- van Zeist, W. and Bottema, S. (Eds.): *Late Quaternary Vegetation of the Near East*, Dr. Ludwig Reichert Verlag, Wiesbaden, 1991.
- Zohary, M. (Ed.): *Geobotanical foundations of the Middle East*, 2 Volumes, Gustav Fischer Verlag and Swets and Zeitlinger, Stuttgart, Amsterdam, 1973.

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Table 1. AMS-radiocarbon dates on core MD01-2430.

Laboratory reference	Depth (cm)	Material dated	¹⁴ C yr BP	Reservoir correction	Calendar age cal yr BP (1 sd)
OS-35404	200	<i>Scrobicularia plana</i>	4780 ± 55	400 ^a	4952–5187
OS-40506	253	<i>Scrobicularia plana</i>	8010 ± 60	400 ^a	8402–8528
OS-40513	337	<i>Arca</i> sp.	10850 ± 65	400 ^a	12230–12601
OS-40514	347	<i>Arca</i> sp.	11050 ± 65	400 ^a	12657–12803
OS-40515	349	<i>Arca</i> sp.	11050 ± 70	400 ^a	12647–12807
OS-40516	384	<i>Turricaspia caspia</i>	13050 ± 75	400 ^a	14742–15066
Y 2/Cape Riva Tephra	691–698	wood	18310 ± 380 ^b	–	21262–22230

^a Reservoir correction following Siani et al. (2000).

^b Mean and standard deviation calculated by Kwiecien et al. (2008) based on original dating series (Eriksen et al., 1990; Pichler and Friedrich, 1976).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.




Fig. 1. Location of the sites discussed. **(a)** NGRIP ice core, North Atlantic (MC52 and VM29-191) and the Near East (speleothem from Soreq Cave). The dashed arrow indicates the Western riding of the Siberian High during winter/spring (Marino et al., 2009). **(b)** Highlight of the sites located in Eastern Mediterranean: sediment core in the Sea of Marmara (MD01-2430), marine cores in the Aegean Sea (SL152 and MNB3), marine core in the Black Sea (GeoB 7622-2), continental core in Greece (Tenaghi Philippon) and speleothem from Sofular Cave. Some archaeological sites are indicated by square. Map **(c)** indicates seasonal rainfall and the movement of depression within the Mediterranean (modify by, Boucher, 1974). Seasons of maximum rainfall are shown only for Turkey and Near East: W (winter), A (autumn), S (spring) and X (indistinct). Figures in the lows indicate annual mean frequency of depressions.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Vegetation dynamics
in the Northeastern
Mediterranean region**

V. Valsecchi et al.




Fig. 2. Depth-age model (line) and 95.4 % confidence intervals (dashed lines). Calibrated radiocarbon dates are represented by dots. Depth-age model by Vidal et al. (2010) is in grey.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.




Fig. 3a. Pollen percentage diagram for core MD01-2430 showing only selected taxa. *Pinus*, aquatics, ferns and reworked pollen are not included in the pollen sum. Mediterranean trees include *Quercus* evergreen type, *Cistus*, *Olea*, *Phillyrea* and *Pistacia*. Steppic plants include *Artemisia*, *Chenopodiaceae*, *Ephedra distachya* type and *E. fragilis* type. Filled black curves show the percentage values of single pollen type; white curves indicate percentages enlarged by ten.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.




Fig. 3b. Pollen influx diagram for core MD01-2430 showing only selected taxa. Gray band indicates the Sapropel 1 (S1).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Pollen percentage, dinocysts percentage and sea surface temperature (SST) for core MD01-2430. Arboreal taxa include all trees and shrubs except *Hippophaë*, *Ephedra distachya* type and *E. fragilis* type. Steppic includes *Artemisia*, *Chenopodiaceae*, *Ephedra distachya* type and *E. fragilis* type. Alkenone derived sea surface temperatures (SST) (Vidal et al., 2010). Thermophilic dinocysts include: *Tuberculodinium vancampoae*, *Tectatodinium pellitum*, *Spiniferites mirabilis*, *Spiniferites delicatus*, *Lingulodinium machaerophorum*, *Polysphaeridium zoharyi*, *Operculodinium israelianum*, and *Stelladinium stellatum*; cool dinocysts include: *Pentaparthosodinium dalei*, *Bitectatodinium tepikiense* s.l., *Caspidinium rugosum* s.l. and *Spiniferites elongatus* s.l. (Londeix et al., 2009). Numbers from 1 to 3 indicate forest set-backs. Grey boxes indicate major climatic shifts: LGM (Last Glacial Maximum) which last from 23–19 cal ka (Mix et al., 2001), HS1 (Heinrich Stadial 1) from ca. 19–15 cal ka (Sanchez Goñi and Harrison, 2010; Stanford et al., 2011), YD (Younger Dryas or GS-1) from ca. 12.7–11.7 cal ka (Steffensen et al., 2008) and S1 from ca. 7–11.5 cal ka (Vidal et al., 2010).

Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.




Fig. 5. Caption on next page.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.

Fig. 5. Selected palaeoclimate records arranged by latitude (north on the top) and paleo proxies for core MD01-2430. **(a–b)** summer and winter insolation values at 30° N, Berger and Loutre, 1991). **(c)** NGRIP $\delta^{18}\text{O}$ (‰ VSMOW) proxy for temperature based on GICC05 timescale (Andersen et al., 2006; Rasmussen et al., 2006; Vinther et al., 2006), black solid curve represents 200 yr (span 0.1) running average smoothing. **(d)** GIPS2 potassium ion (ppb) proxy for the SH (Mayewski et al., 1997), black solid curve represents 200 yr (span 0.01) running average smoothing. **(e)** Ice-rafted hematite-stained grains in the North Atlantic sediment cores MC52 and VM29-191. Numbers from 0 to 8 designate the millennial-scale cycles (Bond et al., 1997, 2001). **(f)** Clay layer frequency record from core GeoB 7622-2 (Black Sea) proxy for winter precipitation in Northern Anatolia (Lamy et al., 2006). **(g)** $\delta^{13}\text{C}$ (‰ VPDB) of stalagmite So-1 from Sofular Cave (Turkey) proxy for type and density of vegetation: values of -12‰ are characteristic for C_3 plants (trees and shrubs) and values of -6‰ for C_4 (grasses and chenopods) plants (Fleitmann et al., 2009), black solid curve represents 200 yr (span 0.01) running average smoothing. **(h)** Arboreal pollen type (%) including temperate and mediterranean plants for core MD01-2430 (this study), numbers from 1 to 3 indicate forest set-backs, B/A and HS1 indicate biostratigraphic zone based on our data. **(i)** Alkenone derived sea surface temperatures (SST) on the core MD01-2430 (Vidal et al., 2010). **(j)** $\delta^{18}\text{O}$ (‰ VPDB) of Soreq Cave (Israel) (Almogi-Labin et al., 2009). For the legend of grey boxes LGM, HS1 and YD see Fig. 4.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vegetation dynamics in the Northeastern Mediterranean region

V. Valsecchi et al.




Fig. 6. Pollen percentage diagram showing selected taxa for core MD01-2430. Filled black curves show the percentage values of single pollen type; white curves indicate percentages enlarged by ten. Gray bands indicate phases of forest withdrawal for anomalies above 20%. Archaeological periods and most important events in the past societies for Northern Anatolia region are shown on the right-side end.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

