

**Iron fluxes to Talos Dome, Antarctica,  
over the past 200 kyr**

P. Vallelonga et al.

**Iron fluxes to Talos Dome, Antarctica,  
over the past 200 kyr**

**P. Vallelonga<sup>1,2</sup>, C. Barbante<sup>2,3,4</sup>, G. Cozzi<sup>2</sup>, J. Gabrieli<sup>2</sup>, S. Schüpbach<sup>3,5</sup>,  
A. Spolaor<sup>3</sup>, and C. Turetta<sup>2</sup>**

<sup>1</sup>Centre for Ice and Climate, Niels Bohr Institute, University of Copenhagen,  
Juliane Maries Vej 30, 2100 Copenhagen, Denmark

<sup>2</sup>Institute for the Dynamics of Environmental Processes (IDPA) – CNR, University of Venice,  
Dorsoduro 2137, 30123 Venice, Italy

<sup>3</sup>Department of Environmental Sciences, University Ca' Foscari of Venice, Dorsoduro 2137,  
30123 Venice, Italy

<sup>4</sup>Accademia Nazionale dei Lincei, Centro B. Segre, via della Lungara 10, 00165, Rome, Italy

<sup>5</sup>Physics Institute, Climate and Environmental Physics, University of Bern, Sidlerstrasse 5,  
3012, Bern, Switzerland

Received: 23 November 2012 – Accepted: 27 November 2012 – Published: 5 December 2012

Correspondence to: P. Vallelonga (ptravis@nbi.ku.dk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Atmospheric fluxes of iron (Fe) over the past 200 kyr are reported for the coastal Antarctic Talos Dome ice core, based on acid leachable Fe concentrations. Fluxes of Fe to Talos Dome were consistently greater than those at Dome C, with the greatest difference observed during interglacial climates. We observe different Fe flux trends at Dome C and Talos Dome during the deglaciation and early Holocene, attributed to a combination of deglacial activation of dust sources local to Talos Dome and reorganization of atmospheric transport pathways with the retreat of the Ross Sea ice shelf. This supports similar findings based on dust particle sizes and fluxes and Rare Earth Element fluxes. We show that Ca and Fe should not be used as quantitative proxies for mineral dust, as they all demonstrate different deglacial trends at Talos Dome and Dome C. Considering that a 20 ppmv decrease in atmospheric CO<sub>2</sub> at the coldest part of the last glacial maximum occurs contemporaneously with the period of greatest Fe and dust flux to Antarctica, we conclude that the maximum contribution of aeolian dust deposition to Southern Ocean sequestration of atmospheric CO<sub>2</sub> is approximately 20 ppmv.

## 1 Introduction

Polar ice cores allow detailed reconstructions of atmospheric composition and aerosol loading which are important for understanding the origins of climate transitions as well as potential feedback effects (Wolff et al., 2006; Fischer et al., 2010). Mineral aerosol (dust) plays a role in several direct and indirect climate feedback processes (Maher et al., 2010) with extremely large (up to 40-fold) changes in dust fluxes at polar and high-latitude regions between glacial and interglacial climates (Fischer et al., 2007). In Central Antarctica, increased dust fluxes arise due to stronger atmospheric coupling with the arid dust-deflation zones of Southern South America in combination with reduced washout of dust (Lambert et al., 2008). Isotopic (Delmonte et al., 2010;

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr**

P. Vallelonga et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Vallelonga et al., 2010) and modelling (Li et al., 2008; Mahowald et al., 2005) investigations indicate changes in dust provenance from glacial to interglacial climates, with a dominant Southern South American dust source during the glacial maxima and enhanced dust entrainment from Australia and local Antarctic ice-free areas during interglacial periods. Marine sediments demonstrate that coupling between Southern South American dust sources and meridional dust transport intensified during the late Pleistocene (Martinez-Garcia et al., 2009).

Records of past atmospheric fluxes of dust and elements such as iron (Fe) are essential for testing the “Iron hypothesis” proposed by Martin (1990), in which CO<sub>2</sub> draw-down in some oceanic zones is controlled not by the availability of macronutrients, such as nitrogen and phosphorus, but by the availability of micronutrients essential for biological growth, particularly Fe. Successful trials of ocean surface Fe addition in High-Nutrient Low-Chlorophyll (HNLC) zones of the Southern and equatorial Pacific Oceans have validated the basis for the Iron hypothesis (Sigman et al., 2010; Smetacek et al., 2012). A complementary mechanism for the transport of nutrients to HNLC zones is wind-driven upwelling of nutrient-rich deep waters: Anderson et al. (2009) identified changes in opal fluxes from a Southern Ocean sediment core which may have been due to shifts in the latitude of Southern westerlies during the last termination ( $T_1$ ). Such shifts consequently alter Southern ocean circulation, the strength of abyssal upwelling and nutrient transport to the subantarctic oceanic zone. Syntheses of terrestrial, marine and ice core dust data are essential for the contribution of aeolian dust deposition to Fe fertilization to be accurately evaluated (Maher et al., 2010).

Records of aeolian Fe inputs to Antarctica are available for two sites: EPICA Dome C (EDC) (Gaspari et al., 2006; Wolff et al., 2006) on the central Eastern plateau and coastal Law Dome (Edwards et al., 2006), showing marked differences in glacial-interglacial Fe fluxes and ratios. EDC samples were leached at pH 1 for > 24 h (identical to this study) while the Law Dome samples were acidified for 1 month, thus allowing for more complete dissolution of acid-labile Fe. An early evaluation of oceanic dust fertilization employed non-sea salt Ca<sup>2+</sup> (nssCa<sup>2+</sup>) as a proxy for Fe (Röthlisberger

et al., 2004). Consequently, the biological pump has been calculated to account for 10 to 50 % of the 80–100 ppmv CO<sub>2</sub> changes observed over glacial-interglacial transitions (Martinez-Garcia et al., 2009; Fischer et al., 2010). Here we present a record of Fe fluxes at Talos Dome (TD) for the past 200 kyr to assess atmospheric dust deposition controls on Southern Ocean paleoproductivity.

## 2 Experimental

Samples were obtained from the TALDICE ice core (159°11' E, 72°49' S; altitude 2315 m a.s.l.; annual mean temperature -41 °C; snow accumulation 80 mm water equivalent yr<sup>-1</sup>; Stenni et al., 2011). The TALDICE-1a chronology (Buiron et al., 2011; Schüpbach et al., 2011) has an uncertainty of 300 yr over T<sub>1</sub> and less than 600 yr over MIS 3. Discrete samples were obtained from a 32 mm × 32 mm section of the inner part of the ice core following Continuous Flow Analysis (Kaufmann et al., 2008). Ice was melted on a gold-coated brass melthead at a rate of 3 cm min<sup>-1</sup>, with discrete samples collected in coulter counter accuvettes at a rate of 0.5 mL min<sup>-1</sup>. Each 15 mL sample integrated a metre of melted ice. Samples were transported frozen to Italy for analysis.

The sample preparation and analytical methods have been reported previously (Barbante et al., 1999). Samples were prepared in a shipping container converted into a particle-free laboratory, with HEPA-filtered overpressured working spaces rated to Class 10 to 100 (US Fed. Std 209E, equivalent to ISO 4 to 5). Melted samples were acidified to pH 1 using sub-boiling distilled HNO<sub>3</sub> (Romil, Cambridge, UK) and analysed at least 24 h later by Inductively Coupled Plasma Sector Field Mass Spectrometry (ICP-SFMS; Finnigan™ ELEMENT2, Thermo Fisher Scientific Inc., Bremen, Germany) coupled to an APEX Q desolvating introduction unit (Elemental Scientific, Omaha, NE, USA). The detection limit, defined as three times the standard deviation of the instrument blank, was 0.029 ngg<sup>-1</sup>, with precisions of 16 % for Holocene samples and 10 % for Last Glacial Maximum (LGM) samples. We report here acid leachable Fe

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concentrations, following the techniques described by Gaspari et al. (2006), in order to compare Talos Dome Fe fluxes directly to those reported for EPICA Dome C. We acknowledge that the Fe fluxes reported here underrepresent total Fe in the samples, as it has been demonstrated that at least one month of acidification is required to completely dissolve Fe-bearing aerosol particles in ice (Edwards et al., 2006). Fluxes were calculated by multiplying Fe concentrations by TALDICE-1a accumulation rates (uncertainty  $\pm 20\%$ ). The time period integrated by each sample varies from about 15 y at 0.6 kyr before present (BP, 1950 CE) to 30 yr at 15 kyrBP and 700 yr by MIS 6 (150 kyrBP). To highlight the trends shown in the figures, we show running average smoothing filters overlaying the data.

### 3 Results and discussion

TALDICE Fe fluxes are shown in Fig. 1, demonstrating a well-described pattern of higher dust and Fe concentrations during colder climate periods, and lower dust and Fe fluxes during warmer climate periods (Wolff et al., 2006). Despite the reduced sampling frequency, it is clearly seen that Fe and dust fluxes to TD and EDC during MIS 6 were less than those of MIS 2. Consistently greater interglacial Fe concentrations were observed at TD compared to EDC during the last two interglacial periods. In addition to greater interglacial Fe fluxes at TD, there was greater variability in the Fe and dust fluxes to TD over the interglacials, compared to EDC. In Table 1, TD Fe concentrations and fluxes are compared with other Antarctic sites for which Fe fluxes are available. These show that Holocene fluxes of Fe at the coastal sites of Talos Dome and Law Dome are an order of magnitude higher than at Dome C, although LGM fluxes are of the same order of magnitude. For Talos Dome, the elevated interglacial dust flux has been attributed to the deglacial activation of proximal dust deflation zones in Victoria Land (Albani et al., 2012), which will be discussed below. For Law Dome, dust fluxes or particle size distributions have not been reported, so the source(s) of interglacial dust cannot be speculated upon. The deposition of dust from local (Antarctic) sources

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



at Law Dome cannot be discounted, as locally-sourced dust has also been observed at Berkner Island, based on the analysis of Sr and Nd isotopic compositions in recent snowpack (Bory et al., 2010).

Deglacial variability of Fe, dust and non-sea salt (nss)  $\text{Ca}^{2+}$  fluxes are shown in Fig. 2, showing a clear decoupling of coastal Antarctica (TD) from Central Antarctica (EDC) from 20 to 10 kyr BP. As mentioned previously, fluxes of these parameters were similar during the LGM, but have been consistently greater at TD since the deglaciation. This deglacial decoupling trend was first observed by Delmonte et al. (2010) for TD dust fluxes and particle size distributions and was attributed to the activation of local (Victoria Land) dust deflation areas during the deglaciation. Albani et al. (2012) reported TD dust particle size distributions at greater temporal resolution (fine/coarse particle size ratios are also shown in Fig. 2), finding a similar deglacial decoupling, which was then followed by a convergence of TD and EDC dust fluxes since 8 kyr BP, as a result of changes to atmospheric transport patterns around the Ross Sea embayment. The retreat of the Ross ice shelf grounding line appears to have led to a preference for the transport of air masses over the Ross Sea, which is not conducive to the deflation of dust in high-altitude areas of Victoria Land.

Dust fluxes to Talos Dome and EDC diverge during transition  $T_1$ , a period of large-scale atmospheric reorganization across Antarctica and the sub-Antarctic region. Post-glacial changes in atmospheric transport and/or provenance have been observed for dust particle sizes (Delmonte et al., 2004) and dust components such as Rare Earth Elements (Wegner et al., 2012; Gabrielli et al., 2010) and Pb isotopes (Vallelonga et al., 2010). The Talos Dome dust record does not demonstrate the “pre-Holocene dust minimum” observed in EDC, DB and Komosmolskaia ice cores in Central East Antarctica (Delmonte et al., 2004), suggesting that different atmospheric transport regimes develop in these different sectors of Antarctica as interglacial conditions develop. Atmosphere-ocean coupled models, supported by stable water isotope data, indicate that the various sectors of Antarctica have differing responses to rapid climate variations such as the deglacial (Buiron et al., 2012). Further, systematic variations

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



have been observed for Rare Earth Element compositions in EDC ice which have been attributed to changing dust provenance during and after the deglaciation (Gabielli et al., 2010).

Such variability can also be observed in the deglacial Fe flux record (Fig. 2), as a distinctive Fe peak in the Antarctic Cold Reversal (ACR) observed at EDC (Gaspari et al., 2006) between 13.0 and 12.7 kyrBP is not present at TD. A similar-size Fe peak is observed later at TD from 12.1 to 11.8 kyrBP. The core chronologies in this period have been synchronized by CH<sub>4</sub> tiepoints, so there is no possibility of attributing this to a misalignment or a dating error. Instead, it appears that these Fe peaks reflect different dust histories for different sectors of Antarctica over the deglacial. Figure 1 shows greater TD Fe/dust ratios during the Holocene compared to the glacial, suggestive of a change in TD dust sources from the LGM to the Holocene.

The high resolution of the TALDICE record also allows centennial-scale variations to be evaluated over the Holocene, as shown in Fig. 3. Iron fluxes varied between higher ( $> 10^{-1} \text{ mg m}^{-2} \text{ a}^{-1}$ ) and lower ( $< 5 \times 10^{-2} \text{ mg m}^{-2} \text{ a}^{-1}$ ) values throughout the Holocene, with highest values during the mid-Holocene (7–10 kyrBP). The period of higher Fe fluxes from 7–10 kyrBP was not matched by consistent changes in dust flux or Fe/dust ratio although there was some tendency toward a greater proportion of fine dust particles at Talos Dome between 7 and 10 kyrBP (Albani et al., 2012). In contrast, there was a consistent change in dust and Fe fluxes and dust size fractions after 7 kyrBP (Fig. 3), which coincided with the retreat of the Ross ice shelf and resulting shifts in regional atmospheric transport patterns away from dust deflation trajectories (Albani et al., 2012). After 7 kyrBP, there are clear decreases in Fe and dust fluxes to Talos Dome, as well as tendency toward deposition of coarser dust. The TD Holocene Fe record displays many spikes with up to 6-fold flux variations in adjacent samples, attributed to the influence of large dust particles emitted from local sources. Delmonte et al. (2010) observed that 5–20  $\mu\text{m}$  diameter dust particles account for 60 to 70 % of total dust mass during the Holocene. By comparison, large particles constitute less than 20 % of total TD dust mass during MIS 2.

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr**

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 2 also shows clearly that deglacial trends of  $\text{nssCa}^{2+}$ , dust and Fe have many individual features on millennial to centennial timescales, which inhibit their applicability as proxies for each other. It is recognized that the common terrestrial sources of dust, Fe and  $\text{nssCa}^{2+}$  is the cause for their similar atmospheric flux patterns over glacial-interglacial time scales, but such commonality cannot be assumed for shorter time periods (Wolff et al., 2006). Comparing the deglacial EDC dust trend to Fe and  $\text{nssCa}^{2+}$  at EDC, we see that the dust minimum at 11.9 kyrBP is not mirrored by minima in Fe or  $\text{nssCa}^{2+}$ . Similarly, deglacial peaks in Fe at EDC (12.9 kyrBP) and Talos Dome (11.9 kyrBP) occur in the absence of corresponding features in deglacial dust fluxes. Some noise may be induced in the  $\text{nssCa}^{2+}$  record because it relies on  $\text{Cl}^-$  fluxes to calculate the contribution of Ca from marine salts. Dust/Fe ratios shown in Fig. 1 change from 0.03 to 0.1, demonstrating that also Fe should not be assumed a 1-to-1 proxy of dust. Variability in dust/Fe ratios could be due to variations in dust sources as well as Fe solubility, both of which are dependent upon dust mineralogy. These examples are sufficient to demonstrate that interpretations should be made cautiously when using one of dust, Fe or  $\text{nssCa}^{2+}$  as a proxy for another, especially when considering changes on millennial or centennial time scales.

In the absence of Fe data, Rothlisberger et al. (2004) evaluated the response of the biological pump to varying dust fluxes by considering changes in EDC  $\text{nssCa}^{2+}$  over the period 40–60 kyrBP. They observed that large  $\text{nssCa}^{2+}$  variations coincided with 20 ppmv  $\text{CO}_2$  variations, and used such a scenario to conclude that changes in Fe fertilization could account for at most 20 ppmv of  $\text{CO}_2$  drawdown. We apply a similar approach for the LGM, but take advantage of the Fe flux records now available for EDC and TD (Fig. 4) and investigate a time period of much greater Fe fluxes. We note that the atmospheric concentration of  $\text{CO}_2$  decreases from 212 ppmv (35–40 kyrBP) to 190.6 ppmv (17–27 kyrBP) at the same time as dust and Fe fluxes increase to their LGM peak. Aeolian contributions to marine primary productivity before 37 kyrBP can be considered minimal because dust and Fe fluxes were close to interglacial values. On this basis, we find a comparable result to those of Rothlisberger et al. (2004), that



approximately 20 ppmv of the glacial-interglacial atmospheric CO<sub>2</sub> change can be attributed to atmospheric transport of mineral dust to the Southern Ocean. Both the Fe and CO<sub>2</sub> data are too sparse to conduct a similar evaluation for the penultimate glacial maximum.

## 4 Conclusions

We show that deglacial Fe, dust and nssCa<sup>2+</sup> fluxes varied across Antarctica, with TALDICE Fe fluxes consistently greater than those of EDC due to the deglacial activation of Antarctic coastal dust deflation sites as well as changing atmospheric transport patterns across Antarctica. Further, we show that fluxes of dust, nssCa<sup>2+</sup> and Fe can vary considerably from each other on millennial to centennial time scales. Based on the similar timing of LGM CO<sub>2</sub> drawdown and heightened Fe fluxes, we follow the technique of Rothlisberger et al. (2004) to attribute 20 ppmv of glacial-interglacial CO<sub>2</sub> variability to aeolian dust deposition over the Southern Ocean. Iron fluxes during MIS 6 were substantially less than those of MIS 2, implying that Fe fertilization contributed less to the similar-magnitude CO<sub>2</sub> increase during the penultimate deglaciation. One of the assumptions of Rothlisberger et al. (2004) was that of constant Fe solubility in dust. Ongoing studies should consider the solubility of Fe deposited to the sub-Antarctic regions: Fe/dust ratios calculated from Table 1 vary from 0.01 to 0.02 (EDC) and from 0.05 to 0.08 (TD), potentially reflecting different dust sources, transport pathways and atmospheric residence times, as well as different chemical properties such as Fe speciation (Spolaor et al., 2012).

*Acknowledgement.* We thank Samuel Albani, Barbara Delmonte, Paolo Gabrielli and Anders Svensson for helpful discussions which improved the manuscript. This work was supported by European Union Marie Curie IIF Fellowship (MIF1-CT-2006–039529, TDICOSO) within the VII Framework Program. The Talos Dome Ice Core Project (TALDICE), a joint European programme, is funded by national contributions from Italy, France, Germany, Switzerland and the UK. Primary logistical support was provided by PNRA at Talos Dome. This is TALDICE publication no. xx.

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## References

- Albani, S., Delmonte, B., Maggi, V., Baroni, C., Petit, J.-R., Stenni, B., Mazzola, C., and Frezzotti, M.: Interpreting last glacial to Holocene dust changes at Talos Dome (East Antarctica): implications for atmospheric variations from regional to hemispheric scales, *Clim. Past*, 8, 741–750, doi:10.5194/cp-8-741-2012, 2012.
- Anderson, R. F., Ali, S., Bradtmiller, L. I., Nielsen, S. H. H., Fleisher, M. Q., Anderson, B. E., and Burckle, L. H.: Wind-driven upwelling in the Southern Ocean and the deglacial rise in atmospheric CO<sub>2</sub>, *Science*, 323, 1443–1448, doi:10.1126/science.1167441, 2009.
- Barbante, C., Cozzi, G., Capodaglio, G., Van de Velde, K., Ferrari, C. P., Boutron, C. F., and Cescon, P.: Trace element determination in alpine snow and ice by double focusing inductively coupled plasma mass spectrometry with microconcentric nebulization, *J. Anal. Atom. Spectrom.*, 14, 1433–1438, doi:10.1039/A901949I, 1999.
- Bory, A., Wolff, E., Mulvaney, R., Jagoutz, E., Wegner, A., Ruth, U., and Elderfield, H.: Multiple sources supply eolian mineral dust to the Atlantic sector of coastal Antarctica: evidence from recent snow layers at the top of Berkner Island ice sheet, *Earth Planet. Sc. Lett.*, 291, 138–148, doi:10.1016/j.epsl.2010.01.006, 2010.
- Buiron, D., Chappellaz, J., Stenni, B., Frezzotti, M., Baumgartner, M., Capron, E., Landais, A., Lemieux-Dudon, B., Masson-Delmotte, V., Montagnat, M., Parrenin, F., and Schilt, A.: TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica, *Clim. Past*, 7, 1–16, doi:10.5194/cp-7-1-2011, 2011.
- Buiron, D., Stenni, B., Chappellaz, J., Landais, A., Baumgartner, M., Bonazza, M., Capron, E., Frezzotti, M., Kageyama, M., Lemieux-Dudon, B., Masson-Delmotte, V., Parrenin, F., Schilt, A., Selmo, E., Severi, M., Swingedouw, D., and Udisti, R.: Regional imprints of millennial variability during the MIS 3 period around Antarctica, *Quaternary Sci. Rev.*, 48, 99–112, doi:10.1016/j.quascirev.2012.05.023, 2012.
- Delmonte, B., Petit, J. R., Andersen, K. K., Basile-Doelsch, I., Maggi, V., and Ya Lipenkov, V.: Dust size evidence for opposite regional atmospheric circulation changes over East Antarctica during the last climatic transition, *Clim. Dynam.*, 23, 427–438, doi:10.1007/s00382-004-0450-9, 2004.
- Delmonte, B., Baroni, C., Andersson, P. S., Schoberg, H., Hansson, M., Aciego, S., Petit, J. R., Albani, S., Mazzola, C., Maggi, V., and Frezzotti, M.: Aeolian dust in the Talos Dome ice core

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr**

P. Vallelonga et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(East Antarctica, Pacific/Ross Sea sector): Victoria Land versus remote sources over the last two climate cycles, *J. Quaternary Sci.*, 25, 1327–1337, doi:10.1002/jqs.1418, 2010.

Edwards, R., Sedwick, P. N., Morgan, V., and Boutron, C.: Iron in ice cores from Law Dome: a record of atmospheric iron deposition for maritime East Antarctica during the Holocene and last glacial maximum, *Geochem. Geophys. Geos.*, 7, doi:10.1029/2006GC001307, 2006.

Fischer, H., Siggaard-Andersen, M.-L., Ruth, U., Röthlisberger, R., and Wolff, E.: Glacial/interglacial changes in mineral dust and sea-salt records in polar ice cores: sources, transport, and deposition, *Rev. Geophys.*, 45, RG1002, doi:10.1029/2005RG000192, 2007.

Fischer, H., Schmitt, J., Luthi, D., Stocker, T. F., Tschumi, T., Parekh, P., Joos, F., Kohler, P., Volker, C., Gersonde, R., Barbante, C., Le Floch, M., Raynaud, D., and Wolff, E.: The role of Southern Ocean processes in orbital and millennial CO<sub>2</sub> variations – a synthesis, *Quaternary Sci. Rev.*, 29, 193–205, doi:10.1016/j.quascirev.2009.06.007, 2010.

Gabrielli, P., Wegner, A., Petit, J. R., Delmonte, B., De Deckker, P., Gaspari, V., Fischer, H., Ruth, U., Kriews, M., Boutron, C. F., Cescon, P., and Barbante, C.: A major glacial-interglacial change in aeolian dust composition inferred from rare earth elements in Antarctic ice, *Quaternary Sci. Rev.*, 29, 265–273, doi:10.1016/j.quascirev.2009.09.002, 2010.

Gaspari, V., Barbante, C., Cozzi, G., Cescon, P., Boutron, C. F., Gabrielli, P., Capodaglio, G., Ferrari, C., Petit, J. R., and Delmonte, B.: Atmospheric iron fluxes over the last deglaciation: climatic implications, *Geophys. Res. Lett.*, 33, L03704, doi:10.1029/2005GL024352, 2006.

Kaufmann, P., Federer, U., Hutterli, M. A., Bigler, M., Schüpbach, S., Ruth, U., Schmitt, J., and Stocker, T. F.: An improved continuous flow analysis system for high-resolution field measurements on ice cores, *Environ. Sci. Technol.*, 42, 8044–8050, doi:10.1021/es8007722, 2008.

Lambert, F., Delmonte, B., Petit, J. R., Bigler, M., Kaufmann, P. R., Hutterli, M. A., Stocker, T. F., Ruth, U., Steffensen, J. P., and Maggi, V.: Dust-climate couplings over the past 800 000 years from the EPICA Dome C ice core, *Nature*, 452, 616–619, doi:10.1038/nature06763, 2008.

Li, F., Ginoux, P., and Ramaswamy, V.: Distribution, transport, and deposition of mineral dust in the Southern Ocean and Antarctica: contribution of major sources, *J. Geophys. Res.*, 113, D10207, doi:10.1029/2007JD009190, 2008.

Lüthi, D., Floch, M. L., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura, K., and Stocker, T. F.: High-resolution carbon diox-

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ide concentration record 650 000–800 000 years before present, *Nature*, 453, 379–382, doi:10.1038/nature06949, 2008.

Maher, B., Prospero, J., Mackie, D., Gaiero, D. M., Hesse, P. P., and Balkanski, Y.: Global connections between aeolian dust, climate and ocean biogeochemistry at the present day and at the last glacial maximum, *Earth-Sci. Rev.*, 99, 61–97, doi:10.1016/j.earscirev.2009.12.001, 2010.

Mahowald, N., Baker, A. R., Bergametti, G., Brooks, N., Duce, R. A., Jickells, T. D., Kubilay, N., Prospero, J. M., and Tegen, I.: Atmospheric global dust cycle and iron inputs to the ocean, *Global Biogeochem. Cy.*, 19, GB4025, doi:10.1029/2004GB002402, 2005.

Martin, J. H., Gordon, R. M., and Fitzwater, S. E.: Iron in Antarctic waters, *Nature*, 345, 156–158, doi:10.1038/345156a0, 1990.

Martinez-Garcia, A., Rosell-Melé, A., Geibert, W., Gersonde, R., Masqué, P., Gaspari, V., and Barbante, C.: Links between iron supply, marine productivity, sea surface temperature, and CO<sub>2</sub> over the last 1.1 Ma, *Paleoceanography*, 24, PA1207, doi:10.1029/2008PA001657, 2009.

Röthlisberger, R., Bigler, M., Wolff, E. W., Joos, F., Monnin, E., and Hutterli, M. A.: Ice core evidence for the extent of past atmospheric CO<sub>2</sub> change due to iron fertilisation, *Geophys. Res. Lett.*, 31, L16207, doi:10.1029/2004GL020338, 2004.

Schilt, A., Baumgartner, M., Schwander, J., Buiron, D., Capron, E., Chappellaz, J., Loulergue, L., Schüpbach, S., Spahni, R., Fischer, H., and Stocker, T. F.: Atmospheric nitrous oxide during the last 140 000 years, *Earth Planet. Sc. Lett.*, 300, 33–43, doi:10.1016/j.epsl.2010.09.027, 2010.

Schüpbach, S., Federer, U., Bigler, M., Fischer, H., and Stocker, T. F.: A refined TALDICE-1a age scale from 55 to 112 ka before present for the Talos Dome ice core based on high-resolution methane measurements, *Clim. Past*, 7, 1001–1009, doi:10.5194/cp-7-1001-2011, 2011.

Sigman, D. M., Hain, M. P., and Haug, G. H.: The polar ocean and glacial cycles in atmospheric CO<sub>2</sub> concentration, *Nature*, 466, 47–55, doi:10.1038/nature09149, 2010.

Smetacek, V., Klaas, C., Strass, V. H., Assmy, P., Montresor, M., Cisewski, B., Savoye, N., Webb, A., d'Ovidio, F., Arrieta, J. M., Bathmann, U., Bellerby, R., Berg, G. M., Croot, P., Gonzalez, S., Henjes, J., Herndl, G. J., Hoffmann, L. J., Leach, H., Losch, M., Mills, M. M., Neill, C., Peeken, I., Rottgers, R., Sachs, O., Sauter, E., Schmidt, M. M., Schwarz, J., Ter-

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



bruggen, A., and Wolf-Gladrow, D.: Deep carbon export from a Southern Ocean iron-fertilized diatom bloom, *Nature*, 487, 313–319, doi:10.1038/nature11229, 2012.

Spolaor, A., Vallelonga, P., Gabrieli, J., Cozzi, G., Boutron, C., and Barbante, C.: Determination of Fe<sup>2+</sup> and Fe<sup>3+</sup> species by FIA-CRC-ICP-MS in Antarctic ice samples, *J. Anal. Atom. Spectrom.*, 27, 310–317, doi:10.1039/C1JA10276A, 2012.

Stenni, B., Buiron, D., Frezzotti, M., Albani, S., Barbante, C., Bard, E., Barnola, J. M., Baroni, M., Baumgartner, M., Bonazza, M., Capron, E., Castellano, E., Chappellaz, J., Delmonte, B., Falourd, S., Genoni, L., Iacumin, P., Jouzel, J., Kipfstuhl, S., Landais, A., Lemieux-Dudon, B., Maggi, V., Masson-Delmotte, V., Mazzola, C., Minster, B., Montagnat, M., Mulvaney, R., Narcisi, B., Oerter, H., Parrenin, F., Petit, J. R., Ritz, C., Scarchilli, C., Schilt, A., Schupbach, S., Schwander, J., Selmo, E., Severi, M., Stocker, T. F., and Udisti, R.: Expression of the bipolar see-saw in Antarctic climate records during the last deglaciation, *Nat. Geosci.*, 4, 46–49, doi:10.1038/ngeo1026, 2011.

Vallelonga, P., Gabrielli, P., Balliana, E., Wegner, A., Delmonte, B., Turetta, C., Burton, G., Vanhaecke, F., Rosman, K. J. R., Hong, S., Boutron, C. F., Cescon, P., and Barbante, C.: Lead isotopic compositions in the EPICA Dome C ice core and Southern Hemisphere Potential Source Areas, *Quaternary Sci. Rev.*, 29, 247–255, doi:10.1016/j.quascirev.2009.06.019, 2010.

Wegner, A., Gabrielli, P., Wilhelms-Dick, D., Ruth, U., Kriews, M., De Deckker, P., Barbante, C., Cozzi, G., Delmonte, B., and Fischer, H.: Change in dust variability in the Atlantic sector of Antarctica at the end of the last deglaciation, *Clim. Past*, 8, 135–147, doi:10.5194/cp-8-135-2012, 2012.

Wolff, E. W., Fischer, H., Fundel, F., Ruth, U., Twarloh, B., Littot, G. C., Mulvaney, R., Rothlisberger, R., de Angelis, M., Boutron, C. F., Hansson, M., Jonsell, U., Hutterli, M. A., Lambert, F., Kaufmann, P., Stauffer, B., Stocker, T. F., Steffensen, J. P., Bigler, M., Siggaard-Andersen, M. L., Udisti, R., Becagli, S., Castellano, E., Severi, M., Wagenbach, D., Barbante, C., Gabrielli, P., and Gaspari, V.: Southern Ocean sea-ice extent, productivity and iron flux over the past eight glacial cycles, *Nature*, 440, 491–496, doi:10.1038/nature04614, 2006.

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

**Table 1.** Deposition of Antarctic Fe and dust before and after the last deglaciation. Talos Dome Fe data compared to EPICA Dome C (EDC) (Gaspari et al., 2006; Delmonte et al., 2010) and Law Dome (Edwards et al., 2006). Dust concentrations and fluxes are also shown for EDC and Talos Dome (Delmonte et al., 2010). Note that Fe and dust concentrations are shown in normal font while fluxes are shown in bold font. Indicative Holocene accumulation rates are also shown for comparative purposes.

|   | Talos Dome        |                   | Law Dome           | Dome C            |                    |
|---|-------------------|-------------------|--------------------|-------------------|--------------------|
| Holocene accumulation<br>(m H <sub>2</sub> O eq. yr <sup>-1</sup> ) | 0.07              |                   | 0.8                | 0.034             |                    |
| Concentrations (ngg <sup>-1</sup> )                                 | Fe                | Dust              | Fe                 | Fe                | Dust               |
| <b>Fluxes (10<sup>-2</sup> mgm<sup>-2</sup> yr<sup>-1</sup>)</b>    |                   |                   |                    |                   |                    |
| Holocene<br>(2–16.5 kyrBP)  | 1.5<br><b>9</b>   | 20<br><b>130</b>  | 0.06<br><b>4.5</b> | 0.2<br><b>0.7</b> | 15<br><b>40</b>    |
| LGM<br>(18–25 kyrBP)  | 14.8<br><b>45</b> | 250<br><b>850</b> | 6.4<br><b>45</b>   | 16<br><b>24</b>   | 750<br><b>1000</b> |
| LGM/Holocene  | 10<br><b>5</b>    | 12<br><b>6</b>    | 107<br><b>10</b>   | 74<br><b>36</b>   | 50<br><b>25</b>    |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 1.** Talos Dome Fe (this work) and dust (Delmonte et al., 2010) fluxes and  $\delta^{18}\text{O}$  (Stenni et al., 2011) compared to Fe (Wolff et al., 2006) and dust (Lambert et al., 2008) fluxes at EPICA Dome C.  $\text{CO}_2$  concentrations were compiled from various Antarctic ice cores for the period 0.2–138 kyrBP (Schilt et al., 2010) and from EDC before 138 kyrBP (Lüthi et al., 2008).

## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






**Fig. 2.** Deglacial changes in Fe, nssCa<sup>2+</sup> and dust at Dome C and Talos Dome. TALDICE  $\delta^{18}\text{O}$  values are from (Stenni et al., 2011). Fine (1–5  $\mu\text{m}$ )/coarse (5–10  $\mu\text{m}$ ) dust particle ratios are from Albani et al. (2012) and dust fluxes are from Delmonte et al. (2010). nssCa<sup>2+</sup> data are shown for TD (this study) and EDC (Röthlisberger et al., 2004) ice cores. Deglacial EDC Fe fluxes are from Gaspari et al. (2006). The vertical dashed lines, located at 11.9 and 12.9 kyr BP, highlight enhanced Fe fluxes which occur independently of dust fluxes at Talos Dome and Dome C, respectively.



## Iron fluxes to Talos Dome, Antarctica, over the past 200 kyr

P. Vallelonga et al.



**Fig. 3.** Holocene Fe (this work) and dust (Delmonte et al., 2010) fluxes and fine (1–5 μm)/coarse (5–10 μm) dust particle ratios (Albani et al., 2012) in the TALDICE ice core (blue lines). EPICA Dome C dust fluxes (Wolff et al., 2006) are also shown (red squares). Note the fine/coarse dust particle ratios are plotted here on a reversed scale.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 4.** Comparison of LGM Fe fluxes and late glacial drawdown of CO<sub>2</sub>. TALDICE δ<sup>18</sup>O paleotemperature values are from (Stenni et al., 2011). CO<sub>2</sub> concentrations were compiled from various Antarctic ice cores (Lüthi et al., 2008; Schilt et al., 2010) while Fe fluxes are shown for TD (this work) and EDC (Wolff et al., 2006; Gaspari et al., 2006) ice cores.