

This discussion paper is/has been under review for the journal *Climate of the Past* (CP).
Please refer to the corresponding final paper in CP if available.

Caspian Sea level changes during the last millennium: historical and geological evidences from the south Caspian Sea

**A. Naderi Beni¹, H. Lahijani², R. Mousavi Harami¹, K. Arpe^{3,4}, S. A. G. Leroy⁴,
N. Marriner⁵, M. Berberian⁶, V. Andrieu-Ponel⁷, M. Djamali⁷, and A. Mahboubi¹**

¹Geology Department, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran

²Marine Geology Department, Iranian National Institute for Oceanography (INIO), Tehran, Iran

³Max Planck Institute for Meteorology, Hamburg, Germany

⁴Institute for the Environment, Brunel University, UB8 3PH Uxbridge, London, UK

⁵Aix-Marseille Université, CNRS, CEREGE, UMR6635 – IRD, CEREGE, UMR 161 –
Collège de France, CEREGE, 13545 Aix en Provence cedex 4, France

⁶School of Mathematics, Science, and Technology, Ocean County College, Toms River,
NJ 08754-2001, USA

⁷Institut Méditerranéen de Biodiversité et d'Ecologie UMR7263 CNRS/Aix-Marseille Université
– Europôle Méditerranéen de l'Arbois – Pavillon Villemin – BP 80, 13545 Aix-en-Provence
Cedex 04, France

1397

Received: 13 February 2013 – Accepted: 28 February 2013 – Published: 12 March 2013

Correspondence to: A. Naderi Beni (amnaderi@inio.ac.ir)

Published by Copernicus Publications on behalf of the European Geosciences Union.

were taken from the morphological zone 1 of Voropaev et al. (1998), which is the most sensitive area to sea level changes. The cores were analysed sedimentologically to correlate with historical documents as well as other geological findings.

The elevation of the coring sites was measured using an Ashtech Mobile Mapper 100. The magnetic susceptibility (MS) of the core samples was measured using a MS2C core-logging scanner from Bartington. The diameter of the susceptibility meter loop was 10 cm and the progression step was 2 cm. The sensitivity of the meter was about 2×10^{-6} SI.

Both of core samples were split and sub-sampled at the laboratory of the Iranian National Institute for Oceanography (INIO) for sedimentological analysis. A Nabertherm P330 furnace was used for loss-on-ignition to measure organic matter and carbonate content based on the methods outlined in Heiri et al. (2001). Grain-size measurements were made using a Horiba Laser Scattering Particle Size Distribution Analyzer LA-950.

Fossil content was identified to aid in characterizing the past depositional environments based on the atlas of the invertebrates of the CS (Birstein et al., 1968).

Two articulated bivalve shells of *Cerastoderma lamarcki*, an indicator of marine environment, were selected and sent to Poznan Radiocarbon laboratory for ^{14}C dating. Calendar ages were obtained from the CALIB Rev 6.0.1 software (Reimer et al., 2009) based on three different databases of IntCal09, Marine09 and Mixed Marine NoHem to compare the results and correlate them with historical findings.

Several historical documents were studied mainly in the library of the Ferdowsi University of Mashhad and related geographical names and positions were extracted based on the "Historical Geography of Cities" (Nahchiri, 1999). We have used the most reliable literature sources covering the last millennium. Historical observations of the CS environments were gathered to compare and contrast with geological records for the same period.

Note that in this study the dates are given in AD, unless otherwise stated. The Persian geographic names and other Persian words are written as they are pronounced and were originally written, with direct and simplified transliteration from Persian to English.

1405

Diacritical marks and special characters are used to differentiate vowel the "A" [short; e.g. ant] from "Ā" [long; e.g. Rāmsar], and Arabic "ain" (also used in Persian) as "ʿA" [e.g. 'Abbās]. Elevations are above or below the present mean sea level. Rud and Daryā mean river and Dāgh means mountains.

5 4 Results and interpretation

4.1 Geological findings

The CSL changes and their impact on coastal evolution have been investigated by several researchers e.g. Kroonenberg et al. (2000, 2007), Lahijani et al. (2009), Kazanci et al. (2004), Leroy et al. (2011), Kakroodi et al. (2012) and Naderi Beni et al. (2013). The results of the geological investigations on the CSL are summarized in Table 1.

In this study we focused on two short cores from the south-eastern flank of the CS to compare the results with other geological findings as well as historical evidence.

The lithology of the cores shows a succession of terrestrial (fluviodeltaic in Kakroodi et al., 2012) and marine environments during the last millennium (Fig. 4). Generally, the marine facies comprise finer-grained materials compared to terrestrial deposits based on the modal grain size and contain marine bivalve fossils of *Cerastoderma lamarcki*. The terrestrial deposits constitute alternations of thin layers of fine sand, silt and clay, without any fossil content but containing gypsum minerals that are concentrated in some horizons. The presence of gypsum minerals in terrestrial sediments could be related to the flat topography of the region and warm climatic conditions that lead to water evaporation during dry seasons. Sea level rise is indicated by a change from the terrestrial facies to marine facies. In the marine facies, organic matter and carbonate contents tend to increase due to higher organic activity in shallow marine environment and presence of calcareous shell bearing organisms (Fig. 4).

The magnetic susceptibility results show lower values in terrestrial deposits, which could be related to the provenance of the grains or the increase in evaporative minerals.

1406

The feeding watercourses have their headwaters in the Kopeh Dāgh that is mantled in calcareous deposits. The higher values of the magnetic susceptibility in the marine facies could be linked to the presence of paramagnetic minerals, e.g. muscovite, flogopite and biotite, which are transported by longshore currents from the southern coast of the CS. The paramagnetic components of the southern coast deposits are provided by igneous and metamorphic rock outcrops of Alborz Mountains (Lahijani and Tavakoli, 2012). According to the presence of some brackish water gastropods such as *Theodoxus palassii*, as well as Charophytes that coexist with marine gastropods *Horatia marina* and *Pyrgohydrobia* sp. and marine bivalve of *Cerastoderma lamarcki* (Birstein et al., 1968), it seems that the marine facies could represent a shallow marine environment and/or an open lagoon that was influenced by fresh-water input. The formation of barrier-lagoon complexes during rapid sea-level rise has been reported by Kroonenberg et al. (2007), Lahijani et al. (2009) and Naderi Beni et al. (2013) in different parts of the Caspian coast and, therefore, it is more probable to link the marine facies to an open lagoon environment which was influenced by fresh water input.

4.2 Radiocarbon dating

The age of the oldest marine facies of cores A and B (Fig. 4) is dependent on the database used to calibrate the radiocarbon data (Table 2). Although almost all of the chronological data are coincident with the LIA in the North Atlantic Ocean and already recorded in the CS by a high-stand (Leroy et al., 2011 in Gilān, and Kakroodi et al., 2012 in Golestān), they could be linked to different sea-level rise episodes between the thirteenth to the seventeenth centuries depending on the databases used for calibration (Table 2).

According to the elevation of the coring site (Table 2), it seems that sea level in dating horizons reached -26.75 m and -25.8 m for core A and B, respectively, as the subsidence of the region is negligible for the last millennium.

1407

4.3 Historical evidence

The CSL variability before the instrumental observations (prior to 1850) was investigated by Brückner (1890) on along the CS coast. He used a wide range of observational evidence, e.g. travel descriptions, navigation maps, and paintings to garner data on CSL changes. Typical examples were walls along the shore with markings of sea levels, reports about buildings that disappeared under the water, and islands that emerged or disappeared. The results of Brückner (1890) are summarized in Table 3.

Fedchina (1980) used the Russian cartographic data from 1556 to 1925 to reconstruct a CSL curve. The same method was followed by Komarova (1980) for 1700 to 1850 time period. The results of Fedchina (1980) and Komarova (1980) are summarized for the 1700 to 1850 time period in Fig. 5. In spite of some differences between their analyses, the results generally show a good agreement in the CSL changes during the time period with high-stands up to -22 m (Fig. 5).

Varushchenko et al. (1987) used a wide range of historical, archaeological and geological evidences to reconstruct CSL changes for the last 2400 yr. Many historical and archaeological documents used by Varushchenko et al. (1987) for the last millennium are the same that Brückner (1890) and Komarova (1980) considered in their works (Table 4). However, their results have many differences especially in the early centuries of the last millennium.

Lithological evidence from coring along the Caspian coastline was investigated by Karpychev (1998) and combined with historical information (Karpychev, 2001) for layers of pebbles, for instance, which are an indication for shores at that level and for which the age was determined by radiocarbon-dating. The results of Karpychev (1998, 2001) are summarized in Table 5.

According to the results of Tables 3, 4 and 5, the CSL curve could be plotted from the tenth to the twentieth centuries (Fig. 6).

The Iranian literary texts are especially rich in statements pertinent to understanding the historical geography and geological events of the Iranian Plateau and its

1408

Abnormal fluctuations were also observed at Baku, Lankarān and the southern Caspian shore, where sea level oscillated within the space of an hour by fifty to more than a hundred centimetres in 1868 and 1960 (Hedin, 1892; Musketov and Orlov, 1893; Ambraseys and Melville, 1982).

5 Large sea-waves were also observed at Āstārā associated with the 1910, M_s 5.4, Moghān earthquake (Kondorskaya and Shebalin, 1977). Large sea-waves were noticed all along the southern Caspian coast from Miānkāleh in the east to Anzali in the west (Ambraseys and Melville, 1982) during the 1890, $M_s \sim 7.2$ Tāsh earthquake in the east Alborz, about 60 km to the SE of the Caspian shore.

10 Sea waves flooded the coast of the Cheleken Island during the 1895 M_s 7.4 Krasnovodsk (Qezel Suyu, Turkmenbāshi) earthquake on the eastern Caspian shore (Kondorskaya and Shebalin, 1977; Ambraseys, 1997). The water in the harbour quickly swelled and reached the railroad tracks. A ship 30 km from Krasnovodsk reported that the sea swelled with the shock and ejected a column of water and smoke (possible submarine mud-volcano eruption).

15 During the 1962 M_w 7.0 Bu'in earthquake, which took place 142 km to the SW of the Caspian shore, some irregularities in the behaviour of the water level of the CS were noticed at the Anzali, Naushahr and Bābolsar ports, where waves followed the earthquake. The tide gauge records at the Anzali port showed abnormal variations in sea level before and after the earthquake; waves with amplitudes of nearly two feet and periods between 15 and 50 min were recorded. At the Naushahr port, a series of swells had been reported but made no noticeable damage. In addition, the morning after the earthquake, the CS was muddy for more than a nautical mile (Ambraseys, 1962, 1963).

20 The 1990 M_w 7.3 Rudbār earthquake (Berberian et al., 1992; Berberian and Walker, 2010), which was much larger than the 1962 M_w 7.0 Bu'in earthquake, and its epicentre was closer to the CS (about 68 km to the SW of the Caspian shore vs. 142 km), should have caused unusual fluctuations in the CS, much greater than the 1962 event. Unfortunately, our efforts in obtaining the hourly/daily records at the south CS failed because the gauge station was broken down during the earthquake.

1415

4.3.5 Other historical findings

Ibn Hawqal (1988, page 141) in tenth century described the Aral Sea (Fig. 1) as a salt lake that had not been freshened even by the Āmu-daryā water: *“Although considerable water from the Jayhun River (Āmu-daryā) discharges into the lake, the area of the lake has not been increased and the water is not freshened and people believe that the lake is connected to the CS via an underground channel”*. Although this is corroborated by the Hudud al-'Alam (1973), Mostowfi (1999) mentioned that the Jayhun River (Āmu-daryā) was diverted from the Aral Sea towards the CS by the sons of Genghis Khan when they surged towards Iran in 1219 (Létolle, 2000).

10 Mar'ashi (1982) quoted that during the invasion of Māzandarān by Uzbeks in 1392, the army invaded Āmol (Fig. 8), then arrested people and after that transferred them to Khārazm by ship from the CS and the Jayhun River. This story indicates that Āmu-daryā was discharging into the CS at that time via the Uzboy waterway. According to Barthold (1984), this waterway was open until the late sixteenth century.

15 In addition to the above-mentioned river avulsion, some reports on other Caspian rivers provide useful information on the south Caspian river courses.

The Sefidrud was reported as the largest river on the south Caspian coast in the Hudud al-'Alam (1973). According to this report, the people of Gilān could be categorized into two groups, the first group settled between the river and the sea, i.e. Rasht; while the other group was situated between the river and the mountains, i.e. Lahijan (Fig. 1). Regarding the East-West direction of the Alborz Mountains (Fig. 1), this categorization is true when the river direction is parallel to the mountains and flow eastward in the plain. Rabino (1917) stated that in 1740 a large tributary separated from the Sefidrud, 6 km south of the river mouth, and discharged into the Anzali Lagoon, and that this tributary was large enough to enable ships to carry passengers from Anzali to Pīrbāzār near Rasht. This tributary could be navigated in 1875 because Farhād Mirzā (1987) used it in his journey to Europe. Barthold (1984) mentioned that the tributary

1416

understanding of the impacts of CSL changes on the environment as well as human societies.

Owing to descriptive nature of many historical sources, some uncertainties in quantifying sea-level variations exist. These uncertainties may be related to observational errors, different standards in describing the events used by different historians and geographers and seismic events. Considering negligible rate of uplifting/subsidence of the region during the last millennium and due to seismicity of the CS region, earthquakes could cause considerable vertical displacement of the shoreline in some places. These local vertical movements should be taken into account when attempting to reconstruct CSL changes based on historical and archaeological evidence.

One of the ways to decrease the uncertainties associated with historical data is the correlation with geological findings as well as the comparison of individual observations with other contemporaneous reports. The multidisciplinary approach leads to a better understanding of past environments and, moreover, to corroborate the accuracy of geological findings.

Some contradictions, however, occur between historical evidence and proxy-based interpretation when determining the relative sea-level position (e.g. historical sea level low-stand from 950 to 1250 a period in which a high-stand was dated by some geoscientists such as Lahijani et al., 2009). It seems that this problem partially comes from uncertainties in dating methods as well as our geological interpretations. Precisely constraining the RE for the CS is a high priority for future studies. Nonetheless, our study on the southeast CS shows that using the Marine09 database for calibration of ^{14}C ages yields more reliable results for this particular region.

As the CS and its watershed area spans the sub-tropics in the southwest to desertic climate in east-northeast, and the humid mid-latitudes in the northwest of the watershed, this could provide a good opportunity for geoscientists to investigate climate change in the Northern Hemisphere. Comparison of the CSL reconstruction for the last millennium with solar irradiance and with the fluctuation of lakes in Middle Asia and Europe, show relatively good agreement between different curves. Based on the

1423

antiphase relationship between the CSL changes and the solar irradiance during the last millennium, it could be concluded that the main historical CSL fluctuations have been modulated by solar activity. However, in some periods no good agreement occurs between the curves, which could be linked to the absence of data from the CSL for certain periods as well as regional irregularities, such as earthquakes.

Despite these problems, this study is able to construct a curve of the CSL variability for the last millennium, which fits to a multitude of observational evidences and can be used for validating simulations with climate models. In the early part of the millennium the absolute values have a wider range of uncertainty but most data agree in the events of increases and decreases of the CSL.

Acknowledgements. This study has been conducted in the framework of a project entitled: "A comparative study of Holocene climate changes in coastal areas of the south-eastern and south-western Caspian Sea based on geological evidence" that is funded by the Iranian National Institute for Oceanography (INIO). This study has also been supported by the Iranian Centre for International Scientific Studies and Collaboration (CISSC), Ministry of Science, Research and Technology of Iran, and its French partner Campus France in the framework of the Franco-Iranian Gundishapour Program. The publication is a contribution to the INQUA Quick-LakeH project (No. 1227). We thank R. Jokar, M. Pourkerman, N. Ghasemi, and M. Hosseindoost for their support with field and laboratory work.

References

- Abbott, K. E.: Notes on Ghilan, P. R. Geogr. Soc. London, 3, 390–395, 1858.
 al-Bakri, A.: Kitāb al-Masālik w-al-Mamālik, 604 AH/1208 AD, edited by: Ghareh Chānlou, H., Motarjem Publication, Tehran, Iran, 1999.
 al-Istakhri: Masalik al Mamalik, 339 AH/951 AD, edited by: Afshar, I., Bongah e Tarjomeh va Nashr-e Ketab, Tehran, 1961.
 al-Maqdisi, M. b. A.: Aḥsan al-taqāsīm fī ma'rifat al-aqālīm, 374 AH/985 AD, Mo'alefān va Motarjemān Publication, Tehran, Iran, 1982.
 al-Mas'ūdī, A. i. a.-H.: Murūj al-dhahab wa-ma'ādin al-jawhar, 335 AH/947 AD, edited by: Payandeh, A., Elmi va Farhangi Publication, Tehran, Iran, 2012.

1424

- Gümilev, N.: History of the Caspian sea level changes during the last 2000 years, in: Humidity Change in Aral-Caspian Region during the Holocene, edited by: Adrianov, B. V., Zorin, L. B., and Nikolaeva, R. V., USSR, Academy of Science, Institute of Water Problem, NAUKA, Moscow, 1980.
- 5 Hedin, S.: Der Demavand nach eigener Beobachtung, Ph.D. Thesis, Berlin, Germany, 304 pp., 1892.
- Heiri, O., Lotter, A. F., and Lemcke, G.: Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results, *J. Paleolimnol.*, 25, 101–110, 2001.
- 10 Hoogendoorn, R. M., Boels, J. F., Kroonenberg, S. B., Simmons, M. D., Aliyeva, E., Babazadeh, A. D., and Huseynov, D.: Development of the Kura delta, Azerbaijan; a record of Holocene Caspian sea level changes, *Mar. Geol.*, 222–223, 359–380, 2005.
- Hudud al-Ālam: Hudud al-Ālam "The regions of the world": a Persian geography, 372 AH/982 AD, edited by: Tehrani, S. J., Majles Publication, Tehran, Iran, 1973.
- 15 ibn Hawqal, M. b. A.: Kitāb sūrat al-arḍ, 366 AH/977 AD, translated from Arabic by: Sho'ar, J., 2nd Edn., Amir Kabir Publication, 400 pp., 1988.
- Jayhāni, M. b. A.: Al Masālik w- al-Mamālik (Ashkāl al Ālam), 279–294 AH/893–907 AD, edited by: Kateb, A. B. A., Āstān Ghods Razavi Publication, Mashhad, Iran, 1989.
- Jovayni, A.-M.: Ta'rīkh-i jahān-gushā, 658 AH/1260 AD, edited by: Qazvini, M. A., Arghavān Publication, Tehran, Iran, 1911.
- 20 Kakroodi, A. A.: Rapid Caspian Sea-level change and its impact on Iranian coasts, Ph.D. Thesis, Department of Geotechnology, Faculty of Civil Engineering and Geosciences, Delft, Netherlands, 121 pp., 2012.
- Kakroodi, A. A., Kroonenberg, S. B., Hoogendoorn, R. M., Mohamm Khani, H., Yamani, M., Ghassemi, M. R., and Lahijani, H. A. K.: Rapid Holocene sea level changes along the Iranian Caspian coast, *Quaternary Int.*, 263, 93–103., 2012.
- Karpychev, Y. A.: Dating of regressive stages in the Caspian Sea using ¹⁴C, *Vodn. Resur.*, 25, 274–278, 1998.
- Karpychev, Y. A.: Variation in the Caspian Sea level in the historic epoch, *Water Resour.*, 1, 1–14, 2001.
- 30 Kazancı, N., Gulbabazadeh, T., Leroy, S. A. G., and Ileri, O.: Sedimentology and environmental characteristics at the Gilān-Mazenderan plain, northern Iran; influence of long and short-term Caspian level fluctuations on geomorphology, *J. Marine Syst.*, 46, 154–168, 2004.

- Kazin, E. A.: Chronology in the Caucasian Mineral Water and Stavropol Plateau, *Bull. of Caucasian Seismic Stations*, Tbilisi, July–December 1973, 1974.
- Knapp, C. C., Knapp, J. H., and Connor, J. A.: Crustal-scale structure of the South Caspian basin revealed by deep seismic profiling, *Mar. Petrol. Geol.*, 21, 1073–1081, 2004.
- 5 Komarova, N. G.: Caspian sea level changes in the Holocene based on historical data, in: Humidity Change in Aral-Caspian Region during the Holocene, edited by: Adrianov, B. V., Zorin, L. B., Nikolaeva, R. V., USSR, Academy of Science, Institute of Water Problem, NAUKA Moscow, 1980.
- Kondorskaya, N. and Shebalin, N.: New Catalog of Strong Earthquakes in the Territory of the USSR from Ancient Times to 1975, Academy of Sciences, Moscow, English translation, updated through 1977, available as Report SE-31, World Data Center A for Solid Earth Geophysics, Boulder, CO, 606 pp., 1977.
- 10 Kousari, S.: Evolution of Sefidrud Delta, *Development in Geological Education*, 1, 31–41, 1986.
- Kroonenberg, S. B., Badyukova, E. N., Storms, J. E. A., Ignatov, E. I., and Kasimov, N. S.: A full sea level cycle in 65 years: barrier dynamics along Caspian shores, *Sediment. Geol.*, 134, 257–274, 2000.
- Kroonenberg, S. B., Abdurakhmanov, G. M., Aliyeva, E. G., Badyukova, E. N., van der Borg, K., Hoogendoorn, R. M., Huseynov, D., Kalashnikov, A., Kasimov, N. S., Rychagov, G. I., Svitoch, A. A., Vonhof, H. B., and Wesselingh, F. P.: Solar-forced 2600 BP and Little Ice Age high-stands of the CS, *Quaternary Int.*, 173–174, 137–143, 2007.
- 20 Lahijani, H. and Tavakoli, V.: Identifying provenance of South Caspian coastal sediments using mineral distribution pattern, *Quaternary Int.*, 261, 128–137, 2012.
- Lahijani, H. A. K., Tavakoli, V., and Amini, A. H.: South Caspian river mouth configuration under human impact and sea level fluctuations, *Environ. Sci.*, 5, 65–86, 2008.
- 25 Lahijani, H. A. K., Rahimpour-Bonab, H., Tavakoli, V., and Hosseindoost, M.: Evidence for Late Holocene high-stands in Central Gilān–East Mazanderan, South Caspian coast, Iran, *Quaternary Int.*, 197, 55–71, 2009.
- Leroy, S. A. G., Marret, F., Giralt, S., and Bulatov, S. A.: Natural and anthropogenic rapid changes in the Kara-Bogaz Gol over the last two centuries reconstructed from palynological analyses and a comparison to instrumental records, *Quaternary Int.*, 150, 52–70, 2006.
- 30 Leroy, S. A. G., Marret, F., Gibert, E., Chalié, F., Reyss, J. L., and Arpe, K.: River inflow and salinity changes in the Caspian Sea during the last 5500 years, *Quaternary Sci. Rev.*, 26, 3359–3383, 2007.

- Leroy, S. A. G., Lahijani, H. A. K., Djamali, M., Naqinezhad, A., Moghadam, M. V., Arpe, K., Shah-Hosseini, M., Hosseindoust, M., Miller, C. S., Tavakoli, V., Habibi, P., and Naderi, M.: Late Little Ice Age palaeoenvironmental records from the Anzali and Amirkola lagoons (south CS): vegetation and sea level changes, *Palaeogeogr. Palaeoecol.*, 302, 415–434, 2011.
- 5 Létolle, R.: Histoire de l'Ouzboï, cours fossile de l'Amou Daria: synthèse et éléments nouveaux, *Studia Iranica*, 29, 195–240., 2000.
- Magny, M., Peyron, O., Gauthier, E., Vannièrè, B., Millet, L., and Vermot-Desroches, B.: Quantitative estimates of temperature and precipitation changes over the last millennium from pollen and lake-level data at Lake Joux, Swiss Jura Mountains, *Quaternary Res.*, 75, 45–54, 2011.
- 10 Mamedov, A. V.: The late Pleistocene-Holocene history of the CS, *Quaternary Int.*, 41–42, 161–166, 1997.
- Mangino, S. and Priestley, K.: The crustal structure of the southern Caspian region, *Geophys. J. Int.*, 133, 630–648, 1998.
- 15 Mar'ashi, Z.: *Tārikh e Tabarestān va Rūyān va Māzandarān, 880 AH/1476 AD*, edited by: Tashibi, M., Shargh Publication, Tehran, Iran, 1982.
- Mayewski, P. A., Rohling, E. E., Curt Stager, J., Karlén, W., Maasch, K. A., Meeker, L. D., Meyerson, E. A., Gasse, F., van Kreveld, S., Holmgren, K., Lee-Thorp, J., Rosqvist, G., Rack, F., Staubwasser, M., Schneider, R. R., and Steig, E. J.: Holocene climate variability, *Quaternary Res.*, 62, 243–255, 2004.
- 20 Mostowfi, H.: *Noz'hat al-Gholub, 740 AH/1340 AD*, edited by: Siaghi, M. D., Tāhā Publication, Qazvin, Iran, 1999.
- Musketov, I. and Orlov, A.: *The Catalog of Earthquakes in the Russian Empire*, *Russ. Geogr. Soc.*, St. Petersburg, 1893.
- 25 Naderi Beni, A., Lahijani, H., Moussavi Harami, R., Leroy, S. A. G., Shah-hosseini, M., Kabiri, K., and Tavakoli, V.: Development of spit-lagoon complexes in response to Little Ice Age rapid sea level changes in the central Gilān coast, South CS, Iran, *Geomorphology*, 187, 11–26, doi:10.1016/j.geomorph.2012.11.026, 2013.
- Nahchiri, A.: *Joghrafiāy e Tārikhī e Shahrhā (Historical geography of cities)*, Madreseh Publication, Tehran, Iran, 1999.
- 30 Nikitin, P. N.: On the problem of seismic activity in the Caspian Sea and its northwestern coast, *Materials of the National Conference on the Problem of the Caspian Sea*, Baku, 1963.
- Nikitin, P. N.: *Chronology of Local Earthquakes in the Northern Caucasus*, Pyatigorsk, 1974.

1429

- Okhravi, R. and Djamali, M.: The missing ancient Lake of Saveh; a historical review, *Iranica Antiqua*, 38, 327–344, 2003.
- Ozyavas, A., Khan, S. D., and Casey, J. F.: A possible connection of Caspian Sea level fluctuations with meteorological factors and seismicity, *Earth Planet. Sc. Lett.*, 299, 150–158, 2010.
- 5 Parodi, S.: *Cose e parole nei Viaggi di Pietro della Valle*, Accademia della Crusca, 1997.
- Priestley, K., Baker, C., and Jackson, J.: Implications of earthquake focal mechanism data for the active tectonics of the South Caspian Basin and surrounding regions, *Geophys. J. Int.*, 118, 111–141, 1994.
- 10 Rabino, H. L.: *Les provinces caspiennes de la Perse: Le Guīlān*, Illustrations, E. Leroux, 1917.
- Rabino, H. L.: *Mazandaran and Astarabad*, Indus Publications, Karachi, 1980.
- Reimer, P., Baillie, M., Bard, E., Bayliss, A., Beck, J., Blackwell, P., Ramsey, C. B., Burr, G., and Edwards, R.: *IntCal09 and Marine09 radiocarbon age calibration curves, 0–50,000 years Cal BP*, *Radiocarbon*, 51, 1111–1150, 2009.
- 15 Rekavandi, H. O., Sauer, E. W., Wilkinson, T., Tamak, E. S., Ainslie, R., Mahmoudi, M., Griffiths, S., Ershadi, M., Van Rensburg, J. J., and Fattahi, M.: An Imperial Frontier of the Sasanian Empire: Further Fieldwork at the Great Wall of Gorgan, Iran, 95–136, 2007.
- Rodkin, M.: Hydroseismicity, new evidence, *J. Geodyn.*, 15, 247–260, 1992a.
- Rodkin, N. V.: Possible correlation of Caspian sea level fluctuations with seismicity, *Izv. Earth Sci.*, 28, 542–549, 1992b.
- 20 Ruddiman, W. F.: *Earth Climate, Past and Future*, W. H. Freeman and Company, New York, 2008.
- Rychagov, G. I.: Holocene oscillations of the Caspian Sea and forecasts based on paleogeographical reconstructions, *Quaternary Int.*, 41–42, 167–172, 1997.
- 25 Shah-hosseini, M., Morhange, C., Naderi Beni, A., Marriner, N., Lahijani, H., Hamzeh, M., and Sabatier, F.: Coastal boulders as evidence for high-energy waves on the Iranian coast of Makran, *Mar. Geol.*, 290, 17–28, 2011.
- Shilo, N. A. and Krivoshey, M. I.: Relation between the Caspian Sea Level Variations and Strain in the Earth's Crust, *Vestnik Russ. Acad. Sci.*, Moscow, 83–90, 1989.
- 30 Sorrel, P., Popescu, S. M., Head, M. J., Suc, J. P., Klotz, S., and Oberhansli, H.: Hydrographic development of the Aral Sea during the last 2000 years based on a quantitative analysis of dinoflagellate cysts, *Palaeogeogr. Palaeoecol.*, 234, 304–327, 2006.

1430

- Storms, J. E. A. and Kroonenberg, S. B.: The impact of rapid sea level changes on recent Azerbaijan ridges, *J. Coastal Res.*, 23, 521–527, 2007.
- Tusi, S. a.-D. M. b. M. A.: 'Ajā'eb al-Makhlūqāt va Ghara'eb al-Mojudat, 555 AH/1160 AD, edited by: Sotudeh, M., Bongāh-e Tarjomeh va Nashr-e Ketāb, Tehran, 1966.
- 5 Ulomov, V. I.: A three-dimensional model of the lithosphere dynamics, seismicity structure, and variations in the Caspian sea level, *Izv.-Phys. Solid Earth*, 39, 353–364, 2003.
- Voropaev, G. V., Krasnozhan, G. F., and Lahijani, H.: Caspian river deltas, *Caspia Bull.*, 1, 23–27, 1998.
- Walker, M. J. C.: *Quaternary Dating Methods*, John Wiley and Sons, Chichester, UK, 2005.

1431

Table 1. Geological findings on the Caspian sea-level (CSL) changes in the Late Holocene. The sea level position is inserted where the data are available.

Location	Event	Age (AD)	CSL	Reference
South CS	Sea-level rise and spit development	1042	–24	Lahijani et al. (2009)
South CS	Sea level rises to –24 m	1289–1403	–24	Kakroodi et al. (2012)
South CS	Relative sea level rise	1311–1445	–	Naderi Beni et al. (2013)
South CS	Sea level rises to –24 m	1335–1446	–24	Kakroodi et al. (2012)
South CS	Inundation of Gorgān Wall	1344–1460	–22	Rekavandi et al. (2007)
West CS	Sea-level rise and barrier formation	1350–1640	–24	Kroonenberg et al. (2007)
South CS	Relative sea-level rise	1408–1514	–	Naderi Beni et al. (2013)
South CS	Sea-level rise	1460	–25	Lahijani et al. (2009)
West CS	Sea-level rise and barrier formation	1590–1710	–24	Kroonenberg et al. (2007)
South CS	Sefidrud avulsion	1600	–	Lahijani et al. (2009)
West CS	Kura River diverted to Qezel Agac Bay	1600–1800	–	Hoogendoorn et al. (2005)
South CS	Age of the core base in Amirkolā	1620	–	Leroy et al. (2011b)
South CS	Relative sea-level rise	1696–1726	–	Naderi Beni et al. (2013)
South CS	Anzali Spit broken into barrier islands	1700–1830	–	Leroy et al. (2011b)
West CS	Development of new Kura Delta	1800	–	Hoogendoorn et al. (2005)
East CS	Sea-level rise	1830	–25.5	Leroy et al. (2006)

1432

Table 6. Historical earthquakes in the Caspian Sea region and the observed consequences on the sea level (Modified after Dotsenko et al., 2002).

Year (AD)	Location	Evidence
918	Derbent	Part of the coast with fortifications was submerged in the sea.
957–972	Derbent	The fall of sea level caused horizontal displacement of the shoreline by around 150 m from its normal position.
958	Ruyān	The Caspian Sea in Gilān was agitated by high tides
1668	Terka	The sea submerged part of the beach. The rise of water level was observed in the delta of the Terek River.
1868	Bāku	Short-term rise and fall of sea level with amplitude about 0.45 m were observed.
1876	Oblivnoy (island)	Unusual sea-level oscillations occurred after strong underwater explosion in conditions of dead calm. Event was observed from the ship.
1890	Tāsh	Large sea waves were noticed along the Iranian Caspian coast
1895	Cheleken Island	Flooding of north and west areas of Uzun-Ada as a result of a rise in water level in the bay. Large waves caused flooding of buildings and the dock. A few wooden houses were washed out to sea. Pipeline was destroyed.
1902	Bāku	Unusual waves resulted in dangerous motion of ships in the port. The event was observed after a destructive earthquake near Shimaha.
1910	Moghān	Large sea waves were observed in Āstārā
1933	Kuuli-Mayak	Sudden rise of sea level up to 1.35 m for 10 min. Fishing boats and equipment were washed out to sea.
1939	Livanov Shoal	The passing of a solitary large wave was observed from two ships that were 15 miles apart.
1960	Bāku	Sea-level oscillations up to 1 m were observed for 2–3 h.
1962	Bu'in	Irregularities were noticed at Anzali, Naushahr and Babolsar ports
1986	Livanov Shoal	Unusual high-frequency sea level oscillations of 2–3 cm amplitude were observed over the earthquake for 1–1.5 min. The event was fixed from the seiner and 45 fishing ships.

Table 7. The Caspian sea level (CSL) during the last millennium based on comparison between historical observations and geological events.

No	Age (AD)	CSL (m)	Historical observation	Reference	Geological event	Reference
1	907	–23	Sea level position in Ahlam, Chālus and Rāmsar	Hudud al-'Alam (1973), ibn Hawqal (1988), Jayhāni (1989)	Progradation of the old Kura Delta	Hoogendoorn et al. (2005)
2	977	–24	Sea level fall	Hudud al-'Alam (1973), ibn Hawqal (1988)	Sea-level fall	Hoogendoorn et al. (2005)
3	982	« –23.8	Bāb Island is in the map	Hudud al-'Alam (1973), ibn Hawqal (1988)	–	–
4	1208	–24	Ābeskun was on the shoreline	Al-Bakri (1999)	–	–
5	1260	> –24	Ābeskun was flooded	Jovayni (1911)	High-stand	Naderi Beni et al. (2013)
6	1304	–19	Rapid sea-level rise	Banāketi (1969), Mostowfi (1999), Marin Sanudo (1320 in Gümilev, 1980), Al-'Umarī (2010)	High-stand	Kakroodi et al. (2012), Naderi Beni et al. (2013), Rekavandi et al. (2007)
7	1587	–28	Construction of Safavid castle in Derbent	Gümilev (1980)	–	–
8	1628	–23	Establishment of ports and structures along the Caspian Sea coast	Parodi (1987), Gümilev (1980)	Sea-level rise and barrier formation	Kroonenberg et al. (2007)
9	1771	–23	Sea-level rise	Abbott (1858), Brückner (1890), Rabino (1980)	Widespread evidence	Leroy et al. (2011), Naderi Beni et al. (2013)
10	1815	–23.5	Sea-level position in Galugāh and Gomishān	Rabino (1980)	Anzali Spit broken into barriers	Leroy et al. (2011)
11	1875	–25	Sea-level rise at Anzali royal tower	Farhād Mirzā (1987)	Karā Bogāz Gol	Leroy et al. (2006)




Fig. 1. Modern map of the Caspian Sea and its seaboard countries. The positions of the Aral Sea, the main rivers as well as the main faults of the region are highlighted. The inset on the top left shows the whole watershed area of the Caspian Sea.

1439




Fig. 2. Caspian Sea-level changes over the last 10 000 yr, uncalibrated radiocarbon ages (Rychagov, 1997). The study period of the present investigation is denoted by the grey shading.

1440




Fig. 3. The Iranian Caspian coast, its prominent coastal landforms and major rivers. The cities mentioned in this paper are marked on the map. The morphological zones of the south Caspian coast (Voropaev et al., 1998) are highlighted. 1: Coasts with gentle slopes on the beach and in the nearshore zone. 2: Coasts with gentle slopes on the beach and steep slopes in the nearshore zone. 3: Coasts with steep slopes on the beach and nearshore zone. 4: Coasts with steep slopes on the beach and gentle slopes in the nearshore zone. Core locations are indicated by the star symbol.

1441




Fig. 4. Lithology of two cores taken near the Hassanqoli Bay in the southeast of the Caspian Sea. The position of the cores is presented in Fig. 3. The stars show the dated horizons and the corresponding radiocarbon ages (BP). GS: Grain Size; MS: Magnetic Susceptibility; TOM: Total Organic Matter. The vertical axis is depth in cm.

1442




Fig. 5. The Caspian sea-level (CSL) changes during the eighteenth century and first half of the nineteenth century (Fedchina, 1980) and the sea-level curve from Komarova (1980) based on Russian cartographic maps.

1443




Fig. 6. The Caspian sea-level curve based on Brückner (1890), Varushchenko et al. (1987) and Karypychev (1998, 2001). The dashed lines connecting the filled symbols are interpolations. The continuous line from 1850 to 2000 shows the instrumental observations. A -26.5 line, the CSL in 1995, was added for ease of comparing the different levels during the millennium. The Medieval Climate Anomaly (MCA) and the Little Ice Age (LIA) are indicated by shaded boxes.

1444




Fig. 7. The tenth century map of the Caspian Sea by Ibn Hawqal (1988). The map was rotated to show the north at the top and some of the names have been translated into English. Ābeskun, Gorgān, Ahlam and Chālus were described as the most important ports of the south Caspian Sea by the tenth century geographer.

1445




Fig. 8. The Iranian Caspian coast and reported historical observations. The ancient names are denoted in parentheses: (1) The possible position of sea level in Dahestān in second half of the tenth century (al-Istakhri, 1961, 1973; Barthold, 1984; ibn Hawqal, 1988). (2) The probable position of Nim-mardān Bay (Mostowfi, 1999; Kakroodi et al., 2012). (3) The probable position of Ābeskun along the Caspian shoreline (Jovayni, 1911; al-Istakhri, 1961; ibn Hawqal, 1988; Mostowfi, 1999; al-Mas'ūdī, 2012). (4) The sea-level position in 1006 in Galougāh (Rabino, 1980). (5) The position of Ashraf port in 1628 (Parodi, 1997). (6) Uzbeks shipped the captives of Māzandarān from Sāri (Mostowfi, 1999). (7) Sea-level position in Ahlam in the ninth century (al-Ya'qūbī, 1968). (8) Sea-level position in Ahlam reported by ibn Hawqal (1988), Mostowfi (1999) and Mar'ashi (1982). (9) Sea-level position in Chālus reported by ibn Hawqal (1988) and Jayhāni (1989) in tenth century. (10) Sea-level position in an area between Daylam and Māzandarān in the tenth century (ibn Hawqal, 1988; Jayhāni, 1989). (11) The Sefidrud or its tributary course in the tenth century (Hudud al-'Alam, 1973). (12) The main course of Sefidrud and its tributary in the eighteenth and nineteenth centuries (Rabino, 1917; Barthold, 1984; Farhād Mirzā, 1987). (13) The Anzali Spit broke into barrier islands in the fifteenth century (Mar'ashi, 1982).

1446

