

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

This discussion paper is/has been under review for the journal Climate of the Past (CP). Please refer to the corresponding final paper in CP if available.

On the effect of orbital forcing on mid-Pliocene climate, vegetation and ice sheets

M. Willeit, A. Ganopolski, and G. Feulner

Potsdam Institute for Climate Impact Research, Potsdam, Germany

Received: 24 March 2013 – Accepted: 25 March 2013 – Published: 3 April 2013

Correspondence to: M. Willeit (willeit@pik-potsdam.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We present results from modeling of the mid-Pliocene warm period (3.3–3 million years ago) using the Earth system model of intermediate complexity CLIMBER-2 analyzing the effect of changes in boundary conditions as well as of orbital forcing on climate.

5 Firstly we performed equilibrium experiments following PlioMIP (Pliocene Model Intercomparison Project) protocol with a CO₂ concentration of 405 ppm, reconstructed mid-Pliocene orography and vegetation and a present day orbital configuration. Simulated global Pliocene warming is about 2.5°C, fully consistent with results of atmosphere-ocean general circulation model simulations performed for the same modeling setup. A factor separation analysis attributes 1.5°C warming to CO₂, 0.3°C to
10 orography, 0.2°C to ice sheets and 0.4°C to vegetation.

Transient simulations for the entire mid-Pliocene warm period with time-dependent orbital forcing as well as interactive ice sheets and vegetation give a global warming varying within the range 1.9–2.8°C. Ice sheet and vegetation feedbacks in synergy
15 act as amplifiers of the orbital forcing transforming seasonal insolation variations into an annual mean temperature signal. The effect of orbital forcing is more significant at high latitudes, especially during summer, when the warming over land varies in the wide range from 0–10°C. The modeled ice sheet extent and vegetation distribution also show significant temporal variations. Modeled and reconstructed data for North-
20 ern Hemisphere sea surface temperatures and vegetation distribution show the best agreement if the reconstructions are assumed to be representative for the “warmest” periods during the orbital cycles. This suggests that low-resolution Pliocene paleoclimate reconstructions can reflect not only the impact of increased CO₂ concentrations and topography changes but also the effect of orbital forcing. Therefore, the climate
25 (Earth system) sensitivity estimates from Pliocene reconstructions which do not account for the effect of orbital forcing can be biased toward high values.

CPD

9, 1703–1734, 2013

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

The time period between 3.3 and 3 million years ago (Mya), traditionally named mid-Pliocene warm period (MPWP), is the most recent interval in the Earth history which is believed to have been significantly warmer than the preindustrial climate. Most paleoproxies suggest that atmospheric CO₂ concentration during the MPWP was in the range from 300 to 400 ppm (Raymo et al., 1996; Pagani et al., 2009; Seki et al., 2010; Bartoli et al., 2011), the global surface air temperature was higher by 2–3 °C (estimate based on model simulations) (Haywood and Valdes, 2004; Jansen et al., 2007; Lunt et al., 2012b) and sea level was up to 25 m above present during the warmest periods of the MPWP (Miller et al., 2012; Foster and Rohling, 2013). If these estimates are sufficiently accurate, this makes the MPWP a useful (although by no means direct) analogue for the future greenhouse world (Salzmann et al., 2009; Haywood et al., 2011b).

A comprehensive set of reconstructions of climate and boundary conditions for the MPWP is available from the PRISM3D (Pliocene Research, Interpretation and Synoptic Mapping) data set (Dowsett et al., 2010). PRISM3D includes global reconstructions for sea surface temperature (SST), vegetation distribution, ice sheet extent and orography. Most of the data are the outcome of a combination of reconstructed data and modeling results. SST reconstructions show little changes in the tropics relative to present-day while significant warming is observed at mid to high latitudes, up to 3–4 °C in the Northern Hemisphere (NH) with major warming occurring in the North Atlantic (Dowsett et al., 2009; Dowsett and Robinson, 2009). There are numerous pieces of evidence from paleodata that the spatial distribution of vegetation during the MPWP was substantially different from today. The main difference was a northward expansion of taiga replacing tundra in the NH. The coast of Antarctica was at least locally covered by some vegetation. The vegetation distribution in mid and low latitudes was essentially similar to present day, although the moister climate favored forest and woodland growth in some regions now covered by steppe or grassland (Thompson and Fleming,

CPD

9, 1703–1734, 2013

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1996). The PRISM3D vegetation reconstruction is based on a combined model-data approach and is described in Salzmann et al. (2008). It combines an internally consistent dataset of reconstructions at 202 palaeobotanical sites with results of simulations of the BIOME4 vegetation model driven by modeled mid-Pliocene climatology.

5 Where data coverage is poor the vegetation reconstruction is mostly based on the model output. Some important regions with low data availability include high northern Eurasia and North American mid-latitudes. This should be kept in mind when comparing model output with reconstructions. More applications of the BIOME4 model to the MPWP can be found in, e.g. Haywood et al. (2002b) and Salzmann et al. (2009).
10 Simulations with the BIOME4 model in offline mode do not account for the feedback of the vegetation on climate. MPWP simulations with a coupled climate-vegetation model showed a generally reasonably good agreement with reconstructed vegetation (PRISM2) but regionally important differences exist, with the model underestimating the northward expansion of coniferous forest and overestimating the extension of arid areas (Haywood and Valdes, 2006). MPWP sea level has been estimated to be up to 25 m higher than present during warmest periods of the MPWP corresponding to the “warm” orbital configurations, which implies significant reduction of the Greenland Ice Sheet and the Antarctic Ice Sheet (Pollard and DeConto, 2009; Miller et al., 2012; Foster and Rohling, 2013). PRISM3D spatially explicit estimation of ice sheet extent
20 is primarily based on modeling approaches (Hill et al., 2007). Time slice modeling experiments with different MPWP orbital configurations suggest that the Greenland ice sheet varied from almost ice-free to nearly present-day state, while Antarctic ice volume showed variations up to 30 % (Dolan et al., 2011). PRISM3D topography, which we use in this study, is more similar to present day than that in PRISM2, with small elevation reductions in the Andes and the Rocky Mountains (Rowley and Garzzone, 2007). Because of a coarse spatial resolution of the CLIMBER-2 model we first aggregate anomalies of MPWP orography from the PRISM3D dataset to the CLIMBER grid and then add them to present day model orography.
25

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Some attempts were made to use the MPWP to derive the so-called Earth System Sensitivity (ESS), i.e. the long-term quasi-equilibrium response of global temperature to radiative forcing. In contrast to the Climate Sensitivity (CS, also called Charney sensitivity) which involves only fast feedbacks, ESS includes also the slow feedbacks, related to ice sheets, vegetation and atmospheric aerosols. The ESS estimates from MPWP range from modest 4.5°C (Lunt et al., 2009) to very high 7.1–8.7°C (Pagani et al., 2009) values. This large range can be explained by uncertainties in reconstructed CO₂ concentrations and model derived global temperatures. Similar attempts have been made to estimate the dependence of global mean sea level on temperature or CO₂ (e.g. Foster and Rohling, 2013). Obviously, any attempt to employ MPWP data to constrain climate models and climate sensitivity depend not only on the accuracy of paleoclimatic reconstructions, but also on the validity of the implicit assumption that reconstructed MPWP climate represents quasi-equilibrium response to elevated level of CO₂ and modified topography. Testing this assumption has important implications for the interpretation of MPWP reconstructions. Even if climatic conditions during the MPWP were relatively stable compared to the more recent past, significant variability is evident in the benthic oxygen isotope ($\delta^{18}\text{O}$) record (Lisiecki, 2005) suggesting a not negligible response of the climate system to orbital forcing. PRISM3D SSTs are well known to represent warmest conditions during the MPWP because of the warm peak averaging procedure involved in the reconstruction (Dowsett, 2007). The temporal resolution of vegetation proxies, mostly fossil pollen, is very coarse and consequently vegetation reconstructions do not resolve orbital variations (Salzmann et al., 2008; Thompson and Fleming, 1996). Where information on vegetation variability in time is available, the biome representing the warmest climatic conditions has been included in the PRISM3D reconstructions (Haywood et al., 2013b). Furthermore, in regions with poor data coverage PRISM3D vegetation is heavily based on the BIOME4 simulation driven by climatology from an atmospheric general circulation model (AGCM) with prescribed PRISM2 boundary conditions, including SSTs (Salzmann et al., 2008). Since

reconstructed SSTs represent warm MPWP conditions, it is possible that vegetation reconstructions are also warm biased in regions with scarce data availability.

Up to now most simulations of MPWP climate represent equilibrium simulations with atmosphere general circulation models with prescribed SSTs (e.g. Chandler et al., 1994; Sloan et al., 1996; Haywood et al., 2000; Jiang, 2005; Haywood et al., 2009) or with coupled atmosphere-ocean climate models (e.g. Haywood and Valdes, 2004; Haywood et al., 2007; Lunt et al., 2007). In both cases, vegetation distribution and ice sheet extent were prescribed. The most recent coordinated efforts as a part of PMIP3/CMIP5 (Paleoclimate Model Intercomparison Project/Coupled Model Intercomparison Project) is PlioMIP (Pliocene Model Intercomparison Project) (Haywood et al., 2010, 2011a). In the case of PlioMIP, the protocol includes present day orbital parameters, CO₂ concentration of 405 ppm with other greenhouse gases (GHGs) at the preindustrial level, as well as orography, ice sheets and vegetation from PRISM3D reconstructions. Results of eight models participating in PlioMIP reveal a wide range of global temperature responses to identical forcings and boundary conditions (1.8–3.6 °C) and similarly wide range (3.5–6.8 °C) for the implied ESS, with some models having an ESS almost identical to the CS, while others have an ESS twice as large as the CS (Haywood et al., 2013a). Typical feature of most of simulations of the MPWP with coupled climate models is an overestimation of warming in low latitudes and underestimation of warming in high latitudes compared to available paleoclimate reconstructions. The discrepancies between data and models in tropical temperatures are present not only in simulations of the MPWP but also other paleoclimate simulations both for warm and cold climates (Lunt et al., 2012a). The reasons for such systematic difference remain debatable. Weaker simulated MPWP warming in high latitudes may have several reasons. One is that climate models underestimate the strength of climate feedbacks in the high latitudes. One such feedback is the climate-vegetation feedback, which in coupled climate models is accounted via prescribed MPWP vegetation. Another possibility is that the standard PlioMIP setup does not account for some orographic changes, such as changes in ocean bathymetry which can affect ocean circulation (Robinson et al.,

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2011). At last, low temporal resolution of most of Pliocene paleoclimate records make it possible that some of the reconstructions are warm-biased and represent climate conditions corresponding to “warm” orbital configurations rather than mean conditions over the entire 3.3–3 Mya period.

5 The effect of orbital forcing has previously been analyzed with time slice experiments using an AGCM coupled to a slab ocean (Haywood et al., 2002a). For different orbital configurations they find small annual changes in SSTs and land surface temperatures, but seasonally more significant variability due to insolation distribution changes. Further they show only minor variations in vegetation cover in offline experiments with the BIOME4 model. However, their model did not include interactive vegetation and ice sheets which can amplify the direct effect of orbital forcing.

10 So far the role of orbital forcing on MPWP climate, vegetation and ice sheets has not been analyzed with a fully coupled model. Here we present results from transient simulations using an Earth system model driven by time-dependent orbital forcing including experiments with interactive vegetation and ice sheets.

2 Model description and experimental setup

2.1 Model

For our analysis we use the Earth system model of intermediate complexity CLIMBER-2 (Petoukhov et al., 2000; Ganopolski et al., 2001). CLIMBER-2 includes a 2.5-dimensional statistical-dynamical atmosphere model, a 3-basin zonally averaged ocean model including sea ice, a dynamic model of the terrestrial vegetation (Brovkin et al., 1997, 2002) and the 3-dimensional polythermal ice sheet model SICOPOLIS based on shallow ice approximation (Greve, 1997). The latter is only applied for the Northern Hemisphere. Different components of the model have different spatial resolution. Atmospheric and land components have latitudinal resolution of 10° and longitudinal of approximately 51°. The ocean model has a latitudinal resolution of 2.5° and 21

CPD

9, 1703–1734, 2013

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



levels vertically. The ice sheet model has a latitudinal resolution of 0.75° , longitudinal resolution of 1.5° and vertically 20 levels in sigma-coordinates. Atmosphere and ice sheets are coupled bi-directionally using a physically based energy balance approach (Calov et al., 2005). Surface runoff and ice calving simulated by the ice sheet model are applied to the ocean as surface freshwater flux. Since CO_2 concentration was prescribed, the marine carbon cycle model was not used in this study. The CLIMBER-2 model in different configurations has been used for numerous studies of past and future climates (e.g. Ganopolski et al., 1998, 2001; Claussen et al., 2006; Schneider von Deimling et al., 2006; Ganopolski and Calov, 2011; Brovkin et al., 2012).

2.2 MPWP boundary conditions

In all MPWP experiments (equilibrium and transient) we prescribed global orography according to PRISM3D dataset. Since our ice sheet model is only applied to the Northern Hemisphere, in all experiments, the Antarctic ice sheet extent and elevation were also prescribed according to PRISM3D. Following the PlioMIP protocol (Haywood et al., 2010), the CO_2 concentration was set to 405 ppm in all experiments. The 405 ppm value is at the high end of concentration estimates (Pagani et al., 2009; Seki et al., 2010) but is chosen to account for unknown additional contributions from non- CO_2 greenhouse gases such as methane. In the set of equilibrium simulations which followed PlioMIP protocol, we also prescribed Greenland ice sheets and vegetation according to PRISM3D reconstructions (Dowsett et al., 2010). PRISM3D data are adjusted and aggregated to the CLIMBER-2 resolution. For vegetation we used the PRISM3D product in its megabiome version. Because in CLIMBER-2 only forest, grass and deserts are distinguished we aggregated the megabiome data to these three classes at the model resolution (Appendix A). As a final result we obtained the fraction of forest, grass and desert cover in each grid cell. Additionally, annual mean values of the leaf area index are derived from the vegetation cover in a semi-interactive vegetation model setup with fixed cover fractions. Orbital parameters in PlioMIP experiments were set to their present day values. In addition to the PlioMIP type of experiment we

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



performed a set of transient experiments through the entire MPWP period with time-dependent orbital forcing computed according to Berger and Loutre (1992). In these experiments the vegetation cover and the NH ice sheets were either interactive or prescribed according to PRISM3D (see below).

2.3 PlioMIP experiment and factor separation analysis

Since CLIMBER-2 is a model of intermediate complexity, to enhance credibility of modeling results, it is useful to compare its results with available results of more complex climate models for identical experiments. Since transient MPWP experiments have not yet been undertaken with coupled AOGCMs, we performed with CLIMBER-2 experiments using the PlioMIP protocol. In addition, to disentangle the effects of CO₂ concentration, orography, vegetation and ice sheets on the mid-Pliocene climate, a full factor separation is performed following the technique described in Lunt et al. (2012b). For four factors this requires running 2⁴ = 16 different experiments, with all possible combination of changes in the boundary conditions. The total global annual MPWP temperature change will be a combination of the contribution from the single factors. The way the single contributions are computed is described in Appendix B.

3 Transient experiments

To estimate the effect of orbital forcing on the MPWP climate, ice sheets and vegetation we perform a suite of transient experiments with time dependent orbital forcing. With respect to both ice sheets and vegetation we performed three types of experiments: the first one with prescribed present-day ice sheets (vegetation), the second with prescribed PRISM3D ice sheets (vegetation) and the third with interactive ice sheets (vegetation).

CPD

9, 1703–1734, 2013

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Temperature and precipitation anomalies simulated for the PlioMIP modeling setup are illustrated in Figs. 1a and 2a, respectively. Our spatial patterns for temperature compare favorably well with the model mean from PlioMIP (Haywood et al., 2013a). For precipitation our results are consistent with the wide range simulated by different models in PlioMIP (Haywood et al., 2013a). The geographical distribution of the separate effect of CO₂, orography, ice sheets and vegetation on annual mean temperature and precipitation are shown in Figs. 1 and 2 together with the synergy between the factors. The main contribution to the total signal is from the higher CO₂ concentration, while the other factors can be equally important but only on a regional scale. Ice sheets have a very local impact limited to Greenland and Antarctica. Regarding temperature, vegetation has a significant effect in high latitudes and over an extended region around northern Africa, Europe and central Asia. All factors are important for the warming in polar regions. Orography changes are also important in south-east Asia and South America. The largest changes in precipitation occur in the tropics and have significant contributions from CO₂, orography and vegetation. The synergy term is small for both temperature and precipitation. Our spatial patterns for the annual temperature factor separation compare favorably well with those from Lunt et al. (2012b) except for orography, where discrepancies are attributable to different versions of PRISM datasets used. As already pointed out for global mean values, also regionally differences exist regarding the amplitude of the warming, with results from Lunt et al. (2012b) showing higher temperature anomalies.

We additionally performed an experiment with PRISM3D boundary conditions but interactive vegetation. Simulated global temperature in experiment with interactive vegetation is 0.2 °C lower compared to the experiment with PRISM3D vegetation, i.e. with interactive vegetation and present day orbital configuration the model can explain only half of the warming attributed to vegetation by the factor separation analysis. A comparison of the modeled forest cover with its PRISM3D counterpart is presented in Fig. 6c,d. Overall the model shows lower forest fractions compared to the reconstructions. The discrepancy is particularly evident in high northern latitudes where modeled forest area

is significantly lower than suggested by MPWP reconstruction. We will discuss the effect of orbital forcing on vegetation in the next section.

4.2 Transient runs

The time series of global mean annual temperature differences from preindustrial for different transient experiments of the MPWP are presented in Fig. 3b. Mid-Pliocene warming of $\Delta T_{\text{PlioMIP}} = 2.5^\circ\text{C}$ simulated in the PlioMIP experiment is shown for comparison. Including time dependent orbital forcing in the experiment with interactive vegetation results in global temperature varying by 0.5°C in the MPWP interval, with values up to 2.7°C during “warmest” orbital configurations. Temperature variability on the orbital time scales is dominated by the precessional component of orbital forcing as was shown already in Claussen et al. (2006). Interestingly enough, the precessional component is only clearly seen during the last 100 kyr (1 kyr = 1000 yr) of the MPWP. To clarify the origin of such nonlinear response to the precessional component of the orbital forcing, we performed an additional experiment where the fraction of vegetation was fixed in the Sahara/Sahel region (between 10°N and 30°N) but interactive elsewhere. Fixing Sahara/Sahel vegetation leads to significant weakening of precessional variability which unambiguously points on surface albedo in Sahara/Sahel region as the main cause for the precessional component of global temperature variability. Since parameterization of surface albedo in the CLIMBER-2 model is rather crude, it is possible that the effect of vegetation on surface albedo is overestimated. On the other hand, the “greening” of Sahara would reduce the atmospheric dust load and, as a result, lower the planetary albedo and warm the NH. This effect is not accounted for in PlioMIP-type simulations. In addition to the precessional cycle, global mean temperature also contains a pronounced 100 kyr cycle. This cycle is not related to the vegetation, because the same variability is observed also in a transient simulation with fixed vegetation (not shown). This 100 kyr cyclicity is explained by corresponding eccentricity variations (Fig. 3a) which causes global annual mean insolation variations of up to 0.5Wm^{-2} (Paillard, 2001). Global temperature in simulations with interactive ice sheets but fixed

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



vegetation is also dominated by the precessional component. This is related to a strong response of the NH ice sheets to the summer insolation (Fig. 4). During periods of high insolation, large portions of the Greenland ice sheet disappeared, while during cold summers Greenland ice sheet occupies the entire island, and a relatively small ice sheet grows over the Arctic archipelago. As it was shown in Ganopolski and Calov (2011), high levels of CO_2 prevent the Earth from falling into the full glacial state. Previous studies have already addressed the effect of orbital forcing on ice sheets in offline time-slice experiments of the MPWP showing very different ice sheets states under different orbital configurations (Dolan et al., 2011). Considerable variations in NH ice volume are also modeled in our coupled transient experiments (Fig. 4). The magnitude of NH ice sheet volume variations corresponds to approximately 15 m sea level equivalent, which is not in conflict with magnitude of benthic $\delta^{18}\text{O}$ variability (Lisiecki, 2005). However, modeled ice-volume variability is dominated by precession, while the benthic $\delta^{18}\text{O}$ record is dominated by obliquity. This fact, at least partly, can be explained by a compensation of precessional variability in the ice volume between the two hemispheres (Raymo et al., 2006; Dolan et al., 2011). The combined effect of interactive ice sheets and vegetation results in further amplification of the precessional forcing. Global mean modeled warming varies now within the even larger range 1.9–2.8°C. Again, the best agreement with $\Delta T_{\text{Pliocene}}$ is found in “warm” orbital intervals.

More detailed information can be inferred from zonal temperature anomalies (Fig. 5). The most striking feature is the large temperature variability in the fully coupled transient experiment both over land and ocean. The variability is particularly significant in summer over land where the temperature anomalies span a 10°C interval, with temperatures ranging from values similar to present day to up to 10°C higher. This has fundamental implications for the interpretation of reconstructions over both ocean and land, since dating accuracy and temporal resolution of MPWP proxy records are generally not high enough to resolve orbital time scales. Again, the zonal temperatures for the Pliocene like experiment are shown for comparison. The mean MPWP zonal temperatures of the fully coupled transient run are generally lower than in the

compared to the preindustrial state everywhere except above 60° N, where forest expands at the expense of tundra. However, the large increase in forest fraction between 70–80° N suggested by paleoclimate reconstruction cannot be explained by the higher CO₂ concentration alone. Here, a generally better agreement between model and data is found in the transient runs during “warm” orbital periods (Fig. 7). This results are apparently in contrast with those from Haywood et al. (2002b), where they show small variations in vegetation cover in time slice experiments with different orbital configurations.

5 Conclusions

Results of our MPWP simulations with PRISM3D boundary conditions and present day orbital parameters are consistent with the results of PlioMIP experiments. The global annual mid-Pliocene warming relative to preindustrial is 2.5 °C, regionally overestimated in the tropics and underestimated in the high latitudes compared with paleoclimate reconstructions. Changes in CO₂ concentrations, orography, vegetation and ice sheets explain 62 %, 18 %, 13 % and 8 % of the warming, respectively. In transient simulations including time-dependent orbital forcing, interactive vegetation and NH ice sheets we found that a warming comparable with the PlioMIP experiment is found only during time intervals with “warm” orbital configurations. Global warming during the MPWP varies in the range from 1.9 °C to 2.8 °C. This results from non-linearities in the climate system. Feedbacks between climate, vegetation and ice sheets amplify the precessional component of the orbital forcing, transforming seasonal insolation variations into an annual mean temperature signal. Regionally, and especially seasonally, the effect of orbital forcing is comparable or even more important than higher CO₂ and orographic changes. The most significant effect of orbital forcing is found in summer over NH land, where temperatures vary by up to 10 °C during the 300 kyr interval. Simulated response of vegetation and ice sheets to the orbital forcing is much larger than to change in CO₂ concentration alone. Only a combination of elevated CO₂ and

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



“optimal” orbital configuration leads to SST and vegetation cover being more consistent with reconstructions.

In summary, we find that orbital forcing plays a fundamental role in controlling modeled MPWP climate- and consequently vegetation distribution and ice sheet extent. If the MPWP reconstructions are warm-biased and therefore contain orbital fingerprints, then a large portion of discrepancies between data and model can be reconciled. This is the case for NH SSTs, boreal vegetation cover and ice sheet extent, suggesting that at least some of the proxy data may reflect not only the effect of increased CO₂ concentration and changed topography, but also the effect of orbital forcing. This has wider implications for the ongoing attempts to estimate climate/Earth system sensitivity with a combined model-reconstructions approach. Our results imply that Earth system sensitivity estimates based on MPWP reconstructions are likely to be biased to high values. As has already been proposed elsewhere (Haywood et al., 2013b), a possible solution to this problem is to use reconstructions from a time-slice with close to present-day orbital configuration instead of the whole MPWP slab, although it will be a challenge to obtain enough data to reconstruct a mid-Pliocene time slice, especially for vegetation.

Our modeling results do not support a notion that the Earth system sensitivity is significantly higher than the classical climate (or Charney) sensitivity. Indeed, results of model simulations (Dolan et al., 2011) in conjunction with the existence of a strong temporal variability of the benthic $\delta^{18}O$ on orbital time scales, suggest that a significant fraction of the ice sheets reduction during the warmest intervals of MPWP was related to the orbital forcing rather than elevated CO₂. As far as vegetation feedback is concerned, it amplifies CO₂ induced temperature change by only 0.2°C in our model (less than 10% of climate sensitivity), which is consistent with previous studies (e.g. O’ishi et al., 2009). Admittedly, there are climate feedbacks which are not included in our model and which can be important for amplification of CO₂ induced warming, such as aerosols (e.g. aeolian dust), atmospheric chemistry, etc. However, the fact that our model is able to simulate MPWP climate anomaly in reasonable agreement with

CPD

9, 1703–1734, 2013

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reconstructions can be interpreted as evidence against the existence of very strong and unaccounted climate feedbacks.

Appendix A

Vegetation aggregation

5 The PRISM3D megabiome vegetation dataset distinguishes 9 plant functional types (PFTs), while CLIMBER-2 differentiates only between forest and grassland. In CLIMBER-2 each gridcell on land is composed by a mosaic of forest, grass and desert with fractional areas computed by the vegetation module. On the other hand, in PRISM3D a unique PFT or desert is assigned to each grid cell. According to the
10 PRISM3D – CLIMBER-2 transfer function between surface types in Table 2 we transform the PRISM3D from 10 to 3 surface types keeping the $2 \times 2^\circ$ resolution. Then for each $10 \times 51^\circ$ CLIMBER-2 cell we count the number of corresponding $2 \times 2^\circ$ cells covered by forest, grass and desert covering the same area, and from that compute the forest, grass and desert share which is then used as boundary condition for CLIMBER-
15 2.

Appendix B

Factor separation

Adopting the notation from Lunt et al. (2012b) we indicate with a subscript the boundary conditions that are prescribed from PRISM3D (c for CO_2 , o for orography, i for ice sheets and v for vegetation). E.g. T_{iv} is the temperature from the experiment with prescribed PRISM3D ice and vegetation, but with present day CO_2 and orography. Assuming that the single factors are independent from each other and including the
20

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



non-linearities in the synergy term S we can write, using Taylor expansion (Lunt et al., 2012b):

$$\Delta T = dT_{\text{CO}_2} + dT_{\text{oro}} + dT_{\text{ice}} + dT_{\text{veg}} + S \quad (\text{B1})$$

where the single terms can be computed from the 16 model simulations with different combinations of boundary conditions as follows:

$$dT_{\text{CO}_2} = \frac{1}{8}((T_c - T) + (T_{\text{co}} - T_o) + (T_{\text{ci}} - T_i) + (T_{\text{cv}} - T_v) + (T_{\text{cov}} - T_{\text{ov}}) + (T_{\text{cio}} - T_{\text{io}}) + (T_{\text{civ}} - T_{\text{iv}}) + (T_{\text{ciov}} - T_{\text{ioiv}})) \quad (\text{B2})$$

$$dT_{\text{oro}} = \frac{1}{8}((T_o - T) + (T_{\text{co}} - T_o) + (T_{\text{oi}} - T_i) + (T_{\text{ov}} - T_v) + (T_{\text{cov}} - T_{\text{cv}}) + (T_{\text{cio}} - T_{\text{ci}}) + (T_{\text{oiv}} - T_{\text{iv}}) + (T_{\text{ciov}} - T_{\text{civ}})) \quad (\text{B3})$$

$$dT_{\text{ice}} = \frac{1}{8}((T_i - T) + (T_{\text{oi}} - T_o) + (T_{\text{ci}} - T_c) + (T_{\text{iv}} - T_v) + (T_{\text{ioiv}} - T_{\text{ov}}) + (T_{\text{cio}} - T_{\text{co}}) + (T_{\text{civ}} - T_{\text{cv}}) + (T_{\text{ciov}} - T_{\text{cov}})) \quad (\text{B4})$$

$$dT_{\text{veg}} = \frac{1}{8}((T_v - T) + (T_{\text{ov}} - T_o) + (T_{\text{iv}} - T_i) + (T_{\text{cv}} - T_c) + (T_{\text{cov}} - T_{\text{co}}) + (T_{\text{ioiv}} - T_{\text{io}}) + (T_{\text{civ}} - T_{\text{ci}}) + (T_{\text{ciov}} - T_{\text{cio}})) \quad (\text{B5})$$

The synergy between the factors is given by:

$$S = T_{\text{ciov}} - T + dT_{\text{CO}_2} + dT_{\text{oro}} + dT_{\text{ice}} + dT_{\text{veg}}. \quad (\text{B6})$$

Acknowledgements. The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7 2007–2013) under grant agreement no. 238366.

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Bartoli, G., Hönisch, B., and Zeebe, R. E.: Atmospheric CO₂ decline during the Pliocene intensification of Northern Hemisphere glaciations, *Paleoceanography*, 26, 1–14, 2011. 1705
- Berger, A. and Loutre, M.: Astronomical solutions for paleoclimate studies over the last 3 million years, *Earth Planet. Sc. Lett.*, 111, 369–382, 1992. 1711, 1730
- Brovkin, V., Ganopolski, A., and Svirezhev, Y.: A continuous climate-vegetation classification for use in climate-biosphere studies, *Ecol. Model.*, 101, 251–261, 1997. 1709
- Brovkin, V., Bendtsen, J., Claussen, M., Ganopolski, A., Kubatzki, C., Petoukhov, V., and Andreev, A.: Carbon cycle, vegetation, and climate dynamics in the Holocene: experiments with the CLIMBER-2 model, *Global Biogeochem. Cy.*, 16, 86.1–86.20, 2002. 1709
- Brovkin, V., Ganopolski, A., Archer, D., and Munhoven, G.: Glacial CO₂ cycle as a succession of key physical and biogeochemical processes, *Clim. Past*, 8, 251–264, doi:10.5194/cp-8-251-2012, 2012. 1710
- Calov, R., Ganopolski, A., Claussen, M., Petoukhov, V., and Greve, R.: Transient simulation of the last glacial inception. Part I: Glacial inception as a bifurcation in the climate system, *Clim. Dynam.*, 24, 545–561, 2005. 1710
- Chandler, M., Rind, D., and Thompson, R.: Joint investigations of the middle Pliocene climate II: GISS GCM Northern Hemisphere results, *Global Planet. Change*, 9, 197–219, 1994. 1708
- Claussen, M., Fohlmeister, J., Ganopolski, A., and Brovkin, V.: Vegetation dynamics amplifies precessional forcing, *Geophys. Res. Lett.*, 33, 1–5, 2006. 1710, 1714
- Dolan, A. M., Haywood, A. M., Hill, D. J., Dowsett, H. J., Hunter, S. J., Lunt, D. J., and Pickering, S. J.: Sensitivity of Pliocene ice sheets to orbital forcing, *Palaeogeogr. Palaeoclimatol.*, 309, 98–110, 2011. 1706, 1715, 1718
- Dowsett, H.: The PRISM palaeoclimate reconstruction and Pliocene sea-surface temperature, in: *Deep-Time Perspectives on Climate Change: Marrying the Signal from Computer Models and Biological Proxies*, edited by: Williams, M., Haywood, A. M., Gregory, F. J., and Schmidt, D. N., The Geological Society, Micropaleontological Society Special Publications, London, 459–480, 2007. 1707
- Dowsett, H. J. and Robinson, M. M.: Mid-Pliocene equatorial Pacific sea surface temperature reconstruction: a multi-proxy perspective., *Philos. T. R. Soc. A*, 367, 109–125, 2009. 1705

CPD

9, 1703–1734, 2013

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dowsett, H. J., Chandler, M. A., and Robinson, M. M.: Surface temperatures of the Mid-Pliocene North Atlantic Ocean: implications for future climate, *Philos. T. R. Soc. A*, 367, 69–84, 2009. 1705
- Dowsett, H., Robinson, M., Haywood, A., Salzmann, U., Hill, D., Sohl, L., Chandler, M., Williams, M., Foley, K., and Stoll, D.: The PRISM3D paleoenvironmental reconstruction, *Stratigraphy*, 7, 123–139, 2010. 1705, 1710
- Foster, G. L. and Rohling, E. J.: Relationship between sea level and climate forcing by CO₂ on geological timescales., *P. Natl. Acad. Sci. USA*, 110, 1209–1214, 2013. 1705, 1706, 1707
- Ganopolski, A. and Calov, R.: The role of orbital forcing, carbon dioxide and regolith in 100 kyr glacial cycles, *Clim. Past*, 7, 1415–1425, doi:10.5194/cp-7-1415-2011, 2011. 1710, 1715
- Ganopolski, A., Kubatzki, C., Claussen, M., Brovkin, V., and Petoukhov, V.: The influence of vegetation-atmosphere-ocean interaction on climate during the mid-holocene, *Science*, 280, 1916–1919, 1998. 1710
- Ganopolski, A., Petoukhov, V., Rahmstorf, S., Brovkin, V., Claussen, M., Eliseev, A., and Kubatzki, C.: CLIMBER-2: a climate system model of intermediate complexity. Part II: Model sensitivity, *Clim. Dynam.*, 17, 735–751, 2001. 1709, 1710
- Greve, R.: Application of a polythermal three-dimensional ice sheet model to the Greenland ice sheet: response to steady-state and transient climate scenarios, *J. Climate*, 10, 901–918, 1997. 1709
- Haywood, A. M. and Valdes, P. J.: Modelling Pliocene warmth: contribution of atmosphere, oceans and cryosphere, *Earth Planet. Sc. Lett.*, 218, 363–377, 2004. 1705, 1708
- Haywood, A. M. and Valdes, P. J.: Vegetation cover in a warmer world simulated using a dynamic global vegetation model for the Mid-Pliocene, *Palaeogeogr. Palaeocl.*, 237, 412–427, 2006. 1706
- Haywood, A., Valdes, P., and Sellwood, B.: Global scale palaeoclimate reconstruction of the middle Pliocene climate using the UKMO GCM: initial results, *Global Planet. Change*, 25, 239–256, 2000. 1708
- Haywood, A., Valdes, P., and Sellwood, B.: Magnitude of climate variability during middle Pliocene warmth: a palaeoclimate modelling study, *Palaeogeogr. Palaeocl.*, 188, 1–24, 2002a. 1709
- Haywood, A. M., Valdes, P. J., Francis, J. E., and Sellwood, B. W.: Global middle Pliocene biome reconstruction: a data/model synthesis, *Geochem. Geophys. Geosy.*, 3, 1–18, 2002b. 1706, 1717

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Haywood, A. M., Valdes, P. J., and Peck, V. L.: A permanent El Niño-like state during the Pliocene?, *Paleoceanography*, 22, PA1213, doi:10.1029/2006PA001323, 2007. 1708
- Haywood, A. M., Chandler, M. A., Valdes, P. J., Salzmann, U., Lunt, D. J., and Dowsett, H. J.: Comparison of mid-Pliocene climate predictions produced by the HadAM3 and GCMAM3 general circulation models, *Global and Planet. Change*, 66, 208–224, 2009. 1708
- Haywood, A. M., Dowsett, H. J., Otto-Bliesner, B., Chandler, M. A., Dolan, A. M., Hill, D. J., Lunt, D. J., Robinson, M. M., Rosenbloom, N., Salzmann, U., and Sohl, L. E.: Pliocene Model Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment 1), *Geosci. Model Dev.*, 3, 227–242, doi:10.5194/gmd-3-227-2010, 2010. 1708, 1710
- Haywood, A. M., Dowsett, H. J., Robinson, M. M., Stoll, D. K., Dolan, A. M., Lunt, D. J., Otto-Bliesner, B., and Chandler, M. A.: Pliocene Model Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment 2), *Geosci. Model Dev.*, 4, 571–577, doi:10.5194/gmd-4-571-2011, 2011a. 1708
- Haywood, A. M., Ridgwell, A., Lunt, D. J., Hill, D. J., Pound, M. J., Dowsett, H. J., Dolan, A. M., Francis, J. E., and Williams, M.: Are there pre-Quaternary geological analogues for a future greenhouse warming?, *Philos. T. R. Soc. A*, 369, 933–56, 2011b. 1705
- Haywood, A. M., Hill, D. J., Dolan, A. M., Otto-Bliesner, B. L., Bragg, F., Chan, W.-L., Chandler, M. A., Contoux, C., Dowsett, H. J., Jost, A., Kamae, Y., Lohmann, G., Lunt, D. J., Abe-Ouchi, A., Pickering, S. J., Ramstein, G., Rosenbloom, N. A., Salzmann, U., Sohl, L., Stepanek, C., Ueda, H., Yan, Q., and Zhang, Z.: Large-scale features of Pliocene climate: results from the Pliocene Model Intercomparison Project, *Clim. Past*, 9, 191–209, doi:10.5194/cp-9-191-2013, 2013a. 1708, 1712, 1713
- Haywood, A., Dolan, A., Pickering, S., Dowsett, H., McClymont, E., Prescott, C., Salzmann, U., Hill, D., Lunt, D., Pope, J., and Valdes, P.: On the identification of a Pliocene time slice for data-model comparison, *Phil. Trans. R. Soc. A.*, in press, 2013b. 1707, 1718
- Hill, D., Haywood, A., Hindmarsh, R., and Valdes, P.: Characterizing ice sheets during the Pliocene: evidence from data and models, in: *Deep-Time Perspectives On Climate Change: Marrying the Signal from Computer Models and Biological Proxies*, edited by: Williams, M., Haywood, A. M., Gregory, F. J., and Schmidt, D. N., The Geological Society, Micropaleontological Society Special Publications, London, 2007. 1706
- Jansen, E., Overpeck, J., Briffa, K., Duplessy, J., Joos, F., Masson-Delmotte, V., Olago, D., Otto-Bliesner, B., Peltier, W., Rahmstorf, S., Ramesh, R., Raynaud, D., Rind, D., Solomina, O., Villalba, R., and Zhang, D.: Palaeoclimate. Chapter 6, in: *Climate Change 2007: The*

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007. 1705

Jiang, D.: Modeling the middle Pliocene climate with a global atmospheric general circulation model, *J. Geophys. Res.*, 110, D14107, doi:10.1029/2004JD005639, 2005. 1708

Lisiecki, L. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records, *Paleoceanography*, 20, 1–17, 2005. 1707, 1715

Lunt, D. J., Valdes, P. J., Haywood, A., and Rutt, I. C.: Closure of the Panama Seaway during the Pliocene: implications for climate and Northern Hemisphere glaciation, *Clim. Dynam.*, 30, 1–18, 2007. 1708

Lunt, D. J., Haywood, A. M., Schmidt, G. A., Salzmann, U., Valdes, P. J., and Dowsett, H. J.: Earth system sensitivity inferred from Pliocene modelling and data, *Nat. Geosci.*, 3, 60–64, 2009. 1707

Lunt, D. J., Dunkley Jones, T., Heinemann, M., Huber, M., LeGrande, A., Winguth, A., Loptson, C., Marotzke, J., Roberts, C. D., Tindall, J., Valdes, P., and Winguth, C.: A model–data comparison for a multi-model ensemble of early Eocene atmosphere–ocean simulations: EoMIP, *Clim. Past*, 8, 1717–1736, doi:10.5194/cp-8-1717-2012, 2012a. 1708

Lunt, D. J., Haywood, A. M., Schmidt, G. A., Salzmann, U., Valdes, P. J., Dowsett, H. J., and Loptson, C. A.: On the causes of mid-Pliocene warmth and polar amplification, *Earth Planet. Sc. Lett.*, 321–322, 128–138, 2012b. 1705, 1711, 1712, 1713, 1719, 1720

Miller, K. G., Wright, J. D., Browning, J. V., Kulpecz, A., Kominz, M., Naish, T. R., Cramer, B. S., Rosenthal, Y., Peltier, W. R., and Sossian, S.: High tide of the warm Pliocene: implications of global sea level for Antarctic deglaciation, *Geology*, 40, 407–410, 2012. 1705, 1706

O’ishi, R., Abe-Ouchi, A., Prentice, I. C., and Sitch, S.: Vegetation dynamics and plant CO_2 responses as positive feedbacks in a greenhouse world, *Geophys. Res. Lett.*, 36, L11706, doi:10.1029/2009GL038217, 2009. 1718

Pagani, M., Liu, Z., LaRiviere, J., and Ravelo, A. C.: High Earth-system climate sensitivity determined from Pliocene carbon dioxide concentrations, *Nat. Geosci.*, 3, 27–30, 2009. 1705, 1707, 1710

Paillard, D.: Glacial cycles: toward a new paradigm, *Rev. Geophys.*, 39, 325–346, 2001. 1714

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Petoukhov, V., Ganopolski, A., Brovkin, V., Claussen, M., Eliseev, A., Kubatzki, C., and Rahmstorf, S.: CLIMBER-2: a climate system model of intermediate complexity. Part I: Model description and performance for present climate, *Clim. Dynam.*, 16, 1–17, 2000. 1709
- Pollard, D. and DeConto, R. M.: Modelling West Antarctic ice sheet growth and collapse through the past five million years, *Nature*, 458, 329–32, 2009. 1706
- Raymo, M., Grant, B., Horowitz, M., and Rau, G.: Mid-Pliocene warmth: stronger greenhouse and stronger conveyor, *Mar. Micropaleontol.*, 27, 313–326, 1996. 1705
- Raymo, M., Lisiecki, L., and Nisancioglu, K.: Plio-Pleistocene ice volume, antarctic climate, and the global $\delta^{18}\text{O}$ record, *Science*, 313, 492–495, 2006. 1715
- Robinson, M. M., Valdes, P. J., Haywood, A. M., Dowsett, H. J., Hill, D. J., and Jones, S. M.: Bathymetric controls on Pliocene North Atlantic and Arctic sea surface temperature and deepwater production, *Palaeogeogr. Palaeoclimatol.*, 309, 92–97, 2011. 1708
- Rowley, D. B. and Garzzone, C. N.: Stable isotope-based paleoaltimetry, *Annu. Rev. Earth Planet. Sc.*, 35, 463–508, 2007. 1706
- Salzmann, U., Haywood, A. M., Lunt, D. J., Valdes, P. J., and Hill, D. J.: A new global biome reconstruction and data-model comparison for the Middle Pliocene, *Global Ecol. Biogeogr.*, 17, 432–447, 2008. 1706, 1707
- Salzmann, U., Haywood, A. M., and Lunt, D. J.: The past is a guide to the future? Comparing Middle Pliocene vegetation with predicted biome distributions for the twenty-first century, *Philos. T. R. Soc. A*, 367, 189–204, 2009. 1705, 1706
- Schneider von Deimling, T., Held, H., Ganopolski, A., and Rahmstorf, S.: Climate sensitivity estimated from ensemble simulations of glacial climate, *Clim. Dynam.*, 27, 149–163, 2006. 1710
- Seki, O., Foster, G. L., Schmidt, D. N., Mackensen, A., Kawamura, K., and Pancost, R. D.: Alkenone and boron-based Pliocene $p\text{CO}_2$ records, *Earth Planet. Sc. Lett.*, 292, 201–211, 2010. 1705, 1710
- Sloan, L., Crowley, T., and Pollard, D.: Modeling of middle Pliocene climate with the NCAR GENESIS general circulation model, *Mar. Micropaleontol.*, 27, 51–61, 1996. 1708
- Thompson, R. and Fleming, R.: Middle Pliocene vegetation: reconstructions, paleoclimatic inferences, and boundary conditions for climate modeling, *Mar. Micropaleontol.*, 27, 27–49, 1996. 1705, 1707

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Table 1. Annual global temperature and precipitation factor separation. The sum of the single contributions and the total changes including the synergy term are shown as well.

	CO ₂	Vegetation	Orography	Ice sheets	Sum	Sum + Synergy
Temperature [°C]	1.54 (62 %)	0.44 (18 %)	0.33 (13 %)	0.19 (8 %)	2.50 (100 %)	2.52
Precipitation [mm day ⁻¹]	0.16 (62 %)	0.06 (23 %)	0.03 (12 %)	0.01 (4 %)	0.26 (100 %)	0.27

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. PRISM3D-CLIMBER-2 surface-type correspondence.

PRISM3D megabiome surface types	CLIMBER-2 surface types
tropical forest warm-temperate forest 50 % savanna and dry woodland temperate forest boreal forest	forest
grassland and dry shrubland tundra dry tundra 50 % savanna and dry woodland	grass
desert	desert

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.




Fig. 1. Annual mean surface air temperature factor separation. **(a)** Total MPWP temperature change, **(b)** synergy, and contributions from changes in **(c)** CO₂, **(d)** vegetation, **(e)** orography and **(f)** ice sheets.

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.




Fig. 2. Annual mean precipitation factor separation. **(a)** Total MPWP precipitation change, **(b)** synergy, and contributions from changes in **(c)** CO_2 , **(d)** vegetation, **(e)** orography and **(f)** ice sheets.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.




Fig. 3. (a) Maximum summer insolation at 65°N computed from Berger and Loutre (1992) (black solid line) and present-day value (black, dashed) and eccentricity computed from Berger and Loutre (1992) (grey solid line) and present-day value (grey, dashed). (b) Annual global temperature anomalies (MPWP-preindustrial) for the fully coupled transient experiment (solid red), transient experiment with fixed PRISM3D vegetation and interactive ice sheets (blue), transient experiment with fixed PRISM3D ice sheets and interactive vegetation with fixed (dashed green) and interactive Sahara/Sahel (solid green). The PlioMIP experiment equilibrium warming is shown for comparison (dashed black).




Fig. 4. Total NH ice sheet volume (blue) and forest area between 60–80° N (green) for the fully coupled experiment (solid lines). Dashed green lines represent forest area between 60–80° N computed from PRISM3D reconstructions.

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 5. Zonal mean temperature anomalies (MPWP-preindustrial). Annual **(a)** and summer **(c)** surface air temperatures over land and annual **(b)** and summer **(d)** sea surface temperatures for the fully coupled transient simulation, showing the time-average over the entire MPWP (solid red line) and the temperature variability range (shaded red) and the PlioMIP experiment equilibrium temperature anomalies (dashed black). The blue solid lines represent PRISM3D SST anomalies.

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.




Fig. 6. Forest fractions. Modelled maximum **(a)** and minimum **(b)** forest fraction (excluding glacier area) in each grid cell during the MPWP for the fully coupled transient simulation. **(c)** Modelled equilibrium forest fraction for the Pliocene with interactive vegetation. **(d)** Forest fraction derived from PRISM3D reconstructions (as described in Appendix A).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. Zonal forest fraction anomalies (MPWP-present-day) for PRISM3D (blue), for the fully coupled transient experiment at the time of maximum summer insolation at 65° N (red) and for the PlioMIP equilibrium simulation with interactive vegetation (yellow).

Effect of orbital forcing on mid-Pliocene climate

M. Willeit et al.

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[⏪](#) | [⏩](#)

[◀](#) | [▶](#)

[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

