



Received: 26 March 2013 – Accepted: 28 March 2013 – Published: 5 April 2013

Correspondence to: N. Combourieu-Nebout (nathalie.nebout@lsce.ipsl.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## CPD

9, 1969–2014, 2013

### Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

To understand the effects of future climate change on the ecology of the central Mediterranean we can look to the impacts of long-term, millennial to centennial-scale climatic variability on vegetation in the basin. Pollen data from the Adriatic Marine core MD 90-917 allows us to reconstruct vegetation and regional climate changes over the south central Mediterranean during the Holocene. Clay mineral ratios from the same core reflect the relative contributions of riverine (illite and smectite) and eolian (kaolinite) contributions to the site, and thus act as an additional proxy with which to test precipitation changes in the Holocene.

Vegetation reconstruction shows vegetation responses to the late-Glacial Preboreal oscillation, most likely driven by changes in seasonal precipitation. Pollen-inferred temperature declines during the early-mid Holocene, but increases during the mid-late Holocene, similar to southern-western Mediterranean climatic patterns during the Holocene. Several short climatic events appear in the record, indicating the sensitivity of vegetation in the region to millennial-scale variability.

Reconstructed summer precipitation shows a regional maximum between 8000 and 7000 cal yr BP similar to the general pattern across southern Europe. Two important shifts in vegetation occur at 7700 and between 7500 and 7000 yr. These vegetation shifts are linked to changes in seasonal precipitation and are correlated to increased river inputs respectively from the north (7700 event) and from the central Adriatic borderlands (7500–7000 event). These results reinforce the strengths of multi-proxy analysis and provide a deeper understanding of the role of precipitation and particularly the seasonality of precipitation in mediating vegetation change in the central Mediterranean during the Holocene.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



# 1 Introduction

The Mediterranean is highly sensitive to climate change, and Mediterranean ecosystems, already widely affected by anthropogenic pressures, are likely to be strongly impacted by future heat and drought stresses. Precipitation changes are of particular concern for the region. Increasing dryness predicted by GCMs for the Mediterranean will induce large water resource deficits and might threaten the region's habitability (Giorgi, 2006; IPCC, 2007; Giorgi and Lionello, 2008). While modern Mediterranean precipitation is controlled by the influences of both subtropical and mid-latitude climatic belts, resulting in dry summers and rainy winters, climate in the Mediterranean has changed during the Holocene, providing an opportunity to study ecosystem response under a range of climate conditions. In particular, past shifts in precipitation may help to envisage the future response of the Mediterranean region to predicted climate change. Reconstructions of past vegetation and climate in the different basins of the Mediterranean will thus produce information that allows us to draw a general framework to understand regional change and the drivers of environmental and climate change across the Mediterranean basin.

The South Adriatic provides an excellent opportunity to record high-resolution vegetation changes in Italy and the Balkans in response to Holocene climate variability. Its central location in the Mediterranean basin should be highly sensitive to the connections and conflicting influences between Northern and Southern atmospheric systems and their relative roles in mediating precipitation and runoff in the central Mediterranean during the Holocene.

Holocene vegetation change in the central Mediterranean has been investigated using continental and marine records (e.g. Watts et al., 1996; Combourieu-Nebout et al., 1998; Jahns and van den Bogaard, 1998; Magri, 1999; Magri and Sadori, 1999; Allen et al., 2002; Oldfield et al., 2003; Piva et al., 2008; Kotthoff et al., 2008, 2011; Di Donato et al., 2008; Di Rita and Magri, 2009; Di Rita et al., 2013). Although these records vary in temporal and taxonomic resolution, they show a general pattern of forest expansion

CPD

9, 1969–2014, 2013

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





For this paper, only 13 <sup>13</sup>C dates are used, providing a reliable age model for the last 13 000 cal yr BP (Table 1).

### 3 Present-day environmental settings

#### 3.1 Climate and atmospheric circulation pattern

5 Atmospheric circulation in the Adriatic is controlled by two dominant climatic systems. In the winter the Eurasian high extends south-westward, retreating northward in the summer when it is replaced by the north-eastward expansion of the Açores high. This pattern results in a typically Mediterranean climate (Fig. 1): winters are generally stormy and cool/cold while summers are warm/hot and dry. Spring climate in the  
10 region is generally long, characterized by fluctuating weather while autumn is very short with an abrupt transition toward winter. The length and intensity of summer dryness increases southwards across the Adriatic; total annual precipitation grades from a maximum in the east to a minimum in the west (Fig. 1). The mountainous regions in the Adriatic basin may show patterns that are dramatically different than their surroundings. In the Apennines or Dinarides temperatures decrease and precipitation increases  
15 according to altitudinal elevation.

A number of winds blow over the central Mediterranean (Fig. 1). The Bora is a cold wind that commonly blows from the North and North-east during winter. The Mistral (Italian Maestral) is a northwest wind that blows from spring to autumn, more often in  
20 summer with decreasing strength to the south. The Sirocco blows from the south from autumn to spring, bringing dust from North Africa. The Ostro is a southerly wind often identified with the Sirocco. The Jugo is an east to south-east dry wind that blows most often in the winter. The west to south-west wind called Libeccio blows year-round but rarely reaches the Adriatic basin.



altitudes, composed of deciduous trees such as *Carpinus*, *Ostrya*, *Fraxinus* and *Quercus*. At higher elevation where summer precipitation is higher, montane forests are mainly dominated by *Fagus*, *Abies alba* and *Picea*.

Natural vegetation in the region has been affected by human pressure in recent times; forest clearance and intensive farming often result in landscapes of bush-wood or open vegetation.

### 3.4 Pollen inputs

Palaeoenvironmental interpretation of the changes depicted in the pollen assemblages is based on the assumption that the pollen signal recorded in the marine core reflects the regional vegetation across an area of several hundred square kilometres (Heusser and Balsam, 1977; Dupont and Wyputta, 2003; Hooghiemstra et al., 1992, 2006). Pollen composition in deep basins of the Adriatic Sea must then be strongly affected by the dominant winds and hydrological currents that bring pollen grains from their source to the site in which they are ultimately carried.

Most pollen grains within the MD90-917 core, and especially arboreal pollen from *Quercus*, *Quercus ilex*, *Carpinus*, *Corylus*, *Fraxinus*, and *Abies* are likely to originate on either coast of the Adriatic basin given similarities between coasts in the modern vegetation community structure (Ozenda, 1975; Horvat et al., 1974). However, natural stands of *Olea* are nearly absent on the Dalmatian coast today and are only found in southern Greece on the Balkan peninsula (Ozenda, 1975), thus *Olea* pollen grains are more likely to arrive from the Italian coast. Nevertheless, olive cultivation by human societies during the last millennia (Mercuri et al., 2012) may contribute to broader *Olea* representation. *Picea* today grows only in the Dalmatian mountains, in the northern Apennines and the Alps, therefore its pollen is likely to be derived from these areas, brought to the Adriatic core site by Po river inflow. *Picea* may also be carried by eastern winds, however, *Picea* pollen grains are considered to be transported relatively short distances by wind in Europe (Hicks, 2001), thus *Picea* pollen is probably of northern origin, deposited to the site by discharge from the Po river.

1976

CPD

9, 1969–2014, 2013

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ultimately, pollen inputs into the Adriatic basin will draw a broad regional picture of Italian and Balkan vegetation from the wide range of habitats extending from the coast to the highlands.

### 3.5 Clay mineral origin

5 The Po River and the eastern Apennine rivers are the main sources of clay-sized particles in the western Adriatic. Although the Apennine rivers have relatively small drainage areas, studies have shown that they have very high sediment contributions relative to their drainage area (Frignani et al., 1992; Milliman and Syvitski, 1992; Alvisi et al., 1996; Bartolini et al., 1996; Sorgente, 1999; Tomadin, 2000). Detrital material in the Po River  
10 comes largely from the Alps, supplying illite, associated with chlorite, to the Adriatic Sea (Chamley, 1989; Alonso and Maldonado, 1990; Tomadin, 2000). Because illite is resistant to degradation and transport it generally represents the relative contribution of physical weathering to sedimentation. As a result, illite dominates the clay mineral sedi-  
15 ment fraction in the deeper parts of the Adriatic (Tomadin, 2000). By contrast, Apennine sediment sources are rich in smectite, which is mainly dispersed southeastward along the coast, with further downslope transport toward the deep basin through seasonal gradient and turbidity currents (Franco et al., 1982; Tomadin, 2000).

Croatian rivers to the east, containing illite and kaolinite with minor smectite (Durn et al., 1999) and Albanian rivers to the south, carrying smectite, supply sediment to the  
20 eastern Adriatic. The contribution of these river inputs is reduced because of very low riverine terrigenous loads and particle trapped along the Adriatic margin (e.g. Tomadin, 2000; Cattaneo et al., 2003).

The eolian contribution to deep-sea sediments is of major importance in the Mediterranean. Massive plumes of desert dust export clay-mineral particles toward the  
25 Mediterranean (Rea et al., 1985; Guerzoni and Chester, 1996) via regional meridian wind systems (Prospero, 1981; Loye-Pilot et al., 1986; Pye, 1987; Bergametti et al., 1989; Tomadin and Lenaz, 1989; Guerzoni and Chester, 1996; Guerzoni et al., 1999; Moulin et al., 1997; Rodriguez et al., 2001; Torres-Padron et al., 2002; Ginoux et al.,

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2004). Illite and kaolinite can also be transported through eolian processes. Kaolinite abundance is maximum in the south-eastern Sahara compared to western Sahara while illite displays the reverse pattern (Avila et al., 1997; Caquineau et al., 1998; Guerzoni et al., 1999). The southerly Sirocco wind most likely supplies eolian kaolinite-rich dust from North Africa to the Adriatic Sea during spring and summer (Goudie and Middleton, 2001).

## 4 Methods

### 4.1 Pollen record

Pollen extractions from marine sediments followed a previously described standard protocol (e.g. Faegri and Iversen, 1964; Combourieu-Nebout et al., 1998). Samples underwent successive treatments with HCl and HF acids and several sievings to obtain pollen concentrated residues. Pollen grains have been counted on slides with an Olympus light microscope with  $\times 500$  magnification. Between 150 and 300 grains were counted in each sample, based on pollen concentration. *Pinus* is over-represented in most marine samples. Because of over-representation of *Pinus* in the MD 90-917 pollen samples, all assemblages had at least 100 non-*Pinus* counts (Heusser and Balsam, 1977; Turon, 1984).

Pollen diagrams use sums excluding *Pinus* pollen. *Pinus* percentages have been calculated using the total sum of identified grains. Pteridophyta is reported using the total sum of all identified pollen grains and Pteridophyta. Pollen zones were visually determined and confirmed using cluster analysis (CONISS software, Grimm, 1987) (Fig. 2a and b).

Fossil pollen assemblages from the MD 90-917 core show similarities to a range of vegetation types from semi-desert to mountain deciduous and coniferous forest. The interpretation of these assemblages follows the modern plant-climate relationships in Eurasia and Northern Africa (Woodward, 1987; Peyron, 1998).

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A pollen record spanning from Last Glacial Maximum to Holocene was published by Combourieu-Nebout et al. (1998). This new analysis includes 51 additional counts, improving the temporal resolution of the Holocene section of this core. The new pollen data corresponds to time sampling intervals of from 15 to 500 yr with an average of 140 yr. Samples of the upper 80 cm were pollen poor and did not provide reliable data, possibly as a result of poor pollen preservation. The proposed pollen record, from 3 to 0.8 m depth, highlights vegetation changes at the end of deglaciation and during most of Holocene, from ~ 13000 to ~ 2500 cal yr BP.

### 4.2 Pollen-based reconstructions

We use the modern analogue technique “MAT” (Hutson, 1980; Overpeck et al., 1985; Guiot, 1990) to reconstruct past climate for this core. MAT has been successfully tested for the Holocene from Mediterranean terrestrial pollen records (e.g. Davis et al., 2003; Pross et al., 2009; Peyron et al., 2011, 2013; Magny et al., 2012; Joannin et al., 2012) and from marine pollen records (e.g. Kotthoff et al., 2008; Dormoy et al., 2009; Peyron et al., 2013; Desprat et al., 2013) with results that are generally supported by secondary proxies and regional records. Like most pollen-based climate reconstruction techniques, MAT aims to quantitatively reconstruct past climate from fossil assemblages based on the present-day environment.

MAT compares past assemblages to modern pollen assemblages without requiring real statistical calibration, using squared-chord distance to determine the degree of similarity between samples with known climate parameters (modern pollen samples) to a sample for which climate parameters are to be estimated (fossil pollen sample). More details on the MAT are given in Peyron et al. (2013).

This method requires a high-quality, taxonomically consistent modern pollen dataset. The pollen-inferred reconstruction here uses an updated modern pollen dataset compiled by Dormoy et al. (2009), containing more than 2000 samples from the Mediterranean region, to which we added 26 modern surface samples from around Lake Trifoglietti (Calabria, southern Italy) collected by Joannin et al. (2012a), and from around

Mount Altesina and Lake Preola (Sicily). Climate data comes from New et al. (2002). *Pinus* pollen was removed from the modern and fossil samples because of its over-representation in marine pollen assemblages. In this study, the quantitative climate reconstructions for the core MD 90-917 were performed for annual, winter and summer temperature (TANN, MTCO and MTWA), and annual, spring, summer, autumn and winter precipitation (PANN, Pspr, Psum, Paut and Pwin) to emphasize changes in seasonality.

### 4.3 Sediment clay fraction

Samples were prepared following standard protocols described in Bout-Roumazeilles et al. (2007). All samples were first decalcified with 0.2 N hydrochloric acid. Excess acid was removed by repeated centrifugations. The clay-sized fraction ( $< 2\mu\text{m}$ ) was isolated by settling, and oriented on glass slides (oriented mounts). Three XRD (X-ray diffraction) determinations were performed: (a) untreated sample; (b) glycolated sample (after saturation for 12 h in ethylene glycol); (c) sample heated at  $490^\circ\text{C}$  for two hours. The analyses were run on a Philips PW 1710 X-ray diffractometer, between  $2.49$  and  $32.5^\circ$ . Each clay mineral is characterized by its layer plus interlayer interval as revealed by XRD analysis. Smectite is characterized by a peak at  $14\text{ \AA}$  on the untreated sample test, which expands to  $17\text{ \AA}$  after saturation in ethylene glycol and retracts to  $10\text{ \AA}$  after heating. Illite presents a basal peak at  $10\text{ \AA}$  on the three tests (natural, glycolated, and heated). Kaolinite is characterized by peaks at  $7$  and  $3.57\text{ \AA}$  on the untreated sample and after saturation in ethylene glycol. Both peaks disappear or are strongly reduced after heating. Semi-quantitative estimation of clay mineral abundances, based on the pseudo-voigt deconvolution for the doublet kaolinite–chlorite ( $3.57\text{--}3.53\text{ \AA}$ ), was performed using the software MacDiff developed by Petschick (2000). Here we use the illite to kaolinite and smectite to kaolinite ratios in order to evaluate the balance between inputs from rivers, respectively Po River (illite) and eastern Apennine rivers (smectite), versus eolian inputs from the south (kaolinite). These ratios are calculated using the semi quantitative estimation of each clay mineral.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 5 Vegetation and climate for core MD 90-917

Palynological investigations using marine sediment cores have confirmed close links between pollen data from marine sediments and regional vegetation, and the ability of marine palynology to reconstruct climate changes at a regional scale (e.g. Heusser and Balsam, 1977; Hooghiemstra et al., 1992, 2006; Combourieu-Nebout et al., 2009). The reconstructed vegetation history for MD 90-917 is similar to other Italian and Balkan sequences for the Holocene. This support for the paleoecological accuracy of the MD 90-1917 core suggests its ability to reveal the regional consequences of the climatic events of the last 13 000 cal yr BP on central Mediterranean vegetation (Table 2, Fig. 2a and b).

The Younger Dryas event (GS-1 event) is characterised by the dominance of semi-desert elements such as *Artemisia*, *Chenopodiaceae* and *Ephedra* (Fig. 2b) which developed extensively around a lowered Adriatic Sea (also described in Combourieu-Nebout et al., 1998). This steppic vegetation is recorded in numerous regional pollen records from the same time (e.g. Watts et al., 1996; Magri and Sadori, 1999; Deneffe, 2000; Allen et al., 2002; Bordon et al. 2009; Kotthoff et al., 2008; Combourieu-Nebout et al., 2009; Fletcher et al., 2010; Desprat et al., 2013). They indicate that cold-arid climate conditions prevailing over the whole Mediterranean basin and especially in Adriatic basin (Fig. 3).

Between 12 000 and 11 700 cal yr BP, semi-desert elements are replaced by arboreal association composed by *Quercus*, *Carpinus*, *Corylus* and *Abies* showing the afforestation driven by climatic warming at the beginning of the Holocene (Fig. 2a). The spread of thermophilous taxa is observed from several pollen sequences across central Mediterranean corresponds to the Preboreal improvement at the Younger Dryas/Holocene transition (Bjork et al., 1996), however this begins at an older age in the marine core (e.g. Watts et al., 1996; Sadori and Narcisi, 2001; Allen et al., 2002; Sadori et al., 2011; Di Rita et al., 2013). Forest expansion is interrupted by a strong drop in arboreal taxa around 11 700 cal yr BP, marked by an increase in herbaceous taxa

CPD

9, 1969–2014, 2013

### Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(*Asteroidae* and *Cichoriodeae*) and semi-desert taxa, excluding *Artemisia* (Fig. 2a and b).

Climate reconstructions using pollen data show a decrease in temperature and precipitation at this time (Figs. 3 and 4). Although lake records from the central Mediterranean are rare, the same pattern can be seen, an increase in temperate forest followed by a slight return of herbs (lago Trifoglietti – Joannin et al., 2012a; lago di Vico – Di Rita et al., 2013). This climate oscillation recorded in the MD90-917 appears to lead the Greenland ice cores  $\delta^{18}\text{O}$  record (Fig. 4; Rasmussen et al., 2007) and probably corresponds to the  $\sim 11\,300$ yr BP Preboreal oscillation (PBO: Björk et al., 2001; Magny et al., 2012).

Climatic general trend of the pollen data in the early Holocene coincides with the alkenone-inferred SST increase at the same time, obtained from MD 90-917 (Sicre et al., 2013). Both pollen-based temperature and alkenone-inferred SST responses precede the foraminifer-based SST from this same core (Siani et al., 2013; Sicre et al., 2013). To explain such discrepancies, Sicre et al. (2013) suggest a local response in alkenone-inferred SST record. The temporal lead shown by MD 90-917 relative to the Trifoglietti and Lake Vico vegetation records should not be related to local behavior, since MD 90-917 should be representing the regional vegetation of the central Mediterranean. Given this, age discrepancies between continental records and MD 90-917 may be due to  $^{13}\text{C}$  date uncertainties. If this is the case, the lead in vegetation response (and alkenone-inferred SST) relative to foraminifer inferred SST from the same core may suggest that the vegetation response to the Preboreal oscillation in the region could be driven by precipitation changes rather than temperature variations.

After 11 000 cal yr BP, a mixed deciduous forest (mainly dominated by *Quercus* with regular occurrence of and accompanied by *Corylus*, *Carpinus*, *Fagus*, *Alnus*, *Betula*) expands (Fig. 2a and b), peaking around 7000 cal yr BP, corresponding to the “Holocene climate optimum”. This expansion occurs in Italy and in the Balkan peninsula at the same time (e.g. Watts et al., 1996; Jahns and van der Bogaard, 1998; Denèfle et al., 2000; Allen et al., 2002). Evidence for a climate optimum at 7000 cal yr BP agrees

CPD

9, 1969–2014, 2013

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Temperate forest dominated by *Quercus* fluctuates between 7000 and 5000 cal yr BP, decreasing after 4500 cal yr BP. *Quercus* becomes less abundant while *Carpinus*, *Corylus* and *Alnus* increase. This marks progressive opening of the forest, corresponding to decreases in summer precipitation, reflected in the pollen-based climate reconstruction. A cold, moist event is inferred from the rise in *Abies*, *Fagus*, and to a lesser extent *Picea* around 4000 cal yr BP (Fig. 3). Increased precipitation would have increased river flows, bringing the poorly dispersed *Abies* pollen to the core location. The upper samples of the core sequence, to 3000 cal yr BP, show an increase in herbaceous taxa (Asteraceae and Poaceae), continuing the trend of declining tree cover, probably influenced by increasing of human impacts in coastal Italy and Balkans. The increase in steppic elements such as *Artemisia* during this period may also indicate regional drying, particularly in the summer, and may represent the establishment of the summer conditions associated with the modern Mediterranean climate.

Mediterranean taxa, primarily *Q. ilex*, are continuously present in the pollen record beginning at 8000 cal yr BP. The Mediterranean forest begins to diversify after 6000 cal yr BP with the continuous presence of *Olea* and *Phillyrea* pollen, and, after 3500 cal yr BP with the presence of *Pistacia*. This coincides with the decreasing abundance of deciduous trees, especially deciduous *Quercus*, and the increase in herbs such as *Asteraceae* and steppic taxa (*Artemisia* excluded) at 3500 cal yr BP. This pattern has been observed elsewhere in the central Mediterranean (Sadori et al., 2011) and marks a shift in the precipitation regime, with decreasing summer precipitation and a progressive establishment of the modern Mediterranean climate (Fig. 3). *Olea*, the emblematic plant of the Mediterranean area, occurs weakly at the beginning of the Holocene and remains sporadic up to 6000 cal yr BP, generally associated only with warming climate. After 6000 cal yr BP *Olea* is continuously present but is most abundant in the upper part of the record, around 3000 cal yr BP, in agreement with the central Mediterranean vegetation history proposed by Sadori et al. (2011) using continental records. *Olea* is present in marine records across the Mediterranean at this time (e.g. Combourieu-Nebout et al., 2009; Desprat et al., 2013) and may also reflect



## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

cold events are associated with declines in *Quercus* pollen, and have been detected in other marine cores from Mediterranean (Fig. 7; e.g. Combourieu Nebout et al., 2009; Schmiel et al., 2010; Fletcher et al., 2010, 2013; Desprat et al., 2013) and may be linked to the millennial-scale cold events recorded in North Atlantic (Bond et al., 1997, 2001; Combourieu-Nebout et al., 2009; Schmiel et al., 2010; Fletcher et al., 2010, 2013; Fletcher and Zielhofer, 2011; Desprat et al., 2013; Magny et al., 2013). The rapid response of the central Mediterranean to millennial scale variability confirms the influence of mid-latitude atmospheric circulation in propagating changes, depending on insolation and ice sheet volume (Desprat et al., 2013). Our reconstruction suggests that the first two events (Younger Dryas and Preboreal Oscillation) were characterized by sharp seasonal contrast with very cold winters; this pattern differs strongly from the modern climate, which established itself progressively after 7700 cal yr BP with warmer winters more favorable to Mediterranean forest communities (Fig. 3).

## 6.2 Precipitation pattern

Pollen-inferred annual precipitation for MD 90-917 increases from 12 000 to 3000 cal yr BP with a maximum between 8000 and 7500 cal yr BP (Fig. 3). This trend is mirrored by springs and autumn precipitation, but, while summer and annual precipitation show similar trends of increasing precipitation from the early Holocene to 7500 cal yr BP, summer precipitation declines more strongly than annual precipitation to 3000 cal yr BP. The Holocene summer precipitation trend inferred from our pollen data is similar to speleothem records in central Italy (Zhorniak et al., 2011) and east-Mediterranean (Bar-Matthews et al., 1998, 2011), although speleothems record features that should be linked to changes in winter precipitation (Fig. 8).

The summer precipitation record fits well with the pattern described by Magny et al. (2012), which suggests that lakes located southward of 40° N experienced high lake-levels, reflecting precipitation maxima along the Early to Mid-Holocene in Italy, while northern Italian lakes were characterized by low lake-levels. The summer precipitation inferred from four Italian pollen records appears to confirm the opposing

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



precipitation regimes for northern and southern sites in the Mediterranean during the Holocene (Peyron et al., 2013). There is strong agreement between the high lake-levels reconstructed in Sicily, the wet summer conditions reconstructed for southern Italy by Peyron et al. (2013) and the summer precipitation inferred from the MD 90-917 pollen record. It is of note that the MD 90-917 summer precipitation curve shows similarities with the increasing trend at the northern lake Ledro during the early Holocene while the Mid-to-Late Holocene precipitation curve fits with the southern Mediterranean summer precipitation trend (Fig. 8, Peyron et al., 2013; Joannin et al., 2012b). Such a signal may illustrate the possible conflict of mid-latitudinal and subtropical monsoonal climatic systems in the Mediterranean area and emphasizes the fact that our record is located at the confluence between northern and southern influences. Climate changes recorded here are clearly driven by insolation changes and likely reflect the increasing influence of westerlies during the second half of Holocene, as demonstrated by the rise in winter precipitation in the upper part of our record.

The summer precipitation based on the MD 90-917 pollen record shows good fit to (i) sea surface salinity changes from the same core, interpreted as increasing runoff into the Adriatic deep sea basin (Siani et al., 2013) and (ii) changes in clay mineral composition (Fig. 8). Clays mineral records, presented as illite/kaolinite and smectite/kaolinite ratios (Fig. 8) reflect contributions from the eastern Apennine/Po rivers and dust input from Africa, respectively. These ratios clearly show a change in detrital input. Wind-blown dust (high kaolinite), supplied from the south by Sirocco winds, is more abundant during the first part of Holocene, while the rise of smectite after 9000 cal yr BP indicates the increasing contribution from Italian rivers to the basin.

A first rise in summer precipitation occurs at around 11 900 cal yr BP. Both I/K and S/K display a major peak at 12 500 cal yr BP, suggesting important discharges from both the Po and the eastern Apennine rivers. A single S/K peak at 12 000 cal yr BP suggests a major discharge from eastern Apennine rivers, synchronous with the precipitation maxima as reconstructed from pollen assemblages, related to the Preboreal event (Fig. 8).



## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



presently supplied to the Adriatic through the Po River and rather originates from Italy slopes through the small but active eastern Apennine rivers flows (Fig. 6). Based on the depth at which the core was obtained, its sedimentation should be mostly under the influence of the illite-rich Po river plume which flows southeastward in the open sea.

5 The smectite-rich southeastward Apennine plumes are generally restricted to shallower depths along the coast. Smectite is transferred toward the deep basin through turbidity and gravity currents when river discharge increases seasonally (Tomadin, 2000). As a consequence the 7000 cal yr BP event likely reflects a local pattern of change, reflecting increases in precipitation affecting the central Italian coast, nearest the core site.  
10 This suggests that decreases in salinity were caused by inflow from local rivers. These inputs persisted throughout the second half of Holocene but it is likely that they were then too weak to induce any significant decrease in salinity in Adriatic Sea surface.

## 7 Conclusions

15 The MD 90-917 marine pollen record provides the regional signal of central Mediterranean vegetation to long-term and millennial climate fluctuations during the late-Glacial and Holocene, and provides an integrated picture of temperature and precipitation changes during that period:

- the early Holocene is clearly expressed in vegetation changes and, despite a short chronological time-lag with the ice core chronology, the Preboreal climate oscillation is recorded in the core by an expansion of central Mediterranean forest taxa, followed by the return of open vegetation. Shifts in vegetation during the Preboreal oscillation are probably driven by precipitation changes, especially summer precipitation, and are correlated with inputs from local and regional rivers since vegetation changes occurred prior to SST increases recorded in the MD 90-917 core.  
20  
25

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Pollen-based temperature reconstructions follow the pattern reconstructed of other proxies in the southern Mediterranean, with an increasing west and north-west climatic influence. Several cold events are recorded and may be linked to millennial scale variability during the Holocene.
- Precipitation reconstructed from the pollen record shows a trend of increasing annual and seasonal (winter, spring and autumn) precipitation during the Holocene. Summer precipitation reflects a trend seen elsewhere in the southern Mediterranean, with a maximum around 7000 cal yr BP and then a decline until 3000 cal yr BP. Two significant peaks in precipitation occur at 7700 cal yr BP and between 7500 and 7000 cal yr BP that are associated with peak runoff from the Po River and from local central Adriatic rivers, respectively.

*Acknowledgements.* This study is a part of the LAMA ANR project (MSHE Ledoux, USR 3124, CNRS) projects financially supported by the French CNRS (National Centre for Scientific Research). The authors received funding from French CNRS and CEA for their researches. SJG is funded through the NSF-Macrosystems program. This research has been supported by the Institut Paul Emile Victor for the Prometée cruise of the R/V *Marion Dufresnes*. Palynological samples were processed by J. P. Cazet and I. Dormoy. This is LSCE contribution no. 4587.



The publication of this article is financed by CNRS-INSU.

## References

- Allen, J. R. M., Watts, W. A., McGee, E., and Huntley, B.: Holocene environmental variability – the record from Lago Grande di Monticchio, Italy, *Quatern. Int.*, 88, 69–80, 2002.
- Alonso, B. and Maldonado, A.: Late Quaternary sedimentation patterns of the Ebro turbidite systems (northwestern Mediterranean): two styles of deep-sea deposition, *Mar. Geol.*, 95, 353–377, 1990.
- Alvisi, F., Beks, J., Frignani, M., Langone, L., Moodley, L., Mowbray, S., Price, N. B., and Ravaioli, M.: Toward a sediment and heavy metal mass balance for the western Adriatic Sea, 2nd Workshop MYP-MAST, Iraklio, Greece, extended abstracts, 59–64, 1996.
- Avila, A., Queralt-Mitjans, I., and Alarcón, M.: Mineralogical composition of African dust delivered by red rains over northeastern Spain, *J. Geophys. Res.*, 102, 21977–21996, 1997.
- Bar-Matthews, M. and Ayalon, A.: Mid-Holocene climate variations revealed by high-resolution speleothem records from Soreq Cave, Israel and their correlations with cultural changes, *Holocene*, 21, 163–172, 2011.
- Bar-Matthews, M., Ayalon, A., and Kaufman, A.: Middle to late Holocene (6500 yr period) paleoclimate in the eastern Mediterranean region from stable isotopic composition of speleothems from Soreq Cave, Israel, in: *Environment and Society in Times of Climate Change*, edited by: Issar, A. and Brown, N., Kluwer Academic, Dordrecht, 203–214, 1998.
- Bartolini, C., Caputo, R., and Pieri, M.: Pliocene-Quaternary sedimentation in the northern Apennine Foredeep and related denudation, *Geol. Mag.*, 133, 255–273, 1996.
- Bergametti, G., Gomes, L., Remoudaki, E., Desbois, M., Martin, D., and Buat-Menard, P.: Present transport and deposition patterns of African dusts to the north-western Mediterranean, in: *Paleoclimatology and Paleometeorology: Modern and Past Patterns of Global Atmospheric Transport*, edited by: Leinen, M. and Sarnthein, M., Kluwer Academic, Dordrecht, 282, 227–251, 1989.
- Björck, S., Kromer, B., Johnsen, S., Bennike, O., Hammarlund, D., Lemdahl, G., Possnert, G., Rasmussen, T. L., Wohlfarth, B., Hammer, C. U., and Spurk, M.: Synchronized terrestrial-atmospheric deglacial records around the North Atlantic, *Science*, 274, 1155–1160, 1996.
- Björck, S., Muscheler, R., Kromer, B., Andresen, C. S., Heinemeier, J., Johnsen, S. J., Conley, D., Koç, N., Spurk, M., and Veski, S.: High-resolution analyses of an early Holocene climate event may imply decreased solar forcing as an important climate trigger, *Geology*, 29, 1107–1110, 2001.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., deMenocal, P., Priore, P., Cullen, H., Hajdas, I., and Bonani, G.: A pervasive millennial-scale cycle in north Atlantic Holocene and Glacial Climates, *Science*, 278, 1257–1266, 1997.
- Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M. N., Showers, W., Hoffman, S., Lotti, R., Hajdas, I., and Bonani, G.: Persistent solar influence on north Atlantic climate during the Holocene, *Science*, 294, 2130–2136, 2001.
- Bordon, A., Peyron, O., Lézine, A., Brewer, S., and Fouache, E.: Pollen-inferred Late-Glacial and Holocene climate in southern Balkans (Lake Maliq), *Quatern. Int.*, 200, 19–30, 2009.
- Bout-Roumazeilles, V., Nebout, N. C., Peyron, O., Cortijo, E., Landais, A., and Masson-Delmotte, V.: Connection between South Mediterranean climate and North African atmospheric circulation during the last 50 000 yr BP north Atlantic cold events, *Quaternary Sci. Rev.*, 26, 3197–3215, 2007.
- Caquineau, S., Gaudichet, A., Gomes, L., Magonthier, M.-C., and Chatenet, B.: Saharan dust: clay ratio as a relevant tracer to assess the origin of soil-derived aerosols, *Geophys. Res. Lett.*, 25, 983–986, 1998.
- Cattaneo, A., Correggiari, A., Langone, L., and Trincardi, F.: The late-Holocene Gargano subaqueous delta, Adriatic shelf: sediment pathways and supply fluctuations, *Mar. Geol.*, 193, 61–91, 2003.
- Chamley, H.: *Clay Sedimentology*, Springer-Verlag, Berlin, 623 pp., 1989.
- Combourieu-Nebout, N., Paterne, M., Turon, J.-L., and Siani, G.: A high-resolution record of the Last Deglaciation in the central Mediterranean sea: palaeovegetation and palaeohydrological evolution, *Quaternary Sci. Rev.*, 17, 303–332, 1998.
- Combourieu-Nebout, N., Peyron, O., Dormoy, I., Desprat, S., Beaudouin, C., Kotthoff, U., and Marret, F.: Rapid climatic variability in the west Mediterranean during the last 25 000 years from high resolution pollen data, *Clim. Past*, 5, 503–521, doi:10.5194/cp-5-503-2009, 2009.
- Davis, B. A. S., Brewer, S., Stevenson, A. C., Guiot, J., and data contributor: The temperature of Europe during the Holocene reconstructed from pollen data, *Quaternary Sci. Rev.*, 22, 1701–1716, 2003.
- Denefle, M., Lézine, A. M. M., Fouache, E., and Dufaure, J. J.: A 12 000 yr pollen record from Lake Maliq, Albania, *Quaternary Res.*, 54, 423–432, 2000.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Desprat, S., Combourieu-Nebout, N., Essallami, L., Sicre, M. A., Dormoy, I., Peyron, O., Siani, G., Bout Roumazeilles, V., and Turon, J. L.: Deglacial and Holocene vegetation and climatic changes in the southern Central Mediterranean from a direct land–sea correlation, *Clim. Past*, 9, 767–787, doi:10.5194/cp-9-767-2013, 2013.
- 5 Di Donato, V., Esposito, P., Russo-Ermolli, E., Scarano, A., and Cheddadi, R.: Coupled atmospheric and marine palaeoclimatic reconstruction for the last 35 ka in the Sele Plain-Gulf of Salerno area (southern Italy), *Quatern. Int.*, 190, 146–157, 2008.
- Di Rita, F. and Magri, D.: Holocene drought, deforestation and evergreen vegetation development in the Central Mediterranean: a 5500 yr record from Lago Alimini Piccolo, Apulia, Southeast Italy, *Holocene*, 19, 295–306, 2009.
- 10 Di Rita, F., Anzidei, A. P., and Magri, D.: A Lateglacial and early Holocene pollen record from Valle di Castiglione (Rome): vegetation dynamics and climate implications, *Quatern. Int.*, 288, 73–80, 2013.
- Dormoy, I., Peyron, O., Combourieu Nebout, N., Goring, S., Kotthoff, U., Magny, M., and Pross, J.: Terrestrial climate variability and seasonality changes in the Mediterranean region between 15 000 and 4000 years BP deduced from marine pollen records, *Clim. Past*, 5, 615–632, doi:10.5194/cp-5-615-2009, 2009.
- Dupont, L. M. and Wyputta, U.: Reconstructing pathways of aeolian pollen transport to the marine sediments along the coastline of SW Africa, *Quaternary Sci. Rev.*, 2, 157–174, doi:10.1016/S0277-3791(02)00032-X, 2003.
- 20 Durn, G., Ottner, F., and Slovenec, D.: Mineralogical and geochemical indicators of the polygenetic nature of terra rossa in Istria, Croatia, *Geoderma*, 91, 125–150, 1999.
- Fægri, K. and Iversen, J.: *Textbook of Pollen Analysis*, Munksgaard, Copenhagen, Denmark, 237 pp., 1964.
- 25 Finné, M., Holmgren, K., Sundqvist, H. S., Weiberg, E., and Lindblom, M.: Climate in the eastern Mediterranean, and adjacent regions, during the past 6000 yr – a review, *J. Archaeol. Sci.*, 38, 3153–3173, doi:10.1016/j.jas.2011.05.007, 2011.
- Fletcher, W. J. and Zielhofer, C.: Fragility of western Mediterranean landscapes during Holocene rapid climate changes, *Catena*, 103, 16–29, 2011.
- 30 Fletcher, W. J., Sanchez Goñi, M. F., Peyron, O., and Dormoy, I.: Abrupt climate changes of the last deglaciation detected in a Western Mediterranean forest record, *Clim. Past*, 6, 245–264, doi:10.5194/cp-6-245-2010, 2010.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Fletcher, W. J., Debret, M., and Sanchez Goñi, M. F.: Mid-Holocene emergence of a low-frequency millennial oscillation in western Mediterranean climate: implications for past dynamics of the North Atlantic atmospheric westerlies, *Holocene*, 23, 153–166, doi:10.1177/0959683612460783, 2013.
- 5 Franco, P., Jeftic, L., Malanotte Rizzoli, P., Michelato, A., and Orlic, M.: Descriptive model of the northern Adriatic, *Oceanol. Acta*, 5, 379–389, 1982.
- Frignani, M., Langone, L., Pacelli, M., and Ravaioli, M.: Input, distribution and accumulation of dolomite in sediments of the Middle Adriatic Sea, *Rap. Comm. Int. Mer Medi.*, 33, p. 324, 1992.
- 10 Frignani, M., Langone, L., Ravaioli, M., Sorgente, D., Alvisi, F., and Albertazzi, S.: Fine-sediment mass balance in the western Adriatic continental shelf over a century time scale, *Mar. Geol.*, 222, 113–133, 2005.
- Ginoux, P., Prospero, J. M., Torres, O., and Chin, M.: Long-term simulation of global dust distribution with the GOCART model: correlation with North Atlantic Oscillation, *Environ. Modell. Softw.*, 19, 113–128, 2004.
- 15 Giorgi, F.: Climate change hot-spots, *Geophys. Res. Lett.*, 33, L08707, doi:10.1029/2006GL025734, 2006.
- Giorgi, F. and Lionello, P.: Climate change projections for the Mediterranean region, *Global Planet. Change*, 63, 90–104, 2008.
- 20 Goudie, A. S. and Middleton, N. J.: Saharan dust storms: nature and consequences, *Earth Sci. Rev.*, 56, 179–204, 2001.
- Grauel, A. L. and Bernasconi, S. M.: Core-top calibration of  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  of *G. ruber* (white) and *U. mediterranea* along the southern Adriatic coast of Italy, *Mar. Micropaleontol.*, 77, 175–186, 2010.
- 25 Grimm, E. C.: CONISS: a FORTRAN 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares, *Comput. Geosci.*, 13, 13–35, 1987.
- Guerzoni, S. and Chester, R.: *The Impact of Desert Dust Across the Mediterranean*, Kluwer Academic, Dordrecht, 1996.
- 30 Guerzoni, S., Chester, R., Dulac, F., Herut, B., Loy -Pilot, M.-D., Measures, C., Migon, C., Molinaroli, E., Moulin, C., Rossini, P., Saydam, C., Soudine, A., and Ziveri, P.: The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea, *Prog. Oceanogr.*, 44, 147–190, 1999.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Guiot, J.: Methodology of the last climatic cycle reconstruction in France from pollen data, *Palaeogeogr. Palaeoclim.*, 80, 49–69, 1990.
- Heusser, L. E. and Balsam, W. L.: Pollen distribution in the NE Pacific ocean, *Quaternary Res.*, 7, 45–62, 1977.
- 5 Hicks, S.: The use of annual arboreal pollen deposition values for delimiting tree-lines in the landscape and exploring models of pollen dispersal, *Rev. Palaeobot. Palynol.*, 117, 1–29, doi:10.1016/S0034-6667(01)00074-4, 2001
- Hooghiemstra, H., Stalling, H., Agwu, C. O. C., and Dupont, L. M.: Vegetational and climatic changes at the northern fringe of the Sahara 250 000–5000 yr BP: evidence from 4 marine pollen records located between Portugal and the Canary Islands, *Rev. Palaeobot. Palynol.*, 10 74, 1–53, 1992.
- Hooghiemstra, H., Leizine, A.-M., Leroy, S. A. G., Dupont, L., and Marret, F.: Late Quaternary palynology in marine sediments: a synthesis of the understanding of pollen distribution patterns in the NW African setting, *Quatern. Int.*, 148, 29–44, 2006.
- 15 Horvat, I., Glavac, V., and Ellenberg, H.: *Vegetation Südosteuropas*, Geobotanica Selecta, 4, Fischer, Stuttgart, 1974.
- Hutson, W.: The Agulhas current during the Late Pleistocene: analysis of modern faunal analogs, *Science*, 207, 64–66, 1980.
- IPCC: *Climate Change 2007: Synthesis Report, Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Geneva, Switzerland, 104 pp., 2007.
- 20 Jahns, S. and van der Bogaard, C.: New palynological and tephrostratigraphical investigations of two salt lagoons on the island of Mljet, south Dalmatia, Croatia, *Veget. Hist. Archaeobot.*, 7, 219–234, 1998.
- 25 Jalut, G., Esteban Amat, A., Bonnet, L., Gauquelin, T., and Fontugne, M.: Holocene climatic changes in the western Mediterranean, from south-east France to south-east Spain, *Palaeogeogr. Palaeoclim.*, 160, 25–290, 2000.
- Jalut, G., Dedoubat, J. J., Fontugne, M., and Otto, T.: Holocene circum-Mediterranean vegetation changes: climate forcing and human impact, *Quatern. Int.*, 200, 4–18, doi:10.1016/j.quaint.2008.03.012, 2009.
- 30

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Joannin, S., Brugiapaglia, E., de Beaulieu, J.-L., Bernardo, L., Magny, M., Peyron, O., Goring, S., and Vanni re, B.: Pollen-based reconstruction of Holocene vegetation and climate in southern Italy: the case of Lago Trifoglietti, *Clim. Past*, 8, 1973–1996, doi:10.5194/cp-8-1973-2012, 2012a.

5 Joannin, S., Vanni re, B., Galop, D., Peyron, O., Haas, J.-N., Gilli, A., Chapron, E., Wirth, S. B., Anselmetti, F., Desmet, M., and Magny, M.: Climate and vegetation changes during the Lateglacial and Early-Mid Holocene at Lake Ledro (southern Alps, Italy), *Clim. Past Discuss.*, 8, 5583–5632, doi:10.5194/cpd-8-5583-2012, 2012b.

10 Kotthoff, U., Pross, J., M ller, U. C., Peyron, O., Schmiedl, G., and Schulz, H.: Climate dynamics in the borderlands of the Aegean Sea during formation of Sapropel S1 deduced from a marine pollen record, *Quaternary Sci. Rev.*, 27, 832–845, 2008.

Kotthoff, U., Koutsodendris, A., Pross, J., Schmiedl, G., Bornemann, A., Kaul, C., Marino, G., Peyron, O., and Schiebel, R.: Impact of late glacial cold events on the northern Aegean region reconstructed from marine and terrestrial proxy data, *J. Quaternary Sci.*, 26, 86–96, 2011.

15 Loye-Pilot, M.-D., Martin, J.-M., and Morelli, J.: Influence of Saharan dust on the rainfall acidity and atmospheric input to the Mediterranean, *Nature*, 321, 427–428, 1986.

20 Magny, M., Peyron, O., Sadori, L., Ortu, E., Zanchetta, G., Vanni re, B., and Tinner, W.: Contrasting patterns of precipitation seasonality during the Holocene in the south- and north-central Mediterranean, *J. Quaternary Sci.*, 27, 290–296, 2012.

Magny, M., Combourieu Nebout, N., Bout-Roumazeilles, V., Desprat, S., Joannin, S., Peyron, O., Sadori, L., Siani, G., Sicre, M. A., Vanni re, B., Wagner, B., de Beaulieu, J. L., Brugiapaglia, E., Chapron, E., Colombaroli, D., Debret, M., Desmet, M., Didier, J., Esselami, L., Galop, D., Gilli, A., Haas, J. N., Kallel, N., Millet, L., Revel, M., Simonneau, A., Stock, A., Tinner, W., Turon, J. L., and Zanchetta, G.: North–south palaeohydrological contrasts in the central Mediterranean during the Holocene: tentative synthesis and working hypotheses, *Clim. Past Discuss.*, in preparation, 2013.

25 Magri, D.: Late Quaternary vegetation history at Lagaccione near Lago di Bolsena (central Italy), *Rev. Palaeobot. Palynol.*, 106, 171–208, 1999.

30 Magri, D. and Sadori, L.: Late Pleistocene and Holocene pollen stratigraphy at Lago di Vico (central Italy), *Veget. Hist. Archaeobot.*, 8, 247–260, 1999.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Mercuri, A. M., Bandini Mazzanti, M., Florenzano, A., Montecchi, M. C., and Rattighieri, E.: *Olea*, *Juglans* and *Castanea*: the OJC group as pollen evidence of the development of human-induced environments in the Italian peninsula, Quatern. Int., doi:10.1016/j.quaint.2013.01.005, in press, 2012.

5 Milliman, J. D. and Syvitski, J. P. M.: Geomorphic/tectonic control of sediment discharge to the ocean: the importance of small mountainous rivers, J. Geol., 100, 525–544, 1992.

Moulin, C., Lambert, C. E., Dulac, F., and Dayan, U.: Control of atmospheric export of dust from North Africa by the north Atlantic Oscillation, Nature, 387, 691–694, 1997.

10 New, M., Hulme, M., and Jones, P. D.: Representing twentieth century space-time climate variability, Part 2: development of 1901–96 monthly grids of terrestrial surface climate, J. Climate, 13, 2217–2238, 2000.

15 Oldfield, F., Asioli, A., Accorsi, C. A., Mercuri, A. M., Juggins, S., Langone, L., Rolph, T., Trincardi, F., Wolff, G., Gibbs, Z., Vigliotti, L., Frignani, M., van der Post, K., and Branch, N.: A high resolution late Holocene paleo environmental record from the central Adriatic Sea, Quaternary Sci. Rev., 22, 319–342, 2003.

Orange, D., Garcia-Garcia, A., Lorenson, T., Nittrouer, C., Milligan, T., Miserocchi, S., Langone, L., Corregiani, A. I., and Trincardi, F.: Shallow gas and flood deposition on the Po Delta, Mar. Geol., 222, 159–177, 2005.

20 Overpeck, J. T., Webb, T., and Prentice, I. C.: Quantitative interpretation of fossil pollen spectra: dissimilarity coefficients and the method of modern analogs, Quaternary Res., 23, 87–108, 1985.

Ozenda, P.: Sur les étages de végétation dans les montagnes du bassin méditerranéen, Documents de Cartographie Ecologique, 16, 1–32, 1975.

25 Palinkas, C. M. and Nittrouer, C. A.: Cliniform sedimentation along the Apennine shelf, Adriatic Sea, Mar. Geol., 234, 245–260, 2007.

Petschick, R.: MacDiff v 4.2.5 (Free Geological Software), Geologisch-Paläontologisches Institut, available at: <http://servermac.geologie.uni-frankfurt.de/Rainer.html>, last access: April 2013, Universität Frankfurt/Main, Frankfurt, Germany, 2001.

30 Peyron, O., Guiot, J., Cheddadi, R., Tarasov, P., Reille, M., de Beaulieu, J. L., Bottema, S., and Andrieu, V.: Climatic reconstruction in Europe for 18 000 yr BP from pollen data, Quaternary Res., 49, 183–196, 1998.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Peyron, O., Goring, S., Dormoy, I., Kotthoff, U., Pross, J., de Beaulieu, J. L., Drescher-Schneider, R., Vanni re, B., and Magny, M.: Holocene seasonality changes in central Mediterranean reconstructed from Lake Accesa and Tenaghi Philippon pollen sequences, *Holocene*, 21, 131–146, 2011.
- 5 Peyron, O., Magny, M., Goring, S., Joannin, S., de Beaulieu, J.-L., Brugiapaglia, E., Sadori, L., Garfi, G., Kouli, K., Ioakim, C., and Combourieu-Nebout, N.: Contrasting patterns of climatic changes during the Holocene in the Central Mediterranean (Italy) reconstructed from pollen data, *Clim. Past Discuss.*, 8, 5817–5866, doi:10.5194/cpd-8-5817-2012, 2012.
- 10 Piva, A., Asioli, A., Trincardi, F., Schneider, R. R., and Vigliotti, L.: Late-Holocene climate variability in the Adriatic sea (central Mediterranean), *Holocene*, 18, 153–167, 2008.
- Pollunin, O.: *Flowers in Greece and the Balkans*, A Field Guide, Oxford Univ. Press, 1980.
- Poulain, P. M.: Adriatic Sea surface circulation as derived from drifter data between 1990 and 1999, *Mar. Syst.*, 29, 3–32, 2001.
- 15 Prospero, J. M.: Arid regions as sources of minerals aerosols in the marine atmosphere, *Geol. Soc. Am. S.*, 186, 71–86, 1981.
- Pross, J., Kotthoff, U., M ller, U. C., Peyron, O., Dormoy, I., Schmiedel, G., Kalaitzidis, S., and Smith, A. M.: Massive perturbation in terrestrial ecosystems of the eastern Mediterranean region associated with the 8.2 kyr BP climatic event, *Geology*, 37, 88–890, 2009.
- Pye, K.: *Aeolian Dust and Dust Deposits*, Academic Press, New York, London, 334 pp., 1987.
- 20 Rasmussen, S. O., Vinther, B. M., Clausen, H. B., and Andersen, K. K.: Early Holocene climate oscillations recorded in three Greenland ice cores, *Quaternary Sci. Rev.*, 26, 1907–1914, 2007.
- Ravazzi, C.: Late Quaternary history of spruce in southern Europe, *Rev. Palaeobot. Palynol.*, 120, 131–177, 2002.
- 25 Rea, D. K., Leinen, M., and Jacenek, T. R.: Geologic approach to the long-term history of atmospheric circulation, *Science*, 227, 721–725, 1985.
- Reimer, P. J., Baillie, M. G. L., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., Buck, C. E., Burr, G. S., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., McCormac, F. G., Manning, S. W., Reimer, R. W., Richards, D. A., Southon, J. R., Talamo, S.,
- 30 Turney, C. S. M., van der Plicht, J., and Weyhenmeyer, C. E.: IntCal09 and Marine09 radiocarbon age calibration curves – 50 000 yr cal BP, *Radiocarbon*, 51, 1111–1150, 2009.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Rodriguez, S., Querol, X., Alastuey, A., Kallos, G., and Kakaliagou, O.: Saharan dust contributions to PM<sub>10</sub> and TSP levels in southern and eastern Spain, *Atmos. Environ.*, 35, 2433–2447, 2001.

Sadori, L. and Narcisi, B.: The postglacial record of environmental history from Lago di Pergusa (Sicily), *Holocene*, 11, 655–671, 2001.

Sadori, L., Jahns, S., and Peyron, O.: Mid-Holocene vegetation history of the central Mediterranean, *Holocene*, 21, 117–129, 2011.

Sadori, L., Ortu, E., Peyron, O., Zanchetta, G., Vanni ere, B., Desmet, M., and Magny, M.: The last 7 millennia of vegetation and climate changes at Lago di Pergusa (central Sicily, Italy), *Clim. Past Discuss.*, in press, 2013.

Schmiidl, G., Kuhnt, T., Ehrmann, W., Emeis, K., Hamann, Y., Kotthoff, U., Dulski, P., and Pross, J.: Climatic forcing of eastern Mediterranean deep-water formation and benthic ecosystems during the past 22 000 yr, *Quaternary Sci. Rev.*, 29, 3006–3020, 2010.

Siani, G., Paterne, M., Michel, E., Sulpizio, R., Sbrana, A., Arnold, M., and Haddad, G.: Mediterranean Sea surface radiocarbon reservoir age changes since the Last Glacial Maximum, *Science*, 294, 1917–1920, doi:10.1126/science.1063649, 2001.

Siani, G., Sulpizio, R., Paterne, M., and Sbrana, A.: Tephrostratigraphy study for the last 18 000 <sup>13</sup>C years in a deep-sea sediment sequence for the South Adriatic, *Quaternary Sci. Rev.*, 23, 2485–2500, 2004.

Siani, G., Paterne, M., and Colin, C.: Late glacial to Holocene planktic foraminifera bioevents and climatic record in the South Adriatic Sea, *J. Quaternary Sci.*, 25, 808–821, 2010.

Siani, G., Magny, M., Paterne, M., Debret, M., and Fontugne, M.: Paleohydrology reconstruction and Holocene climate variability in the South Adriatic Sea, *Clim. Past*, 9, 499–515, doi:10.5194/cp-9-499-2013, 2013.

Sicre, M.-A., Siani, G., Genty, D., Kallel, N., and Essallami, L.: Seemingly divergent sea surface temperature proxy records in the central Mediterranean during the last deglacial, *Clim. Past Discuss.*, 9, 683–701, doi:10.5194/cpd-9-683-2013, 2013.

Sj ogren, P., Knaap, W. V. D., Huusko, A., and Leeuwen, J. F. V.: Pollen productivity, dispersal, and correction factors for major tree taxa in the Swiss Alps based on pollen-trap results, *Rev. Palaeobot. Palynol.*, 152, 200–210, 2008.

Sorgente, D.: Studio della sedimentazione attuale e recente nel medio Adriatico attraverso l'uso di traccianti radioattivi, Ph.D. Thesis, University of Bologna, Bologna, Italy, 178 pp., 1999.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Tomadin, L.: Sedimentary fluxed and different dispersion mechanism of the clay sediments in the Adriatic Basin, *Rend. Fis. Acc. Lincei*, 9, 161–174, 2000.
- Tomadin, L. and Lenaz, R.: Eolian dust over the Mediterranean and their contribution to the present sedimentation, in: *Paleoclimatology and Paleometeorology: Modern and Past Patterns of Global Atmospheric Transport*, edited by: Leinen, M. and Sarnthein, M., Kluwer Academic, Dordrecht, 267–281, 1989.
- Torres-Padrón, M. E., Gelado-Caballero, M. D., Collado-Sánchez, C., Siruela-Matos, V. F., Cardona-Castellano, P. J., and Hernández-Brito, J. J.: Variability of dust inputs to the CANIGO zone, *Deep-Sea Res. Pt. II*, 49, 3455–3464, 2002.
- Turon, J.-L.: Le paynoplanton dans l'environnement actuel de l'Atlantique nordoriental, évolution climatique et hydrologique depuis le dernier maximum glaciaire, Ph.D. thesis, Bordeaux 1 University, Bordeaux, France, 1984.
- van der Knaap, W. O., van Leeuwen, J. F. N., Finsinger, W., Gobet, E., Pini, R., Schweizer, A., Valsecchi, V., and Ammann, B.: Migration and population expansion of *Abies*, *Fagus*, *Picea*, and *Quercus* since 15 000 years in and across the Alps, based on pollen-percentage threshold values, *Quaternary Sci. Rev.*, 24, 645–680, 2005.
- Walker, M. J. C., Berkelhammer, M., Björck, S., Cwynar, L. C., Fisher, D. A., Long, A. J., Lowe, J. J., Newnham, R. M., Rasmussen, S. O., and Weiss, H.: Formal subdivision of the Holocene series/epoch: a discussion paper by a working group of INTIMATE (integration of ice-core, marine and terrestrial records) and the subcommission on Quaternary stratigraphy (International Commission on Stratigraphy), *J. Quaternary Sci.*, 27, 640–659, 2012.
- Walter, W., Harnickell, E., and Mueller-Dombois, D.: *Climate Diagrams*, Springer-Verlag, 1975.
- Watts, W. A., Allen, J. R. M., and Huntley, B.: Vegetation history and palaeoclimate of the Last Glacial period at Lago Grande di Monticchio, southern Italy, *Quaternary Sci. Rev.*, 15, 133–53, 1996.
- Wheatcroft, R. A., Nittrouer, C. A., Miserocchi, S., and Trincardi, F.: A comparison of recent floods and flood deposits of the Eel and Po rivers, in: *American Geophysical Union – Chapman Conference: Formation of Sedimentary Strata on Continental Margins*, Puerto Rico, 17–19 June 2001.
- Woodward, F. I.: *Climate and Plant Distribution*, Cambridge Univ. Press, Cambridge, UK, 1987.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wu, H., Guiot, J., Brewer, S., and Guo, Z.: Climatic changes in Eurasia and Africa at the last glacial maximum and mid-Holocene: reconstruction from pollen data using inverse vegetation modelling, *Clim. Dynam.*, 29, 211–229, doi:10.1007/s00382-007-0231-3, 2007.

Zanchettin, D., Traverso, P., and Tomasino, M.: Po River discharges: a preliminary analysis of a 200-yr tile series, *Climatic Change*, 89, 411–433, 2008.

Zhornyak, L. V., Zanchetta, G., Drysdale, R. N., Hellstrom, J. C., Isola, I., Regattieri, E., Piccini, L., Baneschi, I., and Couchoud, I.: Stratigraphic evidence for a “pluvial phase” between ca. 8200–7100 ka from Renella cave (central Italy), *Quaternary Sci. Rev.*, 30, 409–417, 2011.

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** Age model for core MD 90-917, AMS  $^{13}\text{C}$  ages and corresponding calibrated ages from INTCAL04 (Reimer et al., 2009) integrating a  $^{13}\text{C}$  marine reservoir correction (Siani et al., 2001).

Depth (cm)	Species	$^{13}\text{C}$ age BP (yr)	Error $\pm 1\sigma$	Cal age BP (yr)
0–2	<i>G. bulloides</i>	1010	60	555–609
140–142	<i>G. ruber</i>	4180	70	4082–4290
167–169	<i>G. ruber</i>	4750	70	4855–4986
175–177	<i>G. bulloides</i>	5000	70	5344–5466
190–192	<i>G. bulloides</i>	5680	70	5990–6128
230–232	<i>G. bulloides</i>	6920	90	7413–7511
240–242	<i>G. bulloides</i>	7930	80	8171–8340
250–252	<i>G. ruber</i>	8170	70	8390–8482
275–277	<i>G. bulloides</i>	10 390	90	11 304–11 624
295–297	<i>G. bulloides</i>	10 800	90	12 116–12 399
305–307	<i>G. bulloides</i>	10 830	90	12 225–12 406
315–317	<i>G. bulloides</i>	11 140	90	12 721–12 853
335–337	<i>G. bulloides</i>	11 520	100	12 939–13 114

**Table 2.** Short description of the pollen zonation of the MD 90-917 pollen diagram.

Pollen zone	Interval cm	Age kyr	Pollen zone signature	Main biome	Climate
MD90 917-XII	80–120	2641–3671	<i>Pinus</i> (50%); ↓ <i>Abies</i> (< 4%) and <i>Picea</i> (< 6%) <i>Quercus</i> (45%), <i>Carpinus</i> (6%) <i>Corylus</i> in low percentages, ↑ <i>Fagus</i> (4%) Mediterranean taxa, <i>Q. ilex</i> (↑ 12%), <i>Olea</i> (3–4%) and <i>Pistacia</i> (↑ 4%) ↑ <i>Cichoriodeae</i> ; ↑ <i>Artemisia</i> (15%)	Warm temperate Mediterranean oak forest	Warm and humid stability in T° ↑ seasonal Pmm
MD90 917-XI	128–140	3877–4186	<i>Pinus</i> (50%); ↑ <i>Abies</i> (< 15%), <i>Picea</i> (3%) <i>Quercus</i> (40%), <i>Carpinus</i> (6%), <i>Corylus</i> and <i>Fagus</i> in low percentages, Mediterranean taxa in low percentages, <i>Q. ilex</i> (4%)	Cool mixed forest	Cool and relatively humid
MD90 917-X	142–179	4244–5579	<i>Pinus</i> low (20–40%); Low percentages in <i>Abies</i> and <i>Picea</i> (< 6%) ↓ <i>Quercus</i> (45%), ↑ <i>Carpinus</i> (↑ 10%) and <i>Corylus</i> (↑ 10%) Mediterranean taxa in stable percentages, Herbs in stable percentages; slight increase in steppic taxa (5–10%)	Warm temperate oak forest	Warm and humid stability in T° Pmm slight decrease
MD90 917-IX	180–220	5623–6960	<i>Pinus</i> low (20–40%); low percentages in <i>Abies</i> and <i>Picea</i> (< 6%), Temperate trees stable, ↑ <i>Quercus</i> (↑ 45–65%), ↓ <i>Carpinus</i> (~ 5%) and <i>Corylus</i> (~ 5%), ↑ Mediterranean taxa, <i>Q. ilex</i> (6–7%), <i>Olea</i> , <i>Phillyrea</i> and <i>Pistacia</i> ↑ in herbs; <i>Cichoriodeae</i> (5–10%); steppic taxa low	Warm temperate oak forest	Warm and humid – stability in T° and Pmm slightly reincrease
MD90 917-VIIIb	221–232	7010–7620	<i>Pinus</i> (20–50%), ↓ <i>Abies</i> (< 10%) and <i>Picea</i> (< 5%) decrease, ↑ Temperate trees, <i>Quercus</i> stable (< 45%), <i>Corylus</i> (5–20%), <i>Carpinus</i> (4–10%), <i>Fagus</i> (↑ 5%), <i>Ulmus</i> , (↑ 5%), <i>Alnus</i> (3–8%), Mediterranean taxa stable ( <i>Q. ilex</i> mainly, <i>Olea</i> and <i>Pistacia</i> presence), ↓ Herbs ( <i>Asteraceae</i> < 5%, <i>Poaceae</i> < 5%), steppic taxa low (10%)	Warm temperate oak forest	Warm and humid – increase in T° and Pmm slight decrease

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Table 2. Continued.

Pollen zone	Interval cm	Age kyr	Pollen zone signature	Main biome	Climate
MD90 917-VIIIa	233–234	7700–7779	<i>Pinus</i> (50%), ↑ <i>Abies</i> (↑ 25%) and <i>Picea</i> (↑ 11%), ↓ Temperate trees, <i>Quercus</i> (< 45%), <i>Corylus</i> , <i>Carpinus</i> absent, <i>Fagus</i> (↑ 10%), Mediterranean taxa low, Herbs low, steppic taxa low (slight ↑ <i>Chenopodiaceae</i> ; 2–5%)	Cool mixed forest	Cool and humid
MD90 917-VII	235–238	7859–8058	<i>Pinus</i> (30–60%); <i>Abies</i> and <i>Picea</i> low, Temperate trees: <i>Quercus</i> (30–60%), <i>Corylus</i> (< 5%), <i>Carpinus</i> (4%), <i>Fagus</i> (↑ 10%), Mediterranean taxa low ( <i>Q. Ilex</i> mainly, <i>Olea</i> presence), ↓ Herbs ( <i>Asteraceae</i> < 5%, <i>Poaceae</i> < 5%), steppic taxa nearly absent	Warm temperate oak forest	Warm and humid – Max T° and Pmm slight re-increase
MD90 917-VI	239–250	8157–8456	<i>Pinus</i> (30–60%); <i>Abies</i> and <i>Picea</i> low, Temperate trees: <i>Quercus</i> (40%), <i>Corylus</i> (5–10%), <i>Carpinus</i> (4–10%), <i>Fagus</i> (↑ 8%), Mediterranean taxa ↑ ( <i>Q. Ilex</i> mainly, <i>Olea</i> and <i>Pistacia</i> presence), ↓ Herbs ( <i>Asteraceae</i> < 5%, <i>Poaceae</i> < 5%), steppic taxa low	Warm temperate oak forest	Warm and humid – Max T° and Pmm slight decrease
MD90 917-V	252–258	8678–9445	<i>Pinus</i> (around 50%), <i>Abies</i> and <i>Picea</i> low, ↑ temperate trees, maximum <i>Quercus</i> (45–70%), <i>Corylus</i> (5–10%), <i>Carpinus</i> (2–10%), Mediterranean taxa low ( <i>Q. Ilex</i> only), ↓ Herbs ( <i>Asteraceae</i> < 5%, <i>Poaceae</i> < 5%) and steppic taxa low	Warm temperate oak forest	Warm and humid – Max T° and Pmm
MD90 917-IV	259–269	9546–10 737	Large decrease in <i>Pinus</i> (< 50%), <i>Abies</i> and <i>Picea</i> nearly absent, Increase in temperate trees, <i>Quercus</i> (40%), <i>Corylus</i> (5–10%), <i>Carpinus</i> (2–4%), Mediterranean taxa low ( <i>Q. Ilex</i> and first occurrence in <i>Olea</i> ), ↓ Herbs ( <i>Asteraceae</i> < 12%, <i>Poaceae</i> < 7%), steppic taxa very low	Warm temperate oak forest	Progressive ↑ T° and Pmm

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[I◀](#)

[▶I](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)


[Interactive Discussion](#)

**Table 2.** Continued.

Pollen zone	Interval cm	Age kyr	Pollen zone signature	Main biome	Climate
MD90 917-III	270– 282	10 858– 11 742	<i>Pinus</i> abundant (60–80%), <i>Abies</i> and <i>Picea</i> percentages low, <i>Quercus</i> (20–40%), ↓ <i>Corylus</i> and <i>Carpinus</i> , slight ↑ <i>Alnus</i> , <i>Betula</i> , <i>Tilia</i> , ↑ Herbs ( <i>Asteraceae</i> 20%), ↑ <i>Chenopodiaceae</i> > 10%)	Grassland mixed to open oak forest	Cool and moderately dry
MD90 917-II	284– 288	11 821– 11 980	<i>Pinus</i> abundant (65–75%), <i>Abies</i> and <i>Picea</i> percentages low, ↑ <i>Quercus</i> (15–40%), <i>Corylus</i> (↑ 10%), <i>Carpinus</i> (3%), ↓ Herbs ( <i>Asteraceae</i> < 20%, <i>Poaceae</i> around 10%), ↓ steppic taxa low	Open Warm mixed forest	Slight increase in T° and Pmm
MD90 917-I	290– 350	12 060– 13 257	<i>Pinus</i> abundant (65–90%), <i>Abies</i> and <i>Picea</i> percentages low, Low percentages in temperate trees, <i>Quercus</i> < 20%, <i>Alnus</i> , <i>Betula</i> , <i>Carpinus</i> present, Herbs abundant, <i>Asteraceae</i> 10–20%, <i>Poaceae</i> around 10%, Steppic taxa abundant ( <i>Artemisia</i> 15%, <i>Chenopodiaceae</i> 10%, <i>Ephedra</i> up to 7%)	Steppe	Cold and dry

**Holocene vegetation and climate changes in Mediterranean**

N. Combourieu-Nebout et al.



**Fig. 1.** Location of core MD 90-917. On the map is indicated sea surface circulation in Adriatic Sea and winds blowing over the Adriatic basin. A selection of ombrothermic diagrams are presented on the right for the surrounding borderlands to show the climate of the studied area (Walter et al., 1975; New et al., 2000; New LoCClim software).

[Title Page](#)

[Abstract](#)   [Introduction](#)

[Conclusions](#)   [References](#)

[Tables](#)   [Figures](#)

[⏪](#)   [⏩](#)

[⏴](#)   [⏵](#)

[Back](#)   [Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.



**Fig. 2a.** Pollen diagram of the Core MD 90-917: arboreal taxa. Pollen percentages are calculated on a sum excluding *Pinus*; *Pinus* percentages are calculated on the total pollen sum.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



**Holocene vegetation and climate changes in Mediterranean**

N. Combourieu-Nebout et al.



**Fig. 3.** Selected pollen curves of core MD 90-917 and pollen based reconstruction of precipitation (seasonal – Pspr, Psum, Paut, Pwin – and annual – Pann) and temperature (winter Twin, annual Tann and summer Tsum).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Holocene vegetation and climate changes in Mediterranean


N. Combourieu-Nebout  
et al.



**Fig. 4.** Comparison between two main pollen groups; temperate trees (green) and semi-desert (red) and NGRIP oxygen isotope record (NGRIP members, 2004; Lemieux Dudon, 2011).

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout et al.



**Fig. 5.** Comparison of the total tree percentage curve (green) (this paper) and the  $\delta^{18}\text{O}$  (blue) and  $\delta^{13}\text{C}$  (red) record from the MD 90-917 core (Siani et al., 2013).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Holocene vegetation and climate changes in Mediterranean


N. Combourieu-Nebout et al.



**Fig. 6.** Comparison between MD 90-917 pollen-based winter and summer temperature reconstructions and the combined climate records of south-eastern and south-western Europe (from Davis and Brewer, 2003).

## Holocene vegetation and climate changes in Mediterranean

N. Combourieu-Nebout  
et al.



**Fig. 7.** Comparison of millennial cold events recorded in MD 90-917 with the cold events recorded from vegetation changes in other Mediterranean marine cores (Desprat et al., 2013; Schieldt et al., 2010; Fletcher et al., 2010, 2013; Combourieu Nebout et al., 2009). On the left are pointed the North Atlantic Bond events (Bond et al., 2001) and, on the right, the periods of high lake levels recorded in the Southern Alps (Magny et al., 2012).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 8.** Comparison of the summer precipitation changes reconstructed from MD 90-917 pollen data and high lake levels in the alps (Magny et al., 2012), sea surface salinity reconstructed from MD 90-917  $\delta^{18}\text{O}$  record (Siani et al., 2013), Illite/Kaolinite (I/K) and Smectite/Kaolinite (S/K) ratios from MD 90-917 and  $\delta^{18}\text{O}$  speleothem records from the Renella cave (Italy) (Zhornyak, 2011) and Soreq cave (Israel) (Bar-Matthews et al., 2011).

**Holocene vegetation and climate changes in Mediterranean**

N. Combourieu-Nebout et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

