

**Vegetation  
responses to  
interglacial warming**

A. V. Lozhkin and  
P. M. Anderson

# Vegetation responses to interglacial warming in the Arctic, examples from Lake El'gygytgyn, northeast Siberia

A. V. Lozhkin<sup>1</sup> and P. M. Anderson<sup>2</sup>

<sup>1</sup>Northeast Interdisciplinary Scientific Research Institute, Far East Branch, Russian Academy of Sciences, 16 Portovaya Street, Magadan, 685000, Russia

<sup>2</sup>Earth & Space Sciences and Quaternary Research Center, University of Washington, Seattle, 98195-1310, USA

Received: 28 August 2012 – Accepted: 3 September 2012 – Published: 15 January 2013

Correspondence to: P. M. Anderson (pata@u.washington.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

Palynological data from Lake El'gygytyn reveal responses of plant communities to a range of climatic conditions that can help assess the possible impact of global warming on arctoboreal ecosystems. Vegetation associated with climatic optima suggests two types of interglacial responses: one is dominated by deciduous taxa (the post-glacial thermal maximum (PGTM) and marine isotope stage (MIS5)) and the second by evergreen conifers (MIS11, MIS31). The MIS11 forests show a similarity to *Picea-Larix-Betula-Alnus* forests of Siberia. While dark coniferous forest also characterizes MIS31, the pollen taxa show an affinity to the modern boreal forest of the lower Amur valley in the Russian Far East. Despite vegetation differences during the thermal maximum, all four glacial-interglacial transitions are alike, being dominated by deciduous woody taxa. Initially *Betula* shrub tundra established and was replaced by tundra with tree-sized shrubs (PGTM), *Betula* woodland (MIS5), or *Betula-Larix* (MIS11, MIS31) forest. The consistent occurrence of deciduous forest and/or high shrub tundra in all interglaciations as they approach or achieve maximum warmth underscores the significance of this biome for modeling efforts. The El'gygytyn data also suggest the possible elimination or massive reduction of arctic plant communities under extreme warm-earth scenarios.

## 1 Introduction

Marine and ice cores contain vital data that document fluctuations in past climate and provide insight into the possible causes of the observed paleoclimatic variability. However, such records cannot provide information on the responses of biotic systems to those changes. Most paleo-biotic records do not have the longevity, continuity, or temporal resolution of ocean and ice cores, which limits the comparison of long-term trends in biological and climatic data. Lake El'gygytyn (Lake E), whose basin was created following a meteor impact  $\sim 3.58$  Ma ago, presents a rare exception having yielded a

CPD

9, 245–267, 2013

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



continuous sedimentary record dating back  $\sim 2.8$  Ma (Melles et al., 2012). The location of the lake in arctic Chukotka (Fig. 1) provides a unique look into the response of terrestrial ecosystems at northern high latitudes to a broad range of warm and cold conditions.

Several biological proxies have been examined in cores from Lake E (Brigham-Grette et al., 2007; Melles et al., 2012); we present here interpretations of past vegetation based on palynological data. Previous research has shown that cool, dry intervals are clearly marked in the Lake E record by a dominance of herb pollen (particularly Poaceae) and usually higher values of *Selaginella rupestris* spores (Lozhkin et al., 2007; Matrosova, 2009). These spectra contrast to interglacial assemblages where arboreal pollen is at a maximum, herbaceous taxa are low, and spores are often dominated by *Sphagnum*. Continued research on a newly raised core shows that interglacial assemblages display vegetation and climatic variability both within and between isotope stages. For example, marine isotope stage (MIS) 11.3 and MIS31 have been described as “super interglaciations” with maximum summer temperatures  $\sim 4^\circ$  to  $5^\circ$  C warmer than either MIS5.5 or the postglacial thermal maximum (PGTM; Melles et al., 2012).

These warmer-than-present intervals are of particular interest for understanding possible responses and feedbacks related to increasing global temperatures (e.g., Harrison et al., 1995; Kaplan et al., 2003). Both conceptual and quantitative models have considered two types of vegetation responses at northern high latitudes; one suggests the expansion of evergreen conifer forest northward across North America and much of Eurasia (e.g., Claussen, 1996; Kaplan et al., 2003), whereas the second postulates a dominance of broadleaf deciduous forest in current areas of tundra (e.g., Bonan et al., 1990; Chapin and Starfield, 1997). Paleobotanical data are one means by which these alternative scenarios may be evaluated. Although analysis continues on Lake E, sufficient palynological data are available to compare the vegetation responses during the “super interglaciations” and the more moderate warming represented in MIS5.5 and the PGTM.

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 2 Methods

One to two cm<sup>3</sup> sediment samples were processed for palynological analysis following standard procedures used in preparing organic-poor samples (PALE, 1994). Samples for MIS1-MIS2 and MIS4-MIS6 are from core PG1352, whereas core D1 provided samples from MIS11-MIS12, and MIS 31. Pollen sums for interglacial spectra generally exceed 500 identified pollen grains. Percentages of individual taxa are based on a sum of all arboreal, nonarboreal, and unidentified pollen grains. Subsum percentages on the left of the diagrams (Figs. 2–4) are calculated from a total of pollen and spores. Pollen zonation was done subjectively using changes in percentages of major taxa. Although many palynologists have differentiated tree and shrub *Betula* based on morphology and/or size, blind tests in our laboratories of modern reference materials suggest that these criteria are inconsistent and unreliable; thus we do not separate the *Betula* pollen. *Picea obovata* pollen is classified within *Picea* sect. Eupicea, *Picea ajanensis* within *Picea* sect. Omorica, *Pinus sylvestris* within *Pinus* subg. Diploxylon, and *Pinus pumila* and *Pinus sibirica* within *Pinus* subg. Haploxylon. Following traditional palynological usage, *Duschekia fruticosa* is referred to as shrub *Alnus*. All other plant taxonomy follows Cherepanov (1995).

The age model for the El'gygytyn record is based on paleomagnetic stratigraphy and tuning of additional proxy data to the marine isotope stratigraphy and to variations in regional insolation (see Melles et al., 2012). Because pollen zones do not always correspond to isotope substages, the zones are identified by isotope stage (number) and as a pollen assemblage (letter). Following palynological format the lowest zone is indicated by "a" the next youngest by "b", etc., in contrast to the order of isotope substages.

CPD

9, 245–267, 2013

### Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

### 3 Modern setting

Lake E (67.5° N, 172° E; ~ 500 m a.s.l.) has a diameter of ~ 12 km and a maximum depth of 170 m (Melles et al., 2012). The local vegetation is a mix of discontinuous and continuous herb-lichen tundra (Kozhevnikov, 1993). Shrubs (*Salix krylovii*, *S. alaxensis*, *Betula exilis*) are not abundant and limited to protected sites in valleys and saddles within the crater walls and along the outlet valley. The vegetation in the Lake E catchment is more barren than in the surrounding Chukchi Uplands, because of unusual soil qualities associated with the meteor impact crater. The Uplands are dominated by shrub tundra that includes *Betula exilis*, *Duschekia fruticosa* (shrub *Alnus*), *Pinus pumila*, *Salix*, and Ericales. *Larix gmelinii* treeline is located ~ 150 to the southwest of the lake, but the main body of the forest is found at ~ 300 km distance. Modern pollen studies indicate that the pollen rain in Lake E is more indicative of regional than local vegetation with up to 60 % of the pollen (*Alnus*, *Betula*, *Pinus pumila*) being blown in from the Chukchi Upland (Lozhkin et al., 2001). The over-representation of Upland vegetation, which reflects regional conditions, strengthens the use of the Lake E pollen record for paleoclimatic interpretations. Estimated mean climatic values are: annual temperature -13.5 to -14 °C; July temperature 8.8 to 9 °C; January temperature -31 to -32 °C; annual precipitation 255 mm (Melles et al., 2012).

### 4 Interglacial vegetation of Lake E and the Chukchi Uplands

While interglaciations share similarities in boundary conditions (e.g., high summer insolation, warm sea-surface temperatures; minimal terrestrial ice-cover), variations in the climatic forcings and strengths of feedback mechanisms add a uniqueness to each warm interval (see Melles et al., 2012). The following sections describe the variability of vegetation responses to climates of the PGTM, MIS5, MIS11, and MIS31.

CPD

9, 245–267, 2013

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 4.1 PGTM

High percentages of *Betula* and *Alnus* characterize the late glacial and Early Holocene pollen assemblages at Lake E (Fig. 2; pollen zone 1a), an assemblage that is common across Northeast Siberia (Lozhkin et al., 1993). Statistical comparisons to modern pollen and related climatic parameters reveal mean July temperatures were perhaps 2–4 °C warmer than present between ~8600 and 10 700 <sup>14</sup>CyrBP (Lozhkin et al., 2007). Analog analysis also indicates warm January temperatures and increased July precipitation as compared to modern. These conclusions are based on “possible pollen analogs” to southern Beringia shrub tundra. Macrofossil evidence from the Yana-Indigirka-Kolyma Lowland (Fig. 1) and northern Chukotka documents the occurrence of tree or tree-sized shrubs of *Salix* and *Alnus* beyond present-day treeline (Lozhkin, 1993). Using both pollen and plant macrofossils to enhance understanding of plant functional types, Edwards et al. (2005) concluded that an unusual biome of deciduous forest and forest-tundra characterized Beringia during the PGTM. In the Chukchi Uplands, vegetation was probably a mosaic of forest and tundra communities including gallery forests of tree *Betula* and perhaps *Salix*, and possibly stands of deciduous trees in protected sites within the interfluves. Other areas likely supported high shrub tundra.

## 4.2 MIS5

Because MIS5 vegetation has been described elsewhere (Lozhkin et al., 2007), we focus here on MIS5.5. Like the PGTM, the MIS5 climatic optimum (Fig. 2; pollen zone 5a) is characterized by high percentages of *Betula* and *Alnus* pollen. The dominance of these woody taxa suggests that the regional vegetation during MIS5.5 was also a mix of deciduous forest and high shrub tundra. The reduced herb percentages as compared to the PGTM indicate a more extensive and/or denser population of *Betula*, *Alnus*, and possibly *Salix* in the regional vegetation. The spectra may also indicate a greater establishment of trees/high shrubs within the Lake E catchment as compared to the PGTM.

### Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Plant macrofossils of last interglacial age have been found throughout Northeast Siberia (Lozhkin and Anderson, 1995). Although chronological control is insufficient to assign sites specifically to MIS5.5, current understanding of the valley stratigraphies suggest they are from MIS5. The reconstructions vary from open *Betula* forest (Anadyr basin; Fig. 1), to *Larix* forest with tree *Betula*, tree *Alnus*, and *Pinus pumila* (northern Chukotka, upper Kolyma and Indigirka basins; Yana-Indigirka-Kolyma Lowland) to *Larix* forest with *Picea* (upper Kolyma and Indigirka basins; northern Lowlands; northern Okhotsk sea coast). Although *Picea* may have extended almost to the coast of the East Siberian Sea, the Lake E record raises the possibility that these sites may actually be from MIS11 (see below). Nonetheless, the preponderance of woody remains and nature of the MIS5.5 pollen spectra at Lake E strongly indicate the extensive presence of forest in Northeast Siberia and likely establishment of deciduous forest in the Chukchi Uplands. Analog-based analyses of the pollen data indicate the prevalence of warmer and wetter conditions as compared to modern (Lozhkin et al., 2007; Melles et al., 2012).

### 4.3 MIS11

Three vegetation types characterize MIS11 (Fig. 3), and all types indicate greater summer warmth and annual precipitation as compared to modern (Melles et al., 2012). However, the history of MIS11 diverges dramatically from the vegetation succession seen in MIS5 and MIS1 (i.e., *Betula-Alnus-Pinus pumila*).

The earliest climatic amelioration in MIS11 is indicated by high percentages of *Betula* and decreasing percentages of Poaceae pollen (pollen zone 11a). In contrast to the MIS6/MIS5 and MIS2/MIS1 transitions, *Betula* dominated vegetation appears to persist for a longer time in MIS11 prior to the arrival of *Alnus*. Lower zone 11a, representing the earliest effects of warming, was likely characterized by *Betula* shrub tundra with *Salix* and Ericales. Possibly shrubs achieved large growth forms as occurred in the PGTM. *Betula* pollen increases to 45–60% in the mid-to-upper levels of zone 11a, percentages that generally exceed modern values from Northeast Siberia

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Anderson et al., 2002; Lozhkin et al., 2002). These values more closely approximate modern surface samples from Kamchatka (Lozhkin and Anderson, unpublished data) where forests of *Betula ermanii* are common. In Alaska, highest *Betula* percentages (> 40 %) also occur in interior forests, where *Betula papyrifera* is found in valleys and lower mountain slopes (Anderson and Brubaker, 1993). Inasmuch as we can draw from modern analogs, these data suggest that *Betula* shrub tundra probably was replaced by *Betula* woodland. The presence of *Larix* pollen, even in only a single sample, further suggests the establishment of deciduous woodland. The decreasing percentages of graminoid and Ericales and increases in *Betula* pollen likely indicate a greater presence of wooded landscapes by upper zone 11a.

Zone 11b, representing the climatic optimum and corresponding to MIS11.3, includes some variation within the pollen assemblage. The pollen spectra between 1855–1905 cm: (1) are dominated by *Picea* sect. *Eupicea*, *Alnus*, and *Betula*; (2) show a consistent appearance of *Larix*; and (3) have decreased percentages of herb taxa, particularly Poaceae. *Pinus* subg. Haploxylon pollen appears initially in low amounts but increases in later spectra (1827–1855 cm). This pollen type may represent either *Pinus pumila* or *Pinus sibirica* (see below). The increase in *Pinus* pollen is associated with a slight decline in *Betula* and *Alnus* pollen.

The abrupt increase in *Picea* pollen at the zone 11a-11b boundary suggests a rapid replacement of deciduous woodland by *Picea obovata*-*Larix*-*Betula*-*Alnus* forest. The expansion of *Picea obovata* to Lake E represents a range extension northeastward of ~2000 km (Kolossova, 1980). *Chosenia macrolepis* possibly was present in this forest (see below), but its pollen can not be differentiated from shrub *Salix*. Herb pollen is virtually absent, which is characteristic of modern spectra from closed dark coniferous forest. Although other coniferous taxa (e.g., *Picea* sect. *Omorica*, *Pinus* subg. *Diploxylon*) and *Corylus* appear, these grains likely represent long-distance transport, and the pollen is not indicative of a local presence. Shrub communities of *Betula*, *Alnus*, and *Salix* grew along streams and on moist sites near the lake shore. Shrub tundra perhaps

occupied higher elevations, but the regional vegetation was dominated by dark coniferous forest.

The inferred vegetation for early MIS11.3 (1855–1905 cm) is reminiscent of that found today in the middle reaches of the Yama and Maimandzha Rivers, which is located ~ 150 km to the east of Magadan (Fig. 1) and thought to be a relict forest from a previous warm interval (Rheutt, 1970). Its plant communities contrast to the regional vegetation of light coniferous forest dominated by *Larix gmelinii* and *Pinus pumila*. In the Yama-Maimandzha lowlands, the forest is a mix of *Picea obovata*, *Larix*, tree *Alnus* (*Alnus hirsuta*), and tree *Betula* (*B. platyphylla*, *B. lanata*). *Populus suaveolens* and *Chosenia macrolepis* are limited to floodplains. *Picea* grows to 20<sup>+</sup> m heights, and *Larix* and *Alnus hirsuta* have diameters of 30–50 cm (Lozhkin, personal observation). The percentages of *Pinus* pollen occurring between 1855 and 1905 cm are sufficiently low to suggest *Pinus* shrubs or trees were likely absent or at best minor components of the ancient vegetation. This pattern is consistent with the modern Yama-Maimandzha *Picea* forest, where *Pinus sibirica* is absent and *Pinus pumila* is restricted to other areas of the drainage.

Dark coniferous forest continues to dominate the landscape during later MIS11.3 (1827–1855 cm). Pollen percentages indicate that *Picea obovata*, *Alnus hirsuta*, and *Betula* remain important elements in the forest communities, although the abundance of the latter two species perhaps decreased from earlier in the zone. The most significant shift in the pollen assemblage of late MIS11.3 is the rise in *Pinus* subg. Haploxylon pollen. As mentioned above, this pollen type represents both *Pinus pumila* and *Pinus sibirica*. Modern range distributions of *Picea obovata* and *Pinus pumila* do not overlap in Northeast Siberia (*Pinus pumila* only) or to the west of the Lena drainage (*Picea obovata* only; Kolosova, 1980). However, the two conifers do co-occur in the middle and upper reaches of the Lena catchment, extending southward to the Amur drainage. Here mean July temperatures are usually  $\leq 12^{\circ}\text{C}$  (Mock, 2002). In contrast, *Pinus sibirica* is found from south Eastern Siberia to northern sectors of the Western Siberian Plain, a distribution it shares with *Picea obovata*. In some of this region, mean

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



July temperatures can be as high as 20°C. The three species ranges overlap in the area of Lake Baikal, where their distribution reflects altitudinal gradients. *Larix-Picea-Pinus* forest is found along large flat areas between ~ 1100–1900 m, and *Pinus pumila* is part of a subalpine tundra zone beyond 1900 m (Suslov, 1961).

5 These modern distributions suggest three alternative paleovegetation interpretations. If the pollen is exclusively from *Pinus pumila*, its increase suggests that the *Picea* forest: (1) became sufficiently open to allow for the establishment of *Pinus* shrubs in the understory; and/or (2) the shrubs formed a mid-elevation shrub zone beyond altitudinal treeline. As compared to early MIS11, this vegetation change would perhaps  
10 suggest a slight summer cooling (to open the forest or lower treeline) and/or increased depth/duration of snow fall (required to protect *Pinus pumila* from winter desiccation; Khokhyrakov, 1985). However, the persistence of low percentages of herb pollen does not suggest a significant opening of the forests or decrease in *Picea* cover on mountain slopes. If the pollen represents *Pinus sibirica*, then the Lake E record marks a range  
15 extension of ~ 2500 km (Kolossova, 1980) and possibly an extensive forest cover across northern Chukotka. The addition of *Pinus sibirica* would not require changes in summer conditions as it shares similar requirements with *Picea obovata*, raising questions as to why *Pinus sibirica* was absent in early MIS11.3. If both *Pinus* species are present, then the regional and/or local vegetation probably had a strong altitudinal component  
20 with increased snow fall to aid establishment of *Pinus pumila*.

Quantitative paleoclimatic reconstructions indicate little systematic change in summer temperatures (ranging from ~ 12–16°C) between early and late MIS11.3 (Melles et al., 2012). The absence of *Pinus sibirica* during early MIS11.3 probably was not caused by insufficient summer temperatures, especially given the establishment of *Picea obovata* at the time. Possibly delayed migration from an unknown but distant population  
25 may account for the tree's absence. Mean annual precipitation is estimated to be at its highest during late MIS11.3 (> 600 mm yr<sup>-1</sup> vs. ~ 350 mm yr<sup>-1</sup> during early MIS11.3). If this trend reflects greater seasonal and/or annual snow fall, then this shift in climate would provide a reasonable mechanism for the expansion of *Pinus pumila*. This

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

conclusion, if correct, would indicate that early MIS11.3 experienced drier cool-seasons or seasonal transitions as compared to late MIS11.3. This interpretation would also suggest that *Pinus pumila* was the main *Pinus* species present during late MIS11.3.

Pollen zone 11c represents the final stages of MIS11 and is marked by the sharp decrease in *Picea* pollen, a more gradual decline in *Alnus* pollen, and increases in herb pollen (primarily from Cyperaceae and Poaceae). Areas of graminoid meadows or tundra were more common than previously, suggested by the higher percentages of herb pollen, slightly greater amounts of Ericales and *Salix* pollen, and to a lesser extent the occurrence of higher *Sphagnum* values. Trees were limited on both local and regional landscapes. The loss of *Larix* pollen late in the zone suggests that tundra dominated in the latter part of MIS11.

The forest-tundra of zone 11c perhaps was similar to areas found today in Eastern Siberia (e.g., lower Lena River valley), where *Picea* and *Larix* form latitudinal treeline with isolated stands of *Larix* scattered northward across the tundra. *Pinus pumila* may have persisted from zone 11b, having a similar role as seen in the contemporary vegetation. However, paleoclimatic reconstructions show a sharp decline in mean annual precipitation (Melles et al., 2012), a potentially adverse factor in the survival of *Pinus pumila*. Decreased summer temperatures as indicated by the quantitative climate estimates would have been sufficiently severe to eliminate *Pinus sibirica*, if it was present on the zone 11b landscapes.

#### 4.4 MIS 31

Of all warm periods discussed in this paper, MIS31 presents the most unusual pollen spectra (Fig. 4). While dominated by typical boreal taxa (*Picea*, *Larix*, *Abies*, *Betula*, *Alnus*), the MIS31 assemblages also contain small numbers of more temperate deciduous taxa (e.g., *Corylus*, *Carpinus*, *Acer*, *Quercus*). Not surprisingly given the 1.1 Ma age, these assemblages seem to lack exact analogs in the modern landscapes. Although the pollen diagram is somewhat skeletal, the data are sufficient to determine general characteristics of the MIS31 vegetation.

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The MIS32/MIS31 transition (pollen zone 31a) is similar to that of MIS6/MIS5 (Fig. 2; pollen zone 5a) with spectra dominated by *Betula* and *Alnus* pollen and with very low numbers of herb types. However, differences do exist between the two transitions, including the minor presence of *Larix* pollen and occasional occurrence of the warm taxa *Corylus*, *Carpinus*, and *Myrica* in zone 31a. As during pollen zone 5a, the regional vegetation during early MIS31 probably was dominated by deciduous trees possibly with tree-sized shrubs including *Betula*, *Alnus*, and *Salix*. Unlike MIS5, *Larix* was established in the Lake E watershed and likely across the Upland. The low percentages of herb pollen suggest that a relatively dense deciduous forest rather than open woodland was established at this time. Although pollen from other deciduous trees is the result of long-distance, wind transport, occurrence of these taxa suggests that temperate forests were in closer proximity to north-central Chukotka as compared to today.

The zone 31b assemblage represents the thermal maximum within MIS31 and is characterized by several *Picea* and *Pinus* species, *Larix*, *Abies*, *Betula*, *Alnus*, and < 5% herb pollen. Several partial analogs to this assemblage can be found in Asiatic Russia. The northernmost is the dark coniferous forest of the lower Yenisei valley (northern sector of the Western Siberia Plain), where today *Picea obovata*, *Pinus sibirica*, and *Pinus sylvestris* are found. A southern variant of this forest occurs in the upper Yenisei drainage between ~ 51 to 54° N, an area that is transitional between the Siberian dark coniferous taiga and Mongolian steppes. In this region, taiga dominates in the mountains and forests between ~ 1000 and 2000 m and includes *Larix sibirica*, *Picea obovata*, *Abies sibirica*, *Pinus sibirica*, and occasional *Pinus sylvestris*. A third possible analog is from the western shores of the Sea of Okhotsk (from ~ 51° N southward to the lower Amur valley), where the taiga is a mix of *Picea glehnii*, *Abies nephrolepis*, *Larix gmelinii*, and *Betula ermanii*. *Pinus pumila* is limited to protected sites where it can grow in dense thickets achieving over 2 m height (Suslov, 1961). The more temperate taxa *Corylus mandshurica*, *Quercus mongolica*, *Carpinus cordata*, and *Acer mandshuricum* are present in the Sikhote-Alin Mountains, which border the Amur drainage. Possibly the zone MIS31b vegetation represented a mix of plant communities

from the Siberian dark coniferous forest and the Okhotsk taiga, as no modern vegetation type totally corresponds to taxa represented in the pollen assemblage.

The combined percentages of *Betula* and *Alnus* pollen, as in zone 31a, dominate the MIS31b spectra and demonstrate that deciduous forest remained an important landscape component. The presence of *Corylus* in two samples in minor but significant amounts (2–3%) suggests that the plant was probably present within the region. Today the nearest *Corylus* population (*Corylus mandshurica*) is found in the lower reaches of the Amur drainage and its establishment near Lake E would represent a ~2400 km range extension (Kolossova, 1980). Inasmuch as we can draw on the Okhotsk taiga as analog, then it is likely that during zone 31b the vegetation in the Lake E crater and neighboring Uplands was characterized by a marked altitudinal zonation. *Betula-Larix-Alnus* forest likely dominated river valleys and lower slopes. Possibly the *Salix* pollen represents *Chosenia macrolepis*, a common floodplain tree in modern landscapes. Transitioning to mountain slopes, deciduous forest was replaced by conifer communities of *Picea*, *Pinus*, and *Abies*. *Betula ermanii* is common today in *Picea-Abies* forest, sometimes forming pure stands. This *Betula* species occupies rockier and steeper slopes. *Corylus* was likely an understory shrub that grew on warmer, south-facing slopes. Other temperate taxa (e.g., *Acer*, *Carpinus*), if present, would be restricted to the warmest sites. *Pinus sylvestris*, if the *Pinus* subg. Diploxylon pollen is not the result of long-distance transport, was not abundant. As in the case of MIS11, the *Pinus* subg. Haploxylon pollen may be interpreted in two ways. If from *Pinus pumila*, the species likely grew in protected, open areas within the forest and possibly formed high shrub tundra beyond altitudinal treeline. As in zone 11b, low percentages of herb taxa suggest closed forests that extended far up mountain slopes, making the occurrence of a *Pinus pumila* belt less likely. If the pollen represents *Pinus sibirica*, then the taxon would be a member of the coniferous forest.

Within pollen zones 31c and 31d, the pollen spectra indicate steady vegetation changes implying gradual climatic cooling and increasing aridity. Pollen from evergreen conifers decreases, whereas *Betula* and *Alnus* continue as dominant taxa with a

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



consistent appearance of *Larix* pollen. Deciduous forest probably became the most extensive vegetation type both locally and regionally. The increase in Ericales, Poaceae, and total herb pollen and *Sphagnum* spores indicates an opening of the forest and establishment of tundra communities by late zone 31c and zone 31d.

## 5 Discussion and conclusions

The palynological data presented here and in previous papers (Lozhkin and Anderson, 2007; Matrosova, 2009) are in many senses a preview of the exciting paleoenvironmental information contained within the Lake E archive. For example, our focus on only a small part of the vegetation history has revealed interglaciations with an unexpected magnitude of warming (Melles et al., 2012). The pollen data described herein further indicate that there are at least two types of vegetation responses to interglacial conditions. The vegetation in the first type of thermal optimum is dominated by deciduous tree and shrub species and corresponds to the two most recent warming events (PGTM, MIS5.5): the second type, or “super” interglaciation (MIS11.3, MIS31), indicates climax vegetation dominated by evergreen conifers, particularly *Picea*. These exceptional climatic events experienced warmer summers (+4.5°C) and greater annual precipitation (~300 mm) as compared to the PGTM and MIS5.5 (Melles et al., 2012), which themselves are characterized by summers that are warmer and wetter than present (increases of ~2°C and up to 20 mm for mean July temperature and precipitation, respectively; Lozhkin et al., 2007).

Despite the classification into types, the interglaciations share some commonalities. One of the characteristics of the interglacial intervals at Lake E is that they show greater variability in vegetation types as compared to glaciations. These latter periods display great uniformity in their pollen assemblages both through time as in the Lake E record and also in space, at least in the cases of the latest glaciation when the most paleobotanical data are available (Brigham-Grette et al., 2004). This observation implies that: (1) glacial climate and vegetation were less spatially and temporally variable as

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



compared to interglacial times; and (2) once a “threshold” of coolness and/or dryness is surpassed the vegetation response is limited to herb-dominated tundra or steppe tundra. The variability in vegetation response may also suggest that interglacial forests exerted a greater influence on the climate through such feedbacks as changes in albedo and fluxes of sensible heat (e.g., Foley et al., 1994; Levis et al., 1999; Eugster et al., 2000) as compared to that associated with rather uniform glacial tundra.

In each of the glacial to interglacial transitions the rapid establishment of plant communities dominated by woody taxa occurred. The earliest interglacial vegetation was *Betula* shrub tundra (note: this is not evident in early MIS31, but we expect additional pollen counts will show a parallel trend to the other interglaciations) that quickly was replaced by deciduous forest or woodland. For the PGTM and MIS5, the establishment of *Betula* woodland/high shrub tundra and *Betula-Larix* forest/high shrub tundra during the PGTM and MIS5, respectively, represents maximum summer warming. However, during the older warm periods, *Betula-Larix* (MIS11) or *Betula-Alnus-Larix* (MIS31) forests were transitional as interglacial climates continued to become warmer and wetter. Edwards et al. (2005) noted that the widespread establishment of deciduous forest in Beringia during the PGTM represented a novel biome that differed structurally and functionally from evergreen coniferous forest and tundra that dominate arctoboreal settings today. They also commented that deciduous boreal forest should be included in modeling efforts that explore likely biosphere-atmosphere feedbacks and possible impacts caused by global warming. The new Lake E data underscore the importance of this proposition, as deciduous forest seems to play an essential role in interglacial vegetation history, whether it was as transitional or as climax vegetation.

The different interglacial “modes” also suggest the need to explore the potential impact of a range of forcing mechanisms and feedbacks involved in “warm” and “warmer” scenarios. Melles et al. (2012), for example, suggested that the melting of the West Antarctic Ice Sheet, with consequent changes in ocean and atmospheric circulation, was possibly a major factor contributing to the magnitude of temperature and precipitation increases reconstructed for the “super” interglaciations of MIS11 and MIS31.

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Northeast Asia and Beringia before the Last Glacial Maximum, edited by: Madsen, D., The University of Utah Press, Salt Lake City, UT, USA, 29–62, 2004.

Brigham-Grette, J., Melles, M., and Minyuk, P. (Eds.): Arctic Record of the Last 250 ka from El'gygytgyn Crater Lake, NE Russia, *J. Paleolimnol.*, 37, 162 pp., 2007.

Claussen, M.: Variability of global biome patterns as a function of initial and boundary conditions in a climate model, *Clim. Dynam.*, 12, 371–379, 1996.

Chapin III, F. and Starfield, A.: Time lags and novel ecosystems in response to transient climatic change in arctic Alaska, *Climatic Change*, 35, 449–461, 1997.

Czerepanov, S.: *Vascular Plants of Russia and Adjacent States (the Former USSR)*, Cambridge University Press, Cambridge, UK, 516 pp., 1995.

Edwards, M., Brubaker, L., Lozhkin, A., and Anderson, P.: Structurally novel biomes: a response to past warming in Beringia, *Ecology*, 86, 1696–1703, 2005.

Eugster, W., Rouse, W., Pielke, R., McFadden, J., Baldocchi, D., Kittel, T., Chapin, F., Liston, G., Vidale, P., Vaganov, E., and Chambers, S.: Land-atmosphere energy exchange in arctic tundra and boreal forest: available data and feedbacks to climate, *Global Change Biol.*, 6, 84–115, 2000.

Foley, J., Kutzbach, J., Coe, M., and Levis, S.: Feedbacks between climate and boreal forests during the Holocene epoch, *Nature*, 371, 52–52, 1994.

Harrison, S., Kutzbach, J., Prentice, I., Behling, P., and Sykes, M.: The response of northern hemisphere extratropical climate and vegetation to orbitally induced changes in insolation during the last interglaciation, *Quaternary Res.*, 43, 174–184, 1995.

Kaplan, J., Bigelow, N., Prentice, I., Harrison, S., Bartlein, P., Christensen, T., Cramer, W., Matveyeva, N., McGuire, A., Murray, D., Razzhivin, V., Smth, B., Walker, D., Anderson, P., Andreev, A., Brubaker, L., Edwards, M., and Lozhkin, A.: Climate change and arctic ecosystems: 2. Modeling, paleodata-model comparisons, and future projections, *J. Geophys. Res.*, 108, 8171, doi:10.1029/2002JD002559, 2003.

Khokhyrakov, A.: *Flora of Magadan Oblast*, IBPB, FEB, USSR Academy of Sciences, Vladivostok, Russia, 48 pp., 1985 (in Russian).

Kolosova, L. (Ed.): *Geographic Atlas*, Central Office of Geodesy and Cartography of the Soviet Ministry of the USSR, 4th Edn., Moscow, Russia, 1980 (in Russian).

Kozhevnikov, Y.: Vascular plants near El'gygytgyn Lake, in: *Natural Depression El'gygytgyn Lake (Problems of Study and Protection)*, NEISRI, FEB, RAS, Magadan, Russia, 62–82, 1993 (in Russian).

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Levis, S., Foley, J., and Pollard, D.: Potential high-latitude vegetation feedbacks on CO<sub>2</sub>-induced climate change, *Geophys. Res. Lett.*, 26, 747–750, 1999.
- Lozhkin, A.: Geochronology of late Quaternary events in northeastern Russia, *Radiocarbon*, 35, 429–433, 1993.
- 5 Lozhkin, A. and Anderson, P.: The last interglaciation in northeast Siberia, *Quaternary Res.*, 43, 147–158, 1995.
- Lozhkin, A., Anderson, P., Eisner, W., Ravako, L., Hopkins, D., Brubaker, L., Colinvaux, P., and Miller, M.: Late Quaternary lacustrine pollen records from southwestern Beringia, *Quaternary Res.*, 39, 314–324, 1993.
- 10 Lozhkin, A., Anderson, P., Vartanyan, S., Brown, T., Belaya, B., and Kotov, A.: Reconstructions of late Quaternary paleo-environments and modern pollen data from Wrangel Island (northern Chukotka), *Quaternary Sci. Rev.*, 20, 217–233, 2001.
- Lozhkin, A., Anderson, P., Belaya, B., and Stetsekno, T.: Reflections on modern pollen rain of Chukotka from bottom lake sediments, in: *Quaternary Paleogeography of Beringia*, edited by: Simakov, K., NEISRI FEB RAS, Magadan, Russia, 40–50, 2002 (in Russian).
- 15 Lozhkin, A., Anderson, P., Matrosova, T., and Minyuk, P.: The pollen record from El'gygytgyn Lake: implications for vegetation and climate histories of northern Chukotka since the late middle Pleistocene, *J. Paleolimnol.*, 37, 135–153, 2007.
- Matrosova, T.: Reconstruction of vegetation and climate in northern Chukotka during the last 20 350 thousand years (according to palynological evidence from El'gygytgyn Lake), *Vesnik*, 1, 23–30, 2009 (in Russian).
- Melles, M., Brigham-Grette, J., Minyuk, P., Nowaczyk, N., Wennirch, V., DeConto, R., Anderson, P., Andreev, A., Coletti, A., Cook, T., Haltia-Hovi, E., Kukkonen, M., Lozhkin, A., Rosen, P., Tarasov, P., Vogel, H., and Wagner, B.: 2.8 million years of arctic climate change from Lake El'gygytgyn, NE Russia, *Science*, 337, 315–320, 2012.
- 25 Mock, C. J.: Regional climates of Russia, in: *Late Quaternary Vegetation and Climate of Siberia and the Russian Far East (Palynological and Radiocarbon Database)*, edited by: Anderson, P. and Lozhkin, A., NOAA and Russian Academy of Science, Far East Branch, Magadan, 17–26, 2002.
- 30 PALE Steering Committee: Research Protocols for PALE: Paleoclimatic of Arctic Lakes and Estuaries, PAGES Workshop Report Series 94-1, PAGES Core Project Office, Bern, Switzerland, 53 pp., 1994.

Reutt, A.: Vegetation, in: North of the Far East, edited by: Shilo, N., Nauka, Moscow, Russia, 257–299, 1970 (in Russian).

Suslov, S.: Physical Geography of Asiatic Russia, W.H. Freeman, London, UK, 594 pp., 1961.

CPD

9, 245–267, 2013

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Fig. 1.** Map of northeast Siberia showing location of Lake El'gygytyn.

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson



**Fig. 2.** Percentage diagram of select taxa from MIS1-MIS2 and MIS4-MIS6. Note that labels for the pollen assemblages include both a number, indicating the isotope stage, and a letter, indicative of the pollen zone. Climatic optima are represented by zones 1a and 5a.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Vegetation responses to interglacial warming

A. V. Lozhkin and  
P. M. Anderson



**Fig. 3.** Percentage diagram of select taxa from MIS10-12. The climatic optimum is represented by zone 11b.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

