

**Petrophysical
characterization of
the lacustrine
sediment succession**

A. C. Gebhardt et al.

Petrophysical characterization of the lacustrine sediment succession drilled in Lake El'gygytgyn, Far East Russian Arctic

**A. C. Gebhardt¹, A. Francke², J. Kück³, M. Sauerbrey², F. Niessen¹,
V. Wennrich², and M. Melles²**

¹Alfred Wegener Institute for Polar and Marine Research, Columbusstraße,
27515 Bremerhaven, Germany

²University of Cologne, Institute of Geology and Mineralogy, Zùlpicher Straße 49A,
50674 Cologne, Germany

³German Research Centre for Geosciences, Telegrafenberg, 14473 Potsdam, Germany

Received: 17 December 2012 – Accepted: 1 January 2013 – Published: 18 January 2013

Correspondence to: A. C. Gebhardt (catalina.gebhardt@awi.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Seismic profiles of Far East Russian Lake El'gygytyn which was formed by a meteorite impact some 3.6 million years ago show a stratified sediment succession that can be separated into Subunits Ia and Ib at approximately 167 m below lake floor (= ~ 3.17 Ma). The former is well-stratified, while the latter is acoustically more massive. The sediments are intercalated with frequent mass movement deposits mainly in the proximal parts, while the distal part is almost free of such deposits at least in the upper part. In spring 2009, a long core drilled in the lake center within the framework of the International Continental Scientific Drilling Program (ICDP) penetrated the entire lacustrine sediment succession down to ~ 320 m below lake floor and about 200 m further into the meteorite-impact related bedrock. Downhole logging data down to 390 m below lake floor show that the bedrock and the lacustrine part of the core differ largely in their petrophysical characteristics. The contact between the bedrock and the lacustrine sediments is not abrupt, but rather transitional with a mixture of impact-altered bedrock clasts in a lacustrine matrix with varying percentages. Physical and chemical proxies measured on the cores can be used to divide the lacustrine part into five different clusters. These can be plotted in a redox-condition vs. input type diagram with total organic carbon content and magnetic susceptibility values indicating anoxic or oxic conditions and with the Si/Ti ratio representing more clastic or more biogenic input. Plotting the clusters in this diagram allows identifying clusters that represent glacial phases (Cluster I), super interglacials (Cluster II), and interglacial phases (Clusters III and IV).

1 Introduction

The Arctic region is strongly susceptible to global change and, at the same time, plays a major role in the global climate system through feedback processes in the oceans, the atmosphere, and the cryosphere. It is, thus, of importance to understand past climate changes under different climate-forcing conditions in order to make

CPD

9, 351–391, 2013

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Petrophysical
characterization of
the lacustrine
sediment succession**

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

sedimentation regime as well as the first occurrence of ice rafted debris (IRD) in the Arctic Ocean from about 2.53 Ma, and the onset of large-scale glaciation in Scandinavia (by means of a marked increase in IRD flux) was dated to 2.75 Ma by Fronval and Jansen (1996) and Jansen et al. (2000). Since then, the Arctic realm has experienced several advances and retreats of glaciers and ice sheets. A dropstone which was found in sediments as old as 45 Ma and the frequent occurrence of IRD since the early Miocene in a marine record from the Lomonossov Ridge, however, show that the onset of the transition from a greenhouse world to colder climate with sea ice and icebergs might have begun much earlier than hitherto assumed (Moran et al., 2006).

The El'gygytgyn area was never subject to glacial overprint since its formation (Glushkova and Smirnov, 2007) and, thus, the lake contains an undisturbed climate record of approximately 3.6 million years unique for the terrestrial Arctic realm. This record was drilled within the framework of the International Continental Scientific Drilling Program (ICDP). A permafrost core (ICDP Site 5011-3) was retrieved from the eastern shoreline in late autumn 2008, and during winter/spring 2009, a 517 m long drill core (ICDP Site 5011-1) containing lacustrine sediments and the impact-related bedrock underneath was retrieved from the ice cover of the lake (Melles et al., 2011).

This paper aims at characterizing the lacustrine part of core 5011-1 as well as the transitional zone between the lacustrine sediments and the impact-related bedrock by means of petrophysical parameters such as physical properties and downhole logging measurements. These findings are then compared to the facies description by Melles et al. (2007, 2012) and their interpretation.

2 General settings of the investigation area

2.1 Study area

Lake El'gygytgyn (67° 30' N, 172° 05' E) is located about 100 km north of the Arctic Circle in Central Chukotka, NE Russia (Fig. 1). It was formed by a meteorite impact

that was dated using $^{40}\text{Ar}/^{39}\text{Ar}$ to about 3.6 million years (Layer, 2000; Gurov et al., 1979a, b; Belyi, 1998). The lake's surface lies at about 490 m a.s.l. and the surrounding crater rim reaches altitudes of 900 to almost 1000 m a.s.l.

The lake is roughly circular with a diameter of 12 km. Its catchment is limited to the crater rim (293 km² in total, including lake surface) with about 50 small ephemeral creeks draining into the lake (Nolan and Brigham-Grette, 2007). The Enmyvaan River at the southern edge of the lake is its only outflow (Fig. 1a). The lake has a bowl-shaped form with a flat, central plain of 170 m water depth and flanks that are steepest in the northern and northeastern part. A shelf of 10 to 12 m water depth has developed in the southeastern, southern, and southwestern to western part of the lake (Fig. 1a).

The lake is presently ice-covered during 9–10 months with only a short period of completely open water (Nolan and Brigham-Grette, 2007). During the short summer season, the monomictic and ultra-oligotrophic lake gets mixed completely (Nowaczyk et al., 2002; Nolan and Brigham-Grette, 2007). The catchment vegetation consists mainly of moss-tundra interspersed by few shrub willows; the modern tree line lies about 150 km further south and west (Nowaczyk et al., 2002). The current wind system has a bipolar mode with winds approximately from north and south, respectively (Nolan and Brigham-Grette, 2007).

2.2 Lithological succession

The impact crater shows internal geometries as expected for a crater of its size; a central uplift structure interpreted in the form of a central uplift ring structure was revealed by seismic refraction data; it is overlain by an impact breccia (suevite) (Gebhardt et al., 2006). The lacustrine sediments can be divided into two units by means of refraction data; the upper unit is characterized by a seismic velocity of 1550 ms⁻¹ and a thickness of about 170 m, the lower unit by 1650 ms⁻¹ and a variable thickness of 190 m on top of the uplift ring structure to 290 m in the surrounding basin (Gebhardt et al., 2006). Mass movements (slides, debris flows, and turbidites) are a common feature mainly in proximal parts of the lacustrine sediments. These mass movement

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



packages have already been described as massive, acoustically transparent sediment bodies by Niessen et al. (2007) from high-resolution 3.5 kHz seismic data. Gebhardt et al. (2005) distinguished between two types of mass movement sediments: deposits with a distinct “nose” at their front end, forming a well-defined distal border, and deposits that gradually pinch out, forming a diffuse distal border, poorly visible in the profiles. Melles et al. (2007) report fine-grained turbidites from the pilot core recovered in 1998 (PG1351; Fig. 1). A detailed study of four cores along a transect from the lake border towards the center by Juschus et al. (2009) demonstrated that initial slope failures transform into erosive debris and/or grain flows during their advance and accumulate as non-erosive turbidites in the lake center. This process of suspension settling out is also known from other lakes (e.g. Eyles et al., 2003; Finckh et al., 1984; Van Rensbergen et al., 1998, 1999). Whilst only turbidites and grain flow deposits reached the lake center during the studied ~ 350 ka of pilot cores PG1351 and Lz1024 (Juschus et al., 2009; Frank et al., 2013), investigations of the long 5011-1 cores have revealed several slides, slumps and debrites in deeper core sections (Sauerbrey et al., 2013). Slides, slumps, grain and debris flows can be recognized in the seismic sections, but turbidite sequences are rather thin and, thus, beyond resolution. Mass movement deposits make up approximately one third of the entire lacustrine record drilled in the lake. They were extensively studied by Sauerbrey et al. (2013).

In spring 2009, three drill cores were retrieved from the center of Lake El'gygytgyn (Site 5011-1, cores 5011-1A, 1B, 1C) down to a maximum depth of 517.3 m below lake floor (b.l.f.). A detailed description of all drilling details is given by Melles et al. (2011). The cores were transported to the laboratory facilities at the University of Cologne, Germany, where they were opened and described. Based on the core description together with several proxies, a composite profile was composed (Melles et al., 2012; Nowaczyk et al., 2013). The composite sediment core mainly consists of highly variable silty-clayey pelagic sediments divided into different facies types by Melles et al. (2012) and Brigham-Grette et al. (2013) interfingering with mass movement deposits (Sauerbrey et al., 2013).

**Petrophysical
characterization of
the lacustrine
sediment succession**

A. C. Gebhardt et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

“Facies A” consists of fine clastic laminations of less than 5 mm thickness (average is ~ 0.2 mm). The sediments of facies type A are mainly dark gray to black in color. This points at a stratified water column and anoxic bottom water conditions (Melles et al., 2012). The authors associate this facies type with peak glacial conditions and a perennial ice cover of the lake with mean annual air temperatures of at least $4 (\pm 0.5) ^\circ\text{C}$ less than today. This facies was already described in pilot core PG1351 as subunits 3 (“cold & dry”) and 4 (“cold & moist”), characterized by enhanced amounts of total organic carbon (TOC), medium to low biogenic silica content (Melles et al., 2007), and low magnetic susceptibility due to dissolution of magnetite by anoxic conditions (Nowaczyk et al., 2007). The “cold & dry” subtype is further referred to as A_d , “cold & moist” as A_m . Facies A sediments are limited to the younger part of the sediment record (Brigham-Grette et al., 2013), i.e. the uppermost ~ 124 m (≈ 2.6 Ma).

“Facies B” is the most abundant facies type in the composite profile and mainly consists of olive gray to brown, massive to faintly banded silt with greenish bands of typically 1–3 cm thickness. The sediments are characterized by a lack of sedimentary structures, pointing at bioturbation and oxygenated bottom water (Melles et al., 2012). This implies warmer climate with ice-free summers and a mixed water column. This facies reflects a wide range of glacial to interglacial settings including the modern situation. TOC content is rather low in facies type B due to high organic matter decomposition in oxic bottom water conditions; biogenic silica values are intermediate to high due to enhanced primary productivity, and magnetic susceptibility is high reflecting good preservation of magnetite (Melles et al., 2012).

“Facies C” is the least common facies type found in the composite profile (Melles et al., 2012). It is irregularly distributed and consists of distinctly reddish-brown silt. Melles et al. (2012) suggest oxidation of bottom sediments by a well-ventilated water column as responsible for the distinct reddish color. This facies type was interpreted as “super interglacial” conditions, e.g. during extraordinary warm MIS11 and 31 (Melles et al., 2012). Distinct laminae are found in facies type C, probably pointing at winter stratification and anoxic bottom water conditions under a seasonal ice cover. This is

**Petrophysical
characterization of
the lacustrine
sediment succession**

A. C. Gebhardt et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

further supported by high TOC values. Biogenic silica content is also exceptionally high due to diatom blooms probably caused by enhanced nutrient influx from the catchment. Magnetic susceptibility is rather low both due to dissolution of the magnetic susceptibility signal by the high biogenic silica content and partial dissolution of magnetite during periods (winters) with anoxic bottom water conditions.

“Facies D” is laminated similar to Facies A, but its laminae are significantly larger with an average thickness of up to ~ 1 cm. Laminae are characterized by distinct lower boundaries and a coarsening upward sequence from silt to clay with a higher total clay content than in Facies A. Facies D is mostly gray but has some red and green hues in its oldest parts. The well-preserved lamination points at a lack of bioturbation of the bottom sediment, and the characteristic coarsening upward in each lamina suggests repeated pulses of sediment delivery to the lake probably due to variations in fluvial input (Brigham-Grette et al., 2013). Facies D is limited to the Pliocene part of the record with the youngest occurrence at ~ 141 m b.l.f. (≈ 2.9 Ma).

“Facies E” comprises the transition from the impact-altered bedrock to lacustrine sediments. This transition is more/less gradual with sediments composed of impact breccia and impact melt blocks in a matrix of lacustrine sediments, with the bedrock-related particles being dominant in the lower and the lacustrine sediments in the upper part (Koeberl et al., 2012; Raschke et al., 2012).

“Facies F” comprises a wide variety of mass movement deposits, i.e. turbidites, debrites, slumps, slides and grain flows. A detailed description of the mass movement deposits and their distribution within the record is given by Sauerbrey et al. (2013). Only thin mass movement deposits (< 5 cm in thickness) were sampled in the composite profile of 5011-1, thicker ones were omitted. These thinner mass movement deposits are almost exclusively turbidites.

3 Data acquisition and processing

3.1 Seismic data

Prior to deep drilling, two seismic pre-site surveys were carried out in summers 2000 and 2003. In 2000, a single-channel survey was carried out using a Bolt 600B airgun (82 cm³, 6 s shot interval resulting in approximately 8 m shot distance) with a 20-element single-channel hydrophone streamer (Geoacoustics AE5000) as receiver (Niessen et al., 2007). Single-channel reflection data were bandpass-filtered (100-150-350-450 Hz), and an AGC was used for display. In 2003, 2 single-channel and an additional 8 multi-channel profiles were acquired using a Mini-GI gun triggered in G-gun mode at a pressure of 110 bar (426 cm³, 10 s shot interval resulting in approximately 12 m shot distance). For the multi-channel profiles, a 14-channel streamer with an offset of 130 m and a hydrophone spacing of 10 m was used as receiving array (details are given in Niessen et al., 2007). Multi-channel data were processed in a standard sequence including bandpass filtering (70-90-240-300 Hz), velocity analysis, CMP stacking, and predictive deconvolution. Tracklines are shown in Fig. 1.

3.2 Physical properties development

Physical properties data of cores 5011-1A, 1B and 1C were acquired using a Geotek Multi-Sensor Core Logger (MSCL; Geotek Ltd., UK) both in the field laboratory during the drilling campaign in 2009 (magnetic susceptibility measurements on whole cores) and at the Alfred Wegener Institute (AWI) in Bremerhaven, Germany between October 2009 and January 2011 (density measurements on split cores). The data were complimented with density and magnetic susceptibility data from pilot core Lz1024 which were measured at AWI in March 2004 on whole cores.

“Magnetic susceptibility (MS)” was measured in SI units using a Bartington MS-2 meter equipped with a loop sensor of 80 mm internal diameter. Data correction was done with respect to the specific core and loop sensor diameters according to the Bartington

CPD

9, 351–391, 2013

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Manual (Geotek, 2000). Even though temperature inside the field lab container was sometimes variable due to opening and closing of the door with inside temperatures around +20 °C and outside temperatures between −45 and −20 °C, no severe drifting of the temperature-sensitive sensor could be observed. The small drifting that occurred was significantly lower than the lowest susceptibility readings and, thus, did not affect the data. Both magnetic susceptibility and density data were corrected for outliers and composite profiles were spliced accordingly to the sampling scheme used for the discrete samples (Wennrich et al., 2013)

“Gamma-ray density (GRA)” was measured using a ^{137}Cs source mounted on the Geotek MSCL. For density calibration, standard core-size semi-cylinders consisting of different proportions of aluminum and water were logged prior to the cores according to the method described by Best and Gunn (1999) but modified for split cores. Split cores were only approximately 33 mm thick, which is beyond of what our Geotek MSCL can measure reliably. To convert raw gamma ray attenuation counts to density, however, exact thickness measurements are required. We therefore used the surface scans that were measured by the ITRAX XRF core scanner (COX Analytical Systems, Sweden) at the University of Cologne (see Wennrich et al., this volume) in the course of the XRF measurements on the same split cores. These surface scans were calibrated for thickness using a semi-cylindrical piece with a radius of 33.15 mm to simulate a standard split core, and three pieces that were thicker (+10 mm, +20 mm) or thinner (−10 mm) than the standard to calibrate the entire range of possible sediment thicknesses. GRA was calculated using the method described in the Geotek manual (Geotek, 2000).

3.3 Downhole logging data

While drilling hole 5011-1C, operations were stopped four times to allow for the acquisition of downhole logging data. All data presented here were acquired using slimhole probes manufactured by Antares (Germany). Operation of the probes under the extreme conditions of an Arctic winter drilling campaign went overall well but also took its toll with one probe completely broken (acoustic velocity probe) and another partially

**Petrophysical
characterization of
the lacustrine
sediment succession**

A. C. Gebhardt et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 damaged (caliper probe). For downhole logging sessions, the pipe was pulled out of the hole (except for the uppermost approximately 20 m where the casing was pushed into the sediment for stabilization) leaving a sufficiently stable borehole wall. After downhole logging sessions were finished, the pipes were redeployed, and drilling operations were resumed. For drilling operations, bentonite was used as drilling fluid. Downhole logging was carried out to a maximum depth of 394 m below lake floor. In order to fit the downhole logging depths to the composite profile depths, the entire downhole logging dataset was shifted downwards by 3 m. This results in an apparent discrepancy with depths used by the community that is working on the impact-related bedrock (e.g. Koeberl et al., 2012; Raschke et al., 2012); we therefore also shifted their depths by 3 m downwards for comparison with our data.

Electrical resistivity and magnetic susceptibility data of the uppermost approximately 143 m could not be used as they were disturbed by the pipes of nearby abandoned holes 1A and 1B.

15 “Electrical resistivity (ER)” of the surrounding sediments/rock at two different lateral distances from the borehole wall (deep ~ 60 cm and shallow ~ 20 cm, with the actual penetration depending on rock porosity and the resistivity of fluid and rock) was measured using a dual laterolog probe. The probe has a vertical resolution of approximately 10 cm (electrode length: 8 cm), typical logging speed was 12 m min⁻¹.

20 “Borehole magnetic susceptibility (BMS)” was measured using a probe that basically consists of a receiver coil and a transmitter coil that is located 20 cm above the former inside a non-magnetic pressure housing. The transmitter coil induces a 1 kHz alternating magnetic field. Magnetic susceptibility was corrected for the two different borehole diameters drilled during the Lake El’gygytgyn deep drilling project. Down to 274.33 m composite depth (i.e. 443 m below lake surface), a bit size of 124 mm was used; a correction factor of 1.4 was applied for this part. In the deeper part of the hole, a smaller bit size of 98 mm was used for drilling/coring, and accordingly, a correction factor of 1.25 was used. The vertical resolution is approximately 20 cm (detector spacing), but relative

variations can be identified with a resolution of about 5 cm. Penetration is ~ 20 cm, typical logging speed was $8\text{--}10 \text{ m min}^{-1}$.

The “total amount of naturally occurring radioactive radiation (GR)” was measured using a total natural gamma ray probe. This GR probe was always run with any other probe for depth corrections. One GR curve was chosen as the reference (Master-GR), and all other GR curves with their attached other measurements were shifted to fit the Master-GR. Vertical resolution is approximately 10 cm, typical penetration into the rock is about 10 cm.

The “spectrum of the naturally occurring radioactive radiation (SGR)”, i.e. Uranium, Thorium, and Potassium, was measured using a natural gamma ray probe. Logging speed was not faster than 2 m min^{-1} for the SGR probe to allow gathering of a reliable gamma ray spectrum. Vertical resolution is ~ 10 cm, and penetration into the rock is around 15 cm. Gamma rays penetrate steel casing, therefore both the GR and the SGR probes could be run in cased holes. Corrections were carried out for the casing as well as for the different diameters of the borehole. Th and K values are often used for a first estimate and characterization of clay content in the sediments (e.g. Wonik, 2001; Ruffell and Worden, 2000; Schnyder et al., 2006) assuming that they are almost exclusively present in this grain-size fraction, and that K and Th are present in montmorillonite, illite, and kaolinite in different portions. To estimate the clay content in Lake El'gygytyn sediments, we used the approaches given by Wonik (2001):

$$C_{\text{cl}}(\text{K}) = \frac{K - K_{\text{sand}}}{K_{\text{clay}} - K_{\text{sand}}}, \quad (1)$$

and

$$C_{\text{cl}}(\text{Th}) = \frac{\text{Th} - \text{Th}_{\text{sand}}}{\text{Th}_{\text{clay}} - \text{Th}_{\text{sand}}}, \quad (2)$$

with C_{cl} = clay content (%), K_{sand} and Th_{sand} = K and Th content of sand, K_{clay} and Th_{clay} = K and Th content of clay. K and Th contents of sand are normally very low

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and were set to 0.1 % for K and 0.1 ppm for Th; K and Th contents of clay were set to the maximum K and Th values measured in the record, which is 4.5141 % for K and 22.4249 ppm for Th. K and Th in Eqs. (1) and (2) are the actual readings from the SGR dataset. A third approach uses the GR data as follows (Wonik, 2001):

$$V_{cl}(GR) = 0.33 \cdot (2^{2 \cdot GRI} - 1), \quad (3)$$

$$GRI = \frac{GR - GR_{sand}}{GR_{clay} - GR_{sand}}, \quad (4)$$

with V_{cl} = percentage of the volume of clay, $GR_{sand} = 135$ API and $GR_{clay} = 10$ API; GR in Eq. (4) is the actual reading from the GR dataset.

3.4 Si/Ti, TOC data

Silicia/titanium (Si/Ti) ratios were determined on core halves using an X-ray fluorescence (XRF) core scanner (ITRAX, Cox Ltd., Sweden). Details on the scanner settings and processing of the data are given in Wennrich et al. (2013). Total organic carbon (TOC) content was calculated as the difference between total carbon and total inorganic carbon using a DIMATOC 200 carbon analyzer (Dimatec Corp.) in aqueous suspension.

3.5 Statistical analyses

Statistical analyses were carried out using Matlab[®] and the implemented statistical toolbox (Mathworks Inc., Version 7.14.0.739). In a first step, downhole logging data (magnetic susceptibility, electrical resistivity, U counts, Th counts, K counts) were clustered into 3 groups (Clusters 1 to 3) using k-mean clustering to allow for a first characterization of the entire record. Data < 143 m b.l.f. were omitted due to the disturbed magnetic susceptibility and electrical resistivity signal. In a second step, Si/Ti ratios, magnetic susceptibilities and TOC percentages measured on the composite core down

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



confirmed by the fact that only small mass movement deposits, mainly turbidites, were found in pilot cores PG1351 (~ 13 m length) (Melles et al., 2007) and Lz1024 (~ 16 m length) (Juschus et al., 2009). The turbidites were associated with distant debris flows in a conceptual model by Juschus et al. (2009), and their interdependence is confirmed by findings in the drill cores by Sauerbrey et al. (2013).

The wide shelf at the southeastern part of the lake is characterized by aggrading sequences; seismic data from the western and northwestern shelf are not available due to coarse sediments limiting acoustic penetration in these areas. Subunit Ia forms onlaps against the steep slope at the lake margins in a layer-cake manner (Fig. 2), gradually muting a formerly deeper and steeper relief (Niessen et al., 2007). Subunit Ia conformably overlies Subunit Ib with a clear and distinct boundary in between. Subunit Ib has a massive, acoustically chaotic character and rarely shows internal layering in those parts that are visible in our seismic profiles. Its upper boundary has a hummocky surface probably due to thick, chaotic mass movements in its uppermost parts (Fig. 2). Its lower boundary to Unit II lies below the acoustic multiples and is therefore masked. However, refraction data showed that Subunit Ib drapes the central uplift structure that was observed in the seismic refraction data in Units II and III and that is characteristic for impact craters of this size (Gebhardt et al., 2006).

Faults with a vertical offset of up to several meters were observed in the central part of the northern profiles in Unit I. These show a decreasing offset towards the more recent sediment and are inactive in the upper meters of the lake sediments (Fig. 2); this was also observed in high-resolution subbottom profiles (Niessen et al., 2007). The faults are likely related to the central uplift structure.

4.2 Physical properties from downhole and core measurements

Subunit Ia comprises the uppermost ~ 167 m of the sediment column (Fig. 3), which corresponds to approximately 3.17 Ma (Nowaczyk et al., 2013). This also includes the Pliocene/Pleistocene transition at 123 m.b.l.f. (2.6 Ma). Downhole logging data show that the Pleistocene sediments are characterized by relatively constant K and Th

**Petrophysical
characterization of
the lacustrine
sediment succession**

A. C. Gebhardt et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

counts down to approximately 100 m b.l.f. (2.1 Ma, Nowaczyk et al., 2013); magnetic susceptibilities of the sediment core are highly variable, but fluctuate in a range between ~ 15 and $\sim 200 \times 10^{-4}$ SI (Fig. 3). Similarly to magnetic susceptibility, density is highly variable throughout the entire record, but scatters constantly around a mean value of approximately 1.5 g cm^{-3} in the sediments of Subunit Ia (< 3.17 Ma). Lithologies and associated sedimentary facies are characterized by a rapid change between homogeneous (Facies B) and laminated (Facies A) layers that represent warm and cold phases, respectively (Melles et al., 2007), as well as occasional laminated sediments reflecting peak warm conditions (Facies C). These hemipelagic sediments are intercalated by a large number of mass movement deposits of different types such as debris flows and turbidites (Sauerbrey et al., 2013) that get thicker towards the lower boundary of Subunit Ia (Fig. 3). Below 100 m b.l.f., downhole logging K and Th counts show an increase with increasing depth with the highest values exactly at the Pliocene/Pleistocene boundary and strongly decreasing values in the uppermost part of the Pliocene sediments. Magnetic susceptibility values of the Pliocene part of Subunit Ia show a slight increase in amplitude in comparison to the Pleistocene data.

Subunit Ib comprises all lacustrine sediments between 167 m b.l.f. and the boundary to the underlying bedrock at ~ 320 m b.l.f. While magnetic susceptibility values of the Pleistocene part of Subunit Ia originate from sediment core measurements (MS, light blue in Fig. 3), the values of the Pliocene part were measured in the borehole (BMS, dark blue). The two datasets are not completely comparable in terms of their exact values and amplitudes; at first glance, it seems as if the lower part has much higher amplitudes, this however might be an artefact caused by the different measurement methods. Magnetic susceptibility seems to be more variable in longterm trends in the Pliocene part of the sediments; however, it is unclear if this is a real paleoclimate signal or just a scaling effect. Unfortunately, there is not enough overlap in between the two datasets to tune them to similar amplitudes. Nevertheless, it is obvious that magnetic susceptibility is much more variable between approximately 150 and 220 m b.l.f. than below (220 m corresponds to 3.38 Ma, Nowaczyk et al., 2013). Electrical resistivity is

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rather constant throughout the entire Pliocene sediment succession with exception of the lowermost approximately 20 m where a small maximum occurs at ~ 300 m b.l.f. Density shows an increase with increasing depth from mean values around 1.5 g cm^{-3} in the uppermost part of Subunit 1b to mean values around 1.8 g cm^{-3} with values as high as $> 2.0 \text{ g cm}^{-3}$ in the lowermost part, i.e. in the transitional zone between lacustrine and impact-related units. Lithological description of Subunit 1b is not as detailed as for Subunit 1a due to lower recovery and, thus, larger drilling-related gaps (for details on drillings operations, see Melles et al., 2011). As in the upper part, sediments are alternating between Facies A and B, but the homogeneous facies B is much more dominating in the Pliocene part of the record. Only in the lowermost part, i.e. below ~ 270 m b.l.f. (3.48 Ma, Nowaczyk et al., 2013), laminated sediments are more abundant. As in Subunit 1a, mass movement deposits are intercalated frequently in the hemipelagic sediments.

Uranium values are rather constant throughout the entire record with slightly higher values in the bedrock. Two strong exceptions however are observed in the lacustrine part: (a) between ~ 220 and ~ 244 m b.l.f., U values are slightly enhanced, and (b) a strong double peak is observed between ~ 251 and ~ 262 m b.l.f. The U peaks are confirmed by the independently measured total GR.

Electrical resistivity (deep and shallow), borehole magnetic susceptibility, and natural gamma ray counts of K, U and Th were used for cluster analyses to distinguish between different main units between 143 m and 394 m b.l.f. This includes the boundary between the lacustrine sediments and the brecciated bedrock. Three clusters could be distinguished: (1) Cluster 1 is characterized by high electrical resistivity and enhanced K content values (Fig. 4 upper panel). Magnetic susceptibility is rather variable. (2) Cluster 2 is characterized by low electrical resistivity, variable magnetic susceptibility, and low U and K content. (3) Cluster 3 has low electrical resistivity, high U and intermediate K values (Fig. 4 upper panel). It is clearly different from Cluster 2 in almost all parameters, but coincides with Cluster 2 in terms of low resistivity. Plotting these three clusters against depth (Figs. 3 and 5), it becomes obvious that Cluster 1 clearly describes

the bedrock. Cluster 2 comprises the main part of the lacustrine record. Cluster 3 is part of the lacustrine sediments, but only comprises the section between 254.44 and 259.15 m b.l.f. and between 260.7 and 262.5 m b.l.f. where the strong U double-peak is observed (Fig. 3).

Both pelagic sediments and mass movement deposits in Lake El'gygytyn are part of clusters 2 and 3, which implies that these two sediment types do not differ in their petrophysical characteristics. This confirms that the mass movement deposits mainly consist of reworked lacustrine material (Sauerbrey et al., 2013). Enhanced U values in cluster 3 found in the borehole data could not be measured with the ITRAX XRF core scanner in the according core sections, probably due to the scanner's low capability in measuring U. U is removed from the water column and buried in the sediment during oxic conditions (e.g. Anderson et al., 1989); this would probably point at high bottom water oxygen levels when these layers were accumulated. This, however, is not confirmed by the sediment description that does not differ significantly from above or below these layers. Hence, it is more likely that U-rich rocks were eroded in the lake catchment during these periods and transported to the lake by fluvial/eolian rather than gravitational transport processes.

Natural gamma radiation is often measured and used as an indicator for clay content in sediments, using the fact that K and Th are enriched in different clay minerals. This approach, however, does not work in Lake El'gygytyn sediments where clay K and Th do not correlate with clay or any other grain size (2013) neither with the spectral gamma nor with the total gamma dataset (Eqs. 1–4, respectively). This can best be explained by the lake's location in a small catchment with short transport paths from the source rock to the accumulation site, which prohibits full weathering of all grains. K-bearing feldspar grains that would normally weather into K-bearing clay, so K would be an indicator for clay solely. In the case of a very short distance from source to sink, K-bearing feldspar grains of probably fine sand or silt size would also end up in the sediment along with clay. This would in turn mean that the assumption that sand does not contain K (see Eq. 1, methods chapter) is wrong in our case.

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.3 Physical properties from downhole and core measurements

The most prominent change in the downhole logging data occurs at the boundary between lacustrine sediments and the underlying altered bedrock at ~ 320 m b.l.f. (Figs. 3, 5). While sediments above ~ 313 m b.l.f. are clearly lacustrine with alternating homogeneous and laminated layers, intercalated with frequent mass movement deposits, sediments below ~ 313 m b.l.f. are a mixture between a sedimentary matrix and reworked impact breccia. The boundary in between the lacustrine sediments and the underlying bedrock is thus rather a transitional zone than a sharp boundary. In the upper part of this transitional zone, i.e. ~ 313 and 319.8 m b.l.f. (transitional zone T1 in Fig. 5), lacustrine sediments form the dominant part of the record, while below 319.8 m b.l.f. (T2 and T3) the record contains mainly reworked impact breccia in a sedimentary matrix (Raschke et al., 2012). Therefore, the formal boundary between the lacustrine and the impact part of the drill core was defined at 319.8 m b.l.f., between drill runs 97Q and 98Q. Koeberl et al. (2012) subdivide the part of the transitional zone that lies below this boundary into two subunits, one from 319.8 to 323 (T2 in Fig. 5) and the second from 323 to 331 m b.l.f. (T3) (note that the original depth values from both Raschke et al. (2012) and Koeberl et al. (2012) were shifted downwards by 3 m to match the depth scale used by the lacustrine El'gygytgyn scientific community). Both subunits show similar lithologies with fine sand-sized grains mainly composed of glass fragments, intercalated with impact breccia and impact melt blocks. All three subunits of the transitional zone (T1 above, T2 and T3 below the formal boundary) are shown in light to dark grey tones of facies type E in Fig. 5. The boundary between the matrix-dominated (= lacustrine, T1) and the clast-dominated (= impact-related, T2 and T3) part shows up as a sharp boundary in the electrical resistivity and also in the magnetic susceptibility data. Nevertheless, cluster analyses shows that except of two small bands, the entire transitional zone exhibits characteristics that are more similar to the overlying lacustrine succession. Only below this transitional zone, the sediments are clearly of bedrock-type.

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Petrophysical
characterization of
the lacustrine
sediment succession**

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

explained by the lake's location in a rather small catchment of only 293 km² (including the lake's surface) (Nolan and Brigham-Grette, 2007). This means that during warmer as well as during colder periods the same source rock is eroded, and thus all clastic grains that end up in the lacustrine sediments (probably with exception of some eolian grains; these however are a minor contribution to the sediment, see Francke et al., 2013; Fedorov et al., 2013) originate from the same provenance. Nevertheless, differing erosional processes (i.e. more physically-dominated weathering during colder and more chemically-dominated weathering during warmer periods) in the small hinterland as well as diatom blooms during warmer periods are strong enough to generate highly variable sediment properties (cf. Minyuk et al., 2007) with yet almost similar character in terms of petrophysical characteristics. Magnetic susceptibility, in turn, is highly variable in the lacustrine part, probably reflecting different weathering mechanisms and definitely different modes of palaeohydrological conditions (such as anoxia in the bottom water, see Melles et al., 2007, 2012) along with dilution effects by biogenic material.

In order to detect the variability within the lacustrine succession related to the different modes in paleoenvironmental conditions, we carried out clustering analyses on 5538 data points using similar parameters as in Melles et al. (2007, 2012). With Si/Ti ratio, TOC percentage and magnetic susceptibility from core measurements, we were able to identify four clusters (Tables 1 and 2): "Cluster I" is defined by medium TOC percentages, very low Si/Ti ratios and very low magnetic susceptibility. "Cluster II" shows high TOC percentages along with high Si/Ti ratios and medium magnetic susceptibility. "Cluster III" has low TOC percentages and medium Si/Ti ratios along with high magnetic susceptibility. "Cluster IV" is defined by low TOC percentages and Si/Ti ratios combined with high magnetic susceptibility. Density does not vary much between Clusters I, III and IV, but is considerably lower in Cluster II.

5.2 Description of the lacustrine succession

Melles et al. (2007, 2012) used TOC percentage, Si/Ti content and magnetic susceptibility to identify the oxygenation state of the bottom water and, thus, whether the water column was mixed or stratified, which in turn gives evidence on the duration of an ice cover on the lake. During phases with a perennial ice cover, the water column could not mix, and depletion of oxygen in the bottom water lead to enhanced preservation of organic material, while magnetite underwent dissolution, leading to reduced magnetic susceptibility values. During times with seasonal ice cover, in contrast, mixing of the water column was possible during summer months (as it is today, see Nolan and Brigham-Grette, 2007). Organic carbon was thus consumed in the oxic bottom water, and magnetic minerals were buried without alteration (Melles et al., 2007, 2012). Si/Ti rations can be used to estimate the biogenic vs. clastic input to the lake (Melles et al., 2012; Wennrich et al., 2013). Enhanced Si/Ti values point at high biogenic silica contents, which in the case of Lake El'gygytgyn are produced by enhanced primary productivity (here: mainly diatoms) during warmer times with only seasonal ice cover. Low Si/Ti values point at colder periods with perennial ice cover, thus limitation in light penetration necessary for photosynthesis, along with probably enhanced clastic input through the 50 small ephemeral inlets around the lake (Melles et al., 2007, 2012). During times with a perennial ice cover, clastic input is triggered by seasonal moats and vertical conduits in the ice (as is the case today when snow melt starts in late spring, see Nolan et al., 2003; Asikainen et al., 2007; Francke et al., 2013).

Using this information, we can plot the clusters in a redox-condition vs. input-type diagram (Fig. 6). In such a diagram, the different modes of paleoenvironmental conditions known from earlier studies by Melles et al. (2007, 2012) can be visualized as shown in Figs. 6e, f: the glacial modes of Facies A with perennial ice cover and a stratified water column would plot into in the upper left corner (anoxic conditions, dominated by clastic input or by a relative dominance of clastic material due to the lack of biogenic input); the interglacial mode of Facies B with seasonal ice cover and a mixed water column

CPD

9, 351–391, 2013

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



would show up in the right middle part (oxic conditions with variable, but intermediate contents of clastic and biogenic input), and Facies C – the super interglacial mode – would be found in the lower middle with with variably suboxic and oxic conditions and a dominance in biogenic input.

5 When plotting Clusters I to IV into this diagram (Fig. 6), it becomes obvious that sediments of Facies B, i.e. the interglacial sediments, plot into several clusters (Fig. 6a to d): a high portion of Facies B sediments are found in Cluster IV (62.25 % of all Facies B data points), and another 29.59 % plot in Cluster III. In Cluster I and II, some minor 5.27 % and 2.90 % of Facies B sediments are found. This supports the earlier study by
10 Melles et al. (2012) that describes Facies B sediments as highly variable.

Facies F, i.e. the mass movement deposits, also plot into all clusters with the majority in Cluster IV (80.39 %). Almost equal percentages of 10.98 % and 8.24 % plot into Clusters III and II, and negligible 0.39 % is found in Cluster I. As Facies F is not part of the hemipelagic sediments in Lake El'gygytyn, it was omitted in the pie plots in Fig. 6
15 for better visualization of the distribution of Facies types A to D in the different clusters. The fact that both Facies F and Facies B only have minor parts plotting into Clusters I and II, along with the fact that these clusters only represent 13.01 % and 4.51 % of all data points, suggests that these two clusters might represent sediment endmembers of Lake El'gygytyn.

20 "Glacial" Cluster I: Cluster I (687 data points = 13.01 % of entire dataset, Fig. 6a) plots into a field where sediments of Facies A and some of Facies B would be assumed. It contains equal amounts (30.86 % and 30.13 %) of both "cold & dry" Facies A_d and "cold & moist" Facies A_m . Another 30.42 % of this cluster comprises sediments that were classified as Facies B, i.e. sediments interpreted as accumulated during interglacials, and some 5.82 % were even described as from super interglacials (Facies C). This means that sediments of Facies A show similar characteristics as a certain
25 portion of Facies B sediments, so they could not be statistically separated by means of cluster analysis. Nevertheless, the portion of Facies B data that plot into Cluster I is only 5.27 % of all Facies B data (Fig. 6a) and might even be negligible. In fact, samples used

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for this study are generally 2 cm thick, and we chose the facies type of their average composite depth as representative for the entire 2 cm, neglecting that facies boundaries might occur also within samples. Plotting facies types and clusters versus depth (Fig. 7) reveals that Cluster I quite well captures the cold phases (marked with light and dark blue bars) for example at ~ 52.5 to 52.9 m b.l.f., at ~ 53.4 to 53.5 m b.l.f. and at 63.9 to 64.5 m b.l.f.

“Interglacial” Clusters III and IV: Cluster III (1444 data points = 27.34 % of entire dataset, Fig. 6c) as well as Cluster IV (2912 data points = 55.15 % of the entire dataset, Fig. 6b) have very low TOC contents and high magnetic susceptibility values, pointing at oxic bottom water conditions during their deposition. While in Cluster IV Si/Ti ratios are low, slightly higher Si/Ti ratios in Cluster III suggest some biogenic input into the sediment. A high portion of Cluster IV is composed of Facies B sediment (84.82 %), and equal parts of 6.25 and 6.11 % consist of Facies A_d and C, respectively. In Cluster IV, which is the largest cluster and contains more than half of all data points, also Facies B is clearly dominant with 81.30 %, and 12.74 % are made up of Facies C type sediments along with some 4.02 % of Facies A_d. Both clusters contain the majority of all Facies B data points and confirm that this facies type is rather variable yet similar in being deposited under oxic conditions. The Facies A_d sediments found in these two clusters, however, suggest that even during glacial times, oxic (or at least suboxic) conditions in the bottom water were sometimes encountered at least during periods with “cold & dry” conditions, and some bioproduction leading to enhanced Si/Ti ratios was possible. This is in good agreement with findings by Melles et al. (2007; 2012) who suggested that “cold & dry” Facies A_d represents a perennial ice cover without snow cover. This would allow some light penetration and thus some primary productivity in the water column. In contrast, “cold & moist” Facies A_m was interpreted as representing a perennial ice cover covered by snow, inhibiting any light penetration into the water column, leading to only very limited photosynthetic life in the lake and thus low TOC values and Si/Ti ratios. This, in turn, is confirmed by only negligible 0.58 and 0.07 % of Facies A_m in Clusters III and IV, and 0.84 % in Cluster II.

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



“Super interglacial” Cluster II (238 data points = 4.51 % of the entire dataset; Fig. 6g): cluster II has significantly enhanced TOC and Si/Ti ratios and consists to almost equal parts of Facies C and B sediments (47.48 % and 48.32 %). The negligible remainder is 3.36 % Facies D and 0.84 % Facies A_m.

5 While density is rather variable in Clusters I, III and IV, it is clearly lower than average in Cluster II, which is in good agreement with a high content of biogenic silica. Even though approximately half of Cluster II consists of Facies C type sediments, only approximately one fifth of all Facies type sediments plot into this cluster (21.94 %), while some 35.73 and 43.56 % plot into Clusters III and IV. This might either point at a wider
10 range of TOC percentages, Si/Ti ratio and magnetic susceptibility values within this facies type, or these samples are highly biased by facies changes within the distinct samples that led to a wrong assignment of facies type to a specific sample. When plotting facies and clusters together vs. depth (Fig. 7) it becomes obvious that only some parts of Facies C (red bars) were captured by Cluster II: between ~62 and ~62.7 m b.l.f.,
15 Facies C sediments were visually described, but have rather low Si/Ti content and only slightly enhanced TOC values, so they were statistically gathered into Clusters I and IV; these samples are part of the 5.82 % of Facies C samples that were found in Cluster I and 6.11 % in Cluster IV. On the other hand, sediments of the thick Facies C layer between ~64.7 and ~65.6 m b.l.f. show higher Si/Ti ratios and TOC content
20 and were therefore gathered into Clusters III and II. This would imply that even though Facies C is easily detected by means of visual core description, its basic physical and geochemical properties might not always be significantly different from sediments of other facies types, notably from Facies B. Nevertheless, this is in good agreement with findings by Melles et al. (2012) who report that while primary productivity was highest
25 during these extraordinary phases, there are laminae found in the Facies C sediments that suggest at least seasonally suboxic or anoxic conditions in the bottom waters. This could result in a wide variety of TOC percentages and magnetic susceptibility values in the according sediment and make it difficult to gather these sediments in one single cluster.

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



6 Conclusions

Seismic reflection profiles of Lake El'gygytyn exhibit mostly well-stratified sediments with frequent mass movement deposits intercalated in the more proximal areas. The well-stratified acoustic layers are nicely confirmed by the well-layered sediments of the drill core retrieved during winter/spring 2009 with highly variable facies types changing at high frequency in the core. The lacustrine sediment succession can be separated into two seismic subunits Ia and Ib. While Ia is well-stratified, Ib is acoustically more chaotic. The sediment-bedrock boundary was identified earlier by Gebhardt et al. (2006) at around 320 to 330 m b.l.f. by means of a seismic-refraction data derived depth-velocity model. This was confirmed during drilling with the first bedrock material found at approximately 320 m b.l.f. Downhole logging data down to 394 m b.l.f., i.e. through the entire lacustrine column and some 74 m into the bedrock, show that the lacustrine and bedrock part differ clearly in their petrophysical characteristics: Cluster analysis separates three clusters, two of which comprise the entire lacustrine succession, while the third contains the bedrock. The boundary between the impact-related bedrock and the lacustrine succession is not sharp, but rather a transitional zone with an upwards increasing portion of lacustrine material. Potassium and resistivity values are enhanced in the bedrock part.

In the lacustrine succession, a prominent U peak of unknown origin is visible at around 255 m b.l.f., and slightly enhanced Th and K values mark the Pliocene/Pleistocene transition. The core could be clustered into four different clusters (I to IV) down to approximately 262 m composite depth. The clusters show significant differences in terms of their TOC percentage, Si/Ti ratio and magnetic susceptibility, in some cases also density. This allows plotting the clusters into a redox-condition vs. input-type diagram. In comparison with earlier studies we could conclude that Cluster I contains glacial sediments, III and IV sediments from interglacials, and II comprises the sediments from super interglacials.

CPD

9, 351–391, 2013

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. We thank all expedition members of the El'gygytgyn pre-site surveys in 2000 and 2003 as well as of the deep drilling campaign in winter/spring 2009 for their excellent cooperation and support during work at the lake. Funding for this research was provided by the International Continental Scientific Drilling Program (ICDP), the US National Science Foundation (NSF), the German Federal Ministry of Education and Research (BMBF), Alfred Wegener Institute (AWI) and GeoForschungsZentrum Potsdam (GFZ), the Russian Academy of Sciences Far East Branch (RAS FEB), the Russian Foundation for Basic Research (RFBR), and the Austrian Federal Ministry of Science and Research (BMWF). The Russian GLAD 800 drilling system was developed and operated by DOSECC Inc., the down hole logging was performed by the ICDP-OSG, and LacCore, at the University of Minnesota, handled core curation. Financial support for the laboratory analyses was provided by the German Ministry of Education and Research (BMBF grants no. 03G0586B and no. 03G0642B) and the Deutsche Forschungsgemeinschaft (DFG grant no. GE-1924/1-1).

References

- Anderson, R. F., Fleisher, M. O., and LeHuray, A. P.: Concentration, oxidation state, and particulate flux of uranium in the Black Sea, *Geochim. Cosmochim. Ac.*, 53, 2215–2224, 1989.
- Asikainen, C. A., Francus, P., and Brigham-Grette, J.: Sedimentology, clay mineralogy and grain-size as indicators of 65 ka of climate change from El'gygytgyn Crater Lake, North-eastern Siberia, *J. Paleolimnol.*, 37, 105–122, doi:10.1007/s10933-006-9026-5, 2007.
- Belyi, V. F.: Impactogenesis and volcanism of the El'gygytgyn depression, *Petrology*, 6, 96–110, 1998.
- Best, A. I. and Gunn, D. E.: Calibration of marine sediment core loggers for quantitative acoustic impedance studies, *Mar. Geol.*, 160, 137–146, 1999.
- Brigham-Grette, J., Melles, M., Minyuk, P. S., Andreev, A. A., Tarasov, P., DeConto, R. M., König, S., Nowaczyk, N. R., Wennrich, V., Rosén, P., Haltia-Hovi, E., Cook, T. L., Gebhardt, A. C., Meyer-Jacob, C., Snyder, J., and Herzschuh, U.: Pliocene warmth, extreme polar amplification, and stepped Pleistocene cooling recorded in NE Russia, *Science*, submitted, 2013.
- Dansgaard, W., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N. S., Hammer, C. U., Hvidberg, C. S., Steffensen, J. P., Sveinbjørnsdóttir, A. E., Jouzel, J., and

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bond, G. C.: Evidence for general instability of past climate from a 250-kyr ice-core record, *Nature*, 364, 218–220, 1993.

Eyles, N., Doughty, M., Boyce, J. I., Mullins, H. T., Halfman, J. D., and Koseoglu, B.: Acoustic architecture of glaciolacustrine sediments deformed during zonal stagnation of the Laurentide Ice Sheet; Mazinaw Lake, Ontario, Canada, *Sediment. Geol.*, 157, 133–151, doi:10.1016/S0037-0738(02)00229-4, 2003.

Fedorov, G., Nolan, M., Brigham-Grette, J., Bolshiyarov, D., Schwamborn, G., and Juschus, O.: Lake El'gygytgyn water and sediment balance components overview and its implications for the sedimentary record, *Clim. Past Discuss.*, 8, 3977–4001, doi:10.5194/cpd-8-3977-2012, 2012.

Finckh, P., Kelts, K. R., and Lambert, A.: Seismic stratigraphy and bedrock forms in perialpine lakes, *Geol. Soc. Am. Bull.*, 95, 1118–1128, 1984.

Francke, A., Wennrich, V., Sauerbrey, M., Juschus, O., Melles, M., Brigham-Grette, J., and El'gygytgyn Scientific Party: Quaternary grain-size distributions of Lake El'gygytgyn, NE Siberia: Statistic analysis and climate dependency, *Clim. Past*, in preparation, 2013.

Frank, U., Nowaczyk, N. R., Minyuk, P., Vogel, H., Rosén, P., and Melles, M.: A 350 kyr record of climate change from Lake El'gygytgyn, Far East Russian Arctic: refining the pattern of climate modes by means of cluster analysis, *Clim. Past Discuss.*, 8, 5109–5132, doi:10.5194/cpd-8-5109-2012, 2012.

Fronval, T. and Jansen, E.: Late Neogene paleoclimates and paleoceanography in the Iceland-Norwegian Sea. Evidence from the Iceland and Vøring Plateaus, in: *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by: Thiede, J., Myhre, A. M., Firth, J. V., Johnson, G. L., and Ruddiman, W. F., College Station, Texas, USA, 455–468, 1996.

Gebhardt, A. C., Niessen, F., Melles, M., and Juschus, O.: Shallow seismic survey, in: *The Expedition – El'gygytgyn Lake 2003 (Siberian Arctic)*, edited by: Melles, M., Minyuk, P., Brigham-Grette, J., and Juschus, O., Reports on Polar and Marine Research, Alfred-Wegener-Institut für Polar- und Meeresforschung, Bremerhaven, 116–117, 2005.

Gebhardt, A. C., Niessen, F., and Kopsch, C.: Central ring structure identified in one of the world's best preserved impact craters, *Geology*, 34, 145–148, doi:10.1130/G22278.1, 2006.

Geotek: Multi-Sensor Core Logger, Geotek, available at: <http://www.geotek.co.uk/>, 127 pp., 2000.

**Petrophysical
characterization of
the lacustrine
sediment succession**

A. C. Gebhardt et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Glushkova, O. Y. and Smirnov, V. N.: Pliocene to Holocene Geomorphic Evolution and Paleogeography of the El'gygytgn Lake region, NE Russia, *Journal of Paleolimnology Special Issue*, 37, 37–47, doi:10.1007/s10933-006-9021-x, 2007.
- Grootes, P. M., Stuiver, M., White, J. W. C., Johnsen, S. J., and Jouzel, J.: Comparison of oxygen isotope records from the GISP2 and GRIP Greenland ice cores, *Nature*, 366, 552–554, 1993.
- Gurov, E. P., Gurova, E. P., and Rakitskaya, R. B.: Stishovite and coesite in shock metamorphosed rocks of the El'gygytgn crater in Chukotka (in Russian), *Doklady Akademii Nauk USSR*, 248, 213–216, 1979a.
- Gurov, E. P., Valter, A. A., Gurova, E. P., and Kotlovskaya, F. I.: El'gygytgn impact crater, Chukotka: shock metamorphism of volcanic rocks (abs.), *Lunar and Planetary Science*, 10, 479–481, 1979b.
- Harris, S. A.: Thermal history of the Arctic Ocean environs adjacent to North America during the last 3.5 Ma and a possible mechanism for the cause of the cold events (major glaciations and permafrost events), *Prog. Phys. Geog.*, 29, 1–19, 2005.
- Herman, Y. and Hopkins, D. M.: Arctic Ocean climate in late Cenozoic time, *Science*, 209, 557–562, 1980.
- Hogan, K. A., Dowdeswelle, J. A., and Ó Cofaigh, C.: Glacimarine sedimentary processes and depositional environments in an embayment fed by West Greenland ice streams, *Mar. Geol.*, 311–314, 1–16, doi:10.1016/j.margeo.2012.04.006, 2012.
- Jansen, E., Fronval, T., Rack, F., and Channel, J. E. T.: Pliocene-Pleistocene ice rafting history and cyclicity in the Nordic Seas during the last 3.5 Myr, *Paleoceanography*, 15, 709–721, 2000.
- Juschus, O., Melles, M., Gebhardt, A. C., and Niessen, F.: Late Quaternary mass movement events in Lake El'gygytgn, North-Eastern Siberia, *Sedimentology*, 56, 2155–2174, doi:10.1111/j.1365-3091.2009.01074.x, 2009.
- Koerberl, C., Pitarello, L., Reimold, U., Raschke, U., Brigham-Grette, J., Melles, M., and Minyuk, P.: El'gygytgn impact crater, Chukotka, Arctic Russia: impact cratering aspects of the 2009 ICDP drilling project, *Meteorit. Planet. Sci.*, submitted, 2012.
- Layer, P.: ⁴⁰Ar/³⁹Ar age of the El'gygytgn crater event, Chukotka, Russia, *Meteorit. Planet. Sci.*, 35, 591–599, 2000.
- Lerman, A., Imboden, D., and Gat, J.: *Physics and Chemistry of Lakes*, 2nd Edn., Springer, Berlin, Heidelberg, 334 pp., 1995.

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Melles, M., Brigham-Grette, J., Glushkova, O. Y., Minyuk, P. S., Nowaczyk, N. R., and Hubberten, H.-W.: Sedimentary geochemistry of a pilot core from Lake El'gygytyn – a sensitive record of climate variability in the East Siberian Arctic during the past three climate cycles, *J. Paleolimnol. Special Issue*, 37, 89–104, 10.1007/s10933-006-9025-6, 2007.
- 5 Melles, M., Brigham-Grette, J., Minyuk, P. S., Koeberl, C., Andreev, A. A., Cook, T. L., Fedorov, G., Gebhardt, A. C., Haltia-Hovi, E., Kukkonen, M., Nowaczyk, N. R., Schwamborn, G., Wennrich, V., and the El'gygytyn Scientific Party: The Lake El'gygytyn scientific drilling project – conquering Arctic challenges through continental drilling, *Sci. Drill.*, 11, 29–40, doi:10.2204/iodp.sd.11.03.2011, 2011.
- 10 Melles, M., Brigham-Grette, J., Minyuk, P. S., Nowaczyk, N. R., Wennrich, V., DeConto, R. M., Anderson, P. M., Andreev, A. A., Coletti, A., Cook, T. L., Haltia-Hovi, E., Kukkonen, M., Lozhkin, A. V., Rosén, P., Tarasov, P., Vogel, H., and Wagner, B.: 2.8 million years of Arctic climate change from Lake El'gygytyn, NE Russia, *Science*, 337, 315–320, doi:10.1126/science.1222135, 2012.
- 15 Minyuk, P., Brigham-Grette, J., Melles, M., Borkhodoev, V. Y., and Glushkova, O. Y.: Inorganic geochemistry of El'gygytyn Lake sediments (northeastern Russia) as an indicator of paleoclimatic change for the last 250 kyr, *J. Paleolimnol., Special Issue*, 37, 123–133, doi:10.1007/s10933-006-9027-4, 2007.
- 20 Moran, K., Backman, J., Brinkhuis, H., Clemens, S. C., Cronin, T., Dickens, G. R., Eynaud, F., Gattaceca, J., Jakobsson, M., Jordan, R. W., Kaminski, M., King, J. W., Koc, N., Krylov, A., Martinez, N., Matthiessen, J., McInroy, D., Moore, T. C., Onodera, J., O'Regan, M., Pälike, H., Rea, B., Rio, D., Sakamoto, T., Smith, D. C., Stein, R., St John, K., Suto, I., Suzuki, N., Takahashi, K., Watanabe, M., Yamamoto, M., Farrell, J., Frank, M., Kubik, P., Jokat, W., and Kristoffersen, Y.: The Cenozoic palaeoenvironment of the Arctic Ocean, *Nature*, 441, 601–605, doi:10.1038/nature04800, 2006.
- 25 Myhre, A. M., Thiede, J., Firth, J. V., Ahagon, N., Black, K. S., Bloemendal, J., Brass, G. W., Bristow, J. F., Chow, N., Cremer, M., Davis, L., Flower, B., Fronval, T., Hood, J., Hull, D., Koç, N., Larsen, B., Lyle, M., McManus, J., O'Connell, S., Ostermann, L. E., Rack, F. R., Sato, T., Scherer, R., Spiegler, D., Stein, R., Tadross, M., Wells, S., Williamson, D., Witte, B., and Wolf-Welling, T.: Proceedings of the Ocean Drilling Program, Initial Reports, Leg 151, Ocean Drilling Program, College Station, Texas, 926 pp., 1995.
- 30 NGRIP members: High-resolution record of Northern Hemisphere climate extending into the last interglacial period, *Nature*, 431, 147–151, 2004.

**Petrophysical
characterization of
the lacustrine
sediment succession**

A. C. Gebhardt et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Niessen, F., Gebhardt, A. C., Kopsch, C., and Wagner, B.: Seismic investigation of the El'gygytgyn impact crater lake (Central Chukotka, NE Siberia): preliminary results, *J. Paleolimnol.*, Special Issue, 37, 49–63, doi:10.1007/s10933-006-9022-9, 2007.
- Nolan, M. and Brigham-Grette, J.: Basic hydrology, limnology, and meteorology of modern Lake El'gygytgyn, Siberia, *J. Paleolimnol.*, Special Issue, 37, 17–35, doi:10.1007/s10933-006-9020-y, 2007.
- Nowaczyk, N. R., Minyuk, P., Melles, M., Brigham-Grette, J., Glushkova, O., Nolan, M., Lozhkin, A. V., Stetsenko, T. V., Andersen, P. M., and Forman, S. L.: Magnetostratigraphic results from impact crater Lake El'gygytgyn, Northeastern Siberia: a 300 kyr long high-resolution terrestrial palaeoclimatic record from the Arctic, *Geophys. J. Int.*, 150, 109–126, 2002.
- Nowaczyk, N. R., Melles, M., and Minyuk, P.: A revised age model for core PG1351 from Lake El'gygytgyn, Chukotka, based on magnetic susceptibility variations correlated to northern hemisphere insolation variations, *J. Paleolimnol.*, Special Issue, 37, 65–76, doi:10.1007/s10933-006-9023-8, 2007.
- Nowaczyk, N. R., Haltia-Hovi, E. M., Ulbricht, D., Wennrich, V., Kukkonen, M., Rosén, P., Vogel, H., Francke, A., Meyer-Jacob, C., Andreev, A., and Lozhkin, A. V.: Chronology of Lake El'gygytgyn sediments, *Clim. Past*, in preparation, 2013.
- Raschke, U., Reimold, W., Zaag, P., Pitarello, L., and Koeberl, C.: Lithostratigraphy of the impactite and bedrock section in ICDP drill core D1c from the El'gygytgyn impact crater, Russia, *Meteorit. Planet. Sci.*, submitted, 2012.
- Repenning, C. A. and Brouwers, E. M.: Mid-Pliocene to Late Pleistocene Changes in the Arctic Ocean Borderland Ecosystem, Program with Abstracts, INQUA, Ottawa, 251 pp., 1987.
- Ruffell, A., and Worden, R.: Palaeoclimate analysis using spectral gamma-ray data from the Aptian (Cretaceous) of southern England and southern France, *Palaeogeogr. Palaeocl.*, 155, 265–283, 2000.
- Sauerbrey, M. A., Juschus, O., Gebhardt, A. C., Wennrich, V., Nowaczyk, N. R., and Melles, M.: Mass movement deposits in the 3.6 Ma sediment record of Lake El'gygytgyn, Chukotka, NE Russia: classification, distribution and preliminary interpretation, *Clim. Past*, submitted, 2013.
- Schnyder, J., Ruffell, A., Deconinck, J.-F., and Baudin, F.: Conjunctive use of spectral gamma-ray logs and clay mineralogy in defining late Jurassic-early Cretaceous palaeoclimate change

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Dorset, U. K.), *Palaeogeogr. Palaeoclimatol.*, 229, 303–320, doi:10.1016/j.palaeo.2005.06.027, 2006.

Svensson, A., Bigler, M., Kettner, E., Dahl-Jensen, D., Johnsen, S., Kipfstuhl, S., Nielsen, M., and Steffensen, J. P.: Annual layering in the NGRIP ice core during the Eemian, *Clim. Past*, 7, 1427–1437, doi:10.5194/cp-7-1427-2011, 2011.

Van Rensbergen, P., De Batist, M., Beck, C., and Manalt, F.: High-resolution seismic stratigraphy of late Quaternary fill of Lake Annecy (northwestern Alps): evolution from glacial to interglacial sedimentary processes, *Sediment. Geol.*, 117, 71–96, 1998.

Van Rensbergen, P., De Batist, M., Beck, C., and Chapron, E.: High-resolution seismic stratigraphy of glacial to interglacial fill of a deep glacial lake: Lake Le Bourget, Northwestern Alps, France, *Sediment. Geol.*, 128, 99–129, 1999.

Wennrich, V., Minyuk, P. S., Borkhodoev, V., Francke, A., Ritter, B., Raschke, U., Nowaczyk, N. R., Schwamborn, G., Brigham-Grette, J., Melles, M., and El'gygytgyn Scientific Party: Pliocene to Pleistocene climate and environmental history of Lake El'gygytgyn/NE Russia based on high-resolution inorganic geochemistry data, *Clim. Past*, in preparation, 2013.

Wonik, T.: Gamma-ray measurements in the Kirchrode I and II boreholes, *Palaeogeogr. Palaeoclimatol.*, 174, 97–105, 2001.

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Table 1. Amount of data points of the 6 facies types in the 4 clusters (mass movement deposit sediments and tephra were omitted in this table).

	Facies A _d	Facies A _m	Facies B	Facies C	Facies D	total
Cluster I	212	207	209	40	19	708
Cluster II	0	2	115	113	8	238
Cluster III	58	1	1174	184	27	1444
Cluster IV	182	17	2470	178	65	2912
Total	452	227	3968	515	119	4769

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Table 2. Mean and standard deviation (std dev) of total organic carbon (TOC), density, Si/Ti ratio and magnetic susceptibility (Magn. susc.) of clusters I to IV.

	TOC %		Density (g cm^{-3})		Si/Ti ratio		Magn. susc.	
	mean	std dev	mean	std dev	mean	std dev	mean	std dev
Cluster I	0.8354	0.2460	1.4791	0.1086	0.4927	0.1404	50.1806	55.4048
Cluster II	1.0216	0.3369	1.2735	0.1274	1.2826	0.3100	75.0329	41.8939
Cluster III	0.3215	0.1263	1.4385	0.0949	0.8362	0.1385	122.4036	61.3634
Cluster IV	0.2760	0.1076	1.4900	0.0962	0.5327	0.0934	120.7652	76.9754

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.




Fig. 1. Geographical map of the investigation area. **(a):** location of Lake El'gygytyn in NE Russia, **(b):** aerial photograph of the lake surroundings with lake bathymetry and seismic profiles. The profiles marked in red are shown in Figs. 2 and 3. The orange arrow marks the Enmyvaan River, the only outlet of the lake. Drill site ICDP 5011-1 is shown as an orange circle.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.




Fig. 2. Seismic reflection profiles AWI-20008110 and AWI-20008130. Seismic profiles are shown in (a) and (c); line drawings and interpretations in (b) and (d). Green dashed line marks the intersection of the two lines that was chosen for drill site 5011-1, lilac lines show the faults that are likely related to the central uplift structure of the impact crater (see Gebhardt et al., 2006). Seismic profiles clearly show the boundary between Subunits Ia and Ib. Units II and III are masked by the multiples.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. Profile AWI-20008130 with inset of downhole logging and core physical properties data. Note that the light blue part of the magnetic susceptibility data derive from direct core measurements, while the dark blue part is from downhole logging measurements – i.e. exact values and amplitudes of the two datasets are not directly comparable. Dashed turquoise horizontal lines mark the boundaries of seismic Subunits 1a and 1b; the boundary between Units I and II is taken from Gebhardt et al. (2006). Sedimentary facies comprise mass movement deposits (Facies F; green), massive interglacial sediments (Facies B; ochre), laminated super interglacial sediments (Facies C; red), laminated Quaternary glacial sediments representing “cold & dry” (Facies A_d; light blue) and “cold & moist” climates (Facies A_m; dark blue), laminated Pliocene sediments (Facies D; lilac), and transitional sediments between impact breccia and lake sediments (Facies E; gray). Facies types are used sensu Melles et al. (2007; 2012) and Brigham-Grette (2013). The impact breccia/suevite succession according to Koeberl et al. (2012) is given as follows: yellow: suevite; lilac: upper volcanic rock layer; violet: lower volcanic rock layer, green: ignimbrite. The clusters derived from k-mean clustering of downhole logging data are shown in blue for Cluster 1, red for Cluster 2 and green for Cluster 3. Ages are taken from Nowaczyk et al. (2013).

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.




Fig. 4. Crossplots of clusters vs. geophysical and geochemical parameters. Upper panel: clusters 1 to 3 derived from k-mean clustering of downhole logging data (electrical resistivity (deep and shallow), magnetic susceptibility, U, Th, and K counts). Lower panel: clusters I to IV derived from k-mean clustering of core data (TOC content, Si/Ti ratio, magnetic susceptibility). In the lower panel, mean and standard deviation of parameters are shown for each cluster.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.




Fig. 5. Facies and downhole logging data of the transitional zone between impact-related bedrock and lake sediments. Facies description as in Fig. 3. Clusters derived from k-mean clustering of downhole logging data are shown in blue for Cluster 1 (= bedrock) and red for Cluster 2 (= lacustrine). The impact breccia/suevite succession according to Koeberl et al. (2012) is given as follows: yellow: suevite; lilac: upper volcanic rock layer; violet: lower volcanic rock layer, green: ignimbrite.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.




Fig. 6. Pie plots of core-data derived Clusters I to IV versus facies types known from core description. Facies colors correspond to those shown in Figs. 3 and 4. Clusters are distributed according to their redox conditions and clastic vs. biogenic input ratio. **(e)** shows where different paleoenvironmental conditions would plot in such a redox-condition vs. input-type diagram; **(f)** shows this for the Melles et al. (2007, 2012) and Brigham-Grette (2013) facies types. Percentages in **(a)** to **(d)** are calculated for the facies distribution within each cluster.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏴

⏵

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Petrophysical characterization of the lacustrine sediment succession

A. C. Gebhardt et al.




Fig. 7. Downcore distribution of facies types and clusters for 52 to 56 and 62 to 66 m b.l.f. Facies description as in Fig. 3. Clusters are superimposed on magnetic susceptibility, Si/Ti ratio, density and TOC content and are color-coded as in Fig. 4.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion