

Interactive comment on “Extreme warming, photic zone euxinia and sea level rise during the Paleocene/Eocene Thermal Maximum on the Gulf of Mexico Coastal Plain; connecting marginal marine biotic signals, nutrient cycling and ocean deoxygenation” by A. Sluijs et al.

A. Sluijs et al.

a.sluijs@uu.nl

Received and published: 17 March 2014

Dear Dr. Fluegeman,

Thank you for your evaluation of the biostratigraphy. Your point essentially comes down to the occurrence of *Pseudohastigerina wilcoxensis* in the Bells Landing Marl. As you indicate, Wade et al. (2011), in their review of planktic foraminifer biostratigraphy, indicate that the first occurrence of this taxon as calibrated at the GSSP in Africa is in the

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



earliest Eocene. A simple biostratigraphic implication would be that therefore the Bells Landing Marl must be Eocene, although this is inconsistent with all published biostratigraphic interpretations on the Gulf Coastal Plain (see below). Crucially, Mancini and Oliver (Mancini and Oliver, 1981) note that *P. wilcoxensis* likely evolved from *Planoralites chapmani* during the Late Paleocene, partially based on ‘somewhat atypical’ specimens of *P. wilcoxensis* found in lower Tusahoma marls. If *P. wilcoxensis* evolved in the study region it may not be surprising if its first occurrence was earlier here than in Africa. Indeed, its first occurrence was previously reported to be diachronous with latitude (Pardo et al., 1997).

Below, are sections regarding the age of the upper Tusahoma Fm and Bashi Mb that we will include in the revised version of the manuscript. It shows that 1) based on planktic foraminifer biostratigraphy, it cannot be positively concluded that the Bells Landing Marl is Paleocene, but 2) calcareous nannofossil biostratigraphy is inconsistent with an Eocene age of the Bells Landing Marl and indicates that it represents late Paleocene zone NP9. We also further substantiate that the Bashi represents NP10 and exclude the suggestion by Fluegeman that *Discoaster mohlerii* specimens were reworked into this member. Finally, numerous high-resolution studies have focused on the late Paleocene through early Eocene interval preserved in the deep sea around the world, and none have found evidence for a hyperthermal between the PETM and the ETM2. (e.g., Cramer et al., 2003; Zachos et al., 2010). The recognition of a negative carbon isotope excursion and thus a hyperthermal as described in the manuscript is therefore inconsistent with an age in between the PETM and ETM2 as suggested by Fluegeman.

Sincerely,

Also on behalf of all authors,

Appy Sluijs

The occurrence of the planktic foraminifers *Morozovella subbotinae* and *M. velascoensis* (Mancini and Oliver, 1981) restrict the age of the Bells Landing Marl to the time

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

interval spanning the latest Paleocene (Zone P5 of Wade et al., 2011) to the earliest Eocene (Zone E2). The occurrence of *Pseudohastigerina wilcoxensis* would suggest an earliest Eocene age (Wade et al., 2011). However, diachronicity between the GCP and Africa, where the range of this species was calibrated, cannot be excluded. Mancini and Oliver (1981) note that *P. wilcoxensis* likely evolved from *Planorotalites chapmani* during the Late Paleocene, partially based on the “somewhat atypical” specimens of *P. wilcoxensis* found in lower Tusahoma marls. If *P. wilcoxensis* evolved in the study region it may not be surprising if its first occurrence was earlier here than in Africa. Indeed, its first occurrence was previously reported to be diachronous with latitude (Pardo et al., 1997). Moreover, Gibson et al., (1982) and Siesser (1983) determined that the Bells Landing Formation represents the late Paleocene (calcareous nannofossil zone NP9). This is based on the occurrence of *Discoaster multiratiatus* and the combined absence of *Tribrachiatulus* spp., and *Rhomboaster* spp., which excludes an age younger than NP9. Both *Tribrachiatulus* and *Rhomboaster* are heavily calcified genera that are resistant to dissolution, making it unlikely that preservation accounts for their absence. To our knowledge, there is no report of a nannofossil species in the Bells Landing Marl that would suggest an age younger than late Paleocene (Zone NP9).

Above the Bells Landing Marl, the Tusahoma Fm, particularly within the studied region (Fig. 1), is composed of mudstones with intermittent lignites representing estuarine and lagoonal environments (Ingram, 1991), with little to no carbonate and no calcareous microfossils (Gibson and Bybell, 1994). The uppermost beds of the Tusahoma Fm yield brackish to fresh water palynomorphs, notably pollens and spores of late Paleocene affinities (e.g., Frederiksen et al., 1982; Frederiksen, 1998; Harrington, 2003). Collectively, we agree with the published work that the available information indicates a Paleocene age of the Tusahoma formation (e.g., Gibson and Bybell, 1994; Harrington, 2003; Beard, 2008).

Siesser (1983) determined that the Bashi Marl represents calcareous nannofossil zone

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

NP9, perhaps NP10. The NP9 designation for the Bashi Marl is based on the co-occurrence of *Discoaster multiradiatus* and *D. mohleri*, which is stated to be restricted to the late Paleocene. However, Eocene occurrences of *D. mohleri* are known from Demerara Rise, South Atlantic (Mutterlose et al., 2007), Walvis Ridge, South Atlantic (Monechi and Angori, 2006), and Mexico (Garcia-Cordero and Carreño, 2009). The last occurrence of this species is not well-constrained. Gibson et al. (1982) dated the Bashi Marl as NP10 based on the occurrence of *Tribrachiatulus contortus* and *T. bramlettei*, both of which are restricted to NP10 in the earliest Eocene (Vandenbergh et al., 2012).

References

Beard, K. C.: The oldest North American primate and mammalian biogeography during the Paleocene–Eocene Thermal Maximum, *Proceedings of the National Academy of Sciences of the United States of America*, 105, 3815–3818, 2008. Cramer, B. S., Wright, J. D., Kent, D. V., and Aubry, M.-P.: Orbital climate forcing of $\delta^{13}\text{C}$ excursions in the late Paleocene–early Eocene (chrons C24n–C25n), *Paleoceanography*, 18, 1097, 2003. Frederiksen, N. O., Gibson, T. G., and Bybell, L. M.: Paleocene-Eocene boundary in the eastern Gulf Coast, *Gulf Coast Association of Geological Societies Transactions*, 32, 289–294, 1982. Frederiksen, N. O.: Upper Paleocene and lowermost Eocene angiosperm pollen biostratigraphy of the Eastern Gulf Coast and Virginia, *Micropaleontology* 44, 45–68, 1998. Garcia-Cordero, E., and Carreño, A. L.: Upper lower Eocene calcareous nannoplankton from the Las Pocitas core (Tepetate formation), Baja California Sur, Mexico, *Revista Mexicana de Ciencias Geológicas*, 26, 37–47, 2009. Gibson, T. G., and Bybell, L. M.: Sedimentary Patterns across the Paleocene-Eocene boundary in the Atlantic and Gulf coastal plains of the United States, *Bulletin de la Société Belge de Géologie*, 103, 237–265, 1994. Harrington, G. J.: Wasatchian (Early Eocene) pollen floras from the Red Hot Truck Stop, Mississippi, USA, *Palaeontology*, 46, 725–738, 2003. Ingram, S. L.: The Tusahoma-Bashi section at Meridian, Mississippi: First notice of lowstand deposits above the Paleocene-Eocene TP2/TE1 sequence bound-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

ary, *Mississippi Geology*, 11, 9-14, 1991. Mancini, E. A., and Oliver, G. E.: Planktic foraminifers from the Tuscaloosa Sand (upper Paleocene) of southwest Alabama, *Micropaleontology*, 27, 204-225, 1981. Monechi, S., and Angori, E.: Calcareous nanofossil biostratigraphy of the Paleocene/Eocene Boundary, Ocean Drilling Program Leg 208 Hole 1266C, in: *Proceedings of the Ocean Drilling Program, Scientific Results 208*, edited by: Kroon, D., Zachos, J. C., and Richter, C., Ocean Drilling Program, College Station, TX, 1-9, 2006. Mutterlose, J., Linnert, C., and Norris, R.: Calcareous nanofossils from the Paleocene-Eocene Thermal Maximum of the equatorial Atlantic (ODP Site 1260B): Evidence for tropical warming, *Marine Micropaleontology*, 65, 13-31, 2007. Pardo, A., Keller, G., Molina, E., and Canudo, J.: Planktic foraminiferal turnover across the Paleocene-Eocene transition at DSDP Site 401, Bay of Biscay, North Atlantic, *Marine Micropaleontology*, 29, 129-158, 1997. Siesser, W. G.: Paleogene calcareous nannoplankton biostratigraphy: Mississippi, Alabama and Tennessee, *Mississippi Department of Natural Resources Bureau of Geology Bulletin*, 125, 1-61, 1983. Vandenberghe, N., Speijer, R. P., and Hilgen, F. J.: The Paleogene Period, in: *The Geologic Time Scale 2012*, edited by: Gradstein, F. M., Ogg, J. G., Schmitz, M., and Ogg, G., Elsevier, 855-921, 2012. Wade, B. S., Pearson, P. N., Berggren, W. A., and Pälike, H.: Review and revision of Cenozoic tropical planktonic foraminiferal biostratigraphy and calibration to the geomagnetic polarity and astronomical time scale, *Earth-Science Reviews*, 104, 111-142, 2011. Zachos, J. C., McCarren, H., Murphy, B., Röhl, U., and Westerhold, T.: Tempo and scale of late Paleocene and early Eocene carbon isotope cycles: Implications for the origin of hyperthermals, *Earth and Planetary Science Letters*, 299, 242-249, 2010.

Interactive comment on *Clim. Past Discuss.*, 9, 6459, 2013.

CPD

9, C3651–C3655, 2014

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C3655

