

Interactive comment on “Bering Sea surface water conditions during Marine Isotope Stages 12 to 10 at Navarin Canyon (IODP Site U1345)” by Beth E. Caissie et al.

T. Cronin (Referee)

tcronin@usgs.gov

Received and published: 11 February 2016

This paper reconstructs paleoceanography for an important glacial-interglacial cycle, MIS 12-11 in an important region in the Bering Sea using IODP cores. MIS 11 is an especially important interglacial due to pre industrial-level CO₂ concentrations but higher-than-present sea level and in many regions, significantly warmer air and sea temperatures. The study uses sediment, geochemical and micropaleo proxies (diatoms & calcareous nannofossils) for marine productivity, sea ice and land ice reconstruction. So I think a high-latitude marine record of the MIS12-11 period like this one is sorely needed to go with Lake E, Lake B and others.

Some general issues I think the authors should deal with in a revision:

C1

Full screen / Esc

Printer-friendly version

Discussion paper



1) the introduction tends to be unfocused and too long. Please shorten it giving the main hypotheses to be tested. I also think the methods and results sections tend to be long. In section 4.2.2, in the diatom ecology section, can the main taxa or assemblages used for productivity and sea ice be emphasized? Much of this section is taken care of in the table. In fact, it really constitutes a review of high latitude diatom ecology going back to Sancetta, Koizuma and other pioneers; this is useful but it merits its own paper in a micropaleo journal.

2) The methods section is long and could be put in a supplement, in fact some of it is, but the Supplementary Materials is not cited until Page 28.

Likewise, Section 5 is too descriptive and does not focus on the key patterns that address the hypothesis about suborbital variability.

3) related to # 1, I sense some of the text and references are not quite up to date [interglacial sea level papers, Bering Sea sea-level section 2.2 [where is Keigwin 2006 paper?], modern Bering Sea oceanography, see section 2.3)

4) The NADW discussion does not belong under a section called Bering Sea hydrology. Later in the paper, Section 5, there is again NADW discussion in the context of late MIS 11 Bering Sea reversed flow. In general, I don't think the Bering-N Atlantic links are well established mainly due to chronology/correlation issues, which I believe are discussed in a 2009 paper by L&R on Atlantic Pacific diachroneity of O18 records.

5) I wonder if the suggested correlations of this study's IODP core records with emerged Quaternary marine deposits [this is mentioned in several places] are warranted given age uncertainty of the onshore deposits?

6) The 404 ka ice-rafting event discussed on page 28 seems speculative and not up to date on ice-rafting processes in the Arctic and subarctic. This section evolves into a mechanistic explanation, covering the "snow gun" hypothesis and alternatives [turbidites, sea ice etc]. I think this section should be rethought and rewritten. As with the

[Full screen / Esc](#)[Printer-friendly version](#)[Discussion paper](#)

issue of Bering Sea flow reversal in an earlier section, are these central to the question of patterns and causes of variability within MIS11?

7) The paper uses both cores - U1345 & 1343 - although Kim published on U1343 using different proxies but the same O18 for tuning, is there any way to integrate results from both cores better to provide a more robust pattern of MIS paleoceanography?

Is the main focus of the study on orbital glacial-interglacial timescales or millennial timescales (that is, stadials and interstadials within MIS 11, see section 2.1 on sea level, or abrupt reversals like DO events ? The 15-meter thick MIS 11 record [line 199] ought to allow millennial-scale events to be seen. I have concern with the authors statement, in their discussion of the age model and tuning to LR04 and the other site U1343: “we urge caution when interpreting millennial scale changes at the site or comparing our record to others that examine MIS 11 at millennial scale resolution or finer”. I got the impression in the introduction there would be more definitive conclusions reached on within-interglacial climate variability. Plots in Figures 5-8 don't really show me DO-like or Heinrich-like variability, which could be an important new conclusion, given our ideas on what causes such events in at least the N Atlantic region.

In sum, I rated the paper as accept after minor revisions, some changes I am suggesting might take major text-shortening, but the science presented is sound, it is just not clearly packaged or presented.

Specific Comments

Line 27 comma before however

Line 28: This confuses me as the paper is in the Bering Sea, not the N Atlantic: led to “lowered productivity in both the northern Atlantic and the northern Pacific. “

Line 48 proper citation of IPCC 2013

Lines 55-56. Fix grammar in second part of sentence.

Full screen / Esc

Printer-friendly version

Discussion paper



Line 58 – which coastal region were these glaciers?

Line 70 – do you mean little is known from North Pacific Ocean region incl. Bering Sea?

Line 74. Is this marginal zone sea ice?

Line 114 E Antarctic ice was stable. . .

Line 196 “and” no italics

Section 4.2.1. Authors begin to use “ka” in discussions of diatoms but absolute years were not discussed in the age model-tuning section. So please tell readers earlier in the paper, at Table 2 reference, which should be in age model section, about the ages of MIS12, MIS 11 per the tuning to LR04. See line 586- the section title should say how old oldest sediments are, not “beginning of record”

Section 5.1 includes early deglaciation but section 5.2 is on Termination V, which is the deglaciation. Line 625, the debate about the duration of MIS 11 should be mentioned, embodied in papers by Masson-Delmotte, Ruddiman and others. See line 710 on this topic. I became confused about the MIS11 duration and the number of substages. Many authors do NOT use the substage terminology and the LR04 and the Antarctic ice core records show only two MIS 11 peak warm periods. One reason this is critical is this study of MIS 11 in the Bering Sea is one of the most detailed available. So it should shed light on the issue.

Figure 1 in the caption, mention both U-cores plotted in the map.

Figure 3 is a little complicated but it is critical. Consider dividing into 2 figures. The U1345 curve, red line, certainly looks different from that for U1343 – why so? Cook and Kim age model papers might be summarized in the text, in fact it would be useful to reproduce the O18, tie points data for the entire period covered by their tuning study.

Figures 5-8 are fine and do show what the study sought to accomplish: variability in

[Full screen / Esc](#)[Printer-friendly version](#)[Discussion paper](#)

diversity, lithology and microfossil assemblages. A more succinct treatment of these data in the text and better summary in the conclusions would help readers not familiar with these proxies. Why is MIS 11 split on the left in Fig 7 but not in others? Label horizontal colored panels in the figures for clarity. Figure 7 what is the source of N Atlantic stadials? Are they really relevant to this study?

Interactive comment on Clim. Past Discuss., doi:10.5194/cp-2015-184, 2016.

CPD

Interactive
comment

Full screen / Esc

Printer-friendly version

Discussion paper

