

Interactive comment on “Holocene evolution of the North Atlantic subsurface transport” by Janne Repschläger et al.

Janne Repschläger et al.

jr@gpi.uni-kiel.de

Received and published: 17 February 2017

Response to the reviewer on behalf of all co-authors

Response to Referee #2

We acknowledge the comments made by Referee #2 on our manuscript. While generally positive about the results presented Referee #2 criticizes that the freshwater influence mainly discussed as controlling factor for subtropical subsurface transport is too limited and that also other mechanisms should be addressed as driving factors for Holocene changes in the inter-gyre transport.

We propose to revise the discussion by including a more detailed paragraph describing changes in the atmospheric circulation mainly referring to changes in NAO patterns

C1

and its influence on the STG/SPG system (e.g. Olsen et al., 2012). Following the suggestion of Referee #1, also changes in MOW strength and its potential influence on the AF/AC position will be discussed. Finally, more studies (e.g. Blaschek et al., 2015) about the influence of early Holocene meltwater pathways will be included into the discussion.

Furthermore, Referee #2 proposed to extent the reconstruction of the warm water transport route with cores further south/north or east/west. Unfortunately, high-resolution Holocene studies investigating changes along the warm water route including surface and subsurface temperature and salinity reconstructions are limited. (Farmer et al., 2011) published Holocene surface and subsurface temperature and salinity time series from core MD99-2251 positioned at the eastern flank of Reykjanes Ridge underneath the inflow pathway of the NAC towards core RAPID 12-1 K. The dataset of core MD99-2251 shows the same long-term evolution over early to Mid Holocene evolution as the dataset of (Thornalley et al., 2009) and we thus will mention it in a revised manuscript. In order to highlight the connection to the Labrador Sea, a comparison between our time series and surface water datasets of (Solignac et al., 2004) will be added to one of our figures as proposed in our response to Referee #2. However, the sparse number of high-resolution subsurface temperature and $\delta^{18}\text{O}_w$ reconstructions emphasizes the need of further studies and indicates the novelty of our study.

A further aspect raised by Referee #2 was the strongly shortened methods section in the manuscript, that heavily relied on a previous publication (Repschläger et al., 2015). Indeed the methods section has been shortened very strictly in order to avoid self-plagiarism. However, we agree that it should be possible to follow the new manuscript independently. Thus the missing information will be provided in a revised version of the paper using a revised paragraph also requested by Referee #1:

“For our study we used core GEOFAR KF16 taken at 38°N, south of the Azores Islands at the Mid Atlantic Ridge, from 3060 m water depth. This position in the vicinity

C2

of the AF is ideal to trace changes in STG position and associated varying influence of subtropical and temperate waters (Repschläger et al., 2015). The age model for core GEOFAR KF16 is published in (Schwab et al., 2012). Results are compared to published data from the subtropical (Bahr et al., 2013; Cleroux et al., 2012) and subpolar North Atlantic (Thornalley et al., 2009) along the warm water route. We combine published $\delta^{18}\text{O}$ data of planktonic surface and subsurface dwelling foraminifera *G. ruber w.* (Schwab et al., 2012) and *G. truncatulinoides* (Repschläger et al., 2015) with new Mg/Ca records from the same species. All measurements were conducted on samples of the size fraction 315-400 μm on monospecific samples of 10-25 specimens of *G. ruber w.* and 15 specimens of mixed left and right coiling *G. truncatulinoides*. Stable isotope analyses were carried out at Leibniz Laboratory for Radiometric Dating and Stable Isotope Research at Kiel University. For analyses a Finnigan MAT 523 mass spectrometer coupled with a Kiel I carbon preparation device was used and results were calibrated to the Vienna Pee Dee Belemnite (V-PDB) scale. The 2σ standard deviation (std) obtained from 10 replicates of downcore samples was $\pm 0.11\text{‰}$ for *G. ruber w.* and $\pm 0.12\text{‰}$ for *G. truncatulinoides*. Mg/Ca analyses followed the cleaning procedure of (Martin and Lea, 2002; Repschläger et al., 2015), using reductive and oxidative cleaning. Measurements were carried out with a simultaneous inductively coupled plasma-optical emission spectrometry (ICP-OES) instrument with radial plasma observation. Potential shell contamination or coatings by authigenic minerals were monitored using additional Fe, Al and Mn measurements. Analytical 2σ std was $\pm 0.20\text{ mmol/mol}$ Mg/Ca for *G. ruber w.* and $\pm 0.14\text{ mmol/mol}$ Mg/Ca for *G. truncatulinoides*. To convert Mg/Ca values into water temperature estimates, we used the principle equation format $\text{Mg/Ca} = b \exp(a \cdot T)$ with species-specific variables $a=0.76$ and $b=0.07$ for *G. ruber w.* and $a=0.78$ and $b=0.04$ for *G. truncatulinoides* (equation for mixed subsurface dwellers) published by (Cleroux et al. 2008). The summed analytical and calibration 2σ std is $\pm 1^\circ\text{C}$ for *G. ruber w.* and $\pm 2^\circ\text{C}$ for *G. truncatulinoides*. The calibrated Mg/Ca temperature estimates for both species in our core top samples match well with modern surface and subsurface (200 m depth) temperatures at the

C3

Azores coring site. Seasonal temperature effects on the foraminiferal Mg/Ca signal are assumed to play a subordinate role at our coring site (Cleroux et al., 2008; Repschläger et al., 2015). In addition, we assume that the subsurface temperature signal of *G. truncatulinoides* is predominantly determined by the conditions at the AF and not by the migration of *G. truncatulinoides* to shallower, warmer water depths or by thermocline shoaling (see supplementary information and (Repschläger et al., 2015)). Changes in salinity are reconstructed following the procedure described in Repschläger et al. (2015). The temperature effect, estimated from the Mg/Ca was removed from the foraminiferal $\delta^{18}\text{O}$ carbonate using the general equation of Shackleton (1974) for both, *G. ruber w.* and *G. truncatulinoides*, in order to be consistent with the datasets used for comparison. A correction for VPDB to SMOW was included resulting seawater $\delta^{18}\text{O}$ values discussed in the following. For estimation of $\delta^{18}\text{O}$ uncertainties we followed the approach of Cleroux et al., (2011) and used their equation S8:

$$\sigma(\delta^{18}\text{O}_{\text{w-ivc}}) = ((\sigma(\delta^{18}\text{O}_{\text{foram}}))^2 + (\sigma(\text{Temp}))^2 \cdot (0.23)^2)^{1/2}$$

leading to calculated 2σ std of $\pm 0.35\text{‰}$ and $\pm 0.68\text{‰}$ for *G. ruber w.* and *G. truncatulinoides*, respectively. Given the relative large error in temperature and $\delta^{18}\text{O}$ reconstructions, we only used 5-point average time series for interpretation to investigate longer-term trends in the datasets. The $\delta^{18}\text{O}$ records were corrected for ice volume, using the eustatic sea level curve of (Austermann et al., 2013) and are expressed as $\delta^{18}\text{O}_{\text{w-ivc}}$ values throughout the text. Changes in the AF front were reconstructed using the relative abundance of *G. ruber w.* published by (Weinelt et al., 2015). Abundance counts possess a 2σ std of $\pm 2.5\%$. As *G. ruber w.* is most abundant within the STG (Ottens, 1991; Schiebel et al., 2002; Storz et al., 2009), low/high abundances indicate a southward/northward movement of the AF relative to the coring site. Because the position of the AF is mainly related to changes in the westerly wind belt, the abundances of *G. ruber w.* indicate the relative contribution of subtropical water and can be used as tracer for the position of the northern STG rim and thus of the position of westerlies (see also argumentation in Repschläger et al., 2015) before. “Changes in

C4

the AF front were reconstructed using the relative abundance of *G. ruber w.* published by (Weinelt et al., 2015) These abundance counts have an 2σ STD $\pm 2.5\%$. As *G. ruber w.* is most abundant within the STG (Ottens, 1991; Schiebel et al., 2002; Storz et al., 2009), low/high abundances indicate a southward/northward movement of the AF relative to the coring site. Because the position of the AF is related to changes in the westerly wind belt, the abundances of *G. ruber w.* indicate the relative contribution of subtropical water and can be used as tracer for the position of the northern STG rim and thus of the position of westerlies (see also argumentation in Repschläger et al., 2015). “

Additionally to these 2 major criticisms raised by Referee #2, we will perform all technical corrections and text passages suggested to further improve the manuscript. In a revised version of the manuscript, we will add the following references:

Austermann, J., Mitrovica, J. X., Latychev, K., and Milne, G. A.: Barbados-based estimate of ice volume at Last Glacial Maximum affected by subducted plate, *Nature Geosci.*, 6, 553-557, 2013.

Bahr, A., Nürnberg, D., Karas, C., and Grützner, J.: Millennial-scale versus long-term dynamics in the surface and subsurface of the western North Atlantic Subtropical Gyre during Marine Isotope Stage 5, *Global and Planetary Change*, 111, 77-87, 2013.

Blaschek, M., Renssen, H., Kissel, C., and Thornalley, D.: Holocene North Atlantic Overturning in an atmosphere-ocean-sea ice model compared to proxy-based reconstructions, *Paleoceanography*, 30, 2015PA002828, 2015.

Cleroux, C., Cortijo, E., Anand, P., Labeyrie, L., Bassinot, F., Caillon, N., and Duplessy, J.-C.: Mg/Ca and Sr/Ca ratios in planktonic foraminifera: Proxies for upper water column temperature reconstruction, *Paleoceanography*, 23, PA3214, 2008.

Cleroux, C., Debret, M., Cortijo, E., Duplessy, J.-C., Dewilde, F., Reijmer, J., and Massey, N.: High-resolution sea surface reconstructions off Cape Hatteras over the last 10

C5

ka, *Paleoceanography*, 27, PA1205, 2012.

Farmer, E. J., Chapman, M. R., and Andrews, J. E.: Holocene temperature evolution of the subpolar North Atlantic recorded in the Mg/Ca ratios of surface and thermocline dwelling planktonic foraminifers, *Global and Planetary Change*, 79, 234-243, 2011.

Martin, P. A. and Lea, D. W.: A simple evaluation of cleaning procedures on fossil benthic foraminiferal Mg/Ca, *Geochemistry, Geophysics, Geosystems*, 3, 8401, 2002.

Olsen, J., Anderson, N. J., and Knudsen, M. F.: Variability of the North Atlantic Oscillation over the past 5,200 years, *Nature Geosci.*, 5, 808-812, 2012.

Ottens, J. J.: Planktic foraminifera as North Atlantic water mass indicators, *Oceanologica Acta*, 14, 123-140, 1991.

Repschläger, J., Weinelt, M., Kinkel, H., Andersen, N., Garbe-Schönberg, D., and Schwab, C.: Response of the subtropical North Atlantic surface hydrography on deglacial and Holocene AMOC changes, *Paleoceanography*, 30, 2015.

Schiebel, R., Schmuker, B., Alves, M., and Hemleben, C.: Tracking the Recent and late Pleistocene Azores front by the distribution of planktic foraminifers, *Journal of Marine Systems*, 37, 213-227, 2002.

Schwab, C., Kinkel, H., Weinelt, M., and Repschläger, J.: Coccolithophore paleoproductivity and ecology response to deglacial and Holocene changes in the Azores Current System, *Paleoceanography*, 27, PA3210, 2012.

Shackleton, N. J.: Attainment of isotopic equilibrium ocean water and the benthonic foraminifera genus *Uvigerina*: Isotopic changes in the ocean during the last glacial, *Cent. Natl.Rech. Sci. Colloq. Int.*, 219, 203-209, 1974.

Solignac, S., de Vernal, A., and Hillaire-Marcel, C.: Holocene sea-surface conditions in the North Atlantic-contrasted trends and regimes in the western and eastern sectors (Labrador Sea vs. Iceland Basin), *Quaternary Science Reviews*, 23, 319-334, 2004.

C6

Storz, D., Schulz, H., Waniek, J. J., Schulz-Bull, D. E., and Kuçera, M.: Seasonal and interannual variability of the planktic foraminiferal flux in the vicinity of the Azores Current, *Deep Sea Research Part I: Oceanographic Research Papers*, 56, 107-124, 2009.

Thornalley, D. J. R., Elderfield, H., and McCave, I. N.: Holocene oscillations in temperature and salinity of the surface subpolar North Atlantic, *Nature*, 457, 711-714, 2009.

Weinelt, M., Schwab, C., Kneisel, J., Hinz, M., and Climate and societal change in the western Mediterranean area around 4.2 ka BP. In: 2200 BC – Ein Klimasturz als Ursache für den Zerfall der Alten Welt? 2200 BC – A climatic breakdown as a cause for the collapse of the old world? 7. Mitteldeutscher Archäologentag vom 17. bis 19. Oktober 2013 in Halle (Saale). 7th Archaeological Conference of Central Germany October 23–26, 2014 in Halle (Saale). Meller, H., Arz, H. W., Jung, R., and Risch, R. (Eds.), *Tagungen des Landesmuseums für Vorgeschichte Halle 12 Halle [Saale]* 2015.

Interactive comment on *Clim. Past Discuss.*, doi:10.5194/cp-2016-115, 2016.