

The comments of the two reviewers and M. Löffverström are very helpful and we thank them for this careful reading. They clearly show parts of the manuscript that need to be improved and rewritten.

The main point is probably the main conclusion of the paper and the warm LIG temperature at NEEM.

First, we agree that our results are probably not presented the right way and that our conclusion should be rewritten taking into account in figure 3 the broad ranges of accumulation estimate from modeling and data. This will lower the lowest LIG temperature for the NEEM deposition site by 1°C as noted by E. Wolff.

Second, another important point to take into account for this conclusion is the surprising comparison between NGRIP and NEEM d15N level that leads to different estimates of the LIG temperature. It should be noted that the value of 0.29 permil taken for NGRIP for the LIG in the table 1 was not representative enough of the NGRIP LIG section and we apologize for this mistake. In the previous manuscript, we have taken the deeper individual d15N value from the NGRIP ice core record from the initial low resolution study of Landais et al. (2005) while we should have taken the average of d15N values from the high resolution study of Capron et al. (GRL, 2012) over the bottom part of the NGRIP ice core corresponding to the oldest 1000 year period in the NGRIP ice core. When correcting for this and making an average over 1000 years, we obtain a mean d15N value of 0.275 permil for NGRIP LIG. This corresponds to a LIG mean temperature of -26.5+/-2°C, hence an increase of temperature of 5+/-2°C compared to pre-industrial temperature. Even if this temperature increase is larger than in the previous manuscript, this is still smaller than the temperature increase estimated at the NEEM deposition site. This has been corrected in the figure 3. Then, as suggested by reviewer 1, we have added a discussion about this difference, especially on p. 12 and 13.

Finally, we have also provided some more details on the modeling outputs and how they can be used to document seasonality of precipitation and possible link with d18O. The corresponding additions are in yellow highlights in section 3.3.

We provide an answer to every individual comment below:

Comments by M. Löffverström

1: I am somewhat critical to how you use climate model data in the current version of the paper. First, all models are to varying degree simplified versions of reality so a perfect match with proxy data is not to be expected. Secondly, comparing proxies with data in a single grid cell is arguably a misuse of climate models as they are designed to give an indication of the average conditions over a large region (the size of the region is dependent on the model complexity and grid resolution but a rule of thumb is to use at least a few grid cells). You should perhaps also comment on the range of models used in Fig. 4; the figure presents data from EMICS to full GCMs, which are worlds apart both in terms of complexity and modeling strategy (e.g. data constrained vs. free running, highly simplified vs. very complex, etc).

Indeed the models we used have very different complexities and resolution. We mention that on page 12, line 17 in the previous manuscript and emphasize it further in the next version (section 3.3). We hope that this helps to clarify the model – data comparison.

As for the grid resolution, we would argue to keep the climate model-data comparison for just the NEEM upstream grid box. This grid box is 3.75x2.5 degrees, “regridded” sometimes from lower resolution climate models. For answering this comment, we did the same analysis for the surrounding grid boxes, and the mean over the 9 grid boxes (see answer to comments in the online discussion version). The results change indeed a bit, but not drastically, and the assumed maximum of 2°C from LIG external forcing is still valid.

Another argument for keeping this grid box is that the temperature patterns over Greenland are relatively smooth, so you would also not expect the results to change much when analyzing another grid box. This can already be seen in the temperature maps of the Bakker and Lunt papers quoted in the manuscript.

2: You mention that the $\delta^{18}\text{O}$ signal recorded in ice cores can be influenced by changes in transport pathways and precipitation seasonality. The former is a bit tricky to investigate but you can easily perform a similar analysis as in Pausata and Löffverström (2015) (On the enigmatic similarity in Greenland $\delta^{18}\text{O}$ between the Oldest and Younger Dryas, *Geophys. Res. Letters*, 42, doi:10.1002/2015GL066042) and quantify the importance of precipitation seasonality and cloud temperature for the implied $\delta^{18}\text{O}$ signal in the models.

This comment is not easy to answer within the scope of this study because we do not have monthly mean model results for all models. Moreover, it would make more sense for this question to use models equipped with water isotopes.

Still, we had a look on the NorESM simulations using available surface temperature and precipitation rate at a monthly resolution (unfortunately, cloud temperatures are not available). If we calculate the temperature weighted by the precipitation rate for both pre-industrial and LIG, we find a much more important increase (roughly a factor of two) between pre-industrial and LIG than when using the mean annual temperature. Indeed, both the temperature and accumulation rate seasonalities are different from pre-industrial during LIG at NEEM: summer temperature (accumulation rate) increases by 3.5°C (7 mm/month) and winter temperature (accumulation rate) decreases by 2°C (3 mm/month). This monthly temperature and accumulation rate patterns are however significantly different from one grid box to another and probably from one model to another. We thus propose to include this discussion in a revised manuscript in a short paragraph indicating that this result is strongly variable and that a proper study of the seasonality effect on $\delta^{18}\text{O}$ should be made with a model equipped with water isotopes. The corresponding text is insert on p.4:

“As an example, if we use surface temperature and precipitation rate in monthly resolution from the NorESM model at the NEEM LIG deposition site, we observe a simulated increase in summer temperature (accumulation rate) by 3.5°C (7 mm/month) and a decreased in winter temperature (accumulation rate) by 2°C (3 mm/month).

This seasonality effect indeed leads to a factor of two between the precipitation weighted temperature change and the annual mean temperature change between pre-industrial and LIG.”

3: I would be careful citing unpublished work or papers in open discussion, except of course if the papers are accepted and about to be released. There is never a guarantee that a paper will be accepted only because it is in review and the methodology and conclusions might change significantly when the paper is finally published.

We have removed the reference to Langebroek and Nisancioglu, TCD

- **Comments by Anonymous Referee #1**

Major points:

p11 It is very interesting that the NEEM and NGRIP d15N reconstructions at 120 ka are so different: +8.5 versus +3.2‰. This merits more analysis and attention. In particular, is the 120 ka NGRIP is less affected by uncertainty generated by melt processes? This should be mentioned in the conclusions and in the abstract. Moreover the likely reasons for this discrepancy should be subject to further investigation/discussion and emphasis within this manuscript.

This is indeed important to focus more on the NGRIP ice core which was neglected in the first version. When coming back to these data, we actually realized that the d15N value of 0.29 permil taken for NGRIP for the LIG in the table 1 was not representative enough of the NGRIP LIG section and we apologize for this mistake. In the previous manuscript, we have taken the deeper individual d15N value from the NGRIP ice core record from the initial low resolution study of Landais et al. (2005) while we should have taken the average of d15N values from the high resolution study of Capron et al. (GRL, 2012) over the bottom part of the NGRIP ice core corresponding to the oldest 1000 year period in the NGRIP ice core. When correcting for this and making an average over 1000 years, we obtain a mean d15N value of 0.275 permil for NGRIP LIG. This corresponds to a LIG mean temperature of -26.5±2°C, hence an increase of temperature of 5±2°C compared to pre-industrial temperature. This has been corrected all along the text and in Figure 3. Even if this temperature increase is larger than in the previous manuscript, this is still less than the temperature increase estimated at the NEEM deposition site. As a consequence and as suggested by reviewer 1, a discussion is added on p. 12-13:

“In summary, the NGRIP LIG vs pre-industrial temperature increase (+5.2±2.3°C) is thus on the lower end but still compatible within error bars with the NEEM LIG vs pre-industrial temperature increase (+8±2.5°C at 120 ka using the aforementioned $\delta^{15}\text{N}$ value of 0.256 ‰). Several explanations can explain this discrepancy. First, the dating of the NGRIP and NEEM bottom parts is difficult because of the lack of precise relative and absolute age markers; this limits our confidence that we indeed compare $\delta^{15}\text{N}$ levels of same age on the two different cores. Second, we do not have any evidence of melt layers in the bottom of the NGRIP core, opposite to NEEM. This suggests that the NEEM deposition site was indeed warmer than NGRIP at LIG but also suggests that firn densification may have been affected by this process at NEEM which

would bias our reconstruction. Finally, our temperature reconstruction at NEEM is complicated by the fact that the NEEM LIG deposition site and NEEM drilling site have different surface conditions. In particular, an estimate of the pre-industrial accumulation rate is missing at the NEEM deposition site.”

p12 L7-9 Have the studies by Metz et al not already provided enough information to attempt to constrain these 'other influences' on surface temperature and accumulation rates? And if so, should this not be included within the analysis/uncertainties for this study?

This is a good point, in line with the comments of E. Wolff below and we have included the uncertainty on accumulation in the new determination which lowers the low range of LIG temperature increase at the NEEM deposition site to 6°C (instead of 7°C). Section 3.2.1 has thus been almost entirely rewritten (see text with yellow highlights at the end of this document).

Minor points:

All minor points have been taken into account in the revised manuscript.

Comments by EW Wolff (Referee)

the authors reach a conclusion that I don't think their data justify, especially if one includes a further estimate of the accumulation rate, which I would rate as being at least as valid (perhaps more so) compared to the ones they choose. The critical part of the paper is Figure 3. I would make a first comment that, given how crucial this figure is, it's extremely hard to follow. I'll follow this up later. If I accept all the accumulation estimates shown (but I don't, see below), I reach the conclusion of the authors in section 3.2.3, but not the overall conclusion of the paper. This section seems to conclude:

For Summit, the data can probably not be interpreted as a pure thermal signal; For NEEM, you seem to choose a range between the +20% accumulation (4.5 degrees warming), and the “M-D” approach (8 degrees warming). The authors don't explicitly state how they corrected for the upstream issue but I assume from Table 2 that they added 2.5 degrees. Thus the range they estimate is 7-10.5 degrees. For NGRIP, the best estimate is 3.2+/-0.7 degrees, with no significant upstream correction, and a possible extra 0.5 degrees for the warmest part of the LIG.

So taken at face value, one would conclude that one site gives 7-10.5 degrees and the other gives 2.5-4 degrees, which does not seem like a basis for asserting that the climate was 8 degrees warmer and that the NEEM paradox is confirmed.

We agree that the conclusion should be revised and we certainly did not focus enough on the NGRIP site in the previous manuscript (and too much on the NEEM site). Also, as previously mentioned, we have realized only after we submitted our manuscript that the LIG d15N value taken for NGRIP was not correct. When accounting for the updated LIG d15N value based on Capron et al. 2012 high resolution d15N record, we obtain a LIG temperature increase at the NGRIP deposition site larger than originally calculated. In addition, the arguments developed below are sound as well and we should not discard lowest accumulation rate increases as

performed in the manuscript, especially for NEEM whose deposition site is not very well known. This will be corrected in the new version. The temperature of the LIG will thus be higher by 3-7°C at NGRIP and 6 to 11°C at the NEEM deposition site relative to preindustrial (11°C being highly improbable as already explained in the first manuscript and in agreement with comments below). It is still a bit high compared to the model simulations but the conclusion has been tuned down compared to the first version (see section 4 in the text at the end of this document with many parts removed and parts rewritten in yellow highlights”

However, what Figure 3 really tells us is that there is no good way of estimating the accumulation rate in the LIG, and therefore in the end we can't constrain the temperature this way. I can suggest another very valid way to estimate the accumulation rate. In Kapsner et al (1995), which the authors cite, an accumulation rate-temperature estimate for the early Holocene (pre-Boreal, which seems the most relevant) is made from correlation of annual layer thicknesses with oxygen isotope ratio measurements. They find a rather shallow slope within climate periods such as the pre-Boreal. Because they actually did a regression of accumulation against oxygen isotopes, we can in fact estimate directly (without going through temperature) what should be the acc rate change for an oxygen isotope change of 3.5 permil (Table 2, change at deposition site). Given that they report a slope of 0.9%/K, and they used a calibration of 0.53 permil/K, their acc-isotope slope must have been 1.7%/permil, and a 3.6 permil change would correspond to a 6% change in accumulation rate. This would suggest a 3.5 degree warmer firn column, which adding the 2.5 degrees upstream correction, implies the NEEM region was 6 degrees warmer than present.

This is indeed correct. It is still surprising that the results from Kapsner et al. (1995) for GISP2 are different from the Dahl-Jensen et al. (1993) for GRIP and Buchardt et al. (2012) while the methods are similar. But this goes in line with the other comments that the accumulation rate vs temperature relationship may be very variable in Greenland and we should not have discarded the possible stable accumulation rate scenario for warmer period. This has been corrected. We also understand the need for simplification of figure 3 where we can indeed only show the LIG accumulation rate possible range and remove the accumulation rate vs temperature relationships. Figure 3 has been redrawn together with the text on accumulation rate estimates.

Part of section 3.2.1 rewritten for the Kapsner et al., 1995 reference:

“iii: Based on the GISP2 ice core records over the last deglaciation, Kapsner et al. (1995) showed that the relationship between Greenland accumulation rate and temperature was not stable because of variations in atmospheric circulation. Still, they were able to propose a temporal relationship between accumulation rate based on annual layer counting and $\delta^{18}\text{O}$, or temperature reconstructed from $\delta^{18}\text{O}$ and a calibration based on borehole temperature measurements (leading to a $\delta^{18}\text{O}$ vs temperature slope of $0.53\text{‰}\cdot^{\circ}\text{C}^{-1}$). The inferred sensitivity of snow accumulation rate to temperature change during interglacial period varies from $0.9\text{‰}\cdot^{\circ}\text{C}^{-1}$ (Holocene) to $7.5\text{‰}\cdot^{\circ}\text{C}^{-1}$ (Bølling-Allerød), with an uncertainty encompassing zero.

More recently, Buchardt et al. (2012) followed a similar approach but using numerous ice cores. They used an array of 52 shallow ice cores spanning the last decades to centuries with accumulation rate estimates from annual layer

counting on $\delta^{18}\text{O}_{\text{ice}}$ profiles. They identified different temperature vs accumulation rate relationships from one region to another. In central and north Greenland corresponding to the location of the NEEM, NGRIP, GRIP and GISP2 deep ice cores, the Buchardt approach suggests a sensitivity of 1.5 to 9.4%.°C⁻¹ with an uncertainty encompassing zero. This sensitivity is obtained with a $\delta^{18}\text{O}$ vs temperature sensitivity slope of 0.67‰.°C⁻¹ so that the Kapsner and Buchardt estimates agree on a 0 to 14%.‰⁻¹ accumulation rate vs $\delta^{18}\text{O}$ sensitivity. “

In the rest of this review I will go through the paper in more detail, and comment some more on Figure 3. However my overall suggestion is that, taking into account another realistic way of calculating accumulation rate, and the NGRIP result, the conclusions and abstract should be much more balanced and should not claim to be confirming the very high estimate of NEEM Project Members.

As mentioned above, this has been done both in the abstract and main text (see yellow highlights at the end of the document).

As I already suggested, you really only show in this section that we have no good basis for estimating the accumulation rate that applies to this warm interglacial. Here I just comment on some of the estimates you give: i) The Kapsner paper quite conclusively shows that the thermodynamic approach is not really applicable to a situation like that in Greenland, where most of the precipitation is related to cyclones and storm activity. It's fine to mention it as an option, but it is clearly flawed. When it comes to figure 3, I really don't understand what the grey shaded area is meant to represent, so if you leave it, it needs a better explanation. ii) It seems as if this Benson approach is actually an empirical spatial version of the thermodynamic approach? However, while I could almost justify this approach for single storms tracked (and their integrated effect over a year) spatially across Greenland, there seems no basis for translating that into the temporal domain, as you acknowledge on Page 8, line 22. Note also that Benson 1962 seems missing from the reference list.

For simplicity, we have removed the Benson 1962 outputs (indeed a temporal relationship) from Figure 3. The idea is still to keep it as a reference because it is still used today for validation of accumulation rate estimate in Greenland.

iii) Same applies for the Buchardt approach, but since the blue lines are missing from the plot I cannot assess it. I understand why you prefer the sensitivity Buchardt derived for the NW region but don't understand why you give a range for Greenland of 6.7-9.6%/K, when Buchardt's table gives a value of only 1.5%/K (with an error bar encompassing zero) for the nearby NE region, and even a negative slope for SW Greenland. I do not think you have summarised the Buchardt outcome fairly; in my opinion it gives weak evidence for a 6.7%/K slope, but with an uncertainty that encompasses a zero slope. Note also that it is unnecessary to draw a curve to represent Buchardt; he actually derived an acc rate- $\delta^{18}\text{O}$ slope, so for a measured change in $\delta^{18}\text{O}$, you can pinpoint precisely the range of the change in acc rate and draw them as horizontal lines .

We have included the NW region as well in the discussion of section 3.2.1 and simplified figure 3 as mentioned above (only give the range of possible accumulation rate as for 10Be or chemistry).

Part of section 3.2.1 including the reference to Buchardt 2012:

" More recently, Buchardt et al. (2012) followed a similar approach but using numerous ice cores. They used an array of 52 shallow ice cores spanning the last decades to centuries with accumulation rate estimates from annual layer counting on $\delta^{18}\text{O}_{\text{ice}}$ profiles. They identified different temperature vs accumulation rate relationships from one region to another. In central and north Greenland corresponding to the location of the NEEM, NGRIP, GRIP and GISP2 deep ice cores, the Buchardt approach suggests a sensitivity of 1.5 to 9.4%. $^{\circ}\text{C}^{-1}$ with an uncertainty encompassing zero. This sensitivity is obtained with a $\delta^{18}\text{O}$ vs temperature sensitivity slope of 0.67‰. $^{\circ}\text{C}^{-1}$ so that the Kapsner and Buchardt estimates agree on a 0 to 14%.‰ $^{-1}$ accumulation rate vs $\delta^{18}\text{O}$ sensitivity. "

iv) I have not checked the MD approach specifically, but note that this is the same paper that derived a much larger $\delta^{18}\text{O}$ -temperature slope and therefore a smaller temperature change than NEEM Project Members, so it seems inconsistent to use the same model simulations to derive the opposite result.

The accumulation – temperature relationships given in Masson-Delmotte et al. (2015) have indeed a very high slope for the recent warming. This is not necessary inconsistent with the high $\delta^{18}\text{O}$ vs temperature slope but we understand that it may lead to unnecessary complications. We thus propose to gather the accumulation vs temperature and $\delta^{18}\text{O}$ vs temperature relationship from same source (model outputs, reanalyses, ...) and only give the final accumulation rate vs $\delta^{18}\text{O}$ relationship (hence combining values from Tables 6 and 7 from Masson-Delmotte et al., 2015). This leads to values around 10 %.‰ $^{-1}$, hence a 35% change in accumulation rate for an increase in $\delta^{18}\text{O}$ of 3.5‰ at the NEEM deposition site. This again leads to a temperature reconstruction for the NEEM deposition site at the upper boundary of the 6-11 $^{\circ}\text{C}$ range.

Corrected text in section 3.2.1:

"iv: Masson-Delmotte et al. (2015) used estimates of snow accumulation rate and ice $\delta^{18}\text{O}$ in four shallow cores in the NEEM area together with accumulation rate, temperature and when possible snowfall $\delta^{18}\text{O}$ reconstructions from different models simulation (ECHAM5, Global Climate Model developed by the Max Planck Institute for Meteorology; LMDZ, Global Climate Model developed by the Laboratoire de Météorologie Dynamique; MAR, Regional Atmosphere Model) nudged to available climate reanalyses over the 1979 – 2007 period. In addition to model outputs or temperature gridded reconstruction (Box et al., 2009), the amplitude of temperature increase at NEEM can also be estimated using borehole temperature measurements. Gathering the different sources of information for the strong warming period of 1979-2007 leads to a relatively high slope between accumulation rate and temperature (10 to 15.9 %. $^{\circ}\text{C}^{-1}$, the highest value being obtained using outputs from the MAR model nudged to ERA-40 and ERA-Interim reanalyses (Uppala et al., 2005; Dee et al., 2011)). In this study, the sensitivity of accumulation rate vs. $\delta^{18}\text{O}$ can be estimated through the regression between the NEEM $\delta^{18}\text{O}$ and accumulation rate increases over the period 1979 – 2007 leading to a value of 10 %.‰ $^{-1}$, in agreement with the Buchardt estimate. Another solution is to use the $\delta^{18}\text{O}$ vs temperature estimate based on NEEM $\delta^{18}\text{O}$ measurements vs borehole temperature over the recent warming trend (0.8%. $^{\circ}\text{C}^{-1}$) together with the accumulation rate vs temperature

estimate given above, hence leading to a maximum accumulation rate vs $\delta^{18}\text{O}$ sensitivity of 13 %‰⁻¹, again within the range of the Buchardt estimate . “

v) I have not checked the ^{10}Be estimates, but note only that because deposition in Greenland under interglacial conditions is completely dominated by wet deposition, it seems very unlikely that one can derive accumulation rates from ^{10}Be . I agree with your implication that the change in overall scavenging would have to be accounted for in a sophisticated model, and I don't believe that local accumulation rate would be the dominant control on ^{10}Be concentration or flux. For the same reason, I do not understand at all your statement on Page 9, lines 22-24 that changes in aerosol rule out no change in accumulation: if you want to make such a statement it needs considerably more explanation and analysis, as there seems no basis to estimate the changes in sources well enough to make such a statement.

The argument for the statement that the aerosol chemistry renders the same accumulation in the Eemian and the Holocene as unlikely is as follows: If we compare aerosol species which are dominated by wet deposition such as Na (sea salt aerosol) and NO_3 (lightning activity, biological activity) we see that the concentration in the ice in the Eemian is drastically lower than in the HOL (only about 50%). As these species are mainly wet deposited this is only possible if the atmospheric aerosol concentrations over the ice sheet was also lower by 50% at that time. It is unlikely that both Na and NO_3 (which have completely different sources and transport pathways) have both a 50% reduction of source emissions, which is too large anyway. We can also get a reduction of 50% in atmospheric Na and NO_3 concentrations over the ice, if we increase the precipitation rate along the transport pathway and thus increase wet deposition en route. So what the chemistry points evidences is that the precipitation rate during transport in the Northern Hemisphere was significantly higher in the Eemian than in the HOL. If this is not case on the Greenland ice sheet, it means that the precipitation rate should be higher in the Eemian everywhere else except in Greenland, which is unlikely. In particular as sea salt aerosol transport from the open ocean comes jointly with water vapor transport to the ice sheet through storm events an increased washout of Na en route is expected due to higher precipitation rates and this would also dump more water on the ice sheet.

Finally, note that if we scale the change in precipitation rate en route and in local accumulation rate from the HOL to the Eemian according to the Buchardt formula, no change in Na and NO_3 sources strength is required at all.

The text has been rewritten:

“v: Alternative estimates of accumulation rate at the NEEM deposition site are also provided by ^{10}Be data. Sturevik-Storm et al. (2014) compiled mean Holocene ^{10}Be concentration over eight Greenland sites and determined the spatial relationship against mean accumulation rate estimates. This relationship was then applied to LIG ^{10}Be data from NEEM, leading to the conclusion that accumulation rate was 65-90% higher than today at the NEEM deposition site (“ ^{10}Be -approach”). However, the present-day spatial gradients in ^{10}Be concentration are caused by spatially varying contributions of wet deposition to the overall ^{10}Be deposition, assuming a homogeneous atmospheric ^{10}Be aerosol deposition over Greenland. This latter assumption implies that the atmosphere above

Greenland is well mixed with respect to ^{10}Be after transport from the location of troposphere/stratosphere foldings. The latter are the main entrance pathways of stratospheric aerosols into the extratropical northern hemisphere troposphere. The LIG climate is characterized not only by likely enhanced precipitation above Greenland (accumulation) but also higher wet deposition during aerosol transport to Greenland due to higher precipitation rates. As a result, higher scavenging of ^{10}Be bearing aerosol en route must have led also to a lower atmospheric ^{10}Be concentration over Greenland than today. The LIG accumulation estimate by Sturevik-Storm et al. (2014) is therefore most likely an overestimation and the assumption of Sturevik-Storm et al. (2014) that ^{10}Be concentration is only controlled by accumulation rate at the NEEM site may be challenged. More generally, the use of other chemical aerosol species as accumulation rate tracers is hampered by potential changes in the LIG atmospheric concentrations due to emission changes. Qualitatively, a correction of deposition effects using the Buchardt et al. (2012) approach representative for Northwest Greenland, leads to LIG atmospheric concentrations of all chemical aerosol tracers similar to today. In contrast much higher LIG accumulation rates as estimated by Sturevik-Storm et al. (2014) or no changes in accumulation between the LIG and the Holocene imply an unrealistic change in atmospheric aerosol concentrations for several aerosol tracers. Indeed, if we compare aerosol species that are dominated by wet deposition such as Na^+ (sea salt aerosol) and NO_3^- (lightning activity, biological activity), we see that the concentration in the ice in the LIG is lower than in the Holocene (about 50%). As these species are mainly wet deposited, this is only possible if the atmospheric aerosol concentration was also reduced by 50% at that time. It is unlikely that both Na^+ and NO_3^- (which have completely different sources and transport pathways) have a 50% reduction of source emissions. Another solution to explain the reduction of 50% in atmospheric Na^+ and NO_3^- concentration over the ice is to imply an increase of the precipitation rate along the transport pathway and thus increase of wet deposition en route. The chemistry suggests that the precipitation rate during transport in the Northern Hemisphere was significantly higher during the LIG than during the Holocene and there is no reason why Greenland would not be affected by this general increased accumulation rate. Based on the changes in various chemical tracers in the ice (sea salt aerosol, biogenic aerosol, mineral dust) we thus conclude that the LIG accumulation was likely 20 % higher than pre-industrial and similar to the Buchardt approach for Northwest Greenland.”

vi) It seems that GCMs agree with a small change in accumulation rate. Taking all these estimates together with the one I derived from Kapsner et al, and noting that they all have significant weaknesses, I see no reason to rule out a very small change in accumulation rate, which is suggested by Kapsner et al, approach vi, and the lower range of the Buchardt approach when you include their NE data. It's unfortunate but it means you simply can't constrain accumulation rate this way.

This is correct. It is now included in section 3.2.1 (see above and in the text at the end of the document).

Page 14, line 28. I don't see the relevance of this: at current mean summer temperatures we know we can get melt layers (as we did this year). You certainly don't need the summer mean to be 5 degrees warmer than present to expect significant melt every year.

We were referring to comparison with pre-industrial temperature. In the recent years, melt-layers were observed at NEEM during heat wave as in 2012 with summer temperature of $\sim 5^{\circ}\text{C}$ warmer than pre-industrial summers.

Fig 1. Please remove the age scale for the shaded portion beyond 128 ka, as we have no basis for it. Summer insolation should also be cut at 128 ka as it cannot be compared to the climate curves below it beyond that age.

Done

Figure 3. Please make this figure clearer in the caption. I suggest something like: "The black circles. . .NEEM. The light curves are contours of accumulation rate-temperature combinations for a given value of $\delta\epsilon_{15}\text{N}$, with that value shown in the curve. The coloured examples correspond to the measured LIG $\delta\epsilon_{15}\text{N}$ for GRIP (blue), NGRIP (purple) and NEEM (red). The horizontal lines and darker curves correspond to different estimates of the accumulation rate (for the curves, this is as a function of temperature)." Then having explained the overall point of the diagram, you can explain each curve, but of course I would hope that the Buchardt lines will be shown and will include a wider range than in the text, that the Kapsner line is added, and that the vertical arrows better reflect the range of credible estimates.

The figure 3 has been entirely redrawn.

How warm was Greenland during the last interglacial period?

Amaelle Landais¹, Valérie Masson-Delmotte¹, Emilie Capron^{2,3}, Petra M. Langebroek⁴, Pepijn Bakker⁵, Emma J. Stone⁶, Niklaus Merz⁷, Christoph C. Raible⁷, Hubertus Fischer⁷, Anaïs Orsi¹, Frédéric Prié¹, Bo Vinther², Dorthe Dahl-Jensen².

¹ Laboratoire des Sciences du Climat et de l'Environnement - IPSL, UMR 8212, CEA-CNRS-UVSQ-Université Paris Saclay, Gif-sur-Yvette, France

² Center for Ice and Climate, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, 2100 Copenhagen Ø, Denmark.

³ British Antarctic Survey, High Cross Madingley Road, Cambridge CB3 0ET, UK

⁴ Uni Research Climate, Bjerknes Centre for Climate Research, Nygårdsgaten 112-114, 5008 Bergen, Norway

⁵ College of Earth, Ocean and Atmospheric Sciences, Oregon State University, USA

⁶ BRIDGE, School of Geographical Sciences, University of Bristol, Bristol, UK

⁷ Climate and Environmental Physics, Physics Institute, and Oeschger Centre for Climate Change Research, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

Abstract. The last interglacial period (LIG, ~129-116 thousand years ago) provides the most recent case study for multi-millennial polar warming above pre-industrial level and a respective response of the Greenland and Antarctic ice sheets to this warming, as well as a test bed for climate and ice sheet models. Past changes in Greenland ice sheet thickness and surface temperature during this period were recently derived from the NEEM ice core records, North-West Greenland. The NEEM paradox has emerged from an estimated large local warming above pre-industrial level ($7.5 \pm 1.8^\circ\text{C}$ at the deposition site 126 ka ago without correction for any overall ice sheet altitude changes between the LIG and pre-industrial) based on water isotopes, together with limited local ice thinning, suggesting more resilience of the real Greenland ice sheet than shown in some ice sheet models. Here, we provide an independent assessment of the average LIG Greenland surface warming using ice core air isotopic composition ($\delta^{15}\text{N}$) and relationships between accumulation rate and temperature. The LIG surface temperature at the upstream NEEM deposition site without ice sheet altitude correction is estimated to be warmer by $+8.5 \pm 2.5^\circ\text{C}$ ($+8^\circ\text{C}$ being the most likely estimate according to constraints on past accumulation rate) compared to the pre-industrial period. This temperature estimate is consistent with the $7.5 \pm 1.8^\circ\text{C}$ warming initially determined from NEEM water isotopes but on the upper end of the pre-industrial to LIG temperature difference of $+5.2 \pm 2.3^\circ\text{C}$ obtained at the NorthGRIP site by the same method. Climate simulations performed with present day ice sheet topography lead in general to a warming smaller than reconstructed, but sensitivity tests show that larger amplitudes (up to 5°C) are produced in response to prescribed changes in sea ice extent and ice sheet topography. Still, ice sheet simulations forced by 5°C surface warming lead to large ice sheet decay that are not compatible with existing data. Our new, independent temperature constrain therefore reinforces the NEEM paradox.

1 Introduction

Understanding the magnitude, timing and rate of contributions of the Greenland and/or Antarctic ice sheets to the estimated 5 to 10 m increase in global mean sea level during the last interglacial period (LIG, 129-116 thousand years before 1950, hereafter ka) and therefore ice sheet vulnerability to multi-millennial polar warming remains challenging (Masson-Delmotte et al., 2013; Dutton et al., 2015). Therefore, constraints on past polar climate and ice sheet response are required. Additionally, polar temperature reconstructions provide a benchmark to assess the ability of climate models in capturing feedbacks which amplify the impact of orbital forcing on polar temperatures (Masson-Delmotte et al., 2011; Otto-Bliesner et al., 2013a; Capron et al., 2014). This latter is also relevant for future climate projections.

Since the 1960s, numerous Greenland deep ice core records have provided evidence for layers of ice located near bedrock characterised by high values of water stable isotopes ($\delta^{18}\text{O}_{\text{ice}}$), well above pre-industrial Holocene levels (Johnsen et al., 1997). The climate interpretation of the first records was limited due to poor preservation of deep samples (Camp Century, Dye 3), and the lack of remaining air content preventing any dating by synchronisation with global atmospheric records (i.e. atmospheric $\delta^{18}\text{O}$ of O_2 , hereafter $\delta^{18}\text{O}_{\text{atm}}$ and CH_4) from undisturbed Antarctic

records. This synchronisation method was applied for the LIG interval at Summit, where ice from the LIG was unequivocally identified although not unambiguously datable, but sharp variations in $\delta^{18}\text{O}_{\text{ice}}$ at GRIP and GISP2 were attributed to stratigraphic disturbances (Groote et al., 1993; Landais et al., 2004; Landais et al., 2003; Suwa et al., 2006). At NGRIP, continuous climatic and environmental records cover the last 123 ka (NorthGRIP-community-members, 2004). The Greenland record was recently extended back to 128 ka thanks to a 80 m segment of ice in stratigraphic order found in between disturbed layers at the bottom of the NEEM ice core (NEEM comm. members, 2013). The chronology of this core was tied to an Antarctic ice core age scale, based on common changes in atmospheric composition. The unequivocal matching between the NEEM LIG layer and the Antarctic $\delta^{18}\text{O}_{\text{atm}}$ records rules out stratigraphic disturbance **within this segment** (NEEM comm. members, 2013).

Changes in NEEM air content and $\delta^{18}\text{O}_{\text{ice}}$ were corrected for elevation changes due to the upstream displacement of the deposition site, and combined to infer changes in ice sheet topography, and changes in surface air temperature (NEEM comm. members, 2013). This requires assumptions on NEEM $\delta^{18}\text{O}_{\text{ice}}$ – temperature relationships. While Greenland snow isotopic composition has long been related to temperature due to Rayleigh distillation associated with cooling along air mass pathways (Dansgaard et al., 1964), it has been increasingly documented that $\delta^{18}\text{O}_{\text{ice}}$ -temperature relationships are neither stable in time nor in space (e.g. Jouzel et al., 1999) primarily due to changes in the precipitation intermittency, but also evaporation conditions and atmospheric transport (e.g. Krinner et al., 1997; [Masson-Delmotte et al., 2011](#)).

The initial LIG temperature estimate (NEEM comm. members, 2013) was performed using the average Holocene $\delta^{18}\text{O}_{\text{ice}}$ -temperature relationship established from other central Greenland ice cores through calibration against borehole temperature at $0.5\text{‰}\cdot^{\circ}\text{C}^{-1}$ (Vinther et al., 2009). This relationship was also explored in simulations using isotopically enabled atmospheric general circulation models for climate conditions warmer than pre-industrial, either in response to increasing CO_2 concentration in projections, or in response to changes in orbital forcing. These models produced slopes varying from 0.3 to $0.7\text{‰}\cdot^{\circ}\text{C}^{-1}$ in Greenland, depending on changes in moisture sources driven by changes in sea ice and sea surface temperature patterns (Masson-Delmotte et al., 2011; Sime et al., 2013). Based on these lines of evidence, a slope varying from 0.4 to $0.6\text{‰}\cdot^{\circ}\text{C}^{-1}$ was used to estimate the range of changes in LIG temperature based on NEEM $\delta^{18}\text{O}_{\text{ice}}$ (NEEM community members, 2013). At 126 ka, and at the location of the initial snowfall deposition site (about 205 ± 20 km upstream of the current NEEM site), $\delta^{18}\text{O}_{\text{ice}}$ was estimated at 3.6‰ above local pre-industrial level, which translated into local surface air temperature warming of $7.5 \pm 1.8^{\circ}\text{C}$. After accounting for upstream effects and for Greenland ice sheet elevation change based on air content, this led to an estimate of a $8 \pm 4^{\circ}\text{C}$ warming at the NEEM deposition site at 126 ka (NEEM comm. members, 2013). In parallel, ice sheet simulations forced by different LIG climate scenarios were investigated to select only those compatible with limited change in ice thickness at NEEM, based on air content data. This implied limited Greenland ice sheet deglaciation, with a contribution of 1.4 to 4.3 m to the LIG sea level increase (Masson-Delmotte et al., 2013).

These results led to the “NEEM paradox”, where the Greenland ice sheet appears resilient to large multi-millennial surface warming. This paradox was further enhanced by the difficulty of coupled ocean-atmosphere climate models

to capture such warming (Otto-Bliesner et al., 2013b; Capron et al., 2014), even during the warmest summer months (van de Berg et al., 2013), and by the inconsistency of the Greenland ice sheet retreat simulated by ice sheet models in response to such warming (e.g. Stone et al., 2013; Helsen et al., 2013). When accounting for a reduced Greenland ice sheet and a retreat in sea ice cover in the Nordic Seas, atmospheric simulations can explain up to 5°C annual mean warming with respect to pre-industrial (Merz et al., 2014a; 2016). Moreover, all LIG climate modelling studies cited above strongly enhance summer precipitation seasonality in Greenland, suggesting a summer bias for LIG $\delta^{18}\text{O}_{\text{ice}}$ and weaker annual mean change than the initial estimate of $8\pm 4^\circ\text{C}$ (Masson-Delmotte et al., 2011; Merz et al., 2014b). As an example, if we use surface temperature and precipitation rate in monthly resolution from the NorESM model at the NEEM LIG deposition site, we observe a simulated increase in summer temperature (accumulation rate) by 3.5°C (7 mm/month) and a decreased in winter temperature (accumulation rate) by 2°C (3 mm/month). This seasonality effect indeed leads to a factor of two between the precipitation weighted temperature change and the annual mean temperature change between pre-industrial and LIG.

Recently, new information on climatic controls on NEEM $\delta^{18}\text{O}_{\text{ice}}$ has emerged from present-day water isotope monitoring and multi-decadal trends from shallow ice cores (Steen-Larsen et al., 2011; Steen-Larsen et al., 2014). All these datasets coherently document a surprisingly large present-day response of NEEM $\delta^{18}\text{O}_{\text{ice}}$ to temperature, with a slope of $[0.8\text{-}1.2] \text{‰}\cdot^\circ\text{C}^{-1}$ (Masson-Delmotte et al., 2015). If relationships established from the intra-seasonal to the multi-decadal scale remain valid for earlier warm periods such as the LIG, it also implies that the initially reconstructed temperature change based on NEEM $\delta^{18}\text{O}$ was overestimated.

Here, we present new, independent information on LIG annual mean temperature change for several Greenland drilling sites, using the ice core air isotopic composition $\delta^{15}\text{N}$. These Greenland records are described in Section 2. Section 3 details the temperature reconstructions with their associated uncertainties, with a focus on the NEEM deposition site. These temperature estimates depend on assumptions on the past relationship between temperature and accumulation rate. Section 4 presents a comparison to modeling outputs for discussion before the conclusions.

2 Water and air isotope records of the last interglacial in Greenland

2.1 Records of water stable isotopes from multiple ice cores on a coherent chronology

Figure 1 shows the compilation of the LIG $\delta^{18}\text{O}_{\text{ice}}$ records from NGRIP, GRIP and GISP2 sites on a coherent timescale. NEEM $\delta^{18}\text{O}_{\text{ice}}$ is presented on a parallel depth scale adjusted for the alignment of $\delta^{18}\text{O}_{\text{atm}}$ records over the LIG section. As CH_4 and $\delta^{18}\text{O}_{\text{atm}}$ are globally well-mixed atmospheric tracers, comparable values are measured in the Greenland and Antarctic ice cores at the same time period, accounting for the CH_4 interpolar gradient leading to slightly higher CH_4 levels in Greenland than in Antarctica (e.g. Dällenbach et al., 2000). The synchronization between the records is therefore based on parallel large variations in CH_4 and $\delta^{18}\text{O}_{\text{atm}}$ from measurements in the air trapped in bubbles. For the end of the LIG and the glacial inception, NGRIP records were placed on the AICC2012 timescale (Bazin et al., 2013; Veres et al., 2013) using CH_4 and $\delta^{18}\text{O}_{\text{atm}}$ tie-points between NGRIP and the Antarctic EPICA Dronning Maud Land (EDML) ice core (Capron et al., 2010). However, the AICC2012 NGRIP chronology

is limited since (1) no synchronization points are available for ages older than 118 ka (supplementary online material in Bazin et al., 2013; Veres et al., 2013) and (2) the mean CH₄ level is significantly higher at NGRIP than in the EPICA Dome C (EDC) record (Capron et al. 2012). The latter is interpreted in part to reflect a strong increase in the inter-hemispheric CH₄ gradient, which complicates the alignment of NGRIP and EDC CH₄ records (Capron et al., 2012). Additionally, a slight mismatch is observed between the LIG NGRIP and the recently published EDC $\delta^{18}\text{O}_{\text{atm}}$ records (Figure 1; Landais et al., 2013), suggesting that NGRIP ice chronology may be too young by up to 2 ka at 121 ka, when compared to the AICC2012 chronology.

The dated GRIP and GISP2 $\delta^{18}\text{O}_{\text{ice}}$ records are discontinuous, because of strong stratigraphic disturbances over the bottom 300 m of these Summit ice cores. They were initially placed on the Vostok GT4 timescale (Petit et al., 1999) using identification of $\delta^{18}\text{O}_{\text{atm}}$ /CH₄ pairs and taking into account the inter-polar CH₄ gradient (Landais et al., 2003; Suwa et al., 2006). Here, we have transferred these $\delta^{18}\text{O}_{\text{ice}}$ records on AICC2012 using the correspondence between the Vostok GT4 and AICC2012 chronologies (Figure 1).

Finally, the LIG section of NEEM can only be dated using $\delta^{18}\text{O}_{\text{atm}}$ because its CH₄ record is contaminated by in-situ production, in relationship with local summer melt during the LIG (NEEM comm. members, 2013). Figure 1 displays the NEEM $\delta^{18}\text{O}_{\text{ice}}$ record on its depth scale between 2350 and 2490 m, where the linear alignment of depth with AICC2012 is based on the resemblance between EDC and NEEM $\delta^{18}\text{O}_{\text{atm}}$ records.

The continuous NEEM section spanning the LIG ends just after 128 ka (on the AICC2012 timescale). Indeed, the characteristic abrupt increase of CH₄ and high $\delta^{18}\text{O}_{\text{atm}}$ level identified in Antarctic records at 128 ka is absent from the record. This reveals that the NEEM ice core does not encompass any ice from the penultimate deglaciation at that point, similar to GISP2, GRIP or NGRIP (Figure 1). Whether this hiatus arises from the disappearance of this layer due to melt under warm early LIG conditions or due to specific thinning and flow associated with different physical properties of glacial versus transition ice remains to be fully assessed.

2.2 NEEM air $\delta^{15}\text{N}$ record

Relative to the free atmosphere mean value, the $\delta^{15}\text{N}$ value in air trapped in ice cores is influenced by gravitational fractionation directly related to temperature and to the depth at which bubble lock-in occurs. Changes in firn lock-in depth (LID) can be related to changes in surface accumulation and temperature: an increase in temperature leads to a decrease of the LID because of faster metamorphism, while higher accumulation rates lead to an increase of the LID. The development of firn densification models allows to simulate the LID evolution as a function of surface climatic conditions (e.g. Herron and Langway, 1980; Goujon et al., 2003; Li and Zwally, 2004; Helsen et al., 2008; Arthern et al., 2010; Ligtenberg et al., 2015). While there is still a model-data mismatch in cold sites of central Antarctica (Capron et al., 2013), the comparison between firn models and $\delta^{15}\text{N}$ data can be used for testing quantitative temperature and accumulation rate reconstructions in Greenland and high accumulation sites in Antarctica (Guillevic et al., 2013; Kindler et al., 2014; Buizert et al., 2015). During rapid surface temperature

changes in Greenland (e.g. Dansgaard-Oeschger events), $\delta^{15}\text{N}$ is also influenced by thermal fractionation (Severinghaus et al., 1998). However, no rapid $\delta^{18}\text{O}_{\text{ice}}$ changes are found during the LIG, and overall stable NEEM $\delta^{15}\text{N}$ values are also coherent with gravitational fractionation occurring under stable surface accumulation rate and temperature (Figure 2).

The single exception is a negative spike recorded at 2384 m depth, which coincides with the strongest CH_4 spike, as well as a negative excursion of the ^{10}Be record (Sturevik-Storm et al., 2014). We suggest that this singular event reflects positive surface temperatures, leading to intense surface melt and large in-situ CH_4 production (Orsi et al., 2015). Firm air transport and thus $\delta^{15}\text{N}$ are not expected to be significantly affected by melt layers at the surface (Keegan et al., 2014). This probably explains why most of the CH_4 spikes are not associated with any changes in the $\delta^{15}\text{N}$ signal. Still, for the negative 0.07‰ $\delta^{15}\text{N}$ excursion at 2384 m (corresponding to ~121 ka on the NEEM LIG age scale), we propose that positive surface temperatures have led to a sudden shrinking of the firn by about 15 m using the expression of $\delta^{15}\text{N}$ gravitational settling in the firn column:

$$\delta^{15}\text{N} = \exp\left(\frac{g \times LID}{R \times T_{\text{mean}}}\right) - 1 \cong \frac{g \times LID}{R \times T_{\text{mean}}} \quad (1)$$

where g is acceleration due to gravity, R the ideal gas constant and T_{mean} (K) the mean temperature of the firn when bubbles are isolated.

For the rest of the LIG our $\delta^{15}\text{N}$ record shows only very subtle changes and we take the mean $\delta^{15}\text{N}$ value of 0.251 ‰ over the time interval 122-126 ka to obtain a representative mean $\delta^{15}\text{N}$ value (Table 1).

2.3 Spatial structure of isotopic anomalies

To assess the spatial extent of the isotopic anomalies, the magnitude of $\delta^{15}\text{N}$ are compared to the water stable isotope ($\delta^{18}\text{O}_{\text{ice}}$ and deuterium excess, hereafter d-excess) anomalies recorded in different Greenland ice cores between the LIG and pre-industrial (Table 1).

For $\delta^{18}\text{O}_{\text{ice}}$, the strongest increase from pre-industrial to the LIG is recorded at Summit (+3.4 ‰) and NGRIP (+3.3 ‰), and the smallest increase at NEEM (+ 2.1 ‰). However, the NEEM anomaly must be corrected for upstream effects: due to ice flow, the LIG ice at NEEM originates from a 330 m higher upstream location (at the south-east of NEEM) where mean annual $\delta^{18}\text{O}$ today is approximately -35 ‰ (NEEM comm members, 2013). At this deposition site, the LIG isotopic anomaly is therefore 3.6 ‰, close to the value at Summit. We conclude that changes in $\delta^{18}\text{O}_{\text{ice}}$ are rather homogeneously ~ 3.2-3.6 ‰ higher at the LIG than at pre-industrial in central and north-west Greenland (Table 1).

d-excess is not significantly different between the LIG and pre-industrial period at the different drilling sites listed on Table 1. Pre-industrial d-excess values are also very similar among these different sites. This is the reason why extrapolating the surface d-excess values in the NEEM-NGRIP regions enables us to estimate the pre-industrial d-excess at the upstream NEEM deposition site to a conservative value of 11‰.

Assuming that no abrupt climate change took place from 122 to 126 ka (a hypothesis supported by the relatively flat NEEM $\delta^{18}\text{O}_{\text{ice}}$ record), changes in mean $\delta^{15}\text{N}$ are expected to reflect changes in LID. The spatial structure of the $\delta^{15}\text{N}$ changes differs from the pattern of $\delta^{18}\text{O}_{\text{ice}}$. Indeed, the smallest $\delta^{15}\text{N}$ increase is observed at NGRIP (-0.02 ‰) (but the record ends at 120 ka) and the largest one at GRIP and GISP2 (-0.07 to -0.09 ‰), with an intermediate signal at NEEM (-0.04 ‰), albeit with an inherent uncertainty due to the lack of data for the NEEM deposition site today. Because of stratigraphic disruptions, no continuous record is available at Summit. We thus cannot exclude that low $\delta^{15}\text{N}$ levels observed at GRIP and GISP2 on (122-126 ka) ice sections reflect a temporary $\delta^{15}\text{N}$ decrease caused by thermal fractionation or firn shrinking as for the NEEM $\delta^{15}\text{N}$ value at 2384 m (121 ka). We therefore note regional differences for the different available datasets, but stress their heterogeneities (time span, discontinuity, and lack of present-day reference) preventing any robust conclusion.

3 Temperature reconstructions

3.1 Reconstructions based on $\delta^{18}\text{O}_{\text{ice}}$

Today, NW Greenland accumulation is biased towards summer precipitation (based on regional and general circulation atmospheric models; [Steen-Larsen et al., 2011](#)). NEEM summer $\delta^{18}\text{O}_{\text{ice}}$ was monitored through continuous measurements of surface water vapor isotopic composition in 2010-2012 ([Steen-Larsen et al., 2013](#); [Steen-Larsen et al., 2014](#)), revealing a $\delta^{18}\text{O}_{\text{ice}}$ –temperature slope of $0.85 \text{ ‰} \cdot ^\circ\text{C}^{-1}$ at the intra-summer scale. From 1979 to 2007, the increasing trend of $\delta^{18}\text{O}_{\text{ice}}$ recorded in NEEM shallow ice cores was scaled to simulated and estimated local surface air temperature trends, resulting in a multi-decadal slope of $1.05 \pm 0.2 \text{ ‰} \cdot ^\circ\text{C}^{-1}$ for warming above pre-industrial conditions (Masson Delmotte, 2015). These various estimates suggested that the average Holocene $\delta^{18}\text{O}$ -temperature relationship of $0.5 \text{ ‰} \cdot ^\circ\text{C}^{-1}$ based on the calibration with borehole temperature data at other Greenland ice core sites ([Vinther et al., 2009](#); [Vinther et al., 2010](#)) may not be valid for NEEM. Differences between Greenland locations are expected due to changes in the seasonality of precipitation (summer bias at NEEM but not in central or south Greenland), moisture origin as well as possible changes in boundary layer stability and relationships between surface and temperature relationship.

Applying the multi-decadal temporal slope given above, the LIG $\delta^{18}\text{O}_{\text{ice}}$ anomaly at NEEM deposition site translates into a warming of 2.9-4.2°C, twice smaller than the initial estimate based on Holocene calibrations for other sites. Still, it is difficult to assess whether the present-day calibration can apply for the LIG, marked by a different orbital forcing than today, likely with a reduced sea ice extent and different moisture transport pathways (Sime et al., 2013). The second-order isotopic parameter, the d-excess, can provide information on evaporation conditions. Present-day monitoring studies depict low d-excess values for subtropical moisture, contrasting with high d-excess values for moisture from sea ice margin areas (e.g. Steen-Larsen et al, 2015; Pfahl et al, 2014). The d-excess is also affected by distillation, and will decrease in polar regions if $\delta^{18}\text{O}_{\text{ice}}$ increases without any moisture source change. As noted above, available LIG d-excess data (Table 1) are slightly (insignificantly) above pre-industrial levels. A stable or higher d-excess level together with an $\delta^{18}\text{O}_{\text{ice}}$ increase is therefore understood to reflect

a slight poleward shift of moisture sources. In turn, this would imply reduced distillation for $\delta^{18}\text{O}_{\text{ice}}$ and a reduced slope of the relationship between $\delta^{18}\text{O}_{\text{ice}}$ and temperature (Sime et al, 2013) more in line with the average Holocene calibration.

3.2 Reconstructions based on air $\delta^{15}\text{N}$

In the absence of abrupt surface temperature changes, $\delta^{15}\text{N}$ is only affected by the gravitational signal linked to firn LID; the latter is directly related to changes in temperature and accumulation rate. Thus, if accumulation is known, past temperature changes can be inferred from $\delta^{15}\text{N}$. As neither accumulation nor temperature is independently known for the LIG, we have to constrain the accumulation/temperature relationship in the past based on observation and/or models. We now describe the different steps of our procedure to estimate the temperature of the NEEM deposition site during the LIG from $\delta^{15}\text{N}$ measurements.

3.2.1 Different estimates of the link between temperature and accumulation rate in Greenland

i: Accumulation rate and temperature can be linked through thermodynamic laws and ice sheet topography, despite significant uncertainties associated with atmospheric transport characteristics that lead to regional variability (Kaspner et al., 1995; Merz et al., 2014b). In a first approximation, temperature and moisture content of an air mass are linked through saturation pressure (“ P_{sat} -approach”). This first order relationship between accumulation rate and temperature has long been used for Antarctic ice core chronologies (Lorius et al., 1985; Ritz, 1992) with:

$$A(t) = A(t_0) \times \frac{\left[\left(\frac{\partial P_{\text{sat}}}{\partial T} \right) / (T + 273) \right]_t}{\left[\left(\frac{\partial P_{\text{sat}}}{\partial T} \right) / (T + 273) \right]_{t_0}} \quad (2)$$

where $A(t)$ and $A(t_0)$ are the accumulation rates at time t and t_0 respectively, P_{sat} the saturation pressure over ice and T the temperature in °C.

ii: Empirical relationships between accumulation rate and Greenland temperature have been provided by different methods. The dataset obtained in 1952-1955 by Benson (1962) remains a reference today for evaluating surface accumulation rate reconstructions above the Greenland ice sheet (e.g. Hawley et al., 2014; Munk et al., 2003). Surface accumulation rate and temperature data from 146 sites show an exponential increase of accumulation rate versus temperature. Within the associated 1σ envelope, it encompasses the accumulation rate versus temperature increase deduced from the “ P_{sat} -approach”. This spatial relationship between accumulation rate and temperature is however associated with a very large uncertainty envelop challenging the validity of any relationship between temperature and accumulation in Greenland.

iii: Based on the GISP2 ice core records over the last deglaciation, Kaspner et al. (1995) showed that the relationship between Greenland accumulation rate and temperature was not stable because of variations in atmospheric circulation. Still, they were able to propose a temporal relationship between accumulation rate based on annual layer counting and $\delta^{18}\text{O}$, or temperature reconstructed from $\delta^{18}\text{O}$ and a calibration based on borehole temperature measurements (leading to a $\delta^{18}\text{O}$ vs temperature slope of $0.53\text{‰} \cdot ^\circ\text{C}^{-1}$). The inferred sensitivity of snow

accumulation rate to temperature change during interglacial period varies from $0.9\%.\text{°C}^{-1}$ (Holocene) to $7.5\%.\text{°C}^{-1}$ (Bølling-Allerød), with an uncertainty encompassing zero.

More recently, Buchardt et al. (2012) followed a similar approach but using numerous ice cores. They used an array of 52 shallow ice cores spanning the last decades to centuries with accumulation rate estimates from annual layer counting on $\delta^{18}\text{O}_{\text{ice}}$ profiles. They identified different temperature vs accumulation rate relationships from one region to another. In central and north Greenland corresponding to the location of the NEEM, NGRIP, GRIP and GISP2 deep ice cores, the Buchardt approach suggests a sensitivity of 1.5 to $9.4\%.\text{°C}^{-1}$ with an uncertainty encompassing zero. This sensitivity is obtained with a $\delta^{18}\text{O}$ vs temperature sensitivity slope of $0.67\%.\text{°C}^{-1}$ so that the Kapsner and Buchardt estimates agree on a 0 to $14\%.\text{°}^{-1}$ accumulation rate vs $\delta^{18}\text{O}$ sensitivity.

The results of the “Buchardt approach” lie within the 1σ envelope of the “Benson approach” (Figure 3). For the Northwestern Greenland area, in which the NEEM and NGRIP sites are located, the lower end of the accumulation rate/temperature sensitivity in equation (2) applies (Buchardt et al., 2012). The lower line of the Buchardt approach in Figure 3 is hence the most likely accumulation/temperature sensitivity for the NEEM ice core. Note that one limitation needs to be stressed for the Buchardt and Benson approaches: the application of a spatial accumulation/temperature relationship makes the assumption that this sensitivity can be also applied to a temporal change, an assumption that, similar to the case of the spatial $\delta^{18}\text{O}$ /temperature gradient, must not necessarily hold back in time.

iv: Masson-Delmotte et al. (2015) used estimates of snow accumulation rate and ice $\delta^{18}\text{O}$ in four shallow cores in the NEEM area together with accumulation rate, temperature and when possible snowfall $\delta^{18}\text{O}$ reconstructions from different models simulation (ECHAM5, Global Climate Model developed by the Max Planck Institute for Meteorology; LMDZ, Global Climate Model developed by the Laboratoire de Météorologie Dynamique; MAR, Regional Atmosphere Model) nudged to available climate reanalyses over the 1979 – 2007 period. In addition to model outputs or temperature gridded reconstruction (Box et al., 2009), the amplitude of temperature increase at NEEM can also be estimated using borehole temperature measurements. Gathering the different sources of information for the strong warming period of 1979-2007 leads to a relatively high slope between accumulation rate and temperature (10 to $15.9\%.\text{°C}^{-1}$, the highest value being obtained using outputs from the MAR model nudged to ERA-40 and ERA-Interim reanalyses (Uppala et al., 2005; Dee et al., 2011)). In this study, the sensitivity of accumulation rate vs. $\delta^{18}\text{O}$ can be estimated through the regression between the NEEM $\delta^{18}\text{O}$ and accumulation rate increases over the period 1979 – 2007 leading to a value of $10\%.\text{°}^{-1}$, in agreement with the Buchardt estimate. Another solution is to use the $\delta^{18}\text{O}$ vs temperature estimate based on NEEM $\delta^{18}\text{O}$ measurements vs borehole temperature over the recent warming trend ($0.8\%.\text{°C}^{-1}$) together with the accumulation rate vs temperature estimate given above, hence leading to a maximum accumulation rate vs $\delta^{18}\text{O}$ sensitivity of $13\%.\text{°}^{-1}$, again within the range of the Buchardt estimate . These high slopes are mainly linked to atmospheric model outputs (for temperature or accumulation), and we regard them as an upper limit. By reducing the data used to infer the accumulation rate versus temperature slope to observations (NEEM accumulation rate deduced from shallow

cores, borehole profile inversion, gridded reconstruction or instrumental temperature measurements from the Greenland southwestern coast), the slope is only of $9 \pm 1\% \cdot ^\circ\text{C}^{-1}$ which, in this case, is consistent with the “Buchardt approach”.

v: Alternative estimates of accumulation rate at the NEEM deposition site are also provided by ^{10}Be data. Sturevik-Storm et al. (2014) compiled mean Holocene ^{10}Be concentration over eight Greenland sites and determined the spatial relationship against mean accumulation rate estimates. This relationship was then applied to LIG ^{10}Be data from NEEM, leading to the conclusion that accumulation rate was 65-90% higher than today at the NEEM deposition site (“ ^{10}Be -approach”). However, the present-day spatial gradients in ^{10}Be concentration are caused by spatially varying contributions of wet deposition to the overall ^{10}Be deposition, assuming a homogeneous atmospheric ^{10}Be aerosol deposition over Greenland. This latter assumption implies that the atmosphere above Greenland is well mixed with respect to ^{10}Be after transport from the location of troposphere/stratosphere foldings. The latter are the main entrance pathways of stratospheric aerosols into the extratropical northern hemisphere troposphere. The LIG climate is characterized not only by likely enhanced precipitation above Greenland (accumulation) but also higher wet deposition during aerosol transport to Greenland due to higher precipitation rates. As a result, higher scavenging of ^{10}Be bearing aerosol en route must have led also to a lower atmospheric ^{10}Be concentration over Greenland than today. The LIG accumulation estimate by Sturevik-Storm et al. (2014) is therefore most likely an overestimation and the assumption of Sturevik-Storm et al. (2014) that ^{10}Be concentration is only controlled by accumulation rate at the NEEM site may be challenged. More generally, the use of other chemical aerosol species as accumulation rate tracers is hampered by potential changes in the LIG atmospheric concentrations due to emission changes. Qualitatively, a correction of deposition effects using the Buchardt et al. (2012) approach representative for Northwest Greenland, leads to LIG atmospheric concentrations of all chemical aerosol tracers similar to today. In contrast much higher LIG accumulation rates as estimated by Sturevik-Storm et al. (2014) or no changes in accumulation between the LIG and the Holocene imply an unrealistic change in atmospheric aerosol concentrations for several aerosol tracers. Indeed, if we compare aerosol species that are dominated by wet deposition such as Na^+ (sea salt aerosol) and NO_3^- (lightning activity, biological activity), we see that the concentration in the ice in the LIG is lower than in the Holocene (about 50%). As these species are mainly wet deposited, this is only possible if the atmospheric aerosol concentration was also reduced by 50% at that time. It is unlikely that both Na^+ and NO_3^- (which have completely different sources and transport pathways) have a 50% reduction of source emissions. Another solution to explain the reduction of 50% in atmospheric Na^+ and NO_3^- concentration over the ice is to imply an increase of the precipitation rate along the transport pathway and thus increase of wet deposition en route. The chemistry suggests that the precipitation rate during transport in the Northern Hemisphere was significantly higher during the LIG than during the Holocene and there is no reason why Greenland would not be affected by this general increased accumulation rate. Based on the changes in various chemical tracers in the ice (sea salt aerosol, biogenic aerosol, mineral dust) we thus conclude that the LIG

accumulation was likely 20 % higher than pre-industrial and similar to the Buchardt approach for Northwest Greenland.

vi: Atmospheric general circulation model outputs do not suggest important changes in accumulation rate for the LIG compared to pre-industrial values, in line with the small simulated change in annual mean surface temperature (section 3.3). A comparison of some of the model outputs (presented in Lunt et al., 2013) shows very limited accumulation increase (less than 5%) over central Greenland. Stronger increases in accumulation rate at LIG associated with significantly warmer than pre-industrial temperature were obtained in relation with a reduction of sea ice in the Nordic Seas (10% increase in accumulation, Merz et al., 2016). Finally, it has been shown that the geometry of the Greenland ice sheet and topographic changes can lead to various local accumulation scenarios for the LIG at the upstream NEEM deposition site (Merz et al., 2014b): depending on the prescribed LIG ice sheet topography, the modelled accumulation rate at LIG can be 25% lower to 13% higher than the pre-industrial accumulation rate. The lowest estimate is linked to a change in the trajectory of air mass to the NEEM deposition site, with an increased eastward origin. The validity of such a scenario could be assessed by comparing simulated and measured δ -excess variations. The data presently available shows similar δ -excess levels at the LIG and pre-industrial periods, and therefore does not support a significant change in moisture source and trajectory (Table 1). Often associated with relatively small temperature changes, these modelled accumulation rate scenarios for the LIG are on the lower end of the accumulation rate scenarios discussed above **and encompass the possibility of a scenario with no change in the accumulation rate**, ~~and in contradiction with the recent modelled accumulation evolution depicted by Masson-Delmotte et al. (2015).~~

~~From the limitations of each accumulation rate estimates highlighted above, we conclude that the subset of relationships with the highest accumulation rate temperature sensitivity may overestimate the accumulation response, and are less reliable. Accordingly, we argue the saturation pressure and the lower end of the Buchardt approach to be more likely to bracket the true LIG accumulation rate increase.~~

3.2.2 Measured versus modeled evolution of $\delta^{15}\text{N}$ with respect to temperature and accumulation rate changes

Our $\delta^{15}\text{N}$ data is compared with those simulated using a firn densification model forced by these different accumulation rate versus temperature relationships for the LIG. The firnification model relates LID to accumulation rate and temperature. Here, we use the Goujon et al. (2003) model in steady state to calculate LID, and the barometric equation (1) to translate LID changes into $\delta^{15}\text{N}$ changes.

The model correctly captures the present-day $\delta^{15}\text{N}$ values for NEEM and NGRIP, using the current mean values for accumulation rate and temperature (Figure 3). At NEEM and NGRIP, firn studies have recently provided an accurate determination of the LID and $\delta^{15}\text{N}$ profiles (Guillevic et al., 2013; Buizert et al., 2013). At GISP2, there exists no proper determination of the LID due to discontinuous sampling of air bottles and a large scatter of $\delta^{15}\text{N}$ values measured at the bottom of the firn, ranging between 0.305 and 0.325 ‰ (Bender et al., 2006). An average value of 0.31‰ is obtained from high-resolution $\delta^{15}\text{N}$ measurements over the last 4000 yr on the GISP2 core

(Kobashi et al., 2008). For present-day, our simulation at GISP2 ($\delta^{15}\text{N}$ of 0.325‰) therefore lies at the upper limit of available measurements (Table 1).

3.2.3 Reconstructing Greenland LIG temperature

In order to estimate the LIG firn temperature (at the deposition sites), Figure 3 displays the $\delta^{15}\text{N}$ data points for each ice core site on contours of the simulated $\delta^{15}\text{N}$ values as a function of temperature and accumulation. In addition, we display the different accumulation rate estimates detailed in section 3.2.1. We use the different relationships for accumulation rate vs $\delta^{18}\text{O}$ sensitivity together with the 2.1 ‰ increase between the present-day $\delta^{18}\text{O}$ at NEEM and the LIG $\delta^{18}\text{O}$ at the NEEM deposition site to infer a range of accumulation rates for the NEEM LIG compatible with the approaches of Buchardt et al. (2012), Kapsner et al. (1995) and Masson-Delmotte et al. (2015). We indicate the possibility for no change in accumulation rate. We also display the accumulation rate vs temperature relationships from the P-sat approach and from Masson-Delmotte et al. (2015). Note that the highest accumulation rate vs temperature relationship obtained from outputs of the MAR simulation nudged to ERA reanalyses cannot be reconciled with NEEM $\delta^{15}\text{N}$ measurements at LIG and is thus discarded in the following.

For the NEEM LIG deposition site, we detail below the graphical determination of the temperature of the firn column. The intersection between the $\delta^{15}\text{N}$ contour and the “accumulation rate vs temperature” evolution curve or the horizontal lines showing LIG possible accumulation rate levels gives the range of realistic LIG accumulation rate (y-axis) and temperature (x-axis). Note that the “Benson approach” is scattered, due to local surface topography effects, and therefore not directly relevant for the NEEM LIG temperature reconstruction. We have used this “Benson approach” only for validation of the other accumulation rate vs temperature relationships.

Let us assume that the LIG accumulation rate at the NEEM deposition site was the same as today at NEEM, despite increasing temperature. In this conservative but unrealistic case, our $\delta^{15}\text{N}$ data point to a 3.5°C warmer firn column. Assuming a 20 % accumulation increase at the NEEM deposition site leads to an estimate of 4.5°C surface warming between the NEEM upstream deposition site and the current NEEM firn temperature. From the intersection between the $\delta^{15}\text{N}$ level measured in the NEEM LIG section and the accumulation rate estimates from the “P_{sat}-approach” or the maximum accumulation rate deduced from Buchardt et al. (2012) and Kapsner et al. (1995), we obtain a larger estimate of 6-7°C warming of the firn column at the NEEM LIG deposition site compared to the current NEEM firn temperature. This corresponds to an accumulation rate of 26-30 cm water equivalent yr⁻¹, i.e. 32-50% higher than the present-day accumulation rate at NEEM. The highest LIG warming compatible with the $\delta^{15}\text{N}$ data (almost -20°C, i.e. about 9°C above present-day NEEM values at the upstream NEEM deposition site) corresponds to an accumulation rate of 46 cm water equivalent.yr⁻¹, i.e. 130% higher than the present-day accumulation rate at NEEM, using the slope for the relationship between accumulation rate vs temperature suggested in Masson-Delmotte et al. (2015) from the temperature reconstruction with MAR and accumulation rate from NCEP reanalyses. For the ¹⁰Be approach, which represents an upper limit of the possible accumulation increase we find a LIG temperature 7-8°C warmer than at the current drilling site. When corrected for the change of deposition site, this translates to a 6-11°C higher temperature at the NEEM deposition site for LIG compared to pre-industrial temperature level.

At NGRIP, the same graphical approach leads to an estimated temperature of -28.5 to -24°C at the end of the LIG (120 ka on the AICC2012 timescale) compared to -31.5°C for pre-industrial, i.e. a difference of $+5.2\pm 2.3^{\circ}\text{C}$ between 120 ka and pre-industrial. Even if NGRIP is not on a dome, the upstream effect is quite small: the NGRIP LIG deposition site is estimated to lie 48 km upstream in the direction of Summit with small associated altitude gradients between NGRIP and Summit (Buchardt, 2012). Similar to the NEEM temperature reconstruction, the NGRIP LIG temperature uncertainty range arises from the uncertainty in the accumulation rate vs temperature relationship. The full range of estimates for the accumulation difference at NGRIP from the preindustrial to the LIG is hence estimated to be 0 - 100% of the preindustrial value. This $+5.2\pm 2.3^{\circ}\text{C}$ warming at 120 ka may be an underestimation of the full warming range encompassed during the LIG, because the NGRIP ice core does not extend towards the warmest part of the LIG. At the NEEM deposition site however, the temperature estimated following the graphical method of Figure 3 from the $\delta^{15}\text{N}$ value at 120 ka (0.256 ‰) is only 0.5°C lower than the estimated LIG optimum temperature, hence very comparable to the estimated LIG optimum temperature. In summary, the NGRIP LIG vs pre-industrial temperature increase ($+5.2\pm 2.3^{\circ}\text{C}$) is thus on the lower end but still compatible within error bars with the NEEM LIG vs pre-industrial temperature increase ($+8\pm 2.5^{\circ}\text{C}$ at 120 ka using the aforementioned $\delta^{15}\text{N}$ value of 0.256 ‰). Several explanations can explain this discrepancy. First, the dating of the NGRIP and NEEM bottom parts is difficult because of the lack of precise relative and absolute age markers; this limits our confidence that we indeed compare

$\delta^{15}\text{N}$ levels of same age on the two different cores. Second, we do not have any evidence of melt layers in the bottom of the NGRIP core, opposite to NEEM. This suggests that the NEEM deposition site was indeed warmer than NGRIP at LIG but also suggests that firn densification may have been affected by this process at NEEM which would bias our reconstruction. Finally, our temperature reconstruction at NEEM is complicated by the fact that the NEEM LIG deposition site and NEEM drilling site have different surface conditions. In particular, an estimate of the pre-industrial accumulation rate is missing at the NEEM deposition site.

At Summit, only very high temperatures can be reconciled with LIG $\delta^{15}\text{N}$ values 0.7 to 0.9‰ lower than today: -20°C to -16°C according to graphical determination using Figure 3 compared to a pre-industrial temperature of -31.7°C . Still, without a continuous sequence of interglacial ice at Summit, the true origin of the $\delta^{15}\text{N}$ signal at Summit is doubtful. We cannot yet assess whether this signal is purely gravitational or whether it is dominated by a thermal signal or a firn hiatus effect of ~ 16 m as estimated from the barometric equation (1).

Finally, note that the present-day values in accumulation, temperature and $\delta^{15}\text{N}$ at NGRIP and NEEM nicely align with the accumulation-temperature relationship from the P_{sat} approach (Figure 3). At Summit, the current accumulation rate is significantly higher than expected from this estimate, indicating that other advective moisture pathways come into play, consistent with analyses of spatial influences of weather regimes in Greenland. Indeed, Ortega et al (2014) investigated the influence of the main North Atlantic weather regimes in Greenland, and stressed differences between Summit (affected by NAO+ and NAO- weather regimes) and North / North-West Greenland (more strongly affected by e.g. the North Atlantic Ridge regime). Accordingly, we do not include the Summit values in our conclusions.

3.2.4 Limitations of the $\delta^{15}\text{N}$ based temperature reconstruction at the NEEM sites

In the following, limits inherent to this $\delta^{15}\text{N}$ approach are highlighted, which shall motivate further studies to refine the temperature estimate. First, we have applied a firnification model optimized for present-day central Greenland firn to past periods with different, warmer conditions, outside the range of model validation. For instance, the occurrence of substantial summer melt could accelerate firn densification and produce a smaller close-off depth (and therefore smaller $\delta^{15}\text{N}$ values) than expected from the Goujon model for a given temperature. In principle, the validity of firn models in such temperature range can be tested, if firn studies are performed at Greenland sites which are today warmer than the central deep drilling sites of Summit, NGRIP and NEEM, but no data are yet available. Second, the relative changes in accumulation rates and temperature between the NEEM deposition site and NEEM remain difficult to estimate. ~~While we have gathered all the available information on the spatial and temporal relationships between surface temperature and accumulation rates, other influences need to be ideally considered, such as changes in regional atmospheric circulation associated with a different climate and modifications in ice sheet topography (Merz et al., 2014a,b) and/or sea ice extent (Merz et al., 2016).~~ Finally, we have identified a **negative peak** in the NEEM LIG $\delta^{15}\text{N}$ profile, at 2384 m (121 ka), with no parallel signal in $\delta^{18}\text{O}_{\text{ice}}$ or in chemical records. This signal challenges our attribution of $\delta^{15}\text{N}$ variations solely to changes in accumulation rate and/or temperature, and suggests potential influence of surface melt on firn depth, LID and therefore $\delta^{15}\text{N}$. While the overall stability of the NEEM $\delta^{15}\text{N}$ record over the LIG supports a gravitational / climatic interpretation, a dominant influence of surface melt **explaining a removal of snow of 16 m depth** cannot be excluded for the Summit ice core sections associated with very low $\delta^{15}\text{N}$ (0.23 ‰).

3.3 LIG temperatures in Greenland as estimated by climate models

The LIG climate has been simulated by a suite of climate models of various complexities. Most of these simulations are included in the model intercomparison studies of Lunt et al. (2013) and Bakker et al. (2013, 2014). The former study compared equilibrium (“snap-shot”) simulations, covering time-slices within the early LIG (125-130 ka), to temperature proxy data, whereas the latter studies discussed transient simulations covering the entire LIG. **The equilibrium simulations are all described in detail in Lunt et al. (2013): they comprise 11 General Circulation Models (GCM), and 3 Earth System Models of Intermediate Complexity (EMIC). The resolution varies between T21 (~5.6 x 5.6 degrees) for the EMICs to T85 (~1.4 degrees) for the highest resolution model. All the results are interpolated to grids of 3.75° (longitude) x 2.5° (latitude). The transient simulations are described in details in Bakker et al. (2013 and 2014). These are also a combination of GCMs and EMICs, but the highest resolution is only about 3.75°. The main forcing for all the simulations is though orbital parameters and greenhouse gasses. Note that in the majority of the simulations the land-sea configuration, ice sheet extent and height are kept the same as in their respective pre-industrial simulations.**

The model mean of all equilibrium simulations computes an annual mean temperature increase over Greenland between 0 and 2°C with respect to pre-industrial control simulations (Figure 6a of Lunt et al., 2013). At the

upstream NEEM deposition site the same range of annual mean temperature increase is found when analysing the individual model simulations for the 125 ka time slice (0.2-2.2°C, Figure 4). The majority of the transient models simulate a maximum early LIG temperature increase of similar magnitude as the equilibrium model simulations.

This result is not sensitive to the choice of the grid point in the model: when the same analysis is performed for the grid boxes surrounding the NEEM upstream grid box, the maximum of 2°C from LIG external forcing is still valid.

Two exceptions to this 0-2°C warming are (1) the MPI-UW model that computes an annual mean temperature increase as high as ~4°C around 126 ka and (2) the transient simulation of CCSM3, which consistently simulates lower annual mean temperatures for the LIG compared to pre-industrial.

The LIG temperature and precipitation patterns produced by climate models can be applied as a forcing to ice sheet models simulating the LIG evolution of the Greenland ice sheet (e.g. Otto-Bliesner et al., 2006; Robinson et al., 2011, Born and Nisancioglu, 2012; Stone et al. 2013). Several uncertain parameters within these ice sheet models need to be considered (e.g. basal sliding parameter). This, in combination with uncertainties in the schemes translating the large-scale climate forcing to the local Greenland mass balance schemes, results in a large ensemble of possible melting scenarios for the Greenland ice sheet for each individual ice sheet model study. Present-day observations and paleo proxy data help to reduce this large spread. In particular, the paleo information of limited surface elevation reduction at the central ice core locations strongly constrains the simulated LIG Greenland ice sheet evolution (Masson-Delmotte et al., 2013).

Here, we investigate the annual mean surface temperature anomaly at the upstream depositional site of NEEM for simulations fulfilling the (paleo) data constraints in the ice sheet modelling studies of Stone et al. (2013). For these simulations, the corresponding modelled annual mean surface temperature anomaly at the upstream NEEM depositional site is around 1-4°C above pre-industrial. The annual mean LIG temperature forcing applied in these two studies is similar (Figure 4, HadCM3_Bris vs. NorESM_BCCR), whereas the summer temperature anomaly is about 1°C larger over central Greenland in Stone et al. (2013). Also, the ice sheet models and the methods for calculating the surface mass balance from the climate forcing are different. However, the surface temperature anomaly scenario in these studies is mainly restricted by the limited surface elevation reduction during the LIG implied by NEEM data (NEEM community members, 2013). Assuming this elevation reconstruction to be correct, ice sheet simulations associated with a surface temperature anomaly higher than approximately 4-5°C are interpreted to reflect a too large elevation lowering at the ice core locations, and are therefore rejected.

In the modeling studies described above, the feedback of changes in surface elevations on the climate is either not included (surface topography is kept fixed) or not well resolved due to coarse model resolution (Stone et al., 2013). However, Merz et al. (2014a) showed that a steeper surface slope can cause an additional 1-3°C surface air temperature increase due to an increase in katabatic winds which foster downward flux of sensible heat. In order for this effect to be important for NEEM, the ice sheet geometry needs to change such that its LIG depositional site is closer to the rim of the ice sheet, e.g. with large melting of northeast Greenland.

Confirming the suggestion of Sime et al (2013), Merz et al. (2016) used the CCSM3 and CCSM4 models to demonstrate that the large spread in Greenland SAT change among the LIG equilibrium simulations (Figure 4, Lunt et al. (2013)) is mostly due to differences in simulated sea-ice extent. They showed that surface air temperature and accumulation changes at the NEEM LIG deposition site are particularly sensitive to sea-ice retreat in the Nordic Seas.

In summary, studies performed with atmospheric models suggest that a number of processes may combine to produce larger amplitude of warming as simulated in state-of-the-art coupled climate models. They suggest that Greenland LIG surface air temperature change can be amplified in response to regional sea ice retreat and in response to change in the ice sheet topography. For the NEEM deposition site, LIG annual mean surface temperatures of approximately 5°C above pre-industrial can be obtained: 2°C being due to LIG external forcing (orbital and greenhouse gases), 0.6 to 2.3°C to be attributed to a decrease of sea ice in the Nordic Seas and 1 to 2 °C associated with a moderately smaller GrIS. Higher temperatures might be possible in more extreme scenarios, however then it is unlikely that the ice sheet can maintain surface elevations relatively close to modern for all central deep ice core locations as suggested by the ice core records. The coherency between scenarios of large surface air temperature warming and the plausibility of the Greenland ice sheet response still remains to be fully explored. This includes specific analyses of the seasonal aspects of the surface air temperature change but also scrutinizing the elevation change reconstruction of the Greenland Ice Sheet during the LIG based on air content measurements (NEEM community members, 2013).

4. Conclusions and perspectives

In this study, we have compiled the Greenland ice core data and methods available to quantify LIG temperature change. New estimates of temperature change were provided based on $\delta^{15}\text{N}$ and firn densification modeling, largely independent from water stable isotope data. They imply that the mean annual firn temperature at the LIG deposition site, upstream of the current NEEM site, experienced 6-11°C warming, without correcting for changes in elevation related to ice thickness change. As a comparison, the initial estimate of NEEM community members (2013) based on $\delta^{18}\text{O}$ -temperature relationships calibrated on Holocene data led to a LIG surface air temperature at the upstream NEEM deposition site 5.7-9.3°C warmer than at pre-industrial (NEEM comm. members, 2013). ~~The approach based on water stable isotopes strongly depends on the calibration of the isotope temperature relationship. NEEM (2013) used Holocene calibrations based on central Greenland ice cores (0.5 ‰.°C⁻¹). However, Masson-Delmotte et al (2015) showed that water stable isotopes were twice more sensitive to temperature changes at NEEM during recent decades than inferred for the Holocene in central Greenland. Alternatively, we can independently quantify the isotope temperature relationship using our new data: the $\delta^{15}\text{N}$ temperature reconstruction of a 3.5 to 8.5°C temperature change for a 2.1‰ change between NEEM pre industrial and the NEEM deposition site at LIG (hence 2 different sites at 2 different periods) leads to a slope of the isotope temperature slope (0.25 to 0.6 ‰.°C⁻¹), even smaller than the result of the central Greenland Holocene calibration, and comparable to the LGM vs Holocene temporal slope in Greenland (0.3 ‰.°C⁻¹) (Cuffey et al., 1995; Dahl Jensen et al., 1998). This finding implies that~~

~~the isotope temperature relationship inferred for recent warming (associated with e.g. sea ice retreat in the Baffin Bay area) does not apply for warmer than today LIG conditions. Such a low slope is also captured in some isotopic enabled atmospheric models for LIG conditions (Masson Delmotte et al, 2011), but results appear dependent on patterns of sea surface temperature changes and associated moisture origin changes (Sime et al, 2013). Indirectly, our constraints on the isotope temperature relationships can therefore help to test the realism of large scale climate patterns (including atmospheric transport response to sea surface temperature and sea ice changes) simulated in response to orbital forcing, beyond just Greenland temperature.~~

In addition to the $\delta^{15}\text{N}$ temperature reconstruction, ice core data provide multiple lines of evidence of significantly warmer conditions, at least during summer, at the upstream NEEM deposition site during the LIG than today at the NEEM site such as in-situ CH_4 production likely due to summer melt, firn shrinking suggested by the $\delta^{15}\text{N}$ peak at 2384 m.

However, it shall be noted that our reconstruction is associated with large uncertainties because the Greenland accumulation rate during the LIG is not precisely constrained. In addition, uncertainties are enhanced at NEEM because of the distance between the locations of the NEEM drilling site and the NEEM deposition site. The NGRIP site was less affected by ice flow and the temperature estimate is 3-7.5°C higher during the LIG than for pre-industrial period. This LIG temperature estimate corresponds to the lower limit of the LIG temperature estimate for the NEEM deposition site and suggests that the upper limit of this temperature reconstruction at the NEEM deposition site is not realistic, because of a too high accumulation rate.

The evidence of summer melt in the LIG section of the NEEM core is an additional limitation of our approach for the NEEM temperature reconstruction that shall be taken into account. Melting-refreezing, which accelerates firnification processes, is not included in our firnification model, which, strictly speaking, is only valid for the dry snow zone. Still, the occurrence of extensive summer melt implies that mean summer temperatures at the site of deposition frequently reached the melting point, which is about 5°C higher than pre-industrial mean summer temperatures at NEEM. Accordingly, the LIG surface conditions were very similar to the extraordinary 2012 heat wave over Greenland, which led to substantial surface melting at the NEEM site.

Large warming at the NEEM deposition site is difficult to reconcile with climate simulations in response to orbital forcing and greenhouse gases concentration forcing only, when keeping the LIG ice sheet thickness similar to today's value. State-of-the-art intermediate complexity or fully coupled climate models mostly produce an annual mean temperature increase of less than 2°C above pre-industrial present-day during the LIG in NW Greenland. However, sea ice cover retreat in the Nordic Seas and changes in Greenland ice sheet topography may significantly enhance surface warming (Merz et al., 2014a; 2016) and therefore reduce the gap with our estimate. Vice-versa, relaxing the ice sheet thickness constraint derived from the ice core measurements (NEEM community members, 2013) allows for much stronger surface warming at the LIG deposition sites. ~~However, annual mean warming by 5°C or more leads to simulations of larger ice sheet retreat than implied by data constraints on the LIG ice sheet elevation: the NEEM paradox still needs to be solved, possibly through a stronger focus on summer rather than annual mean temperature change.~~

Further work is required to overcome the unavoidable limitations of firm our temperature estimate. New firm monitoring studies in Greenland areas affected by summer melt and in today's ablation zone are crucially needed to improve firm modeling and interpretation of the $\delta^{15}\text{N}$ signal, especially for the sharp anomaly suggested to reflect 15 m firm shrinking. Similarly, monitoring of water stable isotopes in the Arctic water vapor is also critical to better understand and model the relationships between atmospheric circulation, moisture transport pathways, snow-vapour isotopic exchanges and the isotopic composition above the Greenland ice sheet.

Acknowledgments

NEEM is directed and organized by the Centre of Ice and Climate at the Niels Bohr Institute and US NSF, Office of Polar Programs. It is supported by funding agencies and institutions in Belgium (FNRS-CFB and FWO), Canada (NRCan/GSC), China (CAS), Denmark (FIST), France (IPEV, CNRS/INSU, CEA and ANR), Germany (AWI), Iceland (RannIs), Japan (NIPR), South Korea (KOPRI), The Netherlands (NWO/ALW), Sweden (VR), Switzerland, the United Kingdom (NERC) and the USA (USNSF, Office of Polar Programs) and the EU Seventh Framework programs. The Division for Climate and Environmental Physics, Physics Institute, University of Bern, acknowledges continuing support by the Swiss National Science Foundation as well as access to the infrastructure at the Swiss National Supercomputing Centre. The HadCM3 climate model simulations were carried out using the computational facilities of the Advanced Computing Research Centre, University of Bristol - <http://www.bris.ac.uk/acrc/>. This project benefitted from support from Past4Future 'Climate change - Learning from the past climate', a Collaborative Project under the 7th Framework Programme of the European Commission (grant agreement no. 243908). P.M.L. is funded by the Research Council of Norway through the IceBed project (221598). This work was supported by European Research Council Advanced Grant no. [246815](#) *WATERundertheICE*. The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement n°306045 (COMBINISO) and from Agence Nationale de la Recherche (ANR-07-VULN-0009 and ANR-10-CEPL-0008).

References

- Bakker P., Masson-Delmotte V., Martrat B., Charbit S., Renssen H., Gröger M., Krebs-Kanzow U., Lohmann G., Lunt D.J., Pfeiffer M., Phipps S.J., Prange, M., Ritz S.P., Stenni, B., Stone E.J. and Varma V.: Temperature trends during the Present and Last Interglacial periods – a multi-model-data comparison, *Quaternary Science Reviews*, 99, 224-243, 2014.
- Bakker P., Stone E.J., Charbit S., Chroger M., Krebs-Kanzow U., Ritz S.P., Varma V., Khon V., Lunt D.J., Mikolajewicz U., Prange M., Renssen H., Schneider B. and Schulz M.: Last interglacial temperature evolution - a model inter-comparison, *Climate of the past*, 9, 605–619, 2013.

Bazin L., Landais A., Lemieux-Dudon B., Toyé Mahamadou Kele H., Veres D., Parrenin F., Martinerie P., Ritz C., Capron E., Lipenkov V., Loutre M-F., Raynaud D., Vinther B., Svensson A., Rasmussen S.O., Severi M., Blunier T., Leuenberger M., Fischer H., Masson-Delmotte V., Chappellaz J. and Wolff E.W.: An optimized multi-proxy, multi-site Antarctic ice and gas orbital chronology (AICC2012): 120–800 ka, *Climate of the Past* 9:1715-1731, 2013.

Bender M., Floch G., Chappellaz J., Suwa M., Barnola J-M., Blunier T., Dreyfus G., Jouzel J. and Parrenin F.: Gas age-ice age differences and the chronology of the Vostok ice core, 0-100 ka, *Journal of Geophysical Research* 111, doi:10.1029/2005JD006488, 2006.

Benson C.S., Stratigraphic studies in the snow and firn of the Greenland ice sheet. SIPRE Res. Rep. 70, 76–83, 1962.

Born A. and Nisancioglu K.H.: Melting of Northern Greenland during the last interglaciation. *The Cryosphere* 6, 6, 1239-1250, 2012

Box J.E., Yang L., Bromwich D.H., Bai L-S.: Greenland Ice Sheet Surface Air Temperature Variability: 1840–2007, *Journal of Climate*, 22, 14, 4029-4049, 2009

Buchardt, S. L., Clausen, H. B., Vinther, B. M., and Dahl-Jensen, D.: Investigating the past and recent $\delta^{18}\text{O}$ -accumulation relationship seen in greenland ice cores, *Clim. Past*, 8, 2053-2059, 10.5194/cp-8-2053-2012, 2012.

Buizert, C., Martinerie, P., Petrenko, V. V., Severinghaus, J. P., Trudinger, C. M., Witrant, E., Rosen, J. L., Orsi, A. J., Rubino, M., Etheridge, D. M., Steele, L. P., Hogan, C., Laube, J. C., Sturges, W. T., Levchenko, V. A., Smith, A. M., Levin, I., Conway, T. J., Dlugokencky, E. J., Lang, P. M., Kawamura, K., Jenk, T. M., White, J. W. C., Sowers, T., Schwander, J., and Blunier, T.: Gas transport in firn: multiple-tracer characterisation and model intercomparison for NEEM, Northern Greenland, *Atmos. Chem. Phys.*, 12, 4259-4277, doi:10.5194/acp-12-4259-2012, 2012.

Capron E., Govin A., Stone E.J., Masson-Delmotte V., Mulitza S., Otto-Bliesner B., Rasmussen T.L., Sime L.C., Waelbroeck C. and Wolff E.W.: Temporal and spatial structure of multi-millennial temperature changes at high latitudes during the Last Interglacial. *Quaternary Science Reviews*, 103, 116-133, 2014.

Capron E., Landais A., Chappellaz J., Buiron D., Fischer H., Johnsen S.J., Jouzel J., Leuenberger M., Masson-Delmotte V., Stocker T.F.: A global picture of the first abrupt climatic event occurring during the last glacial inception. *Geophysical Research Letters* 39(L15703), doi:10.1029/2012GL052656, 2012.

Capron, E., Landais, A., Lemieux-Dudon, B., Schilt, A., Masson-Delmotte, V., Buiron, D., Chappellaz, J., Dahl-Jensen, D., Johnsen, S., Leuenberger, M., Loulergue, L., and Oerter, H.: Synchronising EDML and NorthGRIP ice cores using

d18O of atmospheric oxygen and ch4 measurements over MIS5 (80-123 ka), *Quaternary Science Reviews*, 29, 235-246, 2010.

Cuffey K.M. and Clow G.D.: Temperature, accumulation and ice sheet elevation in central Greenland through the last deglacial transition, *Journal of Geophysical Research*, 102, C12, 26383-26396, 1995.

Dahl-Jensen D., Mosegaard K., Gundestrup N., Clow G.D., Johnsen S.J., Hansen A.W. and Balling N.: Past temperatures directly from the Greenland ice sheet, *Science*, 282, 268-271, 1998.

Dällenbach A., Blunier T., Flückiger J., Stauffer B., Chappellaz J. and Raynaud D.: Changes in the atmospheric CH₄ gradient between Greenland and Antarctica during the Last Glacial and the transition to the Holocene, *Geophysical Research Letters*, 27, 7, 1005-1008, 2000.

Dansgaard W.: Stable isotopes in precipitation, *Tellus*, 16, 436-468, 1964

Dee, D.P., Uppala, S.M., Simmons, A.J., Berrisford, P. Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M.A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A.C.M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A.P., Monge-Sanz, B.M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137, 656, 553–597, 2011.

Dutton A., Carlson A.E., Long A.J., Milne G.A., Clark P.U., DeConto R., Horton B.P., Rahmstorf S. and Raymo M.E.: Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, 349, doi: 10.1126/science.aaa4019, 2015.

Fettweis, X., Hanna, E., Lang, C., Belleflamme, A., Ericum, M., and Gallee, H.: Important role of the mid-tropospheric atmospheric circulation in the recent surface melt increase over the greenland ice sheet, *Cryosphere*, 7, 241-248, 10.5194/tc-7-241-2013, 2013.

Goujon C., Barnola J-M. and Ritz C.: Modeling the densification of polar firn including heat diffusion: Application to close-off characteristics and gas isotopic fractionation for Antarctica and Greenland sites. *Journal of Geophysical Research*, 108, D24, 4792, 2003

Groote P.M., Stuiver M., White J.W.C., Johnsen S.J. and Jouzel J., Comparison of the oxygen isotope records from the GISP2 and GRIP Greenland ice cores. *Nature*, 366, 552-554, 1993

Guillevic M., Bazin L., Landais A., Kindler P., Orsi A., Masson-Delmotte V., Blunier T., Buchardt S.L., Capron E., Leuenberger M., Martinerie P., Prié F. and Vinther B.M., Spatial gradients of temperature, accumulation and $\delta^{18}\text{O}$ -ice in Greenland over a series of Dansgaard–Oeschger events, *Climate of the Past* 9, 1029-1051, 2013.

Hawley, R., Courville, Z. R., Kehr, L.M., Lutz, E.R., Osterberg, E.C., Overly, T.B. and Wong, G.J.: Recent accumulation variability in northwest Greenland from ground-penetrating radar and shallow cores along the Greenland Inland Traverse, *Journal of Glaciology*, 60, 220, doi: 10.3189/2014JoG13J141, 2014.

Helsen, M. M., van de Berg, W. J., van de Wal, R. S. W., van den Broeke, M. R., and Oerlemans, J.: Coupled regional climate-ice-sheet simulation shows limited greenland ice loss during the Eemian, *Clim Past*, 9, 1773-1788, 10.5194/cp-9-1773-2013, 2013.

Johnsen, S.J., Clausen, H.B., Dansgaard, W., Gundestrup, N.S., Hammer, C.U., Andersen, U., Andersen, K.K., Dahl-Jensen, D., Steffensen, J.P., Sveinbjörnsdóttir, A.E., White, J., Jouzel, J. and Fisher, D.: The $\delta^{18}\text{O}$ record along the Greenland Ice Core Project deep ice core and the problem of possible Eemian climatic instability, *Journal Geophysical Research*, 102, C12, 26397–26410, 1997.

Jouzel, J.: Calibrating the isotopic paleothermometer, *Science*, 286, 910-911, 1999.

Jouzel J., Masson-Delmotte V., Cattani O., Dreyfus G., Falourd S., Hoffmann G., Minster B., Nouet J., Barnola J-M., Fisher H., Gallet J-C., Johnsen S., Leuenberger M., Loulergue L., Luethi D., Oerter H., Parrenin F., Raisbeck G., Raynaud D., Schilt A., Schwander J., Selmo J., Souchez R., Spahni R., Stauffer B., Steffensen J-P., Stenni B., Stocker T.F., Tison J-L., Werner M. and Wolff E.W.: Orbital and millennial Antarctic climate variability over the past 800,000 years, *Science*, 317, 793-796, 2007.

Kapsner, W.R., Alley, R.B., Shuman, A., Anandakrishnan, S. and Grootes, P.M.: Dominant influence of atmospheric circulation on snow accumulation in Greenland over the past 18,000 years, *Nature*, 373, 52-54, 1995.

Keegan K., Albert M.R. and Baker I.: The impact of ice layers on gas transport through firn at the North Greenland Eemian Ice Drilling (NEEM) site, Greenland, *The Cryosphere* 8, 5, 1801-1806, 2014.

Kindler P., Guillevic M., Baumgartner M., Schwander J., Landais A. and Leuenberger M.: Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core, *Clim. Past* 10, 2, 887-902, 2014.

Kobashi, T., Severinghaus, J.P. and Kawamura, K.: Argon and nitrogen isotopes of trapped air in the GISP2 ice core during the Holocene epoch (0–11,500 B.P.): Methodology and implications for gas loss processes, *Geochim. Cosmochim. Acta*, **72**, 4675–4686, 2008.

Krinner G., Genthon C. and Jouzel J.: GCM analysis of local influences on ice core d signals. *Geophysical Research Letters* 24, 2825-2828, 1997.

Landais A., Dreyfus G., Capron E., Jouzel J., Masson-Delmotte V., Roche D.M., Prié F., Caillon N., Chappellaz J., Leuenberger M., Laurantou A., Parrenin F., Raynaud D. and Teste G.: Two-phase change in CO₂, Antarctic temperature and global climate during Termination II. *Nature Geosciences*, 6, 1062–1065, 10.1038/NGEO1985, 2013.

Landais A., Masson-Delmotte V., Jouzel J., Raynaud D., Johnsen S., Huber C., Leuenberger M., Schwander J. and Minster B., The glacial inception as recorded in the NorthGRIP Greenland ice core: timing, structure and associated abrupt temperature changes, *Climate Dynamics*, 26, 2-3, 273-284, 2006.

Landais, A., Chappellaz, J., Delmotte, M., Jouzel, J., Blunier, T., Bourg, C., Caillon, N., Cherrier, S., Malaize, B., Masson-Delmotte, V., Raynaud, D., Schwander, J., and Steffensen, J.: A tentative reconstruction of the last interglacial and glacial inception in greenland based on new gas measurements in the greenland ice core project (grip) ice core, *J Geophys Res-Atmos*, 108, 10.1029/2002jd003147, 2003.

Landais A., Steffensen, J.P., Caillon, N., Jouzel, J., Masson-Delmotte, V. and Schwander, J.: Evidence for stratigraphic distortion in the Greenland Ice Core Project (GRIP) ice core during Event 5e1 (120 kyr BP) from gas isotopes, *J. Geophys. Res.*, 109, 6, doi : 10.1029/2003JD004193, 2004.

~~Langebroek, P.M. and Nisancioglu, K.H.: Moderate Greenland ice sheet melt during the last interglacial constrained by present-day observations and paleo-ice core reconstructions, *The Cryosphere Discuss.*, doi:10.5194/tc-2016-15, in review, 2016.~~

Lorius, C., Ritz, C., Jouzel, J., Merlivat, L., and Barkov, N. I.: A 150 000-year climatic record from Antarctic ice, *Nature*, 316, 591–596, doi:10.1038/316591a0, 1985.

Lunt, D. J., Abe-Ouchi, A., Bakker, P., Berger, A., Braconnot, P., Charbit, S., Fischer, N., Herold, N., Jungclaus, J. H., Khon, V. C., Krebs-Kanzow, U., Langebroek, P. M., Lohmann, G., Nisancioglu, K. H., Otto-Bliesner, B. L., Park, W., Pfeiffer, M., Phipps, S. J., Prange, M., Rachmayani, R., Renssen, H., Rosenbloom, N., Schneider, B., Stone, E. J., Takahashi, K., Wei, W., Yin, Q., and Zhang, Z. S.: A multi-model assessment of last interglacial temperatures, *Clim. Past*, 9, 699-717, doi:10.5194/cp-9-699-2013, 2013.

Masson-Delmotte, V., Braconnot, P., Hoffmann, G., Jouzel, J., Kageyama, M., Landais, A., Lejeune, Q., Risi, C., Sime, L., Sjolte, J., Swingedouw, D., and Vinther, B.: Sensitivity of interglacial greenland temperature and delta o-18: Ice core data, orbital and increased co2 climate simulations, *Clim Past*, 7, 1041-1059, 10.5194/cp-7-1041-2011, 2011.

Masson-Delmotte, V., Schulz, M., Abe-Ouchi, A., Beer, J., Ganopolski, A., González Rouco, J.F., Jansen, E., Lambeck, K., Luterbacher, J., Naish, T., Osborn, T., Otto-Bliesner, B., Quinn, T., Ramesh, R., Rojas, M., Shao X. and Timmermann, A.: Information from Paleoclimate Archives, in: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 383-464 (Chapter 5), 2013

Masson-Delmotte V., **Landais A.**, Stievenard M., Hoffmann G., Cattani O., Falourd S., Johnsen S. J., Jouzel J., Dahl-Jensen D., Sveinbjornsdottir A., White J. C. W., Popp T., and Fischer H., Holocene climatic changes in Greenland: Different deuterium excess signals at Greenland Ice Core Project (GRIP) and NorthGRIP, *J. Geophys. Res.*, 110, 2005.

Masson-Delmotte V., Steen-Larsen H.C., Ortega P., Swingedouw D., Popp T., Vinther B.M., Oerter H., Sveinbjornsdottir A.E., Gudlaugsdottir H., Box J.E., Falourd S., Fettweis X., Gallée H., Garnier E., Gkinis V., Jouzel J., Landais A., Minster B., Paradis N., Orsi A., Risi C., Werner M. and White J.W.C.: Recent changes in north-west Greenland climate documented by NEEM shallow ice core data and simulations, and implications for past-temperature reconstructions, *The Cryosphere*, 9, 4, 1481-1504, 2015.

Merz, N., Born, A., Raible, C. C., Fischer, H., and Stocker, T. F.: Dependence of Eemian Greenland temperature reconstructions on the ice sheet topography, *Clim. Past*, 10, 1221-1238, 10.5194/cp-10-1221-2014, 2014a.

Merz, N., Gfeller, G., Born, A., Raible, C. C., Stocker, T. F., and Fischer, H.: Influence of ice sheet topography on Greenland precipitation during the Eemian interglacial, *Journal of Geophysical Research*, 119, 10749-10768, 10.1002/2014JD021940, 2014b.

Merz, N., Born, A., Raible, C. C., and Stocker, T. F.: Warm Greenland during the last interglacial: the role of regional changes in sea ice cover, *Clim. Past Discuss.*, doi:10.5194/cp-2016-12, in review, 2016.

Munk, J., Jezek, K.C., Forster and R.R., Gogieni, S.P.: An accumulation map for the Greenland dry-snow facies derived from spaceborn radar, *J. Geophys. Res. Atmosphere*, 108, D9, 10.1029/2002JD002481, 2003.

NEEM comm. members, Eemian interglacial reconstructed from a greenland folded ice core, *Nature*, 493, 489-494, 10.1038/nature11789, 2013.

NorthGRIP-community-members: High resolution climate record of the northern hemisphere reaching into last interglacial period, *Nature*, 431, 147-151, 2004.

Orsi, A. J., Kawamura, K., Fegyveresi, J. M., Headly, M. A., Alley, R. B. and Severinghaus, J. P.: Differentiating bubble-free layers from melt layers in ice cores using noble gases. *Journal of Glaciology*, 61, 227, 585-594, 10.3189/2015JoG14J237, 2015

Ortega P., Swingedouw D., Masson-Delmotte V., Risi C., Vinther B., Yiou P., Vautard R. and Yoshimura K.: Characterizing atmospheric circulation signals in Greenland ice cores: insights from a weather regime approach, *Climate Dynamics*, 43, 2585-2605, 2014

Otto-Bliesner, B. L., Marshall, S. J., Overpeck, J. T., Miller, G. H., Hu, A., and CAPE Last Interglacial Project members: Simulating Arctic Climate Warmth and Icefield Retreat in the Last Interglaciation, *Science*, 311, 1751–1753, 2006

Otto-Bliesner B.L., Park W., Pfeiffer M., Phipps S.J., Prange M., Rachmayani R., Renssen H., Rosenbloom N., Schneider B., Stone E.J., Takahashi E., Wei W., Yin Q. and Zang Z.S.: A multi-model assessment of last interglacial temperatures, *Climate of the past*, 9, 699-717, 2013.

Otto-Bliesner, B. L., Rosenbloom, N., Stone, E. J., McKay, N. P., Lunt, D. J., Brady, E. C., and Overpeck, J. T.: How warm was the last interglacial? New model-data comparisons, *Philosophical Transactions of the Royal Society a- Mathematical Physical and Engineering Sciences*, 371, 10.1098/rsta.2013.0097, 2013.

Petit J.R., Jouzel J., Raynaud D., Barkov N.I., Barnola J-M., Basile I., Bender M., Chappellaz J., Davis M., Delaygue G., Delmotte M., Kotlyakov V.M., Legrand M., Lipenkov V.Y., Lorius C., Pepin L., Ritz C., Saltzman E. and Stievenard M.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399, 6735, 429-436, 1999.

Pfahl, S. and Sodemann, H.: What controls deuterium excess in global precipitation?, *Clim. Past*, 10, 771-781, doi:10.5194/cp-10-771-2014, 2014

Ritz, C.: Un modèle thermo-mécanique d'évolution pour le bassin glaciaire Antarctique Vostok-Glacier Byrd: Sensibilité aux valeurs des paramètres mal connus, Ph.D. thesis, Université Joseph Fourier – Grenoble I, 1992.

Robinson A., Calov R. and Ganopolski A.: Greenland ice sheet model parameters constrained using simulations of the Eemian Interglacial, *Climate of the Past*, 7, 381-396, 2011.

Severinghaus J.P., Sowers T., Brook E.J., Alley R.B. and Bender M.L.: Timing of abrupt climate change at the end of the Younger Dryas interval from thermally fractionated gases in polar ice, *Nature*, 391, 6663, 141-146, 1998.

Sime, L. C., Risi, C., Tindall, J. C., Sjolte, J., Wolff, E. W., Masson-Delmotte, V., and Capron, E.: Warm climate isotopic simulations: What do we learn about interglacial signals in greenland ice cores?, *Quaternary Sci Rev*, 67, 59-80, 2013.

Steen-Larsen, H. C., Masson-Delmotte, V., Sjolte, J., Johnsen, S. J., Vinther, B. M., Breon, F. M., Clausen, H. B., Dahl-Jensen, D., Falourd, S., Fettweis, X., Gallee, H., Jouzel, J., Kageyama, M., Lerche, H., Minster, B., Picard, G., Punge, H. J., Risi, C., Salas, D., Schwander, J., Steffen, K., Sveinbjörnsdóttir, A. E., Svensson, A., and White, J.: Understanding the climatic signal in the water stable isotope records from the neem shallow firn/ice cores in northwest Greenland, *J Geophys Res-Atmos*, 116, D06108, 10.1029/2010jd014311, 2011.

Steen-Larsen, H. C., Johnsen, S. J., Masson-Delmotte, V., Stenni, B., Risi, C., Sodemann, H., Balslev-Clausen, D., Blunier, T., Dahl-Jensen, D., Ellehoj, M. D., Falourd, S., Grindsted, A., Gkinis, V., Jouzel, J., Popp, T., Sheldon, S., Simonsen, S. B., Sjolte, J., Steffensen, J. P., Sperlich, P., Sveinbjörnsdóttir, A. E., Vinther, B. M., and White, J. W. C.: Continuous monitoring of summer surface water vapor isotopic composition above the greenland ice sheet, *Atmos. Chem. Phys.*, 13, 4815-4828, 10.5194/acp-13-4815-2013, 2013.

Steen-Larsen, H. C., Masson-Delmotte, V., Hirabayashi, M., Winkler, R., Satow, K., Prié, F., Bayou, N., Brun, E., Cuffey, K. M., Dahl-Jensen, D., Dumont, M., Guillevic, M., Kipfstuhl, J., Landais, A., Popp, T., Risi, C., Steffen, K., Stenni, B., and Sveinbjörnsdóttir, A.: What controls the isotopic composition of greenland surface snow?, *Clim. Past Discuss.*, 10, 377-392, 10.5194/cpd-9-6035-2013, 2014.

Steen-Larsen H.C., Sveinbjörnsdóttir A.E., Jonsson T., Ritter F., Bonne J-L., Masson-Delmotte V., Sodemann H., Blunier T., Dahl-Jensen D. and Vinther B.: Moisture sources and synoptic to seasonal variability of North Atlantic water vapor isotopic composition. *J. Geophys. Res. Atmos.* 120(12):doi:10.1002/2015JD023234, 2015.

Stone, E. J., Lunt, D. J., Annan, J. D., and Hargreaves, J. C.: Quantification of the greenland ice sheet contribution to last interglacial sea level rise, *Clim Past*, 9, 621-639, 10.5194/cp-9-621-2013, 2013.

Sturevik-Storm, A. S., Aldahan, A., Possnert, G., Berggren, A. M., Muscheler, R., Dahl-Jensen, D., Vinter, B. and Usoskin, I. G.: ^{10}Be climate fingerprints during the Eemian in the NEEM ice core, Greenland, *Scientific Reports*, 4, 6408, 10.1038/srep06408, 2014.

Suwa, M., Fischer, J. C., Bender, M. L., Landais, A., and Brook, E. J.: Chronology reconstruction for the disturbed bottom section of the gisp2 and the grip ice cores : Implications for termination II in Greenland, *J. Geophys. Res.*, 111, D02101, 2006.

Uppala, S.M., Kållberg, P.W., Simmons, A.J., Andrae, U., Bechtold, V.D.C., Fiorino, M., Gibson, J.K., Haseler, J., Hernandez, A., Kelly, G.A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R.P., Andersson, E., Arpe, K., Balmaseda, M.A., Beljaars, A.C.M., Van De Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B.J., Isaksen, I., Janssen, P., Jenne, R., McNally, A.P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N.A., Saunders, R.W., Simon, P., Sterl, A., Trenberth, K., Untch, A., Vasiljevic, D., Viterbo, P. and Woollen, J.: The ERA-40 re-analysis, *Q. J. R. Meteorol. Soc.*, 131, 612, 2961–3012, 2005.

Van de Berg, W. J., van den Broeke, M. R., van Meijgaard, E., and Kaspar, F.: Importance of precipitation seasonality for the interpretation of eemian ice core isotope records from greenland, *Clim Past*, 9, 1589-1600, 10.5194/cp-9-1589-2013, 2013.

Veres D., Bazin L., Landais A., Toyé Mahamadou Kele H., Lemieux-Dudon B., Parrenin F., Martinerie P., Blayo E., Blunier T., Capron E., Chappellaz J., Rasmussen S.O., Severi M., Svensson A., Vinther B. and Wolff E.W.: The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years. 9:1733-1748, 10.5194/cp-1733-2013, 2013.

Vinther, B. M., Buchardt, S. L., Clausen, H. B., Dahl-Jensen, D., Johnsen, S. J., Fisher, D. A., Koerner, R. M., Raynaud, D., Lipenkov, V., Andersen, K. K., Blunier, T., Rasmussen, S. O., Steffensen, J. P., and Svensson, A. M.: Holocene thinning of the greenland ice sheet, *Nature*, 461, 385-388, 10.1038/nature08355, 2009.

Vinther, B. M., Jones, P., Briffa, K., Clausen, H., Andersen, K., Dahl-Jensen, D., and Johnsen, S.: Climatic signals in multiple highly resolved stable isotope records from greenland, *Quaternary Sci Rev*, 29, 522-538, 2010.

	Pre-industrial*					LIG**			
	NEEM	NEEM deposition site	NGRIP	GRIP	GISP2	NEEM deposition site	NGRIP	GRIP	GISP2
$\delta^{15}\text{N}$ (‰) Uncertainty : $\pm 0.007\text{‰}$	0.290 ¹		0.310 ²		0.305- 0.325 ³	0.251	0.275	0.230	0.230
$\delta^{18}\text{O}_{\text{ice}}$ (‰) Uncertainty : $\pm 0.1\text{‰}$	-33.6 ⁴	-35 ⁷	-35.5 ⁴	-35.2 ⁴		-31.5 ⁷	-32.2	-31.8	-31.8
d-excess (‰) Uncertainty : $\pm 1.5\text{‰}$	11 ⁴	11	10.5 ⁴	9.5 ⁴		11.5	9.9 ⁵	10.8 ⁶	
Mean temperature (°C)	-28.5 ⁸	-31 ⁷	-31.5 ²	-31.7 ⁴					
accumulation (m i.e. yr ⁻¹)	0.22 ⁸		0.19 ²	0.23 ²					

Table 1: Characteristics of Greenland deep ice core at pre-industrial and during the LIG.

The uncertainty corresponds to the standard error of the mean. Where data was compiled from previous studies, the references for the number in this table are taken from: 1-Buizert et al. (2012); 2-Guillevic et al. (2013); 3-Kobashi et al. (2008) - Bender et al. (2006); 4- Masson-Delmotte et al. (2005); 5- Capron et al. (2012); 6-Jouzel et al. (2007) - 7- NEEM comm. members (2013)- 8 –Masson-Delmotte et al. (2015)

*: The accumulation rate, $\delta^{18}\text{O}$ and d-excess values attributed to the pre-industrial conditions correspond to averages over the last 200 years. For $\delta^{15}\text{N}$, it corresponds to the value at the bottom of the firn (itself built over the last 200 years). The mean temperature is derived from borehole measurements.

**: For determining the $\delta^{18}\text{O}_{\text{ice}}$, $\delta^{15}\text{N}$ and d-excess values attributed to the LIG, we have taken the average of the corresponding records for NEEM, GRIP and GISP2 between 122 and 126 ka (excluding negative $\delta^{15}\text{N}$ peak at 2384 m at NEEM). At NGRIP, we probably miss the first part of the LIG and probably the optimum of the LIG. The values indicated here correspond to the average over the 1000 oldest years recorded in the NGRIP ice core (119 to 120 ka on the AICC2012 timescale).




Figure 1: Synchronized Greenland $\delta^{18}\text{O}_{\text{ice}}$ records.

From top to bottom: summer solstice insolation at 77°N (black); $\delta^{18}\text{O}_{\text{ice}}$ from NGRIP (light blue line), GRIP (open circles) and GISP2 (triangles) on the AICC2012 timescale (bottom axis) and NEEM $\delta^{18}\text{O}_{\text{ice}}$ (dark blue line) on its depth scale (top axis); CH_4 records from NGRIP (light green line), GRIP (open circles), GISP2 (triangles) and EDC (dashed line) on the AICC2012 timescale and NEEM (dark green line) on its depth axis; $\delta^{18}\text{O}_{\text{atm}}$ records from NGRIP (light blue), GRIP (open circles), GISP2 (triangles) and EDC (dashed line) on the AICC2012 timescale and NEEM $\delta^{18}\text{O}_{\text{atm}}$ (turquoise) on its depth axis. The shaded grey rectangle highlights the deepest part of the NEEM records, where no gas synchronization with Antarctic ice core records from the penultimate glaciation is feasible.




Figure 2: NEEM $\delta^{18}\text{O}_{\text{ice}}$ (top), $\delta^{15}\text{N}$ (middle) and CH_4 (bottom) spanning the LIG.

The hatched rectangle indicates the stratigraphic disturbance identified at the upper part of the LIG section. The light grey rectangle highlights the single negative peak in $\delta^{15}\text{N}$ at NEEM during the LIG, corresponding to the strongest positive peak in CH_4 , both peaks being identified at 2384 m depth. The grey and bold horizontal segment indicates the depth range over which the $\delta^{15}\text{N}$ average has been done.




Figure 3: Contour plot showing the evolution of modeled $\delta^{15}\text{N}$ with respect to accumulation rate estimates for LIG at NEEM.

The black circles indicate the accumulation rate vs temperature for pre-industrial conditions for the 3 Greenland sites discussed here, Summit, NGRIP and NEEM. The grey zone corresponds to the repartition of the accumulation rate vs temperature data with a 1 σ envelop from Benson et al. (1962). The dashed horizontal black line indicate the pre-industrial level of NEEM accumulation rate; the khaki horizontal solid line stands for the maximum accumulation rate estimate from Buchardt et al. (2012) or Kaspner et al. (1995); the purple horizontal solid line stands for the NEEM deposition site LIG accumulation rate best guess estimate from chemical aerosol species; the dashed and solid blue horizontal lines show the minimum and maximum accumulation rate estimate for the NEEM deposition site at LIG proposed by Sturevik-Storm et al. (2014) on the basis of the ^{10}Be concentration; the red dashed curve and the green dashed curves indicates the relationship between accumulation and temperature deduced, respectively from the "Psat-approach" and the "Masson-Delmotte approach". The red contour line indicates the NEEM deposition site LIG $\delta^{15}\text{N}$ level excluding the peak at 2384 m. The two vertical arrows indicate the highest and lowest NEEM LIG temperature as determined by the graphical determination (see text). The thin pink contour line indicates the NGRIP LIG $\delta^{15}\text{N}$ level and the blue contour line indicates the GRIP/GISP2 LIG $\delta^{15}\text{N}$ level. The blue circle illustrates the range of possible present-day values for GISP2 $\delta^{15}\text{N}$.




Figure 4: Annual mean temperature anomaly relative to pre-industrial control simulations for the upstream NEEM depositional site extracted from transient and equilibrium climate model simulations.

Note the similarity for the annual mean temperature increase at 125 and 130ka as simulated by HadCM3_Bris and NorESM_BCCR. For more information on the climate models and the simulations themselves we refer to the compilation studies and references therein (Bakker et al., 2013, 2014 and Lunt et al., 2013).