

Interactive comment on “Changes in Holocene meridional circulation and poleward Atlantic flow: the Bay of Biscay as a nodal point” by Yannick Mary et al.

Y. Mary

yannick.mary@u-bordeaux.fr

Received and published: 4 May 2016

We are very grateful for your positive comments and interesting suggestions.

The reference to the article of Mojtahid et al., (2013) is actually mentioned in the reference list (L454), although not at the correct position. This will be corrected in the final version. We apologize for the mistake and thank you for spotting it.

Regarding the influence of solar forcing, short-lived cold spells recorded in the SST signal of core PP10-07 at 8.2, 7, 4, 2.9 and 1.7 ka BP indeed show similarity with the so-called "Bond cycles", at ca 8, 6, 4.5, 3, 1.8 and 0.5ky (Bond et al., 2001). However, the very short duration of these events in the Bay of Biscay calls for a derived phe-

[Printer-friendly version](#)

[Discussion paper](#)



nomenon rather than a direct influence of solar forcing on SST oscillations. The same idea is indirectly suggested in our paper when we refer to the millennial-scale storminess maxima reconstruction (Figure n°3 in the manuscript) of Sorrel et al., (2012). These authors concluded that the solar forcing was not a primary trigger for storminess maxima but did not exclude its possible influence as a weak external driver.

Though, comparing the SST signal of PP10-07 core with Bond cycle proxies, such as drifted ice indices, or directly with solar irradiance signal is a challenging suggestion. We will definitively try such approach (see the preliminary Figure R1) and include it in the final revised version, if possible. For information, this comparison was done and discussed at the scale of the last 2 ka BP in our 2015 paper (Mary et al., 2015).

Moreover, Morley et al., (2014) suggest that the strength of the Latitudinal Thermal Gradient (LTG), driven by contrasting distribution of insolation between polar and tropical latitudes, impacts meridional heat transport by oceanic systems and associated teleconnections. A sharp increase of the LTG occurs around 2000 BP. Such forcing may enhance NAC inflow toward northern latitude, which may explain the large, multi-millennial scale anomalies visible on the Bay of Biscay.

References:

Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M.N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I., Bonani, G. (2001): Persistent solar influence on North Atlantic climate during the Holocene. *Science* 294, 2130–2136.

Mary, Y., Eynaud, F., Zaragosi, S., Malaizé, B., Cremer, M. and Schmidt, S. (2015): High frequency environmental changes and deposition processes in a 2 kyr-long sedimentological record from the Cap-Breton canyon (Bay of Biscay), *The Holocene*, 25, 348–365, doi:10.1177/0959683614558647.

Mojtahid, M., Jorissen, F.J., Garcia, J., Schiebel, R., Michel, E., Eynaud, F., Gillet, H., Cremer, M., Diz Ferreiro, P., Siccha, M., Howa, H. (2013). High resolu-

[Printer-friendly version](#)[Discussion paper](#)

tion Holocene record in the southeastern Bay of Biscay: Global versus regional climate signals. *Palaeogeography, Palaeoclimatology, Palaeoecology* 377, 28–44. doi:10.1016/j.palaeo.2013.03.004

Morley, A., Rosenthal, Y., deMenocal, P. (2014): Ocean-atmosphere climate shift during the mid-to-late Holocene transition. *Earth and Planetary Science Letters* 388, 18–26. doi:10.1016/j.epsl.2013.11.039

Roth, R. and Joos, F. (2013): A reconstruction of radiocarbon production and total solar irradiance from the Holocene ^{14}C and CO_2 records: implications of data and model uncertainties, *Clim. Past*, 9, 1879-1909, doi:10.5194/cp-9-1879-2013. data available from CP at <http://www.clim-past.net/9/1879/2013/>

Sorrel, P., Debret, M., Billeaud I., Jaccard S.L., McManus J.F., Tessier B. (2012): Persistent non-solar forcing of Holocene storm dynamics in coastal sedimentary archives, *Nature Geoscience* 12, 892–896. doi:10.1038/ngeo1619, 2012.

Interactive comment on *Clim. Past Discuss.*, doi:10.5194/cp-2016-32, 2016.

CPD

Interactive
comment

Printer-friendly version

Discussion paper






Fig. 1. Comparison of SST data of the Bay of Biscay with reconstructed solar irradiance data (after Roth and Joos, 2013) and reconstruction of Holocene storm periods (after Sorrel et al., 2012).

Printer-friendly version

Discussion paper

