

1 **Supplementary Information Figure Legends**

2 **Figure S1.** Results of the preheat plateau test on single aliquots of the Lake Little Llangothlin
3 sample L-EVA1231 (field code LL4). There does not appear to be a dependence of equivalent
4 dose with preheat temperature, although more inter-aliquot scatter was observed for the preheat
5 temperature of 280°C. Therefore a higher preheat temperature (260°C) was chosen for
6 subsequent SAR measurements; the cutheat temperature was 220°C.

7 **Figure S2.** Dose recovery test results for L-EVA1228, illustrated as a radial plot. The black line
8 corresponds to the administered dose and the shaded line to 2σ on either side of the central age.

9 **Table S1.** Results of the Finite Mixture Modelling. The dominant age populations used for age
10 calculation are highlighted in italics.

11

12 **Supporting Information Legends (File S1)**13 **Table S1.** Overdispersion values based on dose distributions from the LLL OSL samples.

Sample code	Overdispersion (%)
L-EVA1228 (LL1)	79.9
L-EVA1229 (LL2)	44.4
L-EVA1230 (LL3)	63.9
L-EVA123 (LL4)	61.4

14


15

16 **Table S2.** Results from the Finite Mixture Modelling. The dominant age populations used for
17 age calculation are highlighted in italics; the resulting ages for the dominant and second major
18 age populations are also given for comparison.

Sample code	Number of components	De (Gy)	% population	Age (ka)	BIC
L-EVA1228 (LL1)	4	1.4±0.1 2.8±0.3 <i>6.1±0.6</i> 13.4±1.6	22.4 28.7 32.5 16.5	1.2 ± 0.1 2.3 ± 0.3 <i>5.1 ± 0.5</i> 11.1 ± 1.6	155
L-EVA1229 (LL2)	2	5.7±0.3 <i>19.2±0.4</i>	12.8 87.2	5.6 ± 0.5 <i>18.9 ± 1.2</i>	101
L-EVA1230 (LL3)	3	4.5±0.3 11.8±0.5 <i>26.9±0.9</i>	12.2 39.0 48.8	9.1 ± 0.7 20.6 ± 1.4	234
L-EVA1231 (LL4)	2	7.8±0.5 22.9±1.2	43.2 56.8	8.0 ± 0.7 <i>23.4 ± 1.8</i>	128

19

20




21

22 **Figure S1.** Results of the preheat plateau test on single aliquots of the Lake Little Llangothlin
 23 sample L-EVA1231 (field code LL4). There does not appear to be a dependence of equivalent
 24 dose with preheat temperature, although more inter-aliquot scatter was observed for the preheat
 25 temperature of 280°C. Therefore a higher preheat temperature (260°C) was chosen for
 26 subsequent SAR measurements; the cutheat temperature was 220°C.

27

28



29

30 **Figure S2.** Dose recovery test results for L-EVA1228, illustrated as a radial plot. The black line
 31 corresponds to the administered dose and the shaded line to 2σ on either side of the central age.

32

33

34 Supplementary Information - References

35

36 Botter-Jensen, L., Bulur, E., Duller, G.A.T., Murray, A.S., 2000. Advances in luminescence
37 instrument systems. *Radiation Measurements* 32, 523-528.

38 Bowler, J.M., 1968. Australian landform example no.11: lunette. *Australian Geographer* 10, 402-
39 404.

40 Bowler, J.M., 1973. Clay dunes: their occurrence, formation and environmental significance.
41 *Earth Science Reviews* 9, 315-338.

42 Bowler, J.M., 1983. Lunettes as indices of hydrologic change: A review of the Australian
43 evidence. *Proceedings of the Royal Society of Victoria* 95, 147-168.

44 Fitzsimmons, K.E., Stern, N., Murray-Wallace, C.V., 2014. Depositional history and
45 archaeology of the central Lake Mungo lunette, Willandra Lakes, southeast Australia. *Journal of*
46 *Archaeological Science* 41, 349-364.

47 Murray, A.S., Wintle, A.G., 2000. Luminescence dating of quartz using an improved single-
48 aliquot regenerative-dose protocol. *Radiation Measurements* 32, 57-73.

49 Murray, A.S., Wintle, A.G., 2003. The single aliquot regenerative dose protocol: potential for
50 improvements in reliability. *Radiation Measurements* 37, 377-381.

51 Prescott, J.R., Hutton, J.T., 1994. Cosmic ray contributions to dose rates for luminescence and
52 ESR dating: Large depths and long term variations. *Radiation Measurements* 23, 497-500.

53

54