

Dear Editor,

Thank you very much for reading this paper. We took account of your minor revision in the last version of the manuscript.

The text has been modified following your recommendations. Only for the “radiocarbon dating” section we did not add information about the interpolation as we write later in the “result” section : *“The age model for the upper 1.2 m of the core SU92-33 was based on 7 AMS-14C age measurements and a linear interpolation between these ages (Table 3, Fig. 2). For the lower portion of the core, a control point was established at the onset of the last deglaciation, which is coeval in the western and central Mediterranean Sea at ~17 cal ka BP (Sierro et al., 2005; Melki et al., 2009; Siani et al., 2001)”*. (lines 303-306).

All the references in preparation have been replaced by personal communication.

Figure 3 has been modified following your recommendations. The y-axis for the  $\epsilon\text{Nd}$  record from the Balearic-Alboran Seas has been resized. Plot a) is now renamed “Balearic Sea” and plot b is now renamed “Alboran and Balearic Seas”

# Hydrological variations of the intermediate water masses of the western Mediterranean Sea during the past 20 ka inferred from neodymium isotopic composition in foraminifera and cold-water corals

Quentin Dubois-Dauphin<sup>1</sup>, Paolo Montagna<sup>2,3</sup>, Giuseppe Siani<sup>1</sup>, Eric Douville<sup>4</sup>, Claudia Wienberg<sup>5</sup>, Dierk Hebbeln<sup>5</sup>, Zhifei Liu<sup>6</sup>, Nejb Kallel<sup>7</sup>, Arnaud Dapoigny<sup>4</sup>, Marie Revel<sup>8</sup>, Edwige Pons-Branchu<sup>4</sup>, Marco Taviani<sup>2,9</sup>, Christophe Colin<sup>1\*</sup>

<sup>1</sup>Laboratoire Geosciences Paris-Sud (GEOPS), Université de Paris Sud, Université Paris-Saclay, 91405 Orsay, France.

<sup>2</sup>ISMAR-CNR, via Gobetti 101, 40129 Bologna, Italy.

<sup>3</sup>Lamont-Doherty Earth Observatory, Columbia University, 61 Route 9W, Palisades, NY 10964, USA

<sup>4</sup>Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France.

<sup>5</sup>MARUM-Center for Marine Environmental Sciences, University of Bremen, Leobener Strasse, 28359 Bremen, Germany.

<sup>6</sup>State Key Laboratory of Marine Geology, Tongji University, Shanghai 200092, China.

<sup>7</sup>Laboratoire Georessources, Matériaux, Environnements et Changements Globaux, LR13ES23, Faculté des Sciences de Sfax, Université de Sfax, BP1171, 3000 Sfax, Tunisia.

<sup>8</sup>Geoazur, UNS, IRD, OCA, CNRS, 250 rue Albert Einstein, 06500 Valbonne, France

<sup>9</sup>Biology Department, Woods Hole Oceanographic Institution, 266 Woods Hole Road, Woods Hole, MA 02543, USA.

Correspondence to: Christophe Colin ([christophe.colin@u-psud.fr](mailto:christophe.colin@u-psud.fr))

**Abstract.** We present the neodymium isotopic composition ( $\epsilon\text{Nd}$ ) of mixed planktonic foraminifera species from a sediment core collected at 622 m water depth in the Balearic Sea, as well as  $\epsilon\text{Nd}$  of scleractinian cold-water corals (CWC; *Madrepora oculata*, *Lophelia pertusa*) retrieved ~~at 280-414 m water depth~~ in the Alboran Sea (280-414 m water depth) and the south Sardinian continental margin (414 m water depth). The aim is to constrain hydrological variations at intermediate depths in the western Mediterranean Sea during the last 20 kyr. Planktonic (*Globigerina bulloides*) and benthic (*Cibicidoides pachyderma*) foraminifera from the Balearic Sea were also analyzed for stable oxygen ( $\delta^{18}\text{O}$ ) and carbon ( $\delta^{13}\text{C}$ ) isotopes. The foraminiferal and coral  $\epsilon\text{Nd}$  values from the Balearic and Alboran Sea are comparable over the last ~13 kyr, with mean values of  $-8.94 \pm 0.26$  ( $1\sigma$ ;  $n=24$ ) and  $-8.91 \pm 0.18$  ( $1\sigma$ ;  $n=25$ ), respectively. Before 13 ka BP, the foraminiferal  $\epsilon\text{Nd}$  values are slightly lower ( $-9.28 \pm 0.15$ ) and tend to reflect ~~a~~ higher mixing between intermediate and deep waters, which ~~is~~ are characterized by more unradiogenic  $\epsilon\text{Nd}$  values. The slight  $\epsilon\text{Nd}$  increase after 13 ka BP is associated to a decoupling in the benthic foraminiferal  $\delta^{13}\text{C}$  composition between intermediate and deeper depths, which started at ~16 ka BP. This suggests an earlier stratification of the water masses and a subsequent reduced contribution of unradiogenic  $\epsilon\text{Nd}$  from deep waters. The CWC from the Sardinia Channel show a much larger scattering of  $\epsilon\text{Nd}$  values, from  $-8.66 \pm 0.30$  to  $-5.99 \pm 0.50$ , and a lower average ( $-7.31 \pm 0.73$ ;  $n=19$ ) compared to the CWC and foraminifera from the Alboran and Balearic Sea, indicative of intermediate waters sourced from the Levantine basin. At the time of sapropel S1 deposition (10.2 to 6.4 ka), the  $\epsilon\text{Nd}$  values of the Sardinian CWC become more unradiogenic ( $-8.38 \pm 0.47$ ;  $n=3$  at ~8.7 ka BP), suggesting a significant contribution of intermediate waters

47 | originated from the western basin. ~~Accordingly, we~~ We propose that western Mediterranean intermediate waters  
48 | replaced the Levantine Intermediate Water (LIW), which ~~was~~ would be strongly reduced during the mid-  
49 | sapropel (~8.7 ka BP). This observation supports a notable change of Mediterranean circulation pattern centered  
50 | on sapropel S1 that needs further investigations to be confirmed.

51

## 52 | 1. Introduction

53 | The Mediterranean Sea is a mid-latitude semi-enclosed basin, characterized by evaporation exceeding  
54 | precipitation and river runoff, where the inflow of fresh and relatively warm surface Atlantic water is  
55 | transformed into saltier and cooler (i.e. denser) intermediate and deep waters. Several studies have demonstrated  
56 | that the Mediterranean thermohaline circulation was highly sensitive to both the rapid climatic changes  
57 | propagated into the basin from high latitudes of the Northern Hemisphere (Cacho et al., 1999, 2000, 2002;  
58 | Moreno et al., 2002, 2005; Paterne et al., 1999; Martrat et al., 2004; Sierro et al., 2005; Frigola et al., 2007,  
59 | 2008) and orbitally-forced modifications of the eastern Mediterranean freshwater budget mainly driven by  
60 | monsoonal river runoff from the ~~south~~ subtropics (Rohling et al., 2002; 2004; Bahr et al., 2015). A link between  
61 | the intensification of the Mediterranean Outflow Water (MOW) and the intensity of the Atlantic Meridional  
62 | Overturning Circulation (AMOC) was proposed (Cacho et al., 1999, 2000, 2001; Bigg and Wadley, 2001; Sierro  
63 | et al., 2005; Voelker et al., 2006) and recently supported by new geochemical data in sediments of the Gulf of  
64 | Cádiz (Bahr et al., 2015). In particular, it has been suggested that the intensity of the MOW and, more generally,  
65 | the variations of the thermohaline circulation of the Mediterranean Sea could play a significant role in triggering  
66 | a switch from a weakened to an enhanced state of the AMOC through the injection of saline Mediterranean  
67 | waters in the intermediate North Atlantic at times of weak AMOC (Rogerson et al., 2006; Voelker et al., 2006;  
68 | Khélifi et al., 2009). The Mediterranean intermediate waters, notably the Levantine Intermediate Water (LIW),  
69 | which represent today up to 80 % in volume of the MOW (Kinder and Parilla, 1987) are considered an important  
70 | driver of MOW-derived salt into the North Atlantic. Furthermore, the LIW also plays a key role in controlling  
71 | the deep-sea ventilation of the Mediterranean basin, being strongly involved in the formation of deep waters in  
72 | the Aegean Sea, Adriatic Sea, Tyrrhenian Sea and Gulf of Lions (Millot and Taupier-Letage, 2005). It is  
73 | hypothesized that a reduction of intermediate and deep-water formation as a consequence of surface hydrological  
74 | changes in the eastern Mediterranean basin acted as a precondition for the sapropel S1 deposition by limiting the  
75 | oxygen supply to the bottom waters (De Lange et al., 2008; Rohling et al., 2015; Tachikawa et al., 2015).  
76 | Therefore, it is crucial to gain a more complete understanding of the variability of the Mediterranean  
77 | intermediate circulation in the past and its impact on the MOW outflow and, in general, on the Mediterranean  
78 | thermohaline circulation.

79 | Previous studies have mainly focused on the glacial variability of the deep-water circulation in the western  
80 | Mediterranean basin (Cacho et al., 2000, 2006; Sierro et al., 2005; Frigola et al., 2007, 2008). During the Last  
81 | Glacial Maximum (LGM), strong deep-water convection took place in the Gulf of Lions, producing cold, well-  
82 | ventilated western Mediterranean Deep Water (WMDW) (Cacho et al., 2000, 2006; Sierro et al., 2005), while  
83 | the MOW flowed at greater depth in the Gulf of Cádiz (Rogerson et al., 2005; Schönfeld and Zahn, 2000). With  
84 | the onset of the Termination 1 (T1) at about 15 ka, the WMDW production declined until the ~~transition-onset of~~  
85 | ~~to~~ the Holocene due to the rising sea level, with a relatively weak mode during the Heinrich Stadial 1 (HS1) and

86 the Younger Dryas (YD) (Sierro et al., 2005; Frigola et al., 2008), that led to the deposition of the Organic Rich  
87 Layer 1 (ORL1; 14.5-8.2 ka BP; Cacho et al., 2002).

88 Because of the disappearance during the Early Holocene of specific epibenthic foraminiferal species, such as  
89 *Cibicoides* spp., which are commonly used for paleohydrological reconstructions, information about the  
90 Holocene variability of the deep-water circulation in the western Mediterranean are relatively scarce and are  
91 mainly based on grain size analysis and sediment geochemistry (e.g. Frigola et al., 2007). These authors have  
92 identified four distinct phases representing different deep-water overturning conditions in the western  
93 Mediterranean basin during the Holocene, as well as centennial- to millennial-scale abrupt events of overturning  
94 reinforcement.

95 Faunal and stable isotope records from benthic foraminifera located at intermediate depths in the eastern basin  
96 reveal ~~uninterrupted~~-well-ventilated LIW during the last glacial period and deglaciation (Kuhnt et al., 2008;  
97 Schmiedl et al., 2010). Similarly, a grain-size record obtained from a sediment core collected within the LIW  
98 depth range (~500 m water depth) at the east Corsica margin also documents enhanced bottom currents during  
99 the glacial period and for specific time intervals ~~during-of~~ the deglaciation, such as HS1 and YD (Toucanne et  
100 al., 2012). The Early Holocene is characterized by a collapse of the LIW (Kuhnt et al., 2008; Schmiedl et al.,  
101 2010; Toucanne et al., 2012) synchronous with the sapropel S1 deposition (10.2 – 6.4 cal ka BP; Mercone et al.,  
102 2000). Proxies for deep-water conditions reveal the occurrence of episodes of deep-water overturning  
103 reinforcement in the eastern Mediterranean basin at 8.2 ka BP (Rohling et al., 1997, 2015; Kuhnt et al., 2007;  
104 Abu-Zied et al., 2008, Siani et al., 2013; Tachikawa et al; 2015), responsible for the interruption of the sapropel  
105 S1 in the eastern Mediterranean basin (Mercone et al., 2001; Rohling et al., 2015).

106 Additional insights into Mediterranean circulation changes may be ~~obtained-gained~~ using radiogenic isotopes,  
107 such as neodymium, that represent reliable tracers for constraining water-mass mixing and sources (Goldstein  
108 and Hemming, 2003, and references therein). It has recently been shown that the neodymium (Nd) isotopic  
109 composition, expressed as  $\epsilon Nd = \left( \frac{(^{143}Nd/^{144}Nd)_{sample}}{(^{143}Nd/^{144}Nd)_{CHUR}} - 1 \right) \times 10000$  (CHUR: Chondritic  
110 Uniform Reservoir [Jacobsen and Wasserburg, 1980]) of living and fossil scleractinian CWC faithfully traces  
111 intermediate and deep-water mass provenance and mixing of the ocean (e.g. van de Flierdt et al., 2010; Colin et  
112 al., 2010; López Correa et al., 2012; Monterro-Serrano et al., 2011, 2013; Copard et al., 2012). Differently from  
113 the CWC, the  $\epsilon Nd$  composition of fossil planktonic foraminifera is not related to the ambient seawater at  
114 calcification depths but reflects the bottom and/or pore water  $\epsilon Nd$ , due to the presence of authigenic Fe-Mn  
115 coatings precipitated on their carbonate shell ~~after deposition onto the sediment~~ (Roberts et al., 2010; Elmore et  
116 al., 2011; Piotrowski et al., 2012; Tachikawa et al., 2014; Wu et al., 2015). Therefore, the  $\epsilon Nd$  composition of  
117 planktonic foraminiferal tests can be used as a useful tracer of deep-water circulation changes in the past,  
118 although the effect of pore water on foraminiferal  $\epsilon Nd$  values could potentially complicate the interpretation  
119 (Tachikawa et al., 2014).

120 In the Mediterranean Sea, modern seawater  $\epsilon Nd$  values display a large range from ~-11 to ~-5, and a clear  
121 vertical and longitudinal gradient, with more radiogenic values encountered in the eastern basin and typically at  
122 intermediate and deeper depths (Spivack and Wasserburg 1988; Henry et al., 1994; Tachikawa et al., 2004;  
123 Vance et al., 2004). Considering this large  $\epsilon Nd$  contrast,  $\epsilon Nd$  recorded in fossil CWC and planktonic  
124 foraminifera from the Mediterranean offers great potential to trace intermediate and deep-water mass exchange

125 between the two basins, especially during periods devoid of key epibenthic foraminifera, such as the sapropel S1  
126 or ORL1 events.

127 Here, the  $\epsilon\text{Nd}$  of planktonic foraminifera from a sediment core collected in the Balearic Sea and CWC samples  
128 from the Alboran Sea and the Sardinia Channel was investigated to establish past changes of the seawater  $\epsilon\text{Nd}$  at  
129 intermediate depths and constrain hydrological variations of the LIW during the last ~20 kyr. The  $\epsilon\text{Nd}$  values  
130 have been combined with stable oxygen ( $\delta^{18}\text{O}$ ) and carbon ( $\delta^{13}\text{C}$ ) isotope measurements of benthic (*Cibicidoides*  
131 *pachyderma*) and planktonic (*Globigerina bulloides*) foraminifera and sea-surface temperature estimates by  
132 modern analogue technique (MAT). Results reveal significant  $\epsilon\text{Nd}$  variations at intermediate depths in the  
133 western basin interpreted as a drastic reduction of the hydrological exchanges between the western and eastern  
134 Mediterranean Sea and the subsequent higher proportion of intermediate water produced in the Gulf of Lions  
135 during the time interval corresponding to the sapropel S1 deposition.

136  
137

## 138 2. Seawater $\epsilon\text{Nd}$ distribution in the Mediterranean Sea

139 The Atlantic Water (AW) enters the Mediterranean Sea as surface inflow through the Strait of Gibraltar with an  
140 unradiogenic  $\epsilon\text{Nd}$  signature of ~-9.7 in the strait (Tachikawa et al., 2004) and ~-10.4 in the Alboran Sea  
141 (Tachikawa et al., 2004, Spivack and Wasserburg, 1988) for depths shallower than 50 m. During its eastward  
142 flowing, AW mixes with upwelled Mediterranean Intermediate Water forming the Modified Atlantic Water  
143 (MAW) that spreads within the basin (Millot and Taupier-Letage, 2005) (Fig.1). The surface water  $\epsilon\text{Nd}$  values  
144 (shallower than 50 m) range from -9.8 to -8.8 in the western Mediterranean basin (Henry et al., 1994; [Montagna  
145 et al., in prep](#)[Montagna, pers. comm., 2016](#)) and -9.3 to -4.2 in the eastern basin, with seawater off the Nile delta  
146 showing the most radiogenic values (Tachikawa et al., 2004; Vance et al., 2004; [Montagna, pers. comm.,  
147 2016](#)[Montagna et al., in prep](#)). The surface waters in the eastern Mediterranean basin become denser due to  
148 strong mixing and evaporation caused by cold and dry air masses flowing over the Cyprus-Rhodes area in  
149 winter, and eventually sink leading to the formation of LIW (Ovchinnikov, 1984; Lascaratos et al., 1993, 1998;  
150 Malanotte-Rizzoli et al., 1999; Pinardi and Masetti, 2000). The LIW spreads throughout the entire Mediterranean  
151 basin at depths between ~150-200 m and ~600-700 m, and is characterized by more radiogenic  $\epsilon\text{Nd}$  values  
152 ranging from -7.9 to -4.8 (average value  $\pm 1\sigma$ :  $-6.6 \pm 1$ ) in the eastern basin and from -10.4 to -7.58 ( $-8.7 \pm 0.9$ )  
153 in the western basin (Henry et al., 1994; Tachikawa et al., 2004; Vance et al., 2004; [Montagna, pers. comm.,  
154 2016](#)[Montagna et al., in prep](#)). The LIW acquires its  $\epsilon\text{Nd}$  signature mainly from the partial dissolution of Nile  
155 River particles (Tachikawa et al., 2004), which have an average isotopic composition of -3.25 (Weldeab et al.,  
156 2002), and the mixing along its path with overlying and underlying water masses with different  $\epsilon\text{Nd}$  signatures.  
157 The LIW finally enters the Atlantic Ocean at intermediate depths through the Strait of Gibraltar with an average  
158  $\epsilon\text{Nd}$  value of  $-9.2 \pm 0.2$  (Tachikawa et al., 2004; [Montagna, pers. comm., 2016](#)[Montagna et al., in prep](#)).

159 The WMDW is formed in the Gulf of Lions due to winter cooling and evaporation followed by mixing between  
160 ~~the relative fresh~~ surface waters and the more saline LIW and spreads into the Balearic basin and Tyrrhenian Sea  
161 between ~2000 m and 3000 m (Millot, 1999; Schroeder et al., 2013) (Fig. 1). The WMDW is characterized by an  
162 average  $\epsilon\text{Nd}$  value of  $-9.4 \pm 0.9$  (Henry et al., 1994; Tachikawa et al., 2004; [Montagna, pers. comm.,  
163 2016](#)[Montagna et al., in prep](#)). Between the WDMW and the LIW (from ~700 to 2000 m), the Tyrrhenian Deep  
164 Water (TDW) has been found (Millot et al., 2006), which is produced by the mixing between WMDW and

165 | Eastern Mediterranean Deep Water (EMDW) that cascades in the Tyrrhenian Sea after entering ~~from~~ through the  
166 | Strait of Sicily (Millot, 1999, 2009; Astraldi et al., 2001). The TDW has an average  $\epsilon\text{Nd}$  value of  $-8.1 \pm 0.5$   
167 | ([Montagna, pers. comm., 2016](#)~~Montagna et al., in prep~~).

168

### 169 | 3. Material and methods

#### 170 | 3.1. Cold-water coral and foraminifera samples

171 | Forty-four CWC samples belonging to the species *Lophelia pertusa* and *Madrepora oculata* collected from the  
172 | Alboran Sea and the Sardinia Channel were selected for this study (Fig. 1). Nineteen fragments were collected at  
173 | various core depths from a coral-bearing sediment core (RECORD 23; 38°42.18' N; 08°54.75' E; Fig. 1)  
174 | retrieved from 414 m water depth in the "Sardinian Cold-Water Coral Province" (Taviani et al., 2015) during the  
175 | R/V Urania cruise "RECORD" in 2013. The core contains well-preserved fragments of *M. oculata* and *L.*  
176 | *pertusa* embedded in a brownish muddy to silty carbonate-rich sediment. The Sardinian CWC samples were  
177 | used for U-series dating and Nd isotopic composition measurements. For the southern Alboran Sea, twenty-five  
178 | CWC samples were collected at water depths between 280 and 442 m in the "eastern Melilla Coral Province"  
179 | (Fig. 1) during the R/V Poseidon cruise "POS-385" in 2009 (Hebbeln et al., 2009). Eleven samples were  
180 | collected at the surface of two coral mounds (New Mound and Horse Mound) and three coral ridges (Brittlestar  
181 | ridges I, II and III), using a box corer and a remotely operated vehicle (ROV). In addition, fourteen CWC  
182 | samples were collected from various core depths of three coral-bearing sediment cores (GeoB13728, 13729 and  
183 | 13730) retrieved from the Brittlestar ridge I. Details on the location of surface samples and cores collected in the  
184 | southern Alboran Sea and details on the radiocarbon ages obtained from these coral samples are reported in Fink  
185 | et al. (2013). Like the CWC sample set from the Sardinia Channel, the dated Alboran CWC samples were also  
186 | used for further Nd isotopic composition analyses in this study.

187 | In addition, a deep-sea sediment core (barren of any CWC fragments) was recovered southwest of the Balearic  
188 | Sea at 622 m water depth during the R/V Le Suroît cruise "PALEOCINAT II" in 1992 (SU92-33; 35°25.38' N;  
189 | 0°33.86' E; Fig. 1). The core unit, which consists of 2.1 m of grey to brown carbonaceous clays, was sub-  
190 | sampled continuously at 5-10 cm intervals for a total number of 24 samples used for  $\delta^{18}\text{O}$ ,  $\delta^{13}\text{C}$  and  $\epsilon\text{Nd}$   
191 | analyzes.

192

#### 193 | 3.2. Analytical procedures on cold-water coral samples

##### 194 | 3.2.1. U/Th dating

195 | The nineteen CWC samples collected from the sediment core RECORD 23 (Sardinia Channel) were analysed for  
196 | uranium and thorium isotopes to obtain absolute dating using a Thermo Scientific™ Neptune<sup>plus</sup> MC-ICPMS  
197 | installed at the Laboratoire des Sciences du Climat et de l'Environnement (LSCE, Gif-sur-Yvette, France). Prior  
198 | to analysis, the samples were carefully cleaned using a small diamond blade to remove any visible contamination  
199 | and sediment-filled cavities. The fragments were examined under a binocular microscope to ensure against the  
200 | presence of bioeroded zones and finally crushed into a coarse-grained powder with an agate mortar and pestle.  
201 | The powders (~60-100 mg) were transferred to acid cleaned Teflon beakers, ultrasonicated in MilliQ water,  
202 | leached with 0.1N HCl for ~ 15 s and finally rinsed twice with MilliQ water. The physically and chemically  
203 | cleaned samples were dissolved in 3-4 ml dilute HCl (~10%) and mixed with an internal triple spike with known  
204 | concentrations of  $^{229}\text{Th}$ ,  $^{233}\text{U}$  and  $^{236}\text{U}$ , calibrated against a Harwell Uraninite solution (HU-1) assumed to be at

205 secular equilibrium. The solutions were evaporated to dryness at 70°C, redissolved in 0.6 ml 3N HNO<sub>3</sub> and then  
206 loaded into 500 µl columns packed with Eichrom UTEVA resin to isolate uranium and thorium from the other  
207 major and trace elements of the carbonate matrix. The U and Th separation and purification followed a  
208 procedure slightly modified from Douville et al. (2010). The U and Th isotopes were determined following the  
209 protocol recently revisited at LSCE (Pons-Branchu et al., 2014). The <sup>230</sup>Th/U ages were calculated from  
210 measured atomic ratios through iterative age estimation (Ludwig and Titterton, 1994), using the <sup>230</sup>Th, <sup>234</sup>U  
211 and <sup>238</sup>U decay constants of Cheng et al. (2013) and Jaffey et al. (1971). Due to the low <sup>232</sup>Th concentration (< 1  
212 ng/g; see Table 1), no correction was applied for the non-radiogenic <sup>230</sup>Th fraction.

213

### 214 3.2.2 Nd isotopic composition analyses on cold-water coral fragments

215 Sub-samples of the CWC fragments from the Sardinia Channel used for U-series dating in this study (Table 1) as  
216 well as sub-samples of the twenty-five CWC fragments originating from the Alboran Sea, which were already  
217 radiocarbon-dated by Fink et al. (2013) (Table 2), were used for further Nd isotopic composition analyses. The  
218 fragments (350 to 600 mg) were subjected to a mechanical and chemical cleaning procedure. The visible  
219 contaminations, such as Fe-Mn coatings and detrital particles, were carefully removed from the inner and  
220 outermost surfaces of the coral skeletons using a small diamond blade. The physically cleaned fragments were  
221 ultrasonicated for 10 min with 0.1 N ultra-clean HCl, followed by several MilliQ water rinses and finally  
222 dissolved in 2.5 N ultraclean HNO<sub>3</sub>. Nd was separated from the carbonate matrix using Eichrom TRU and LN  
223 resins, following the analytical procedure described in detail in Copard et al. (2010).

224 The <sup>143</sup>Nd/<sup>144</sup>Nd ratios of all purified Nd fractions were analyzed using the ThermoScientific Neptune<sup>Plus</sup> Multi-  
225 Collector Inductively Coupled Plasma Mass Spectrometer (MC-ICP-MS) hosted at LSCE. The mass-  
226 fractionation correction was made by normalizing <sup>146</sup>Nd/<sup>144</sup>Nd to 0.7219 and applying an exponential law.  
227 During each analytical session, samples were systematically bracketed with analyses of JNdi-1 and La Jolla  
228 standard solutions, which are characterised by accepted values of 0.512115±0.000006 (Tanaka et al., 2000) and  
229 0.511855±0.000007 (Lugmair et al., 1983), respectively. Standard JNdi-1 and La Jolla solutions were analysed  
230 at concentrations similar to those of the samples (5-10 ppb) and all the measurements affected by instrumental  
231 bias were corrected, when necessary, using La Jolla standard. The external reproducibility (2σ) for time resolved  
232 measurement, deduced from repeated analyses of La Jolla and JNdi-1 standards, ranged from 0.1 to 0.5 εNd  
233 units for the different analytical sessions. The analytical error for each sample analysis was taken as the external  
234 reproducibility of the La Jolla standard for each session. Concentrations of Nd blanks were negligible compared  
235 to the amount of Nd of CWC investigated in this study.

236

## 237 3.3. Analyses on sediment of core SU92-33

### 238 3.3.1. Radiocarbon dating

239 Radiocarbon dating was measured at UMS-ARTEMIS (Pelletron 3MV) AMS (CNRS-CEA Saclay, France).  
240 Seven AMS radiocarbon (<sup>14</sup>C) dating were performed in [first 1.2 m of the](#) core SU92-33 on well-preserved  
241 calcareous tests of the planktonic foraminifera *G. bulloides* in the size fraction >150 µm (Table 3). The age  
242 model for the core was derived from the calibrated planktonic ages by applying a mean reservoir effect of ~400  
243 years (Siani et al., 2000, 2001). All <sup>14</sup>C ages were converted to calendar years (cal. yr BP, BP = AD 1950) by

244 using the INTCAL13 calibration data set (Reimer et al., 2013) and the CALIB 7.0 program (Stuiver and Reimer,  
245 1993).

246

### 247 3.3.2. Stable isotopes

248 Stable oxygen ( $\delta^{18}\text{O}$ ) and carbon ( $\delta^{13}\text{C}$ ) isotope measurements were performed in core SU92-33 on well-  
249 preserved (clean and intact) samples of the planktonic foraminifera *G. bulloides* (250-315  $\mu\text{m}$  fraction) and the  
250 epibenthic foraminifera *C. pachyderma* (250-315  $\mu\text{m}$  fraction) using a Finnigan MAT-253 mass spectrometer at  
251 the State Key Laboratory of Marine Geology (Tongji University). Both  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values are presented  
252 relative to the Pee Dee Belemnite (PDB) scale by comparison with the National Bureau of Standards (NBS) 18  
253 and 19. The mean external reproducibility was checked by replicate analyses of laboratory standards and is better  
254 than  $\pm 0.07\text{‰}$  ( $1\sigma$ ) for  $\delta^{18}\text{O}$  and  $\pm 0.04\text{‰}$  for  $\delta^{13}\text{C}$ .

### 255 3.3.3 Nd isotope measurements on planktonic foraminifera

256 Approximately 25 mg of mixed planktonic foraminifera species were picked from the  $>63$   $\mu\text{m}$  size fraction of  
257 each sample already used for stable isotope measurements (Table 4). The samples were gently crushed between  
258 glass slides under the microscope to ensure that all chambers were open, and ultrasonicated with MilliQ water.  
259 Samples were allowed to settle between ultrasonication steps before removing the supernatant. Each sample was  
260 rinsed thoroughly with MilliQ water until the solution was clear and free of clay. The cleaned samples were  
261 dissolved in 1N acetic acid and finally centrifuged to ensure that all residual particles were removed, following  
262 the procedure described in Roberts et al. (2010). Nd was separated following the analytical procedure reported in  
263 Wu et al. (2015). For details on the measurement of Nd isotopes see the section above.

264

### 265 3.3.4. Modern analogue technique (MAT)

266 The palaeo-sea surface temperatures (SST) were estimated using the modern analogue technique (MAT)  
267 (Hutson, 1980; Prell, 1985), implemented by Kallel et al. (1997) for the Mediterranean Sea. This method directly  
268 measures the difference between the faunal composition of a fossil sample with a modern database, and it  
269 identifies the best modern analogues for each fossil assemblage (Prell, 1985). Reliability of SST reconstructions  
270 is estimated using a square chord distance test (dissimilarity coefficient), which represents the mean degree of  
271 similarity between the sample and the best 10 modern analogues. When the dissimilarity coefficient is lower than  
272 0.25, the reconstruction is considered to be of good quality (Overpeck et al., 1985; Kallel et al., 1997). For core  
273 SU92-33, good dissimilarity coefficients are  $<0.2$ , with an average value of  $\sim 0.13$  (varying between 0.07 and  
274 0.19) (Fig. 2a). The calculated mean standard deviation of SST estimates observed in core MD90-917 are  $\sim 1.5$   
275  $^{\circ}\text{C}$  from the late glacial period to the Younger Dryas and  $\sim 1.2$   $^{\circ}\text{C}$  for the Holocene.

276

## 277 4. Results

### 278 4.1. Cold-water corals

279 The good state of preservation for the CWC samples from the Sardinia Channel (RECORD 23; Fig. 1) is attested  
280 by their initial  $\delta^{234}\text{U}$  values (Table 1), which is in the range of the modern seawater value ( $146.8 \pm 0.1$ ; Andersen  
281 et al., 2010). If the uncertainty of the  $\delta^{234}\text{U}_i$  is taken into account, all the values fulfill the so-called “strict”  $\pm 4$   
282  $\text{‰}$  reliability criterion and the U/Th ages can be considered strictly reliable. The coral ages range from  
283  $0.091 \pm 0.011$  to  $10.904 \pm 0.042$  ka BP (Table 1), and reveal three distinct clusters of coral age distribution during



284 the Holocene representing periods of sustained coral occurrence. These periods coincide with the Early Holocene  
285 encompassing a 700-years-lasting time interval from ~10.9 to 10.2 ka BP, the very late Early Holocene at ~8.7  
286 ka BP, and the Late Holocene starting at ~1.5 ka BP (Table 1).

287 Radiocarbon ages obtained for CWC samples collected in the Alboran Sea were published by Fink et al. (2013)  
288 (Table 2). They also document three periods of sustained CWC occurrence coinciding with the Bølling–Allerød  
289 (B-A) interstadial (13.5–12.9 cal ka BP), the Early Holocene (11.2–9.8 cal ka BP) and the Mid- to Late Holocene  
290 (5.4–0.3 cal ka BP).

291 The  $\epsilon\text{Nd}$  record obtained from the CWC samples from the Alboran Sea displays a narrow range from  $-9.22\pm 0.30$   
292 to  $-8.59\pm 0.3$ , which is comparable to the  $\epsilon\text{Nd}$  record of the planktonic foraminifera from the Balearic Sea over  
293 the last 13.5 kyr (Table 2, Fig. 3b). Most of the CWC  $\epsilon\text{Nd}$  values are similar within [the analytical](#) error and the  
294 record does not reveal any clear difference over the last ~13.5 kyr.

295 On the contrary, the CWC samples from the Sardinia Channel display a relatively large  $\epsilon\text{Nd}$  range, with values  
296 varying from  $-5.99\pm 0.50$  to  $-7.75\pm 0.10$  during the Early and Late Holocene, and values as low as  $-8.66\pm 0.30$   
297 during the the mid-sapropel S1 deposition (S1a) at ~8.7 ka BP (Table 1, Fig. 3c).

298

299

#### 300 **4.2 Core SU92-33**

301 The stratigraphy of core SU92-33 was derived from the  $\delta^{18}\text{O}$  variations of the planktonic foraminifera  
302 *G. bulloides* (Fig. 2b). The last glacial/interglacial transition and the Holocene encompasses the upper 2.1 m of  
303 the core (Fig. 2b). The  $\delta^{18}\text{O}$  record of *G. bulloides* shows higher values (~3.5 ‰) during the late glacial  
304 compared to the Holocene (from ~1.5 to 0.8 ‰) exhibiting a pattern similar to those observed in nearby deep-sea  
305 cores from the Western Mediterranean Sea (Sierro et al., 2005; Melki et al., 2009).

306 The age model for the upper 1.2 m of the core SU92-33 was based on 7 AMS- $^{14}\text{C}$  age measurements and a  
307 linear interpolation between these ages (Table 3, Fig. 2). For the lower portion of the core, a control point was  
308 established at the onset of the last deglaciation, which is coeval in the western and central Mediterranean Sea at  
309 ~17 cal ka BP (Sierro et al., 2005; Melki et al., 2009; Siani et al., 2001). Overall, the upper 2.1 m of core SU92-  
310 33 spans the last 19 kyr, with an estimated average sedimentation rate ranging from ~15 cm ka<sup>-1</sup> during the  
311 deglaciation to ~10 cm ka<sup>-1</sup> during the Holocene.

312 April-May SST reconstruction was derived from MAT to define the main climatic events recorded in  
313 core SU92-33 during the last 19 kyr. SST vary from 8.5°C to 17.5°C with high amplitude variability over the last  
314 19 kyr BP (Fig. 2a). The LGM (19-18 ka BP) is characterized by SST values centered at around 12°C. Then, a  
315 progressive decrease of ~4°C between 17.8 ka and 16 ka marks the Heinrich Stadial 1 (HS1) (Fig. 2a). A  
316 warming phase (~14°C) between 14.5 ka BP and 13.8 ka BP coincides with the B-A interstadial and is followed  
317 by a cooling (~11°C) between 13.1 ka BP and 11.8 ka BP largely corresponding to the YD (Fig. 2a). During the  
318 Holocene, SST show mainly values of ~16°C, with one exception between 7 ka BP and 6 ka BP pointing to an  
319 abrupt cooling of ~3°C (Fig. 2a). From the late glacial to the Holocene, SST variations show a similar pattern to  
320 that previously observed in the Gulf of Lions and Tyrrhenian Sea (Kallel et al., 1997; Melki et al., 2009) as well  
321 as in the Alboran Sea (Martrat et al., 2014; Rodrigo-Gámiz et al., 2014). They are globally synchronous for the  
322 main climatic transitions to the well dated South Adriatic Sea core MD90-917 (Siani et al., 2004) confirming the  
323 robustness of the SU92-33 age model (Fig. 2a).

324 The  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  records obtained from the benthic foraminifera *C. pachyderma* display significant variations  
325 at millennial time scales (Figs. 2c and 2d). The  $\delta^{18}\text{O}$  values decrease steadily from  $\sim 4.5$  ‰ during the LGM to  
326  $\sim 1.5$  ‰ during the Holocene, without showing any significant excursion during HS1 and the YD events (Fig.  
327 2c), in agreement with results obtained from the neighbor core MD99-2343 (Sierro et al., 2005).

328 The  $\delta^{13}\text{C}$  record of *C. pachyderma* shows a decreasing trend since the LGM with a low variability from  $\sim 1.6$  ‰  
329 to  $\sim 0.6$  ‰ (Fig. 2d). The heaviest  $\delta^{13}\text{C}$  values are related to the LGM ( $\sim 1.6$  ‰) while the lightest values ( $\sim 0.6$   
330 ‰) characterize the Early Holocene and in particular the period corresponding to the sapropel S1 event in the  
331 eastern Mediterranean basin (Fig. 2d).

332 The  $\epsilon\text{Nd}$  values of planktonic foraminifera of core SU92-33 from the Balearic Sea vary within a relatively  
333 narrow range between  $-9.50 \pm 0.30$  and  $-8.61 \pm 0.30$ , with an average value of  $-9.06 \pm 0.28$  (Table 2, Fig. 3b). The  
334 record shows a slight increasing trend since the LGM, with the more unradiogenic values (average  $-9.28 \pm 0.15$ ;  
335  $n=7$ ) being observed in the oldest part of the record (between 18 and 13.5 ka BP), whereas Holocene values are  
336 generally more radiogenic (average  $-8.84 \pm 0.22$ ;  $n=17$ ) (Fig. 3b).

337

### 338 5. Discussion

339 Overall, the CWC and foraminiferal  $\epsilon\text{Nd}$  values measured in this study point to a pronounced dispersion at  
340 intermediate depth in terms of absolute values and variability in Nd isotopes during the Holocene between the  
341 Alboran and Balearic Seas and the Sardinia Channel. Furthermore, the foraminiferal  $\epsilon\text{Nd}$  record reveals an  
342 evolution towards more radiogenic values at intermediate water depth in the Balearic Sea over the last  $\sim 19$  kyr  
343 (Fig. 3).

344 A prerequisite to properly interpret such  $\epsilon\text{Nd}$  differences and variations through time consists in characterizing  
345 first the present-day  $\epsilon\text{Nd}$  of the main water-mass end-members ~~flowing-present~~ in the western Mediterranean  
346 basin. It is also necessary to evaluate the temporal changes in  $\epsilon\text{Nd}$  of the end-members since the LGM, and  
347 assess the potential influences of lithogenic Nd input and regional exchange between the continental margins and  
348 seawater (“boundary exchange”; Lacan and Jeandel, 2001, 2005) on the  $\epsilon\text{Nd}$  values of intermediate water  
349 masses.

350 During its westward flow, the LIW continuously mixes with surrounding waters with different  $\epsilon\text{Nd}$  signatures  
351 lying above and below. For the western Mediterranean basin, these water masses are the MAW/Western  
352 Intermediate Water (WIW) and the TDW/WMDW, ~~respectively. Accordingly, a well defined and~~ As a result, a  
353 gradual  $\epsilon\text{Nd}$  gradient exists at intermediate depth between the eastern and western Mediterranean basins, with  
354 LIW values becoming progressively more unradiogenic towards the Strait of Gibraltar, from  $-4.8 \pm 0.2$  at 227 m  
355 in the Levantine basin to  $-10.4 \pm 0.2$  at 200 m in the Alboran Sea (Tachikawa et al., 2004). Such an  $\epsilon\text{Nd}$  pattern  
356 implies an effective vertical mixing with more unradiogenic water masses along the E-W LIW trajectory ruling  
357 out severe isotopic modifications of the LIW due to the local exchange between the continental margins and  
358 seawater. Unfortunately, no information exists on the potential temporal variability in  $\epsilon\text{Nd}$  of the Mediterranean  
359 water-mass end-members since the LGM.

360 It has been demonstrated that eolian dust input can modify the surface and sub-surface  $\epsilon\text{Nd}$  distribution of the  
361 ocean in some areas (Arsouze et al., 2009). The last glacial period was associated with an aridification of North  
362 Africa (Sarnthein et al., 1981; Hooghiemstra et al., 1987; Moreno et al., 2002; Wienberg et al., 2010) and higher  
363 fluxes of Saharan dust to the NE tropical Atlantic (Itambi et al., 2009) and the western Mediterranean Sea

364 characterized by unradiogenic  $\epsilon\text{Nd}$  values (between  $-11\pm 0.4$  and  $-14\pm 0.4$ ; see synthesis in Scheuvens et al.,  
365 2013). Bout-Roumazeilles et al. (2013) documented a dominant role of eolian supply in the Siculo-Tunisian  
366 Strait during the last 20 ka, with the exception of a significant riverine contribution (from the Nile River) and a  
367 strong reduction of eolian input during the sapropel S1 event. Such variations in the eolian input to the  
368 Mediterranean Sea are not associated to a significant change in the seawater  $\epsilon\text{Nd}$  record obtained for the Balearic  
369 Sea (core SU92-33) during the sapropel S1 event (Fig. 3). Furthermore, the  $\epsilon\text{Nd}$  signature of the CWC from the  
370 Sardinia Channel (core RECORD 23) shifts to more unradiogenic values ( $-8.66\pm 0.30$ ) during the sapropel S1  
371 event, which is opposite to what ~~would be expected if it was related to~~ a strong reduction of eolian sediment  
372 input. In a recent study, Rodrigo-Gámiz et al. (2015) have documented variations in the terrigenous provenance  
373 from a sediment record in the Alboran Sea (core 293G;  $36^{\circ}10.414'\text{N}$ ,  $2^{\circ}45.280'\text{W}$ , 1840 m water depth) since  
374 the LGM. Radiogenic isotopes (Sr, Nd, Pb) point to changes from North African dominated sources during the  
375 glacial period to European dominated source during the Holocene. Nevertheless, the major Sr-Nd-Pb excursions  
376 documented by Rodrigo-Gámiz et al. (2015) and dated at ca. 11.5, 10.2, 8.9-8.7, 5.6, 2.2 and 1.1. ka cal BP do  
377 not seem to affect the  $\epsilon\text{Nd}$  values of our foraminifera and coral records.

378 Taken together, these results suggest that changes of eolian dust input since the LGM ~~were not responsible for~~  
379 ~~cannot explain~~ the observed  $\epsilon\text{Nd}$  variability at intermediate water depths.

380 Consequently, assuming that the Nd isotopic budget of the western Mediterranean Sea has not been strongly  
381 modified since the LGM, the reconstructed variations of the E-W gradient of  $\epsilon\text{Nd}$  values in the western  
382 Mediterranean Sea for the past and notably during the sapropel S1 event (Fig. 3) are indicative of a major  
383 reorganization of intermediate water circulation.

384

### 385 ***5.1 Hydrological changes in the Alboran and Balearic Seas since the LGM***

386 The range in  $\epsilon\text{Nd}$  for the CWC from the Alboran Sea (from  $-9.22\pm 0.30$  to  $-8.8.59\pm 0.30$ ; Table 2) is very close to  
387 the one obtained for the planktonic foraminifera from the Balearic Sea (from  $-9.50\pm 0.30$  to  $-8.61\pm 0.30$ ; Table 4,  
388 Fig. 3c), suggesting that both sites are influenced by the same intermediate water masses at least for the last 13.5  
389 kyr BP. Today, LIW occupies a depth range between  $\sim 200$  and  $\sim 700$  m in the western Mediterranean basin  
390 (Millot, 1999; Sparnocchia et al., 1999). More specifically, the salinity maximum corresponding to the core of  
391 LIW is found at around 400 m in the Alboran Sea (Millot, 2009) and up to 550 m in the Balearic Sea (López-  
392 Jurado et al., 2008). The youngest CWC sample collected in the Alboran Sea with a rather "recent" age of 0.34  
393 cal ka BP (Fink et al. 2013) displays an  $\epsilon\text{Nd}$  value of  $-8.59\pm 0.30$  (Table 2) that is similar to the present-day value  
394 of the LIW at the same site ( $-8.3\pm 0.2$ ) (Dubois-Dauphin et al., submitted) and is significantly different from the  
395 WMDW  $\epsilon\text{Nd}$  signature in the Alboran Sea ( $-10.7\pm 0.2$ , 1270 m water depth; Tachikawa et al., 2004). Considering  
396 the intermediate depth range of the studied CWC and foraminifera samples, we can reasonably assume that  
397 samples from both sites, in the Balearic Sea (622 m water depth) and in the Alboran Sea (280 to 442 m water  
398 depth), record  $\epsilon\text{Nd}$  variations of the LIW. The  $\epsilon\text{Nd}$  record obtained from planktonic foraminifera generally  
399 displays more unradiogenic and homogenous values before  $\sim 13$  cal ka BP (range from  $-9.46$  to  $-9.12$ ) compared  
400 to the most recent part of the record (range from  $-9.50$  to  $-8.61$ ), with the highest value of  $-8.61\pm 0.3$  in the Early  
401 and Late Holocene.

402 The SST record displays values centered at around 12°C during the LGM with a subsequent rapid SST decrease  
403 towards 9°C, highlighting the onset of the HS1 (Fig. 2a). These values are well comparable to recent high-  
404 resolution SST data obtained in the Alboran Sea (Martrat et al., 2014; Rodrigo-Gámiz et al., 2014).

405 The  $\delta^{18}\text{O}$  record obtained on *G. bulloides* indicates an abrupt 1‰ excursion towards lighter values centered at  
406 about 16 cal ka BP (Table 4), synchronous with the HS1 (Fig. 2b), which is similar to the  $\delta^{18}\text{O}$  shift reported by  
407 Sierro et al. (2005) for a core collected at 2391 m water depth NE of the Balearic Islands (MD99-2343; Fig. 1).  
408 As the Heinrich events over the last glacial period are characterized by colder and fresher surface water in the  
409 Alboran Sea (Cacho et al., 1999; Pérez-Folgado et al., 2003; Martrat et al., 2004, 2014; Rodrigo-Gámiz et al.,  
410 2014) and dry climate on land over the western Mediterranean Sea (Allen et al., 1999; Combourieu-Nebout et  
411 al., 2002; Sanchez Goni et al., 2002; Bartov et al., 2003), lighter  $\delta^{18}\text{O}$  values of planktonic *G. bulloides* are  
412 thought to be the result of the inflow of freshwater derived from the melting of icebergs in the Atlantic Ocean  
413 into the Mediterranean Sea (Sierro et al., 2005; Rogerson et al., 2008).

414 During this time interval, the  $\delta^{13}\text{C}$  record of *C. pachyderma* from the Balearic Sea (core SU92-33) displays a  
415 decreasing  $\delta^{13}\text{C}$  trend after ~16 cal ka BP (from 1.4 ‰ to 0.9 ‰; Table 4; Fig. 4a). Moreover, the  $\delta^{13}\text{C}$  record  
416 obtained on benthic foraminifera *C. pachyderma* from the deep Balearic Sea (core MD99-2343) reveals similar  
417  $\delta^{13}\text{C}$  values before ~16 cal ka BP suggesting well-mixed and ventilated water masses during the LGM and the  
418 onset of the deglaciation (Sierro et al., 2005).

419 The slightly lower foraminiferal  $\epsilon\text{Nd}$  values before ~13 cal ka BP could reflect a stronger influence of water  
420 masses deriving from the Gulf of Lions as WMDW ( $\epsilon\text{Nd}$ :  $-9.4 \pm 0.9$ ; Henry et al., 1994; Tachikawa et al., 2004;  
421 [Montagna, pers. comm., 2016](#)~~Montagna et al., in prep~~). This is in agreement with  $\epsilon\text{Nd}$  results obtained by  
422 Jiménez-Espejo et al. (2015) from planktonic foraminifera collected from deep-water sites (1989 m and 2382 m)  
423 in the Alboran Sea (Fig. 4c). Jiménez-Espejo et al. (2015) documented lower  $\epsilon\text{Nd}$  values (ranging from -  
424  $10.14 \pm 0.27$  to  $-9.58 \pm 0.22$ ) during the LGM, suggesting an intense deep-water formation. This is also associated  
425 to an enhanced activity of the deeper branch of the MOW in the Gulf of Cádiz (Rogerson et al., 2005; Voelker et  
426 al., 2006) linked to the active production of the WMDW in the Gulf of Lions during the LGM (Jiménez-Espejo  
427 et al., 2015).

428 The end of the HS1 (14.7 cal ka BP) is concurrent with the onset of the B-A warm interval characterized by  
429 increased SST up to 14°C in the Balearic Sea (SU92-33; Fig. 3a), also identified for various sites in the  
430 Mediterranean Sea (Cacho et al., 1999; Martrat et al., 2004, 2014; Essallami et al., 2007; Rodrigo-Gámiz et al.,  
431 2014). The B-A interval is associated with the so-called melt-water pulse 1A (e.g. Weaver et al., 2003) occurring  
432 at around 14.5 cal ka BP. This led to a rapid sea-level rise of about 20 m in less than 500 years and large  
433 freshwater discharges in the Atlantic Ocean due to the melting of continental ice sheets (Deschamps et al., 2012),  
434 resulting in an enhanced Atlantic inflow across the Strait of Gibraltar. Synchronously, cosmogenic dating of  
435 Alpine glacier retreat throughout the western Mediterranean hinterland suggests maximum retreat rates (Ivy-  
436 Ochs et al., 2007; Kelly et al., 2006). Overall, these events are responsible for freshening Mediterranean waters  
437 and reduced surface water density, and hence, weakened ventilation of intermediate (Toucanne et al., 2012) and  
438 deep-water masses (Cacho et al., 2000; Sierro et al., 2005). Similarly, lower benthic  $\delta^{13}\text{C}$  values obtained for the  
439 Balearic Sea (Fig. 4a) point to less ventilated intermediate water relative to the late glacial. In addition, a  
440 decoupling in the benthic  $\delta^{13}\text{C}$  values is observed between deep (MD99-2343) and intermediate (core SU92-33)  
441 | waters after ~16 cal ka BP (Sierro et al. 2005), suggesting an enhanced stratification of the water masses (Fig.

442 4a). At this time, the shallowest  $\epsilon\text{Nd}$  record from the deep Alboran Sea (core 300G) shifted towards more  
443 radiogenic values, while the deepest one (core 304G) remained close to the LGM values (Jimenez-Espejo et al.,  
444 2015) (Fig. 4c). Furthermore, results from the UP10 fraction (particles  $> 10 \mu\text{m}$ ) of the MD99-2343 sediment  
445 core (Fig. 4d) indicate a declining bottom-current velocity at 15 ka BP (Frigola et al., 2008). Rogerson et al.  
446 (2008) have hypothesized that during deglacial periods the sinking depth of dense waters produced in the Gulf of  
447 Lions was shallower resulting in new intermediate water (WIW) rather than new deep-water (WMDW) as  
448 observed today during mild winters (Millot, 1999; Schott et al., 1996). Therefore, intermediate depths of the  
449 Balearic Sea could have been isolated from the deep-water with the onset of the T1 (at  $\sim 15$  ka BP). The reduced  
450 convection in the deep western Mediterranean Sea together with the shoaling of the nutricline (Rogerson et al.,  
451 2008) led to the deposition of the ORL 1 (14.5 to 8.2 ka B.P; Cacho et al., 2002) and dysoxic conditions below  
452 2000 m in agreement with the absence of epibenthic foraminifera such as *C. pachyderma* after 11 cal ka BP in  
453 MD99-2343 (Sierro et al., 2005) (Fig. 4a).

454 After 13.5 ka BP, planktonic foraminifera  $\epsilon\text{Nd}$  values from the Balearic Sea (core SU92-33) become more  
455 radiogenic and are in the range of CWC  $\epsilon\text{Nd}$  values from the Alboran Sea (Fig. 4b). These values may reveal a  
456 stronger influence of the LIW in the Balearic Sea during the Younger Dryas, as also supported by the sortable  
457 silt record from the Tyrrhenian Sea (Toucanne et al., 2012) (Fig. 4e). Deeper depths of the Alboran Sea also  
458 record a stronger influence of the LIW with an  $\epsilon\text{Nd}$  value of  $-9.1 \pm 0.4$  (Jimenez-Espejo et al., 2015). In addition,  
459 a concomitant activation of the upper MOW branch, as reconstructed from higher values of Zr/Al ratio in  
460 sediments of the Gulf of Cádiz, can be related to the enhanced LIW flow in the western Mediterranean Sea (Fig.  
461 4f) (Bahr et al., 2015).

462 The time of sapropel S1 deposition (10.2 – 6.4 ka) is characterized by a weakening or a shutdown of  
463 intermediate- and deep-water formation in the eastern Mediterranean basin (Rossignol-Strick et al., 1982; Cramp  
464 and O'Sullivan, 1999; Emeis et al., 2000; Rohling et al., 2015). At this time, planktonic foraminifera  $\epsilon\text{Nd}$  values  
465 from intermediate water depths in the Balearic Sea (core SU92-33) remain high (between  $-9.15 \pm 0.3$  and -  
466  $8.61 \pm 0.3$ ) (Fig. 4b). On the other hand, the deeper Alboran Sea provides a value of  $-9.8 \pm 0.3$  pointing to a  
467 stronger contribution of WMDW (Jimenez-Espejo et al., 2015), coeval with the recovery of deep-water activity  
468 from core MD99-2343 (Frigola et al., 2008).

469

470

### 5.2 Hydrological changes in the Sardinia Channel during the Holocene

471 The present-day hydrographic structure of the Sardinia Channel is characterized by four water masses, with the  
472 surface, intermediate and deep-water masses being represented by MAW, LIW and TDW/WMDW, respectively  
473 (Astraldi et al., 2002a; Millot and Taupier-Lepage, 2005). In addition, the WIW, flowing between the MAW and  
474 the LIW, has also been observed along the Channel (Sammari et al., 1999). The core of the LIW is located at  
475 400-450 m water depth in the Tyrrhenian Sea (Hopkins, 1988; Astraldi et al., 2002b), which is the depth range of  
476 CWC samples from the Sardinia Channel (RECORD 23; 414 m) (Taviani et al., 2015). The youngest CWC  
477 sample dated at  $\sim 0.1$  ka BP has an  $\epsilon\text{Nd}$  value of  $-7.70 \pm 0.10$  (Table 1, Fig. 5), which is similar within error to the  
478 value obtained from a seawater sample collected at 451 m close to the coral sampling location ( $-8.0 \pm 0.4$ ;  
479 [Montagna, pers. comm., 2016](#) [Montagna et al., in prep](#)).

480 The CWC dating from the Sardinia Channel shows three distinct periods of sustained coral occurrence in this  
481 area during the Holocene, with each displaying a large variability in  $\epsilon\text{Nd}$  values. CWC from the Early Holocene

482 (10.9-10.2 ka BP) and the Late Holocene (<1.5 ka BP) exhibit similar ranges of  $\epsilon\text{Nd}$  values (ranging from -  
483  $5.99\pm 0.50$  to  $-7.75\pm 0.20$ ; Table 1, Fig 5c). Such variations are within the present-day  $\epsilon\text{Nd}$  range being  
484 characteristic for intermediate waters in the eastern Mediterranean Sea ( $-6.6\pm 1.0$ ; Tachikawa et al., 2004; Vance  
485 et al., 2004). However, the CWC  $\epsilon\text{Nd}$  values are more radiogenic than those observed at mid-depth in the  
486 present-day western basin (ranging from  $-10.4\pm 0.2$  to  $-7.58\pm 0.47$ ; Henry et al., 1994; Tachikawa et al., 2004;  
487 [Montagna, pers. comm., 2016](#)~~Montagna et al., in prep~~), suggesting a stronger LIW component in the Sardinia  
488 Channel during the Early and Late Holocene. The Sardinian CWC  $\epsilon\text{Nd}$  variability also reflects the sensitivity of  
489 the LIW to changes in the eastern basin such as rapid variability of the Nile River flood discharge (Revel et al.,  
490 2014; 2015; Weldeab et al., 2014) or a modification through time in the proportion between the LIW and the  
491 Cretan Intermediate Water (CIW). Today, the intermediate water outflowing from the Strait of Sicily is  
492 composed by ~66 to 75 % of LIW and 33 to 25 % of CIW (Manca et al., 2006; Millot, 2014). As the CIW is  
493 formed in the Aegean Sea, this intermediate water mass is generally more radiogenic than LIW (Tachikawa et  
494 al., 2004; [Montagna, pers. comm., 2016](#)~~Montagna et al., in prep~~). Following this hypothesis, a modification of  
495 the mixing proportion between the CIW and the LIW may potentially explain values as radiogenic as about -6 in  
496 the Sardinia Channel during the Early and Late Holocene (Fig. 5c). However, a stronger LIW and/or a CIW  
497 contribution cannot be responsible for  $\epsilon\text{Nd}$  values as low as  $-8.66\pm 0.30$  observed during the sapropel S1 event at  
498 8.7 ka BP (Table 1, Fig. 5c). Considering that such unradiogenic value is not observed at intermediate depth in  
499 the modern eastern Mediterranean basin, the most plausible hypothesis suggested here is that the CWC were  
500 influenced by a higher contribution of intermediate water from the western basin.

501

### 502 *5.3 Hydrological implications for the intermediate water masses of the western Mediterranean Sea*

503 The  $\epsilon\text{Nd}$  records of the Balearic Sea, Alboran Sea and Sardinia Channel document a temporal variability of the  
504 east-west gradient in the western Mediterranean basin during the Holocene. The magnitude of the gradient  
505 ranges from ~1.5 to ~3  $\epsilon$  units during the Early and Late Holocene and it is strongly reduced at 8.7 ka BP ([from](#)  
506 [0 to ~0.5  \$\epsilon\$  unit](#)), coinciding with the sapropel S1 event affecting the eastern Mediterranean basin (Fig. 5). Such  
507 variations could be the result of a modification of the Nd isotopic composition of intermediate water masses due  
508 to changes of the LIW production through time and a higher contribution of the western-sourced intermediate  
509 water towards the Sardinia Channel coinciding with the sapropel S1 event.

510 The LIW acquires its radiogenic  $\epsilon\text{Nd}$  [signature](#) in the Mediterranean Levantine basin mainly from Nd exchange  
511 between seawater and lithogenic particles originating mainly from Nile River (Tachikawa et al., 2004). A higher  
512 sediment supply from the Nile River starting at ~15 ka BP was documented by a shift to more radiogenic  $\epsilon\text{Nd}$   
513 values of the terrigenous fraction obtained from a sediment core having been influenced by the Nile River  
514 discharge (Revel et al., 2015) (Fig. 5e). Others studies pointed to a gradual enhanced Nile River runoff as soon  
515 as 14.8 ka BP and a peak of Nile discharge from 9.7 to 8.4 ka recorded by large increase in sedimentation rate  
516 from 9.7 to 8.4 ka ( $>120$  cm/ka) (Revel et al., 2015; Weldeab et al., 2014; Castaneda et al., 2016). Similarly,  
517 enhanced Nile discharge at ~9.5 cal kyr B.P was inferred based on  $\delta^{18}\text{O}$  in planktonic foraminifera from a  
518 sediment core in the southeast Levantine Basin (PS009PC (32°07.7'N, 34°24.4'E; 552 m water depth)  
519 (Hennekam et al., 2014). This increasing contribution of the Nile River to the eastern Mediterranean basin has  
520 been related to the African Humid Period (14.8–5.5 ka BP; Shanahan et al., 2015), which in turn was linked to  
521 the precessional increase in Northern Hemisphere insolation during low eccentricity (deMenocal et al., 2000;

522 Barker et al., 2004; Garcin et al., 2009). An increasing amount of radiogenic sediments dominated by the  
523 Blue/Atbara Nile River contribution (Revel et al., 2014) could have modified the  $\epsilon\text{Nd}$  of surface water towards  
524 | more radiogenic values (~~Revel, pers. comm., 2016~~~~Revel et al., in prep~~). Indeed, planktonic foraminifera  $\epsilon\text{Nd}$   
525 values as high as  $\sim -3$  have been documented in the eastern Levantine Basin (ODP site 967;  $34^{\circ}04.27'\text{N}$ ,  
526  $32^{\circ}43.53'\text{E}$ ; 2553 m water depth) during the sapropel S1 event as a result of enhanced Nile flooding (Scrivner et  
527 al., 2004). The radiogenic signature was likely transferred to intermediate depth as a consequence of the LIW  
528 formation in the Rhodes Gyre, and it might have been propagated westwards towards the Sardinia Channel.  
529 Therefore, considering the more unradiogenic value of the CWC samples from the Sardinia Channel during the  
530 sapropel S1a event, it is very unlikely that eastern-sourced water flowed at intermediate depth towards the  
531 Sardinia Channel. A possible explanation could be the replacement of the radiogenic LIW that was no longer  
532 produced in the eastern basin (Rohling, 1994) by less radiogenic western intermediate water (possibly WIW).  
533 | Such a scenario could even support previous hypotheses ~~that invoke of~~ a potential circulation reversal in the  
534 eastern Mediterranean from anti-estuarine to estuarine during sapropel formation (Huang and Stanley, 1972;  
535 Calvert, 1983; Sarmiento et al., 1988; Buckley and Johnson, 1988; Thunell and Williams, 1989). An alternative  
536 hypothesis would be that reduced surface water densities in the eastern Mediterranean during sapropel S1  
537 | resulted in the LIW sinking to shallower depths than at present. ~~As a result of this shoaling~~~~In this case~~, CWC  
538 from the Sardinia Channel would have been bathed by underlying Western Intermediate Water during the  
539 sapropel S1a event.

540

## 541 **6. Conclusions**

542 The foraminiferal  $\epsilon\text{Nd}$  record from intermediate depths in the Balearic Sea reveals a relatively narrow range of  
543  $\epsilon\text{Nd}$  values varying between -9.50 and -8.61 since the LGM ( $\sim 20$  ka). Between 18 and 13.5 cal ka BP, the more  
544 unradiogenic  $\epsilon\text{Nd}$  values support a vigorous deep overturning in the Gulf of Lions while  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values  
545 indicate a stratification of the water masses after 16 cal ka BP. The stratification together with a decrease of the  
546 deep-water intensity led to more radiogenic values after  $\sim 13$  cal ka BP. The foraminiferal  $\epsilon\text{Nd}$  record, supported  
547 by  $\epsilon\text{Nd}$  values from CWC in the Alboran Sea, shows only minor changes in neodymium isotopes from 13.5 cal  
548 ka BP to 0.34 cal ka BP, suggesting that the westernmost part of the western Mediterranean basin is not very  
549 sensitive to hydrological variations of the LIW.

550 On the contrary, CWC located at the depth of the LIW in the Sardinia Channel exhibit large  $\epsilon\text{Nd}$  variations  
551 (between  $-7.75 \pm 0.10$  and  $-5.99 \pm 0.50$ ) during the Holocene, suggesting either the role of the Nile River in  
552 changing the  $\epsilon\text{Nd}$  of the LIW in the eastern Mediterranean basin or a variable LIW/CIW mixing of the water  
553 outflowing from the Strait of Sicily. At the time of the sapropel S1 event at  $\sim 8.7$  ka BP, CWC display a shift  
554 | toward lower values ( $-8.66 \pm 0.30$ ), similar to those ~~obtained~~~~found~~ at intermediate depths in the westernmost part  
555 of the western basin. This suggests that western-sourced intermediate water likely filled mid-depth of the  
556 southern Sardinia, replacing LIW that was no longer produced (or heavily reduced) in the eastern basin. These  
557 results could potentially support a reversal of the Mediterranean circulation, although this assumption needs  
558 further investigation to be confirmed.

559

## 560 **Acknowledgements**

561 | The research leading to this study has received funding from [the MISTRALS/PALEOMEX/COFIMED](#), the  
562 | French National Research Agency “Investissement d’Avenir” (n°ANR-10-LABX-0018), the HAMOC project  
563 | ANR-13-BS06-0003, ~~the MISTRALS/PALEOMEX/COFIMED~~ and ENVIMED/Boron Isotope and Trace  
564 | Elements project. This work contributes to the RITMARE project. We thank Hiske Fink for selecting and kindly  
565 | providing the cold-water corals samples from the Alboran Sea. We further thank François Thil and Louise  
566 | Bordier for their support with Nd isotopic composition analyses. Paolo Montagna is grateful for financial support  
567 | from the Short Term Mobility Program (CNR). Thanks are also extended to the captains, crews, chief scientists,  
568 | and scientific parties of research cruises RECORD (R/V Urania), POS-385 (R/V Poseidon) and PALEOCINAT  
569 | II (R/V Le Suroît).

570

## 571 | **References**

572 | Abu-Zied, R. H., Rohling, E. J., Jorissen, F. J., Fontanier, C., Casford, J. S. L. and Cooke, S.: Benthic  
573 | foraminiferal response to changes in bottom-water oxygenation and organic carbon flux in the eastern  
574 | Mediterranean during LGM to Recent times, *Mar. Micropaleontol.*, 67(1-2), 46–68,  
575 | doi:10.1016/j.marmicro.2007.08.006, 2008.

576 | Allen, J. R. M., Huntley, B., Brandt, U., Brauer, A., Hubberten, H., Keller, J., Kraml, M., Mackensen, A.,  
577 | Mingram, J., Negendank, J. F. W., Nowaczyk, N. R., Oberhänsli, H., Watts, W. A., Wulf, S. and Zolitschka, B.:  
578 | Rapid environmental changes in southern Europe during the last glacial period, *Nature*, 400(6746), 740–743,  
579 | doi:10.1038/23432, 1999.

580 | Andersen, M. B., Stirling, C. H., Zimmermann, B. and Halliday, A. N.: Precise determination of the open ocean  
581 |  $^{234}\text{U}/^{238}\text{U}$  composition, *Geochemistry, Geophys. Geosystems*, 11(12), Q12003, doi:10.1029/2010GC003318,  
582 | 2010.

583 | Arsouze, T., Dutay, J.-C., Lacan, F. and Jeandel, C.: Reconstructing the Nd oceanic cycle using a coupled  
584 | dynamical – biogeochemical model, *Biogeosciences*, 6(12), 2829–2846, doi:10.5194/bg-6-2829-2009, 2009.

585 | Astraldi, M., Gasparini, G. P., Gervasio, L. and Salusti, E.: Dense Water Dynamics along the Strait of Sicily  
586 | (Mediterranean Sea), *J. Phys. Oceanogr.*, 31(12), 3457–3475, doi:10.1175/1520-  
587 | 0485(2001)031<3457:DWDATS>2.0.CO;2, 2001.

588 | Astraldi, M., Gasparini, G. P., Vetrano, A. and Vignudelli, S.: Hydrographic characteristics and interannual  
589 | variability of water masses in the central Mediterranean: A sensitivity test for long-term changes in the  
590 | Mediterranean Sea, *Deep. Res. Part I Oceanogr. Res. Pap.*, 49(4), 661–680, doi:10.1016/S0967-0637(01)00059-  
591 | 0, 2002a.

592 | Astraldi, M., Conversano, F., Civitarese, G., Gasparini, G. P., Ribera d’Alcalà, M. and Vetrano, a.: Water mass  
593 | properties and chemical signatures in the central Mediterranean region, *J. Mar. Syst.*, 33-34, 155–177,  
594 | doi:10.1016/S0924-7963(02)00057-X, 2002b.

595 | Bahr, A., Kaboth, S., Jiménez-Espejo, F. J., Sierro, F. J., Voelker, A. H. L., Lourens, L., Röhl, U., Reichert, G.



596 J., Escutia, C., Hernández-Molina, F. J., Pross, J. and Friedrich, O.: Persistent monsoonal forcing of  
597 Mediterranean Outflow Water dynamics during the late Pleistocene, *Geology*, 43(11), 951–954,  
598 doi:10.1130/G37013.1, 2015.

599 Barker, P. A., Talbot, M. R., Street-Perrott, F. A., Marret, F., Scourse, J. and Odada, E. O.: Late Quaternary  
600 climatic variability in intertropical Africa, in *Past Climate Variability through Europe and Africa*, pp. 117–138,  
601 Springer Netherlands, Dordrecht., 2004.

602 Bartov, Y., Goldstein, S. L., Stein, M. and Enzel, Y.: Catastrophic arid episodes in the Eastern Mediterranean  
603 linked with the North Atlantic Heinrich events, *Geology*, 31(5), 439, doi:10.1130/0091-  
604 7613(2003)031<0439:CAEITE>2.0.CO;2, 2003.

605 Bigg, G. R. and Wadley, M. R.: Millennial-scale variability in the oceans: an ocean modelling view, *J. Quat.*  
606 *Sci.*, 16(4), 309–319, doi:10.1002/jqs.599, 2001.

607 Bout-Roumazielles, V., Combourieu-Nebout, N., Desprat, S., Siani, G., Turon, J. L. and Essallami, L.: Tracking  
608 atmospheric and riverine terrigenous supplies variability during the last glacial and the Holocene in central  
609 Mediterranean, *Clim. Past*, 9(3), 1065–1087, doi:10.5194/cp-9-1065-2013, 2013.

610 Buckley, H. A. and Johnson, L. R.: Late pleistocene to recent sediment deposition in the central and western  
611 Mediterranean, *Deep Sea Res. Part A. Oceanogr. Res. Pap.*, 35(5), 749–766, doi:10.1016/0198-0149(88)90028-  
612 3, 1988.

613 Cacho, I., Pelejero, C., Grimalt, J. O., Calafat, A. and Canals, M.: C37 alkenone measurements of sea surface  
614 temperature in the Gulf of Lions (NW Mediterranean), *Org. Geochem.*, 30(7), 557–566, doi:10.1016/S0146-  
615 6380(99)00038-8, 1999.

616 Cacho, I., Grimalt, J. O., Sierro, F. J., Shackleton, N. and Canals, M.: Evidence for enhanced Mediterranean  
617 thermohaline circulation during rapid climatic coolings, *Earth Planet. Sci. Lett.*, 183(3-4), 417–429,  
618 doi:10.1016/S0012-821X(00)00296-X, 2000.

619 Cacho, I., Grimalt, J. O., Canals, M., Sbaiffi, L., Shackleton, N. J., Schönfeld, J. and Zahn, R.: Variability of the  
620 western Mediterranean Sea surface temperature during the last 25,000 years and its connection with the Northern  
621 Hemisphere climatic changes, *Paleoceanography*, 16(1), 40–52, doi:10.1029/2000PA000502, 2001.

622 Cacho, I., Grimalt, J. O. and Canals, M.: Response of the Western Mediterranean Sea to rapid climatic variability  
623 during the last 50,000 years: a molecular biomarker approach, *J. Mar. Syst.*, 33-34, 253–272,  
624 doi:10.1016/S0924-7963(02)00061-1, 2002.

625 Cacho, I., Shackleton, N., Elderfield, H., Sierro, F. J. and Grimalt, J. O.: Glacial rapid variability in deep-water  
626 temperature and  $\delta^{18}\text{O}$  from the Western Mediterranean Sea, *Quat. Sci. Rev.*, 25(23-24), 3294–3311,  
627 doi:10.1016/j.quascirev.2006.10.004, 2006.

628 Calvert, S. E.: Geochemistry of Pleistocene sapropels and associated sediments from the Eastern Mediterranean,  
629 *Oceanol. Acta*, 6(3), 255–267, 1983.

630 Castañeda, I. S., Schouten, S., Pätzold, J., Lucassen, F., Kasemann, S., Kuhlmann, H. and Schefuß, E.:  
631 Hydroclimate variability in the Nile River Basin during the past 28,000 years, *Earth Planet. Sci. Lett.*, 438, 47–  
632 56, doi:10.1016/j.epsl.2015.12.014, 2016.

633 Cheng, H., Lawrence Edwards, R., Shen, C.-C., Polyak, V. J., Asmerom, Y., Woodhead, J. D., Hellstrom, J.,  
634 Wang, Y., Kong, X., Spötl, C., Wang, X. and Calvin Alexander, E.: Improvements in  $^{230}\text{Th}$  dating,  $^{230}\text{Th}$  and  
635  $^{234}\text{U}$  half-life values, and U–Th isotopic measurements by multi-collector inductively coupled plasma mass  
636 spectrometry, *Earth Planet. Sci. Lett.*, 371–372, 82–91, doi:10.1016/j.epsl.2013.04.006, 2013.

637 Colin, C., Frank, N., Copard, K. and Douville, E.: Neodymium isotopic composition of deep-sea corals from the  
638 NE Atlantic: implications for past hydrological changes during the Holocene, *Quat. Sci. Rev.*, 29(19–20), 2509–  
639 2517, doi:10.1016/j.quascirev.2010.05.012, 2010.

640 Combourieu-Nebout, N., Turon, J. L., Zahn, R., Capotondi, L., Londeix, L. and Pahnke, K.: Enhanced aridity  
641 and atmospheric high-pressure stability over the western Mediterranean during the North Atlantic cold events of  
642 the past 50 k.y, *Geology*, 30(10), 863–866, doi:10.1130/0091-7613(2002)030<0863:EAAAHP>2.0.CO;2, 2002.

643 Copard, K., Colin, C., Douville, E., Freiwald, A., Gudmundsson, G., De Mol, B. and Frank, N.: Nd isotopes in  
644 deep-sea corals in the North-eastern Atlantic, *Quat. Sci. Rev.*, 29(19–20), 2499–2508,  
645 doi:10.1016/j.quascirev.2010.05.025, 2010.

646 Copard, K., Colin, C., Henderson, G. M., Scholten, J., Douville, E., Sicre, M.-A. and Frank, N.: Late Holocene  
647 intermediate water variability in the northeastern Atlantic as recorded by deep-sea corals, *Earth Planet. Sci. Lett.*,  
648 313–314, 34–44, doi:10.1016/j.epsl.2011.09.047, 2012.

649 Cramp, A. and O’Sullivan, G.: Neogene sapropels in the Mediterranean: a review, *Mar. Geol.*, 153(1–4), 11–28,  
650 doi:10.1016/S0025-3227(98)00092-9, 1999.

651 De Lange, G. J., Thomson, J., Reitz, A., Slomp, C. P., Principato, M. S., Erba, E. and Corselli, C.: Synchronous  
652 basin-wide formation and redox-controlled preservation of a Mediterranean sapropel, *Nat. Geosci.*, 1(9), 606–  
653 610, 2008.

654 DeMenocal, P., Ortiz, J., Guilderson, T. and Sarnthein, M.: Coherent High- and Low-Latitude Climate  
655 Variability During the Holocene Warm Period, *Science* (80-. ), 288(5474), 2198–2202,  
656 doi:10.1126/science.288.5474.2198, 2000.

657 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., Henderson, G. M., Okuno, J. and  
658 Yokoyama, Y.: Ice-sheet collapse and sea-level rise at the Bølling warming 14,600 years ago, *Nature*,  
659 483(7391), 559–564, doi:10.1038/nature10902, 2012.

660 Douville, E., Sallé, E., Frank, N., Eisele, M., Pons-Branchu, E. and Ayrault, S.: Rapid and accurate U–Th dating  
661 of ancient carbonates using inductively coupled plasma-quadrupole mass spectrometry, *Chem. Geol.*, 272(1–4),  
662 1–11, doi:10.1016/j.chemgeo.2010.01.007, 2010.

663 Dubois-Dauphin, Q., Colin, C., Bonneau, L., Montagna, P., Wu, Q., Van Rooij, D., Reverdin, G., Douville, E.,  
664 Thil, F., Waldner, A., Frank, N.: Fingerprinting North-east Atlantic water masses using Neodymium isotopes,  
665 [EPSL-GCA, Submitted](#)  
666

667 Elmore, A. C., Piotrowski, A. M., Wright, J. D. and Scrivner, A. E.: Testing the extraction of past seawater Nd  
668 isotopic composition from North Atlantic deep sea sediments and foraminifera, *Geochemistry, Geophys.*  
669 *Geosystems*, 12(9), doi:10.1029/2011GC003741, 2011.

670 Emeis, K.-C., Sakamoto, T., Wehausen, R. and Brumsack, H.-J.: The sapropel record of the eastern  
671 Mediterranean Sea — results of Ocean Drilling Program Leg 160, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*,  
672 158(3-4), 371–395, doi:10.1016/S0031-0182(00)00059-6, 2000.

673 Fink, H. G., Wienberg, C., De Pol-Holz, R., Wintersteller, P. and Hebbeln, D.: Cold-water coral growth in the  
674 Alboran Sea related to high productivity during the Late Pleistocene and Holocene, *Mar. Geol.*, 339, 71–82,  
675 doi:10.1016/j.margeo.2013.04.009, 2013.

676 Frigola, J., Moreno, A., Cacho, I., Canals, M., Sierro, F. J., Flores, J. a., Grimalt, J. O., Hodell, D. a. and Curtis,  
677 J. H.: Holocene climate variability in the western Mediterranean region from a deepwater sediment record,  
678 *Paleoceanography*, 22(2), n/a–n/a, doi:10.1029/2006PA001307, 2007.

679 Frigola, J., Moreno, A., Cacho, I., Canals, M., Sierro, F. J., Flores, J. A. and Grimalt, J. O.: Evidence of abrupt  
680 changes in Western Mediterranean Deep Water circulation during the last 50kyr: A high-resolution marine  
681 record from the Balearic Sea, *Quat. Int.*, 181(1), 88–104, doi:10.1016/j.quaint.2007.06.016, 2008.

682 Garcin, Y., Junginger, A., Melnick, D., Olago, D. O., Strecker, M. R. and Trauth, M. H.: Late Pleistocene–  
683 Holocene rise and collapse of Lake Suguta, northern Kenya Rift, *Quat. Sci. Rev.*, 28(9-10), 911–925,  
684 doi:10.1016/j.quascirev.2008.12.006, 2009.

685 Hebbeln, D., Wienberg, C., Beuck, L., Freiwald, A., Wintersteller, P. and cruise participants (2009) Report and  
686 preliminary results of R/V POSEIDON Cruise 385 "Cold-water corals of the Alboran Sea (western  
687 Mediterranean Sea)", Faro - Toulon, 29.5. - 16.6.2009. Reports of the Department of Geosciences at the  
688 University of Bremen, No. 273. Department of Geosciences, Bremen University. urn:nbn:de:gbv:46-  
689 ep000106508.

690 Hennekam, R., Jilbert, T., Schnetger, B. and De Lange, G. J.: Solar forcing of Nile discharge and sapropel S1  
691 formation in the early to middle Holocene eastern Mediterranean, *Paleoceanography*, 29(5), 343–356,  
692 doi:10.1002/2013PA002553, 2014.

693 Henry, F., Jeandel, C., Dupré, B. and Minster, J.-F.: Particulate and dissolved Nd in the western Mediterranean  
694 Sea: Sources, fate and budget, *Mar. Chem.*, 45(4), 283–305, doi:10.1016/0304-4203(94)90075-2, 1994.

695 Hooghiemstra, H., Bechler, A. and Beug, H.-J.: Isopollen maps for 18,000 years B.P. of the Atlantic offshore of  
696 northwest Africa: Evidence for paleowind circulation, *Paleoceanography*, 2, 561–582,  
697 doi:10.1029/PA002i006p00561, 1987.

- 698 Hopkins, T. S.: Recent observations on the intermediate and deep water circulation in the Southern Tyrrhenian  
699 Sea, *Oceanol. Acta*, (Special issue), 41–50, 1988.
- 700 Huang, T. C. and Stanley, D. J.: Western Alboran sea: sediment dispersal, pouncing and reversal of currents, in  
701 *The Mediterranean Sea: A Natural Sedimentation Laboratory*, pp. 521–559, Dowden, Hutchinson & Ross,  
702 Stroudsburg, PA., 1972.
- 703 Hutson, W. H.: The Agulhas Current During the Late Pleistocene: Analysis of Modern Faunal Analogs, *Science*  
704 (80-. ), 207(4426), 64–66, doi:10.1126/science.207.4426.64, 1980.
- 705 Itambi, a. C., von Dobeneck, T., Mulitza, S., Bickert, T. and Heslop, D.: Millennial-scale northwest African  
706 droughts related to Heinrich events and Dansgaard-Oeschger cycles: Evidence in marine sediments from  
707 offshore Senegal, *Paleoceanography*, 24(1), PA1205, doi:10.1029/2007PA001570, 2009.
- 708 Ivy-Ochs, S., Kerschner, H. and Schlüchter, C.: Cosmogenic nuclides and the dating of Lateglacial and Early  
709 Holocene glacier variations: The Alpine perspective, *Quat. Int.*, 164-165, 53–63,  
710 doi:10.1016/j.quaint.2006.12.008, 2007.
- 711 Jacobsen, S. B. and Wasserburg, G. J.: Sm-Nd isotopic evolution of chondrites, *Earth Planet. Sci. Lett.*, 50(1),  
712 139–155, doi:10.1016/0012-821X(80)90125-9, 1980.
- 713 Jaffey, A. H., Flynn, K. F., Glendenin, L. E., Bentley, W. C. and Essling, A. M.: Precision measurements of half-  
714 lives and specific activities of <sup>235</sup>U and <sup>238</sup>U, *Phys. Rev. C*, 4(5), 1889–1906, doi:10.1103/PhysRevC.4.1889,  
715 1971.
- 716 Jiménez-Espejo, F. J., Pardos-Gené, M., Martínez-Ruiz, F., García-Alix, A., van de Flierdt, T., Toyofuku, T.,  
717 Bahr, A. and Kreissig, K.: Geochemical evidence for intermediate water circulation in the westernmost  
718 Mediterranean over the last 20kyrBP and its impact on the Mediterranean Outflow, *Glob. Planet. Change*, 135,  
719 38–46, doi:10.1016/j.gloplacha.2015.10.001, 2015.
- 720 Kallel, N., Paterne, M., Labeyrie, L., Duplessy, J.-C. and Arnold, M.: Temperature and salinity records of the  
721 Tyrrhenian Sea during the last 18,000 years, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 135(1-4), 97–108,  
722 doi:10.1016/S0031-0182(97)00021-7, 1997.
- 723 Kelly, M. A., Ivy-Ochs, S., Kubik, P. W., Von Blanckenburg, F. and Schlüchter, C.: Chronology of deglaciation  
724 based on <sup>10</sup>Be dates of glacial erosional features in the Grimsel Pass region, central Swiss Alps, *Boreas*, 35(4),  
725 634–643, doi:10.1111/j.1502-3885.2006.tb01169.x, 2006.
- 726 Khelifi, N., Sarnthein, M., Andersen, N., Blanz, T., Frank, M., Garbe-Schonberg, D., Haley, B. a., Stumpf, R.  
727 and Weinelt, M.: A major and long-term Pliocene intensification of the Mediterranean outflow, 3.5-3.3 Ma ago,  
728 *Geology*, 37(9), 811–814, doi:10.1130/G30058A.1, 2009.
- 729 Kinder, T. H. and Parrilla, G.: Yes, some of the Mediterranean water does come from great depth, *J. Geophys.*  
730 *Res.*, 92, 2901–2906, doi:10.1029/JC092iC03p02901, 1987.

- 731 Kuhnt, T., Schmiedl, G., Ehrmann, W., Hamann, Y. and Hemleben, C.: Deep-sea ecosystem variability of the  
732 Aegean Sea during the past 22 kyr as revealed by Benthic Foraminifera, *Mar. Micropaleontol.*, 64(3-4), 141–  
733 162, doi:10.1016/j.marmicro.2007.04.003, 2007.
- 734 Kuhnt, T., Schmiedl, G., Ehrmann, W., Hamann, Y. and Andersen, N.: Stable isotopic composition of Holocene  
735 benthic foraminifers from the Eastern Mediterranean Sea: Past changes in productivity and deep water  
736 oxygenation, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 268(1-2), 106–115, doi:10.1016/j.palaeo.2008.07.010,  
737 2008.
- 738 Lacan, F. and Jeandel, C.: Tracing Papua New Guinea imprint on the central Equatorial Pacific Ocean using  
739 neodymium isotopic compositions and Rare Earth Element patterns, *Earth Planet. Sci. Lett.*, 186(3-4), 497–512,  
740 doi:10.1016/S0012-821X(01)00263-1, 2001.
- 741 Lacan, F. and Jeandel, C.: Neodymium isotopes as a new tool for quantifying exchange fluxes at the continent–  
742 ocean interface, *Earth Planet. Sci. Lett.*, 232(3-4), 245–257, doi:10.1016/j.epsl.2005.01.004, 2005.
- 743 Lascaratos, A. and Nittis, K.: A high-resolution three-dimensional numerical study of intermediate water  
744 formation in the Levantine Sea, *J. Geophys. Res.*, 103(C9), 18497, doi:10.1029/98JC01196, 1998.
- 745 Lascaratos, A., Williams, R. G. and Tragou, E.: A mixed-layer study of the formation of Levantine intermediate  
746 water, *J. Geophys. Res.*, 98(C8), 14739, doi:10.1029/93JC00912, 1993.
- 747 López Correa, M., Montagna, P., Joseph, N., Rüggeberg, A., Fietzke, J., Flögel, S., Dorschel, B., Goldstein, S.  
748 L., Wheeler, A. and Freiwald, A.: Preboreal onset of cold-water coral growth beyond the Arctic Circle revealed  
749 by coupled radiocarbon and U-series dating and neodymium isotopes, *Quat. Sci. Rev.*, 34, 24–43,  
750 doi:10.1016/j.quascirev.2011.12.005, 2012.
- 751 López-Jurado, J. L., Marcos, M. and Monserrat, S.: Hydrographic conditions affecting two fishing grounds of  
752 Mallorca island (Western Mediterranean): during the IDEA Project (2003-2004), *J. Mar. Syst.*, 71(3-4), 303–  
753 315, doi:10.1016/j.jmarsys.2007.03.007, 2008.
- 754 Ludwig, K. R. and Titterton, D. M.: Calculation of  $^{230}\text{Th}/\text{U}$  isochrons, ages, and errors, *Geochim.  
755 Cosmochim. Acta*, 58(22), 5031–5042, doi:http://dx.doi.org/10.1016/0016-7037(94)90229-1, 1994.
- 756 Lugmair, G. W., Shimamura, T., Lewis, R. S. and Anders, E.: Samarium-146 in the Early Solar System:  
757 Evidence from Neodymium in the Allende Meteorite, *Science* (80-. ), 222(4627), 1015–1018,  
758 doi:10.1126/science.222.4627.1015, 1983.
- 759 Malanotte-Rizzoli, P., Manca, B. B., D’Alcala, M. R., Theocharis, A., Brenner, S., Budillon, G. and Ozsoy, E.:  
760 The Eastern Mediterranean in the 80s and in the 90s: the big transition in the intermediate and deep circulations,  
761 *Dyn. Atmos. Ocean.*, 29(2-4), 365–395, doi:10.1016/S0377-0265(99)00011-1, 1999.
- 762 Manca, B., Ibello, V., Pacciaroni, M., Scarazzato, P. and Giorgetti, A.: Ventilation of deep waters in the Adriatic  
763 and Ionian Seas following changes in thermohaline circulation of the Eastern Mediterranean, *Clim. Res.*, 31,  
764 239–256, doi:10.3354/cr031239, 2006.

765 Martrat, B., Grimalt, J. O., Lopez-Martinez, C., Cacho, I., Sierro, F. J., Flores, J. A., Zahn, R., Canals, M.,  
766 Curtis, J. H. and Hodell, D. a: Abrupt temperature changes in the Western Mediterranean over the past 250,000  
767 years., *Science* (80-. ), 306(5702), 1762–1765, doi:10.1126/science.1101706, 2004.

768 Martrat, B., Jimenez-Amat, P., Zahn, R. and Grimalt, J. O.: Similarities and dissimilarities between the last two  
769 deglaciations and interglaciations in the North Atlantic region, *Quat. Sci. Rev.*, 99, 122–134,  
770 doi:10.1016/j.quascirev.2014.06.016, 2014.

771 Melki, T., Kallel, N., Jorissen, F. J., Guichard, F., Dennielou, B., Berné, S., Labeyrie, L. and Fontugne, M.:  
772 Abrupt climate change, sea surface salinity and paleoproductivity in the western Mediterranean Sea (Gulf of  
773 Lion) during the last 28 kyr, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 279(1-2), 96–113,  
774 doi:10.1016/j.palaeo.2009.05.005, 2009.

775

776 Mercone, D., Thomson, J., Croudace, I. W., Siani, G., Paterne, M. and Troelstra, S.: Duration of S1, the most  
777 recent sapropel in the eastern Mediterranean Sea, as indicated by accelerator mass spectrometry radiocarbon and  
778 geochemical evidence, *Paleoceanography*, 15(3), 336–347, doi:10.1029/1999PA000397, 2000.

779 Mercone, D., Thomson, J., Abu-Zied, R. H., Croudace, I. W. and Rohling, E. J.: High-resolution geochemical  
780 and micropalaeontological profiling of the most recent eastern Mediterranean sapropel, *Mar. Geol.*, 177(1-2),  
781 25–44, doi:10.1016/S0025-3227(01)00122-0, 2001.

782 Millot, C.: Circulation in the Western Mediterranean Sea, *J. Mar. Syst.*, 20(1-4), 423–442, doi:10.1016/S0924-  
783 7963(98)00078-5, 1999.

784 Millot, C.: Another description of the Mediterranean Sea outflow, *Prog. Oceanogr.*, 82(2), 101–124,  
785 doi:10.1016/j.pocean.2009.04.016, 2009.

786 Millot, C.: Heterogeneities of in- and out-flows in the Mediterranean Sea, *Prog. Oceanogr.*, 120, 254–278,  
787 doi:10.1016/j.pocean.2013.09.007, 2014.

788 Millot, C. and Taupier-Letage, I.: Circulation in the Mediterranean Sea, in *Environmental Chemistry*, vol. 5,  
789 edited by A. Saliot, pp. 29–66, Springer Berlin Heidelberg, Heidelberg., 2005.

790 Millot, C., Candela, J., Fuda, J.-L. and Tber, Y.: Large warming and salinification of the Mediterranean outflow  
791 due to changes in its composition, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 53(4), 656–666,  
792 doi:10.1016/j.dsr.2005.12.017, 2006.

793 Montero-Serrano, J.-C., Frank, N., Colin, C., Wienberg, C. and Eisele, M.: The climate influence on the mid-  
794 depth Northeast Atlantic gyres viewed by cold-water corals, *Geophys. Res. Lett.*, 38(19),  
795 doi:10.1029/2011GL048733, 2011.

796 Montero-Serrano, J.-C., Frank, N., Tisnérat-Laborde, N., Colin, C., Wu, C., Lin, K., Shen, C., Copard, K.,  
797 Orejas, C., Gori, A., De Mol, L., Van Rooij, D., Reverdin, G. and Douville, E.: Decadal changes in the mid-  
798 depth water mass dynamic of the Northeastern Atlantic margin (Bay of Biscay), *Earth Planet. Sci. Lett.*, 364,  
799 134–144, doi:10.1016/j.epsl.2013.01.012, 2013.

800 Moreno, A., Cacho, I., Canals, M., Prins, M. a., Sánchez-Goñi, M.-F., Grimal, O. J. and Weltje, G. J.: Saharan  
801 Dust Transport and High-Latitude Glacial Climatic Variability: The Alboran Sea Record, *Quat. Res.*, 58, 318–  
802 328, doi:10.1006/qres.2002.2383, 2002.

803 Moreno, A., Cacho, I., Canals, M., Grimalt, J. O., Sánchez-Goñi, M. F., Shackleton, N. and Sierro, F. J.: Links  
804 between marine and atmospheric processes oscillating on a millennial time-scale. A multi-proxy study of the last  
805 50,000yr from the Alboran Sea (Western Mediterranean Sea), *Quat. Sci. Rev.*, 24(14-15), 1623–1636,  
806 doi:10.1016/j.quascirev.2004.06.018, 2005.

807 Myers, P. G., Haines, K. and Rohling, E. J.: Modeling the paleocirculation of the Mediterranean: The Last  
808 Glacial Maximum and the Holocene with emphasis on the formation of sapropel S1, *Paleoceanography*, 13(6),  
809 586–606, doi:10.1029/98PA02736, 1998.

810 Ovchinnikov, I. M.: The formation of intermediate water in the Mediterranean, *Oceanology*, 24, 168–173, 1984.

811 Overpeck, J. T., Webb, T. and Prentice, I. C.: Quantitative interpretation of fossil pollen spectra: Dissimilarity  
812 coefficients and the method of modern analogs, *Quat. Res.*, 23(1), 87–108, doi:10.1016/0033-5894(85)90074-2,  
813 1985.

814 Paterne, M., Kallel, N., Labeyrie, L., Vautravers, M., Duplessy, J.-C., Rossignol-Strick, M., Cortijo, E., Arnold,  
815 M. and Fontugne, M.: Hydrological relationship between the North Atlantic Ocean and the Mediterranean Sea  
816 during the past 15-75 kyr, *Paleoceanography*, 14(5), 626–638, doi:10.1029/1998PA900022, 1999.

817 Pérez-Folgado, M., Sierro, F. J., Flores, J. A., Cacho, I., Grimalt, J. O., Zahn, R. and Shackleton, N.: Western  
818 Mediterranean planktonic foraminifera events and millennial climatic variability during the last 70 kyr, *Mar.*  
819 *Micropaleontol.*, 48(1-2), 49–70, doi:10.1016/S0377-8398(02)00160-3, 2003.

820 Pinardi, N. and Masetti, E.: Variability of the large scale general circulation of the Mediterranean Sea from  
821 observations and modelling: a review, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 158(3-4), 153–173,  
822 doi:10.1016/S0031-0182(00)00048-1, 2000.

823 Piotrowski, A. M., Galy, A., Nicholl, J. a. L., Roberts, N. L., Wilson, D. J., Clegg, J. a. and Yu, J.:  
824 Reconstructing deglacial North and South Atlantic deep water sourcing using foraminiferal Nd isotopes, *Earth*  
825 *Planet. Sci. Lett.*, 357-358, 289–297, doi:10.1016/j.epsl.2012.09.036, 2012.

826 Pons-Branchu, E., Douville, E., Roy-Barman, M., Dumont, E., Branchu, P., Thil, F., Frank, N., Bordier, L. and  
827 Borst, W.: A geochemical perspective on Parisian urban history based on U–Th dating, laminae counting and  
828 yttrium and REE concentrations of recent carbonates in underground aqueducts, *Quat. Geochronol.*, 24, 44–53,  
829 doi:10.1016/j.quageo.2014.08.001, 2014.

830 Prell, W. L.: Stability of low-latitude sea-surface temperatures: an evaluation of the CLIMAP reconstruction  
831 with emphasis on the positive SST anomalies. Final report, Providence, RI (USA)., 1985.

832 Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., Grootes, P. M.,  
833 Guilderson, T. P., Hafliðason, H., Hajdas, I., HattĹ, C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G., Hughen, K.

834 A., Kaiser, K. F., Kromer, B., Manning, S. W., Niu, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon,  
835 J. R., Staff, R. A., Turney, C. S. M., & van der Plicht, J. (2013). IntCal13 and Marine13 Radiocarbon Age  
836 Calibration Curves 0-50,000 Years cal BP. *Radiocarbon*, 55(4).

837 Revel, M., Colin, C., Bernasconi, S., Combourieu-Nebout, N., Ducassou, E., Grousset, F. E., Rolland, Y.,  
838 Migeon, S., Bosch, D., Brunet, P., Zhao, Y. and Mascle, J.: 21,000 Years of Ethiopian African monsoon  
839 variability recorded in sediments of the western Nile deep-sea fan, *Reg. Environ. Chang.*, 14(5), 1685–1696,  
840 doi:10.1007/s10113-014-0588-x, 2014.

841 Revel, M., Ducassou, E., Skonieczny, C., Colin, C., Bastian, L., Bosch, D., Migeon, S. and Mascle, J.: 20,000  
842 years of Nile River dynamics and environmental changes in the Nile catchment area as inferred from Nile upper  
843 continental slope sediments, *Quat. Sci. Rev.*, 130, 200–221, doi:10.1016/j.quascirev.2015.10.030, 2015.

844 Roberts, N. L., Piotrowski, A. M., McManus, J. F. and Keigwin, L. D.: Synchronous deglacial overturning and  
845 water mass source changes., *Science*, 327(2010), 75–78, doi:10.1126/science.1178068, 2010.

846 Rodrigo-Gámiz, M., Martínez-Ruiz, F., Rampen, S. W., Schouten, S. and Sinninghe Damsté, J. S.: Sea surface  
847 temperature variations in the western Mediterranean Sea over the last 20 kyr: A dual-organic proxy (U K' 37 and  
848 LDI) approach, *Paleoceanography*, 29(2), 87–98, doi:10.1002/2013PA002466, 2014.

849 Rodrigo-Gámiz, M., Martínez-Ruiz, F., Chiaradia, M., Jiménez-Espejo, F. J. and Ariztegui, D.: Radiogenic  
850 isotopes for deciphering terrigenous input provenance in the western Mediterranean, *Chem. Geol.*, 410, 237–250,  
851 doi:10.1016/j.chemgeo.2015.06.004, 2015.

852 Rogerson, M., Rohling, E. J., Weaver, P. P. E. and Murray, J. W.: Glacial to interglacial changes in the settling  
853 depth of the Mediterranean Outflow plume, *Paleoceanography*, 20(3), doi:10.1029/2004PA001106, 2005.

854 Rogerson, M., Rohling, E. J. and Weaver, P. P. E.: Promotion of meridional overturning by Mediterranean-  
855 derived salt during the last deglaciation, *Paleoceanography*, 21(4), 1–8, doi:10.1029/2006PA001306, 2006.

856 Rogerson, M., Cacho, I., Jimenez-Espejo, F., Reguera, M. I., Sierro, F. J., Martinez-Ruiz, F., Frigola, J. and  
857 Canals, M.: A dynamic explanation for the origin of the western Mediterranean organic-rich layers,  
858 *Geochemistry, Geophys. Geosystems*, 9(7), n/a–n/a, doi:10.1029/2007GC001936, 2008.

859 Rohling, E. J.: Review and new aspects concerning the formation of eastern Mediterranean sapropels, *Mar.*  
860 *Geol.*, 122(1-2), 1–28, doi:10.1016/0025-3227(94)90202-X, 1994.

861 Rohling, E. J., Jorissen, F. J. and De stichter, H. C.: 200 Year interruption of Holocene sapropel formation in the  
862 Adriatic Sea, *J. Micropalaeontology*, 16(2), 97–108, doi:10.1144/jm.16.2.97, 1997.

863 Rohling, E. J., Mayewski, P. a, Abu-Zied, R. H., Casford, J. S. L. and Hayes, A.: Holocene atmosphere-ocean  
864 interactions: records from Greenland and the Aegean Sea, *Clim. Dyn.*, 18(7), 587–593, doi:10.1007/s00382-001-  
865 0194-8, 2002.

866 Rohling, E. J., Sprovieri, M., Cane, T., Casford, J. S. ., Cooke, S., Bouloubassi, I., Emeis, K. C., Schiebel, R.,



- 867 Rogerson, M., Hayes, A., Jorissen, F. . and Kroon, D.: Reconstructing past planktic foraminiferal habitats using  
868 stable isotope data: a case history for Mediterranean sapropel S5, *Mar. Micropaleontol.*, 50(1-2), 89–123,  
869 doi:10.1016/S0377-8398(03)00068-9, 2004.
- 870 Rohling, E. J., Marino, G. and Grant, K. M.: Mediterranean climate and oceanography, and the periodic  
871 development of anoxic events (sapropels), *Earth-Science Rev.*, 143, 62–97, doi:10.1016/j.earscirev.2015.01.008,  
872 2015.
- 873 Rossignol-Strick, M., Nesteroff, W., Olive, P. and Vergnaud-Grazzini, C.: After the deluge: Mediterranean  
874 stagnation and sapropel formation, *Nature*, 295(5845), 105–110, doi:10.1038/295105a0, 1982.
- 875 Sammari, C., Millot, C., Taupier-Letage, I., Stefani, A. and Brahim, M.: Hydrological characteristics in the  
876 Tunisia–Sardinia–Sicily area during spring 1995, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 46(10), 1671–1703,  
877 doi:10.1016/S0967-0637(99)00026-6, 1999.
- 878 Sarmiento, J. L., Herbert, T. and Toggweiler, J. R.: Mediterranean nutrient balance and episodes of anoxia,  
879 *Global Biogeochem. Cycles*, 2(4), 427–444, doi:10.1029/GB002i004p00427, 1988.
- 880
- 881 Sánchez-Goñi, M., Cacho, I., Turon, J. L., Guiot, J., Sierro, F. J., Peypouquet, J., Grimalt, J. O. and Shackleton,  
882 N. J.: Synchronicity between marine and terrestrial responses to millennial scale climatic variability during the  
883 last glacial period in the Mediterranean region, *Clim. Dyn.*, 19(1), 95–105, doi:10.1007/s00382-001-0212-x,  
884 2002.
- 885 Sarnthein, M., Tetzlaff, G., Koopmann, B., Wolter, K. and Pflaumann, U.: Glacial and interglacial wind regimes  
886 over the eastern subtropical Atlantic and North-West Africa, *Nature*, 293, 193–196, doi:10.1038/293193a0,  
887 1981.
- 888 Scheuven, D., Schütz, L., Kandler, K., Ebert, M. and Weinbruch, S.: Bulk composition of northern African dust  
889 and its source sediments — A compilation, *Earth-Science Rev.*, 116, 170–194,  
890 doi:10.1016/j.earscirev.2012.08.005, 2013.
- 891 Schmiedl, G., Kuhnt, T., Ehrmann, W., Emeis, K. C., Hamann, Y., Kotthoff, U., Dulski, P. and Pross, J.:  
892 Climatic forcing of eastern Mediterranean deep-water formation and benthic ecosystems during the past 22 000  
893 years, *Quat. Sci. Rev.*, 29(23-24), 3006–3020, doi:10.1016/j.quascirev.2010.07.002, 2010.
- 894 Schönfeld, J. and Zahn, R.: Late Glacial to Holocene history of the Mediterranean outflow. Evidence from  
895 benthic foraminiferal assemblages and stable isotopes at the Portuguese margin, *Palaeogeogr. Palaeoclimatol.*  
896 *Palaeoecol.*, 159(1-2), 85–111, doi:10.1016/S0031-0182(00)00035-3, 2000.
- 897 Schott, F., Visbeck, M., Send, U., Fischer, J., Stramma, L. and Desaubies, Y.: Observations of Deep Convection  
898 in the Gulf of Lions, Northern Mediterranean, during the Winter of 1991/92, *J. Phys. Oceanogr.*, 26(4), 505–524,  
899 doi:10.1175/1520-0485(1996)026<0505:OODCIT>2.0.CO;2, 1996.
- 900 Schroeder, K., Millot, C., Bengara, L., Ben Ismail, S., Bensi, M., Borghini, M., Budillon, G., Cardin, V.,  
901 Coppola, L., Curtil, C., Drago, A., El Moumni, B., Font, J., Fuda, J. L., García-Lafuente, J., Gasparini, G. P.,

- 902 Kontoyiannis, H., Lefevre, D., Puig, P., Raimbault, P., Rougier, G., Salat, J., Sammari, C., Sánchez Garrido, J.  
903 C., Sanchez-Roman, A., Sparnocchia, S., Tamburini, C., Taupier-Letage, I., Theocharis, A., Vargas-Yáñez, M.  
904 and Vetrano, A.: Long-term monitoring programme of the hydrological variability in the Mediterranean Sea: a  
905 first overview of the HYDROCHANGES network, *Ocean Sci.*, 9(2), 301–324, doi:10.5194/os-9-301-2013,  
906 2013.
- 907 Scrivner, A. E., Vance, D. and Rohling, E. J.: New neodymium isotope data quantify Nile involvement in  
908 Mediterranean anoxic episodes, *Geology*, 32(7), 565, doi:10.1130/G20419.1, 2004.
- 909 Shanahan, T. M., McKay, N. P., Hughen, K. A., Overpeck, J. T., Otto-Bliesner, B., Heil, C. W., King, J., Scholz,  
910 C. A. and Peck, J.: The time-transgressive termination of the African Humid Period, *Nat. Geosci.*, 8(2), 140–144,  
911 doi:10.1038/ngeo2329, 2015.
- 912 Siani, G., Paterne, M., Arnold, M., Bard, E., Metivier, B., Tisnerat, N. and Bassinot, F.: Radiocarbon reservoir  
913 ages in the Mediterranean Sea and Black Sea, *Radiocarbon*, 42(2), 271–280 [online] Available from: <Go to  
914 ISI>://000089971000010, 2000.
- 915 Siani, G., Paterne, M., Michel, E., Sulpizio, R., Sbrana, A., Arnold, M. and Haddad, G.: Mediterranean Sea  
916 surface radiocarbon reservoir age changes since the last glacial maximum., *Science* (80-. ), 294(5548), 1917–  
917 1920, doi:10.1126/science.1063649, 2001.
- 918 Siani, G., Sulpizio, R., Paterne, M. and Sbrana, A.: Tephrostratigraphy study for the last 18,000 C years in a  
919 deep-sea sediment sequence for the South Adriatic, *Quat. Sci. Rev.*, 23(23-24), 2485–2500,  
920 doi:10.1016/j.quascirev.2004.06.004, 2004.
- 921 Siani, G., Magny, M., Paterne, M., Debret, M., Fontugne, M. (2013) - Paleohydrology reconstruction and  
922 Holocene climate variability in the South Adriatic Sea. *Climate of the Past*, 9, 499-515.
- 923 Sierro, F. J., Hodell, D. A., Curtis, J. H., Flores, J. A., Reguera, I., Colmenero-Hidalgo, E., Bárcena, M. A.,  
924 Grimalt, J. O., Cacho, I., Frigola, J. and Canals, M.: Impact of iceberg melting on Mediterranean thermohaline  
925 circulation during Heinrich events, *Paleoceanography*, 20(2), n/a–n/a, doi:10.1029/2004PA001051, 2005.
- 926 Sparnocchia, S., Gasparini, G. P., Astraldi, M., Borghini, M. and Pistek, P.: Dynamics and mixing of the Eastern  
927 Mediterranean outflow in the Tyrrhenian basin, *J. Mar. Syst.*, 20(1-4), 301–317, doi:10.1016/S0924-  
928 7963(98)00088-8, 1999.
- 929 Spivack, A. J. and Wasserburg, G. J.: Neodymium isotopic composition of the Mediterranean outflow and the  
930 eastern North Atlantic, *Geochim. Cosmochim. Acta*, 52(12), 2767–2773, doi:10.1016/0016-7037(88)90144-5,  
931 1988.
- 932 Stratford, K., Williams, R. G. and Myers, P. G.: Impact of the circulation on Sapropel Formation in the eastern  
933 Mediterranean, *Global Biogeochem. Cycles*, 14(2), 683–695, doi:10.1029/1999GB001157, 2000.
- 934 Stuiver, M., Reimer, P. J. and Reimer, R.: CALIB 7.0, *Radiocarb. Calibration Progr.*, 2005.
- 935 Tachikawa, K., Roy-Barman, M., Michard, A., Thouron, D., Yeghicheyan, D. and Jeandel, C.: Neodymium

- 936 isotopes in the Mediterranean Sea: comparison between seawater and sediment signals, *Geochim. Cosmochim.*  
937 *Acta*, 68(14), 3095–3106, doi:10.1016/j.gca.2004.01.024, 2004.
- 938 Tachikawa, K., Piotrowski, A. M. and Bayon, G.: Neodymium associated with foraminiferal carbonate as a  
939 recorder of seawater isotopic signatures, *Quat. Sci. Rev.*, 88, 1–13, doi:10.1016/j.quascirev.2013.12.027, 2014.
- 940 Tanaka, T., Togashi, S., Kamioka, H., Amakawa, H., Kagami, H., Hamamoto, T., Yuhara, M., Orihashi, Y.,  
941 Yoneda, S., Shimizu, H., Kunimaru, T., Takahashi, K., Yanagi, T., Nakano, T., Fujimaki, H., Shinjo, R.,  
942 Asahara, Y., Tanimizu, M. and Dragusanu, C.: JNdi-1: a neodymium isotopic reference in consistency with  
943 LaJolla neodymium, *Chem. Geol.*, 168(3-4), 279–281, doi:10.1016/S0009-2541(00)00198-4, 2000.
- 944 Tachikawa, K., Vidal, L., Cornuault, M., Garcia, M., Pothin, A., Sonzogni, C., Bard, E., Menot, G. and Revel,  
945 M.: Eastern Mediterranean Sea circulation inferred from the conditions of S1 sapropel deposition, *Clim. Past*,  
946 11(6), 855–867, doi:10.5194/cp-11-855-2015, 2015.
- 947 Taviani, M., Angeletti, L., Canese, S., Cannas, R., Cardone, F., Cau, A., Cau, A. B., Follesa, M. C., Marchese,  
948 F., Montagna, P. and Tessarolo, C.: The “Sardinian cold-water coral province” in the context of the  
949 Mediterranean coral ecosystems, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, doi:10.1016/j.dsr2.2015.12.008,  
950 2015.
- 951 Thunell, R. C. and Williams, D. F.: Glacial–Holocene salinity changes in the Mediterranean Sea: hydrographic  
952 and depositional effects, *Nature*, 338(6215), 493–496, doi:10.1038/338493a0, 1989.
- 953 Toucanne, S., Jouet, G., Ducassou, E., Bassetti, M. A., Dennielou, B., Angue Minto’o, C. M., Lahmi, M.,  
954 Touyet, N., Charlier, K., Lericolais, G. and Mulder, T.: A 130,000-year record of Levantine Intermediate Water  
955 flow variability in the Corsica Trough, western Mediterranean Sea, *Quat. Sci. Rev.*, 33, 55–73,  
956 doi:10.1016/j.quascirev.2011.11.020, 2012.
- 957 Vance, D., Scrivner, A. E. and Beney, P.: The use of foraminifera as a record of the past neodymium isotope  
958 composition of seawater, *Paleoceanography*, 19(2), PA2009, doi:10.1029/2003PA000957, 2004.
- 959 van de Flierdt, T., Robinson, L. F. and Adkins, J. F.: Deep-sea coral aragonite as a recorder for the neodymium  
960 isotopic composition of seawater, *Geochim. Cosmochim. Acta*, 74(21), 6014–6032,  
961 doi:10.1016/j.gca.2010.08.001, 2010.
- 962 Voelker, A. H. L., Lebreiro, S. M., Schönfeld, J., Cacho, I., Erlenkeuser, H. and Abrantes, F.: Mediterranean  
963 outflow strengthening during northern hemisphere coolings: A salt source for the glacial Atlantic?, *Earth Planet.*  
964 *Sci. Lett.*, 245, 39–55, doi:10.1016/j.epsl.2006.03.014, 2006.
- 965 Weaver, A. J., Saenko, O. a., Clark, P. U. and Mitrovica, J. X.: Meltwater Pulse 1A from Antarctica as a Trigger  
966 of the Bolling-Allerod Warm Interval, *Science* (80-. ), 299(5613), 1709–1713, doi:10.1126/science.1081002,  
967 2003.
- 968 Weldeab, S., Emeis, K.-C., Hemleben, C. and Siebel, W.: Provenance of lithogenic surface sediments and  
969 pathways of riverine suspended matter in the Eastern Mediterranean Sea: evidence from  $^{143}\text{Nd}/^{144}\text{Nd}$  and

970  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios, *Chem. Geol.*, 186(1-2), 139–149, doi:10.1016/S0009-2541(01)00415-6, 2002.

971 Weldeab, S., Menke, V. and Schmiedl, G.: The pace of East African monsoon evolution during the Holocene,  
972 *Geophys. Res. Lett.*, 41, 1724–1731, doi:10.1002/2014GL059361. Received, 2014.

973 Wienberg, C., Frank, N., Mertens, K. N., Stuut, J.-B. W., Marchant, M., Fietzke, J., Mienis, F. and Hebbeln, D.:  
974 Glacial cold-water coral growth in the Gulf of Cádiz: Implications of increased palaeo-productivity, *Earth*  
975 *Planet. Sci. Lett.*, 298(3-4), 405–416, doi:10.1016/j.epsl.2010.08.017, 2010.

976 Wu, Q., Colin, C., Liu, Z., Thil, F., Dubois-Dauphin, Q., Frank, N., Tachikawa, K., Bordier, L. and Douville, E.:  
977 Neodymium isotopic composition in foraminifera and authigenic phases of the South China Sea sediments:  
978 Implications for the hydrology of the North Pacific Ocean over the past 25 kyr, *Geochemistry, Geophys.*  
979 *Geosystems*, 16(11), 3883–3904, doi:10.1002/2015GC005871, 2015.

## 980 **Table captions**

981

982 **Table 1.** U-series ages and  $\epsilon\text{Nd}$  values obtained for cold-water coral samples collected from sediment core RECORD 23  
983 (Sardinia Channel).

984

985 **Table 2.**  $\epsilon\text{Nd}$  values obtained for cold-water corals from the southern Alboran Sea. The AMS  $^{14}\text{C}$  ages published by Fink et  
986 al. (2013) are also reported as Median probability age (ka BP).

987

988 **Table 3.** AMS  $^{14}\text{C}$  ages of samples of the planktonic foraminifer *G. bulloides* from ‘off-mound’ sediment core SU92-33. The  
989 AMS  $^{14}\text{C}$  ages were corrected for  $^{13}\text{C}$  and a mean reservoir age of 400 yrs, and were converted into calendar years using the  
990 INTCAL13 calibration data set (Reimer et al., 2013) and the CALIB 7.0 program (Struiver et al., 2005).

991

992 **Table 4.** Multiproxy data obtained for the upper 2.1 m of sediment core SU92-33 (Balearic Sea). Stable oxygen and carbon  
993 isotopes were measured on benthic (*C. pachyderma*) and planktonic (*G. bulloides*) foraminifera;  $\epsilon\text{Nd}$  values were obtained on  
994 mixed planktonic foraminifera samples. The age results from a combination of 7 AMS- $^{14}\text{C}$  age measurements for the upper  
995 1.2 m of the core and by a linear interpolation between these ages as well as the  $\delta^{18}\text{O}$  variations of the planktonic  
996 foraminifera *G. bulloides*.

997

## 998 **Figure captions**

999

1000 **Figure 1.** Map of the western Mediterranean Sea showing the locations of samples investigated in this study. Yellow dot  
1001 indicates the sampling location of the sediment core from the Balearic Sea (SU92-33); yellow stars indicate the locations of  
1002 the CWC-bearing cores from the Sardinia Channel (RECORD 23) and the southern Alboran Sea (for further details on the  
1003 CWC from the Alboran Sea refer also to Fink et al., 2013). The cores discussed in this paper (Gulf of Cádiz: IODP site  
1004 U1387, Balearic Sea: MD09-2343, northern Tyrrhenian Sea: MD01-2472, Adriatic Sea: MD90-917) are indicated by black  
1005 dots, and seawater stations are marked by open squares. Arrows represent the main oceanographic currents. The black line  
1006 shows the general trajectory of the Modified Atlantic Water (MAW) flowing at the surface from the Atlantic Ocean toward  
1007 the western and eastern Mediterranean. The orange line represents the Levantine Intermediate Water (LIW) originating from  
1008 the eastern basin. The black dashed line shows the trajectory of the Western Mediterranean Deep Water (WMDW) flowing  
1009 from the Gulf of Lions toward the Strait of Gibraltar.

1010

1011 **Figure 2.** (a) Sea Surface Temperature (SST) records of cores SU92-33 (red line) and MD90-917 (green line; Siani et al.,  
1012 2004), (b)  $\delta^{18}\text{O}$  record obtained on planktonic foraminifer *G. bulloides* for core SU92-33, (c)  $\delta^{18}\text{O}$  record obtained on benthic  
1013 foraminifer *C. pachyderma* for core SU92-33, (d)  $\delta^{13}\text{C}$  record obtained on benthic foraminifer *C. pachyderma* for core SU92-  
1014 33. LGM: Last Glacial Maximum; HS1: Heinrich Stadial 1; B-A: Bølling-Allerød; YD: Younger Dryas. Black triangles  
1015 indicate AMS  $^{14}\text{C}$  age control points.

1016  
1017 **Figure 3.** (a) Sea Surface Temperature (SST) record of core SU92-33 (red line), (b)  $\epsilon\text{Nd}$  records obtained on mixed  
1018 planktonic foraminifers from core SU92-33 (open circles) and from cold-water coral fragments collected in the Alboran Sea  
1019 (red squares), (c)  $\epsilon\text{Nd}$  values of cold-water corals from core RECORD 23 (Sardinia Channel).

1020  
1021 **Figure 4.** (a)  $\delta^{13}\text{C}$  records obtained on benthic foraminifer *C. pachyderma* for cores SU92-33 (red line) and MD99-2343  
1022 (blue line; Siero et al., 2005). (b)  $\epsilon\text{Nd}$  records obtained on mixed planktonic foraminifers from core SU92-33 (open circles)  
1023 and from cold-water coral fragments collected in the Alboran Sea (red squares). Modern  $\epsilon\text{Nd}$  values for LIW (orange dashed  
1024 line) and WMDW (blue dashed line) are also reported for comparison. (c)  $\epsilon\text{Nd}$  values obtained for planktonic foraminifera  
1025 with Fe-Mn coatings at sites 300G (36°21.532' N, 1°47.507' W; 1860 m; open dots) and 304G (36°19.873' N, 1°31.631' W;  
1026 2382 m; black dots) in Alboran Sea (Jimenez-Espejo et al., 2015). (d) UP10 fraction (>10  $\mu\text{m}$ ) from core MD99-2343  
1027 (Frigola et al., 2008). (e) Sortable silt mean grain-size of core MD01-2472 (Toucanne et al., 2012). (f) Ln Zr/Al ratio at IODP  
1028 site U1387 (36°48.3' N 7°43.1' W; 559 m) (Bahr et al., 2015).

1029  
1030 **Figure 5.** (a)  $\delta^{18}\text{O}$  record obtained on planktonic foraminifer *G. bulloides* for core SU92-33, (b)  $\delta^{13}\text{C}$  records obtained on  
1031 benthic foraminifer *C. pachyderma* for core SU92-33, (c)  $\epsilon\text{Nd}$  values of cold-water corals from core RECORD 23 (Sardinia  
1032 Channel), (d)  $\epsilon\text{Nd}$  values records obtained on mixed planktonic foraminifera from core SU92-33 (open circles) and from  
1033 cold-water coral fragments collected in the Alboran Sea (red squares), (e)  $\epsilon\text{Nd}$  values obtained on terrigenous fraction of  
1034 MS27PT located close the Nile River mouth in the eastern Mediterranean basin (Revel et al., 2015)

