



Climatic variability in Princess Elizabeth Land (East Antarctica) over the last 350 years

Alexey A. Ekaykin^{1,2}, Diana O. Vladimirova^{1,2*}, Vladimir Ya. Lipenkov¹ and Valérie Masson-Delmotte³

¹Climate and Environmental Research Laboratory, Arctic and Antarctic Research Institute, St. Petersburg, 199397, Russia

5 ²Institute of Earth Sciences, Saint Petersburg State University, St. Petersburg, 199178, Russia

³Laboratoire des Sciences du Climat et de l'Environnement - IPSL, UMR 8212, CEA-CNRS-UVSQ-Université Paris Saclay, Gif sur Yvette, F-91198, France

*now at Center for Ice and Climate, Niels Bohr Institute, University of Copenhagen, Copenhagen Ø, 2100, Denmark

Correspondence to: Alexey A. Ekaykin (ekaykin@aari.ru)

10 **Abstract.** We use isotopic composition (δD) data from 6 sites in Princess Elisabeth Land (PEL) in order to reconstruct the air temperature variability in this sector of East Antarctica for the last 350 years. First, we use the present-day instrumental mean annual surface air temperature data to demonstrate that the studied region (between Russian research stations Progress, Vostok and Mirny) is characterized by uniform temperature variability. We thus construct the stacked record of the temperature anomaly for the whole sector for the period 1958-2015. A comparison of this series with the Southern Hemisphere climatic indices shows that the short-term inter-annual temperature variability is primarily governed by Antarctic Oscillation (AAO) and Interdecadal Pacific Oscillation (IPO) modes of atmospheric variability. However, the low-frequency temperature variability (with period > 27 years) is mainly related to the anomalies of Indian Ocean Dipole (IOD) mode. Then we construct the stacked record of δD for the PEL for the period 1654-2009 from individual normalized and filtered isotopic records obtained at 6 different sites ('PEL2016' stacked record). We use a significant relationship between this record and the stacked PEL temperature record (with an apparent slope of $9 \text{ } \mu\text{C}^{-1}$) to convert 'PEL2016' into temperature scale. Analysis of 'PEL2016' shows a $1 \text{ } ^\circ\text{C}$ warming in this region over the last three centuries, with a particularly cold period from mid-18th to mid-19th century. A peak of cooling occurred in the 1840s - a feature previously observed in other Antarctic records. We reveal that 'PEL2016' correlates with a low-frequency component of IOD. We suggest that the IOD mode influences the Antarctic climate by modulating the activity of cyclones that bring heat and moisture to Antarctica. We also compare 'PEL2016' with other Antarctic stacked isotopic records. This work is a contribution to PAGES and IPICS Antarctica 2k projects.

1 Introduction

While understanding the behavior of Antarctic climate system is crucial in context of the present-day global environmental changes, key gaps arise from limited observations. The only source of climatic data prior to the International Geophysical Year



epoch (1955-1957) is ice core records. Deep ice cores have provided a wealth of climatic and environmental information covering glacial-interglacial variations of the past 800,000 years (EPICA community members, 2004). However, the spatio-temporal characteristics of Antarctic climate variability of the most recent centuries remains poorly known or understood (Jones et al., in press; PAGES 2k Consortium, 2013).

- 5 The network of ice core records spanning the last centuries is moreover unevenly distributed, reflecting heterogeneous efforts for extracting these records. A quite extensive coverage of some regions of Antarctica, such as West Antarctica (Kaspari et al., 2004) or Dronning Maud Land (Altnau et al., 2015; Oerter et al., 2000) contrasts with other regions that still remain white spots. As a result, attempts to reconstruct the climatic variability of the whole Antarctic continent (Jones et al., in press; PAGES 2k Consortium, 2013; Schneider et al., 2006; Frezzotti et al., 2013) are limited by the lack of available data.
- 10 In our previous work we summarized available isotopic data for the vicinity of Vostok Station in order to construct a robust stack climatic record over the past 350 years (Ekaykin et al., 2014). Here we present a new stacked climate record for Princess Elisabeth Land (PEL), the territory located between the Russian stations of Progress, Vostok and Mirny, East Antarctica. This record is based on water stable isotope data from 6 sites, and spans the last 350 years (Figure 1). We evidence a close relationship between the stacked isotopic record and regional surface air temperature variations, and highlight significant
- 15 relationships between regional climate and large-scale modes of variability of the Southern Hemisphere. Classically, Section 2 describes our data and methods, and Section 3 is focused on these results and their discussion, before a conclusion in Section 4.

2 Methods

2.1 Individual records

- 20 In this study we use data from 6 individual records obtained in Princess Elisabeth Land (Figure 1, Table 1).
“105 km” (67.433 °S and 93.383 °E, time interval 1757-1987) is a 727 m ice core drilled in 1988 by specialists of St. Petersburg Mining Institute, about 105 km inland from Mirny station. The isotopic content was measured late in the 1980s at Laboratoire des Sciences du Climat et de l’Environnement (LSCE) with resolution of 1 m. In 2013, the upper 109 m of the core were re-measured at Climate and Environmental Research Laboratory (CERL), with a depth resolution of 5 cm. The core was dated
- 25 by layer counting, and the initial dating was adjusted using the reference horizon of the 1816 Tambora volcanic eruption, identified from Electrical Conductivity Measurements (ECM) (Vladimirova and Ekaykin, 2014). As a result, a record of annual accumulation rate is available.
“400 km” (69.95 °S and 95.617 °E, 1254-1987) refers to an ice core drilled in 1988 at the 400th km from Mirny station, down to 150 m depth. Isotopic measurements were performed at LSCE on 1 m samples. The core was dated according to the simple



Nye depth-age model, taking into account the average accumulation rate at the drilling site (Lipenkov et al., 1998) and the density profile of the core. As a result, no record of annual accumulation rate is available.

“VRS 2013” (78.467 °S and 106.84 °E, 1654-2010) is a stack of 15 individual isotopic records from snow pits and shallow cores recovered in the vicinity of Vostok Station (Ekaykin et al., 2014). The data on temporal variability of snow accumulation rate data is also available for this site.

“NVFL-1” (77.11 °S and 95.072 °E, 1711-1944) is a 18.3 m firn core drilled from the bottom of a 2.5 m snow pit in 2008 close to the Dome B. The chronology was established using the firn density data and the 1816 Tambora volcano ECM peak as a reference horizon.

“NVFL-3” (76.405 °S and 102.167 °E, 1978-2009) is a 3.1 m snow pit dug in 2010 in the northern part of subglacial Lake Vostok. It is dated based on snow stratigraphy and identification of 1993 Pinatubo volcano peak in SO₄²⁻ vertical profile. Chemical measurements were performed at Limnological Institute of Russian Academy of Sciences, Irkutsk, Russia.

“PV-10” (72.805 °S and 79.934 °E, 1976-2009) is a 7.55 m firn core drilled in 2010 about 400 km inland from Progress Station. It was dated using firn density data and taking into account the ECM peak associated with the 1993 deposition from the Pinatubo eruption.

We estimated the dating uncertainty by comparing age calculated using only firn density data and average snow accumulation rate for a given site with age of the reference age markers and came to a conclusion that the age errors do not exceed 10 %. For the reference years (1816 and 1993, where we have absolute dating), the error tends to zero. The largest error is expected for the “400 km” series, where we do not have reference age markers. However, if we use the prominent 1840 cold event (see Section 3.3), observed in all records, as such a marker, then we may estimate a relative dating error for this series as < 6%.

We also use the accumulation data from the site “200 km” (Fig. 1), spanning the period 1640-1987, as published in (Ekaykin et al., 2000).

2.2 Stacked records

Figure 2 displays the individual δD time-series from all 6 sites. Differences between mean values reflect well-known differences in isotopic distillation along a gradient of inland elevation (e.g. Masson-Delmotte et al., 2008). In order to investigate temporal variations only, we calculated normalized values for each series using interval 1757-1944 as a reference period. The short series (NVFL-3 and PV-10) were normalized over 1978-2009 period, and then the normalized values were reduced in terms of mean and STD to the corresponding period of the long series, in order to avoid an artificial dominating of the short records in the stacked series.

We then applied a low-pass filter to cut off the variability with periodicities lower than 27 years. Note that all spectral analyses and filtering was performed with the use of Analyseries software (Paillard et al., 1996). This is motivated by the fact that one



single record in inland Antarctica cannot provide reliable climatic information on a short-term time scale, due to a very low signal-to-noise ratio (Ekaykin et al., 2014). Moreover, the latter study also highlighted multi-decadal climatic variability in this sector of central Antarctica, with a period of 30-50 years.

The normalized and filtered time series are displayed in Figure 3. Despite common features, this comparison clearly shows significant discrepancies between individual records. One reason for such mismatches may lie in age scale uncertainties. However, this hypothesis is ruled out by the comparison of individual series around 1816 and 1993 (dates of firn layers containing Tambora and Pinatubo volcanic eruption debris, denoted by vertical dashed lines in Fig. 3), when the relative dating error tends to zero: observed discrepancies do not only arise from chronological problems. Alternatively, this mismatch may arise from a significant level of noise even in the filtered series.

10 In order to isolate the climatic signal from the noise, we constructed a stacked climatic record for the PEL region, hereafter named PEL2016 (grey line in Fig. 3). For a given year, the value of this record consists of the average of the values of individual records available for this year; the standard deviation within individual records is also reported on Fig. 3.

2.3 Meteorological information available for the studied sector

A number of research stations have been established in the PEL area, as indicated in Fig. 1. Unfortunately, most of them have very short (if any) meteorological records. Relatively long records are available only for 5 stations: Australian station Davis (1957-1964 and 1969-2015), Chinese station Zhongshan (1989-2015), Russian stations Progress (1989, 1991 and 2003-2015), Mirny (1956-2015) and Vostok (1958-2015 with gaps in 1962, 1994, 1996 and 2003). The data were downloaded from <https://legacy.bas.ac.uk/met/READER/> (Turner et al., 2004).

20 The correlation between Progress, Zhongshan and Davis annual mean temperature datasets, located very close to each other, is 0.96-0.98 (note that only statistically significant correlation coefficients with a confidence level > 95 % are reported in the paper). Hereafter, we only used data from the station with the longest record (Davis).

We also use data from automatic weather station (AWS) LGB59 located at the slope of the Antarctic ice sheet inland from Progress station (Fig. 1), available for the period from 1994 to 1999, as well as surface air temperature data from Casey and Mawson.

25 2.4 Climatic indices of Southern Hemisphere

In order to investigate possible relationships between PEL climate multi-decadal variations and large-scale modes of variability, we use data on the indices of the Antarctic Oscillation (AAO), the Interdecadal Pacific Oscillation (IPO) and the Indian Ocean Dipole (IOD).



AAO index is defined as a mean latitudinal difference of sea level pressure at 40 °S and 65 °S, and is considered as a prevailing mode of Atmospheric circulation in the Southern Hemisphere (Marshall, 2003). The data on AAO index is available here: http://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily_ao_index/ao/monthly.ao.index.b79.current.ascii.table (since 1979). Also, at the site of British Antarctic Survey there is AAO data since 1957:

5 <http://www.antarctica.ac.uk/met/gjma/sam.html>,

although data for the 1957-1978 period is considered to be less robust.

IPO is defined as a sea surface temperature (SST) anomaly over the Pacific Ocean. The positive phase of IPO is characterized by relatively warm central and eastern tropical Pacific, and relatively cold north-western and south-western Pacific (Henley et al., 2015; Dong and Dai, 2015). IPO index is closely related to PDO (Pacific Decadal Oscillation), but PDO better characterizes
10 Northern Pacific, while IPO is better applicable to the whole Pacific region. We use IPO data because in the previous study a teleconnection between PDO mode and central Antarctic climate was discovered (Ekaykin et al., 2014).

The data on IPO index since 1870 is available here:

<http://www.esrl.noaa.gov/psd/data/timeseries/IPOTPI/>

IOD is characterized by Dipole Mode Index (DMI) that is defined as the SST gradient between the western equatorial Indian
15 Ocean (50 °E - 70 °E and 10 °S - 10 °N) and the south eastern equatorial Indian Ocean (90 °E - 110 °E and 10 °S - 0 °N). Thus, IOD is an analogue of SOI, but for Indian Ocean. The data on DMI index since 1870 could be found at:

http://www.jamstec.go.jp/frsgc/research/d1/iod/iod/dipole_mode_index.html.

3 Results and discussion

20 3.1 Surface air temperature variability in the Princess Elisabeth Land during the period of instrumental observations (1958-2015)

Here, we first consider the variability of surface air temperature recorded at the meteorological stations in Princess Elisabeth Land, to assess whether uniform climate variability pattern is monitored, and to provide a reference regional temperature record for comparison with the δD stacked record.

25 Correlation coefficients between annual mean surface air temperature data at Vostok, Mirny and Davis vary between 0.6 and 0.9 (Table 2). Correlation coefficients between Automatic Weather Station LGB59 (located between Davis and Vostok, Fig. 1) and these 3 stations vary between 0.86 and 0.96. Despite the short record at LGB59, they are also significant at 95% confidence level. These results demonstrate that the region encompasses between these 3 stations has experienced similar climatic variability. This is further confirmed by a cluster analysis of surface air temperature data from 12 Antarctic stations



(see Supplementary Figure S1), showing that Vostok, Mirny, Casey, Mawson and Davis data form a single cluster in terms of climatic variability.

Interestingly, the correlation coefficient between Mirny and Vostok data is significantly weaker in 1958-1976 ($R=0.53$) than in 1976-2015 ($R=0.74$). This suggests that, before the so-called “1976 climate shift” (Giese et al., 2002) Vostok experienced a higher influence from the Pacific sector of the Southern Ocean (Ekaykin et al., 2014) not encompassed at Mirny. Indeed, the correlation coefficient between temperature data from Vostok and Mc Murdo Station (located in the Pacific sector) was higher before the 1976 shift ($R=0.46$) than after 1976 ($R=0.35$).

During the whole period of instrumental observations, the strongest relationships observed for temperature at Vostok were with temperature data at Mirny and Mawson coastal stations from the Indian Ocean sector, and more precisely the sector between Davis Sea and Cooperation Sea.

As a result, Figure 4a shows the average temperature anomaly from Vostok, Mirny and Davis stations. Hereafter, we use this stacked temperature record as an estimate of the temperature anomaly for the whole PEL sector.

We now compare the low frequency variations in these various temperature records, using the 27-year low pass filter (Figure S2). Both Vostok and Mirny have a 30-year periodicity, maxima in the late 1970s and the late 2000s, and demonstrate a very high similarity at low frequency. While Davis data have the same periodicity, their maxima are shifted to the early 1970s and early 2000s. If we consider other Antarctic stations, we see a complex behavior of air temperature in different sectors of Antarctica: most stations also show a 30-year cycle, but with a significant phase shift relative to PEL region.

In the Indian Ocean sector, temperature peaks appear more and more delayed when moving from west to east. For example, the first maximum occurred late in the 1960s at Mawson, early in the 1970s at Davis, in the second half of the 1970s at Mirny, and late in the 1970s at Casey. This feature may reflect a low-frequency component of the Antarctic Circumpolar Wave (Carril and Navarra, 2001).

With respect to multi-decadal trends, contrasted patterns emerge: some stations (Esperanza, Novolazarevskaya, Davis, Vostok, Mirny, McMurdo) display warming trend, while a cooling trend emerges at Halley or Dumont d’Hurville (Fig. S2).

This comparison of instrumental temperature records highlights different patterns of multi-decadal variability across different sectors of Antarctica, which is important for interpreting paleoclimate records, and for combining various proxy records for temperature reconstructions (Jones et al., in press). Our analysis nevertheless demonstrates coherency within Princess Elisabeth Land, where we will use the stacked temperature record from Vostok, Mirny and Davis as a reference regional signal (hereafter named PEL temperature anomaly) for calibration of δD records.



3.2 Relationships between Princess Elisabeth Land instrumental temperature records and Southern Hemisphere modes of variability

Here, we compare the PEL temperature anomaly with indices that characterize climatic variability in the Southern Hemisphere. First, as expected, a very strong negative relationship with the AAO index ($r = -0.68$) is observed in 1979-2015 (Fig. 4b). The Antarctic Oscillations is the predominant mode of climatic variability in Antarctica: a strong AAO index reflects a large pressure gradient between low and high latitudes, associated with a more zonal circulation around Antarctica, and colder conditions in East Antarctica. We note that no correlation between PEL and AAO is identified prior to 1979, which could be an artifact due to poor estimate of AAO before 1979, when few instrumental records are assimilated in atmospheric reanalyses. The correlation coefficient of PEL temperature anomaly with the IPO index is weak (Fig. 4c), but the residuals of the PEL temperature regression with AAO are negatively correlated with IPO index ($r = -0.47$).

A multiple linear regression approach leads to the conclusion that combined variations in AAO and IPO explain 59% of the temperature variance, at the inter-annual scale. While such tele-connection between Pacific and central Antarctic climate had previously been reported from Vostok data (Ekaykin et al., 2014), the underlying mechanism is not known. Finally, no significant correlation was identified between PEL temperature and the IOD index (Fig. 4d).

However, different results emerge when considering the low-pass filtered time series. At multi-decadal time scales, a strong positive correlation ($r = 0.8$) relates PEL temperature and the AAO (Fig. 4a and 4b), and a very strong positive correlation appears between PEL temperature and the IOD index ($r = 0.93$). We suggest that the Indian Ocean Dipole affects the Antarctic climate through a modulation of cyclonic activity. This is indirectly confirmed by a negative correlation ($r = -0.56$) between the IOD index and the pressure anomaly at Mirny and Davis (not shown). The positive relationship between AAO and temperature in the low frequency band could then be an “induced correlation” caused by a very strong positive correlation between AAO and IOD ($r = 0.8-0.9$) at these time-scales.

3.3 Climatic variability in Princess Elisabeth Land over the last 350 years

The stacked δD record (built from low-pass filtered individual records) is now compared with the filtered PEL temperature composite (Fig. 4a). A significantly positive correlation is observed ($r = 0.66$). We note a discrepancy in the timing of the most recent maximum (2000 according to the isotopic series and 2008 according to the temperature record), while the late 1970s maximum is in phase in the both records. This discrepancy could be explained by biases in the stack isotopic series caused by “edge effect” at the end margin of the record due to varying number of individual records used in the stacked one; or by phase shifts in climatic variability in different sectors of Antarctica (as reported for Davis and Mirny, Fig. S2).

Assuming that the stacked δD record is a proxy of surface air temperature in the PEL region, we estimate the corresponding calibration coefficient by the ratio of the standard deviation of the δD composite record to the standard deviation of the PEL



low-pass filtered temperature record, which allows us to assign a temperature scale to the isotopic record. The apparent isotope-temperature gradient, obtained as a standard deviation of isotopic values divided by standard deviation of temperature values is $13,8 \pm 2,5 \text{ ‰ } ^\circ\text{C}^{-1}$ (the uncertainty is due to different standard deviation of isotopic values in individual records). Such an approach implicitly suggests a perfect correlation between the compared series. If we correct the apparent slope by the observed correlation coefficient, 0.66, it becomes $9 \text{ ‰ } ^\circ\text{C}^{-1}$. The latter value is still considerably higher than the corresponding slopes observed in other regions of Antarctica (see a review in Stenni et al., 2016), but corresponds nicely to an isotope-condensation temperature slope predicted by simple isotope model (Salamatin et al., 2004). Actually, low apparent isotope-temperature slopes obtained based on ice-core data may be due to significant amount of noise in the isotopic records, while in our case we considerably removed noise by filtering and constructing the stacked record.

5 The temperature reconstruction is displayed in Figure 5b as a temperature anomaly relative to the 1980-2009 period. We also show the instrumentally obtained air temperature anomaly in Fig. 5b on the same temperature scale.

Following (Ekaykin et al., 2014), who reported a closer relationship between Vostok isotopic data and summer temperature than with annual mean temperature, we performed additional analyses of relationships between our stacked isotope record and other temperature time series (e.g. monthly or seasonal temperature anomalies), but this did not improve the isotope-temperature correlation.

15 Despite discrepancies in the individual isotopic records (Fig. 3), common signal identified in the stacked record lead to several conclusions about PEL climate variability over the past 350 years. During this time interval, regional surface air temperature shows a long-term increasing trend, and an overall warming by about $1 \pm 0.2 \text{ } ^\circ\text{C}$. Superimposed on this multi-centennial trend, quasi-periodical variability occurs with periods of 30-40 and about 60 years. A colder period is identified in 1750-1860, corresponding to the end of the “Little Ice Age” (PAGES 2k Consortium, 2013).

20 A remarkable cold phase is observed during the 1840s, during which PEL temperature could fall $1.2 \text{ } ^\circ\text{C}$ below present-day (defined as the average value of the last 30 years). As seen in Fig. 3, this event is a robust feature, observed in all 4 individual records available for this time interval. This minimum was also identified in an Antarctic temperature stack record (Schneider et al., 2006) – see Fig. 5d, as well as an ice core drilled in the Ross Sea sector (Rhodes et al., 2012).

25 Further studies are needed to understand whether such remarkable cold conditions arise from internal variability or are driven by the response of regional climate to an external perturbation. A possible candidate could be a response to volcanic forcing (Sigl et al., 2015). A moderate event is associated with the eruption of Cosigüina in 1835. According to the inventory of volcanic events recorded in the Vostok firn cores (Osipov et al., 2014), there was an eruption of an unknown volcano in 1840; however, the amount of deposited sulfate was about 15% of that of Tambora, so it is not expected to have a major effect on climate system. So far, the influence of volcanic forcing on Antarctic climate, and the response time remains poorly known.

30 By contrast, recent studies have stressed the delayed response of the North Atlantic Oscillation (Ortega et al., 2015) to major



volcanic eruptions, as well as their role as pace-makers of bidecadal variability in the North Atlantic (Swingedouw et al., 2015).

The period before 1700 is probably the coldest part of the record, but this is not a robust result as the 2 records spanning this time interval show somewhat different behaviors (Fig. 3). However, another stack of 5 East Antarctic cores from PAGES2k (Fig. 5e) also highlights that the 1690s could have been the coldest decade of the last 350 years.

We also compare the PEL2016 record with other Antarctic temperature reconstructions. Schneider et al. (2006) used high-resolution isotopic records from 5 Antarctic sites (a stack of Law Dome records, Siple Station, a stack of Dronning Maud Land records, and two ITASE sites from West Antarctica). His record is statistically significantly correlated with PEL2016 ($r = 0.36$). This suggests that multi-decadal temperature variability of the relatively small PEL region has some common features with the whole Antarctic continent (warming in the 1820s and 1890s, cold events in the 1840s and 1900s, etc.).

We also investigated the similarities between PEL2016 and the filtered stack normalized isotopic East Antarctic record based on 5 East Antarctic ice cores (Fig. 5e; data are available in Supplementary materials of PAGES 2k Consortium, 2013). The correlation with PEL2016 is weak ($r = 0.13$) but significant. The correlation with the stack from Schneider et al (2006) is again 0.36.

The main difference between our PEL2016 record and the other isotopic stacked records for the whole Antarctica (Fig. 5d) and for East Antarctica (Fig. 5e) appears for long-term trends, with a long-term increase in PEL2016 but no similar feature in the other reconstructions. We suggest that contrasted regional long-term trends may disappear in continental-scale reconstructions (see Fig. S2).

Finally, we compare our PEL2016 record with an IOD time-series since 1870, also processed with a low-pass filter. The strong correlation coefficient ($r = 0.79$) confirms the tight relationship between multi-decadal variations in surface air temperature in this sector of Antarctica and IOD. The Indian Dipole Ocean oscillation appears as the predominant climatic mode affecting multi-decadal climate variability in this part of East Antarctica. While the exact mechanisms underlying this relationship are not known, the IOD is expected to affect the inland Antarctic climate by modulating the cyclonic activity that brings heat and moisture to Antarctic continent.

3.4 Snow accumulation rate variability

We now investigate the low-pass filtered values of snow accumulation rate, available at sites “105 km”, “200 km” and Vostok (the latter is a stack curve from 3 deep snow pits), normalized over the period from 1952 to 1981 (Figure 6). All of them exhibit a negative trend, more prominent for “200 km” series. This result contradicts the stacked Antarctic snow accumulation rate record (Frezzotti et al., 2013) showing an overall increase of the accumulation rate during the last 200 years. Our finding is also not supported by the accurate assessment of average accumulation rate change between successive reference horizons at



Vostok, showing a slight but significant increase of snow accumulation rate since 1816 (Ekaykin et al., 2004). Our results moreover stress the fact that, during the last centuries, opposite long-term trends may have occurred in temperature and accumulation. This is counter-intuitive with respect to atmospheric thermodynamics and to the expected co-variation of heat and moisture advection towards inland Antarctica.

- 5 Processes other than snowfall deposition may however affect the ice core records. In the vicinity of “105 km”, large “transversal” snow dunes have recently been evidenced (Vladimirova and Ekaykin, 2014). Such features may lead to a strong non-climatic variability in the snow accumulation rate in a given point, due to dune propagation effects. Blowing snow events may also have a significant influence on mass balance in the coastal zone of Antarctica (Scarchilli et al., 2010), potentially introducing additional post-deposition noise.
- 10 As a result, we are not confident that the datasets reported in Fig. 6 can be interpreted in terms of climate (snowfall) variations, and further work is needed to decipher the large-scale climate effect (snowfall deposition) from the non-climatic effects potentially associated with post-deposition (wind erosion, dune propagation etc.).

4 Conclusion

In this paper, we presented an analysis of the recent variability in snow isotopic composition (δD) data from 6 snow pits and
15 ice cores recovered in the region of Princess Elisabeth Land (PEL), East Antarctica.

To interpret this data, we have investigated the present-day mean annual surface air temperature variability using the instrumental temperature measurements at stations Mirny, Davis and Vostok located at the margins of the studied sector. It was shown that inter-annual climatic variability strongly covariates at these three stations. Cluster analysis has evidenced coherent variations for these stations, together with the nearby stations of Casey and Mawson. However, we have stressed
20 phase shifts between multi-decadal temperature variations along the coastal stations: temperature maxima and minima at Vostok and Mirny are delayed by a few years compared to those at Davis. At a broader geographical scale, temperature records from different sectors of Antarctica exhibit different climatic variability at decadal scale in terms of periodicities, phasing and trends.

We then compared recent temperature variability in the PEL region with indices of Southern Hemisphere modes of variability, and highlight the importance of the Annular Antarctic Oscillation and the Interdecadal Pacific Oscillation that in total explain
25 59% of the temperature variance in this Antarctic region. At the multi-decadal time-scale, however, temperature variations appear most closely related with the Indian Ocean Dipole mode, understood to modulate the cyclonic activity bringing heat and moisture to Princess Elisabeth Land.



Given limitations of ice core data for inter-annual variations, we have processed our isotopic time-series with a low-pass filter to cut off variability expressed at timescales <27 years. Both common features and significant discrepancies emerge from individual filtered time-series. These differences may arise from true differences in regional climate variations, and/or by non-climatic noise.

5 In order to improve the signal-to-noise ratio, we constructed a stacked isotopic record for the Princess Elisabeth Land based on data from all 6 sites. We then used the significant correlation between this record and instrumentally obtained air temperature record in order to convert the isotopic composition scale into air temperature scale. The apparent isotope-temperature slope is $9 \text{ ‰ } ^\circ\text{C}^{-1}$.

The newly obtained temperature reconstruction covers the period from 1654 to 2009. During this period, temperature appears
10 to have gradually increased by about $1 \text{ }^\circ\text{C}$, from a relatively cold period evidenced from the mid-17th to mid-19th centuries ("Little Ice Age"). The coldest decade is identified in the 1840s, a feature common to several Antarctic isotopic composite signals. By contrast, long-term temperature trends were not identified previously in pan-Antarctic stacked records, possibly due to averaging effects of different regional trends. We found a weak, though significant, positive correlation of our temperature reconstruction with reconstructions previously obtained for the whole Antarctic continent and/or East Antarctica.

15 Finally, our PEL record appears closely related to the low-frequency component of the Indian Ocean Dipole mode.

The three accumulation time series depict decreasing long-term trends and large inter-site differences. Further investigations of non-climatic drivers (including wind erosion and dune effects) are needed prior to confident climatic interpretation.

Our time-series is provided as supplementary information to this manuscript. Understanding the cause for the reconstructed changes will require to compare the PEL record with other regional Antarctic records, expanding the work of Jones et al (in
20 press), and combining simulations and reconstructions in order to better understand the mechanisms of regional climate multi-decadal to centennial variations, and to explore the potential response of Antarctic climate to external forcing factors (e.g. volcanic eruptions).

This study finally stresses the importance of obtaining a dense network of highly resolved ice core records in order to document the complexity of spatio-temporal variations in Antarctic climate, a key focus of the Antarctica 2K project (<http://www.pages-igbp.org/ini/wg/antarctica2k/intro>).
25

Acknowledgement

This work is a contribution to PAGES and IPICS "Antarctica 2k" project. We are grateful to all the participants of Russian Antarctic Expedition (RAE) and drillers from St. Petersburg Mining University for providing us with the high-quality ice cores. We thank RAE for logistical support of our works in Antarctica. The Russian-French collaboration in the field of ice



cores and paleoclimate studies is carried out in the frames of LIA "Vostok". We thank the CERL's staff for the isotopic analyses. The chemical analyses of the samples were performed at Irkutsk's Limnological Institute of RAS in frames of Russian Foundation for Basic Research grant 15-55-16001. One of the authors (VMD) was supported by Agence Nationale de la Recherche in France, grant ANR-14-CE01-0001.

- 5 This study was completed with a financial support from Russian Science Foundation, grant 14-27-00030.

References

- Altnau, S., Schlosser, E., Isaksson, E., and Divine, D. V.: Climatic signals from 76 shallow firn cores in Dronning Maud Land, East Antarctica, *The Cryosphere*, 9, 925-944, 2015.
- Carril, A. F., and Navarra, A.: Low-frequency variability of the Antarctic Circumpolar Wave, *Geophys. Res. Lett.*, 28, 4623-4626, 2001.
- 10 Dong, B., and Dai, A.: The influence of the Interdecadal Pacific Oscillation on temperature and precipitation over the globe, *Clim. Dyn.*, 15, doi:10.1007/s00382-015-2500-x, 2015.
- Ekaykin, A. A., Lipenkov, V. Y., Barkov, N. I., Petit, J. R., and Stievenard, M.: The snow accumulation variability over the last 350 years at the slope of Antarctic ice sheet at 200 km from the Mirny observatory (in Russian), *Kriosfera Zemli*, 4, 57-66, 2000.
- 15 Ekaykin, A. A., Lipenkov, V. Y., Kuzmina, I. N., Petit, J. R., Masson-Delmotte, V., and Johnsen, S.: The changes in isotope composition and accumulation of snow at Vostok station over the past 200 years, *Ann. Glaciol.*, 39, 569-575, 2004.
- Ekaykin, A. A., Kozachek, A. V., Lipenkov, V. Y., and Shibaev, Y. A.: Multiple climate shifts in the Southern Hemisphere over the past three centuries based on central Antarctic snow pits and core studies, *Ann. Glaciol.*, 55, 259-266, 2014.
- 20 EPICA community members: Eight glacial cycles from an Antarctic ice core, *Nature*, 429, 623-628, 2004.
- Frezzotti, M., Scarchilli, C., Becagli, S., Proposito, M., and Urbini, S.: A synthesis of the Antarctic surface mass balance during the last 800 yr, *The Cryosphere*, 7, 303-319, doi:10.5194/tc-7-303-2013, 2013.
- Giese, B. S., Urizar, S. C., and Fuckar, N. S.: Southern hemisphere origins of the 1976 climate shift, *Geophys. Res. Lett.*, 29, 11-4, 2002.
- 25 Henley, B. J., Gergis, J., Karoly, D. J., Power, S., Kennedy, J., and Folland, C. K.: A tripole index for the Interdecadal Pacific Oscillation, *Clim. Dyn.*, 15, doi:10.1007/s00382-015-2525-1, 2015.
- Jones, J. M., Gille, S. T., Goosse, H., Abram, N. J., Canziani, P. O., Charman, D. J., Clem, K. R., Crosta, X., de Lavergne, C., Eisenman, I., England, M. H., Fogt, R. L., Frankcombe, L. M., Marshall, G. J., Masson-Delmotte, V., Morrison, A. K., Orsi,



- A. J., Raphael, M. N., Renwick, J. A., Schneider, D. P., Simpkins, G. R., Steig, E. J., Stenni, B., Swingedouw, D., and Vance, T. R.: Assessing recent trends in high-latitude Southern Hemisphere surface climate, *Nature Climate Change*, in press.
- Kaspari, S., Mayewski, P. A., Dixon, D. A., Spikes, V. B., Sneed, S. B., Handley, M. J., and Hamilton, G. S.: Climate variability in West Antarctica derived from annual accumulation-rate records from ITASE firn/ice cores, *Ann. Glaciol.*, 39, 585-594, 5 2004.
- Lipenkov, V. Y., Ekaykin, A. A., Barkov, N. I., and Pourchet, M.: On the relation of surface snow density in Antarctica to wind speed (in Russian), *Materialy Glyatsiologicheskikh Issledovaniy*, 85, 148-158, 1998.
- Marshall, G. J.: Trends in the Southern Annular Mode from observations and reanalysis, *J. Clim.*, 16, 4134-4143, 2003.
- Masson-Delmotte, V., Hou, S., Ekaykin, A. A., Jouzel, J., Aristarain, A., Bernardo, R. T., Bromwich, D., Cattani, O., Delmotte, 10 M., Falourd, S., Frezzotti, M., Gallee, H., Genoni, L., Isaksson, E., Landais, A., Helsen, M., Hoffmann, G., Lopez, J., Morgan, V., Motoyama, H., Noone, D., Oerter, H., Petit, J. R., Royer, A., Uemura, R., Schmidt, G. A., Schlosser, E., Simoes, J. C., Steig, E., Stenni, B., Stievenard, M., van den Broeke, M., van de Wal, R., van den Berg, W.-J., Vimeux, F., and White, J. W. C.: A review of Antarctic surface snow isotopic composition: Observations, atmospheric circulation and isotopic modelling, *J. Clim.*, 21, 3359-3387, 2008.
- 15 Oerter, H., Wilhelms, F., Jung-Rothenhausler, F., Goktas, F., Miller, H., Graf, W., and Sommer, S.: Accumulation rates in Dronning Maud Land, Antarctica, as revealed by dielectric-profiling measurements of shallow firn cores, *Ann. Glaciol.*, 30, 27-34, 2000.
- Ortega, P., Lehner, F., Swingedouw, D., Masson-Delmotte, V., Raible, C. C., Casado, M., and Yiou, P.: A model-tested North Atlantic Oscillation reconstruction for the past millennium, *Nature*, 523, 71-77, doi:10.1038/nature14518, 2015.
- 20 Osipov, E. Y., Khodzher, T. V., Golobokova, L. P., Onischuk, N. A., Lipenkov, V. Y., Ekaykin, A. A., Shibaev, Y. A., and Osipova, O. P.: High-resolution 900 year volcanic and climatic record from the Vostok area, East Antarctica, *The Cryosphere*, 8, 843-851, 2014.
- PAGES 2k Consortium: Continental-scale temperature variability during the past two millennia, *Nature Geoscience*, 6, 339-346, doi:10.1038/ngeo1797, 2013.
- 25 Paillard, D., Labeyrie, L., and Yiou, P.: Macintosh program performs time-series analysis, *Eos Trans. AGU*, 77, 379, 1996.
- Rhodes, R. H., Bertler, N. A. N., Baker, J. A., Steen-Larsen, H. C., Sneed, S. B., Morgenstern, U., and Johnsen, S. J.: Little Ice Age climate and oceanic conditions of the Ross Sea, Antarctica from a coastal ice core record, *Clim. Past*, 8, 1223-1238, doi:10.5194/cp-8-1223-2012, 2012.
- Salamatin, A. N., Ekaykin, A. A., and Lipenkov, V. Y.: Modelling isotopic composition in precipitation in central Antarctica, 30 *Materialy Glyatsiologicheskikh Issledovaniy*, 97, 24-34, 2004.



- Scarchilli, C., Frezzotti, M., P., G., De Silvestri, L., Agnoletto, L., and Dolci, S.: Extraordinary blowing snow transport events in East Antarctica, *Clim. Dyn.*, 34, 1195-1206, doi:10.1007/s00382-009-0601-0, 2010.
- Schneider, D. P., Steig, E., Van Ommen, T., Dixon, D. A., Mayewski, P. A., Jones, J. M., and Bitz, C. M.: Antarctic temperatures over the past two centuries from ice cores, *Geophys. Res. Lett.*, 33, 1-5, 2006.
- 5 Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., Buntgen, U., Caffee, M., Chellman, N., Dahl-Jensen, D., Fischer, H., Kipfstuhl, S., Kostick, C., Maselli, O. J., Mekhaldi, F., Mulvaney, R., Muscheler, R., Pasteris, D. R., Pilcher, J. R., Salzer, M., Schupbach, S., Steffensen, J. P., Vinther, B. M., and Woodruff, T. E.: Timing and climate forcing of volcanic eruption for the past 2,500 years, *Nature*, 1-7, doi:10.1038/nature14565, 2015.
- Stenni, B., Scarchilli, C., Masson- Delmotte, V., Schlosser, E., Ciardini, V., Dreossi, G., Grigioni, P., Bonazza, M., Cagnati, A., Karlicek, D., Risi, C., Udisti, R., and Valt, M.: Three-year monitoring of stable isotopes of precipitation at Concordia station, East Antarctica, *The Cryosphere Discussions*, 1-30, doi:10.5194/tc-2016-142, 2016.
- 10 Swingedouw, D., Ortega, P., Mignott, J., Guilyardi, E., Masson- Delmotte, V., Butler, P. G., Khodri, M., and Seferian, R.: Bidecadal north Atlantic ocean circulation variability controlled by timing of volcanic eruptions, *Nature Communications*, 6, 1-12, doi:10.1038/ncomms7545, 2015.
- 15 Turner, J., Colwell, S. R., Marshall, G. J., Lachlan-Cope, T. A., Carleton, A. M., Jones, P. D., Lagun, V., Reid, P. A., and Iagovkina, S.: The SCAR READER project: Toward a high-quality database of mean Antarctic meteorological observations, *J. Clim.*, 17, 2890-2898, doi:10.1175/1520-0442(2004)017<2890:TSRPTA>2.0.CO;2, 2004.
- Vladimirova, D. O., and Ekaykin, A. A.: Climatic variability in Davis Sea sector (East Antarctica) for the last 250 years based on geochemical investigations of "105 km" ice core, *Probl. Arktiki i Antarktiki*, 1, 102-113, 2014.

**Table 1: Information on sites where individual time-series were obtained.**

Site series /	Coordinates		Alt., m a.s.l.	Time interval, years AD	Acc. rate, mm w.e.	Sample resolution, cm	δD measurements	Accumulation record available	Reference
	Lat., °S	Long., °E							
105 km	67.433	93.383	1407	1757-1987	310	5	LSCE, mass spectrometry; CERL, laser spectroscopy	Yes	(Vladimirova and Ekaykin, 2014)
400 km	69.95	95.617	2777	1254-1987	170	100	LSCE, mass spectrometry	No	this work
VRS 2013 stack (Vostok)	78.467	106.84	3490	1654-2010	21	1-7	LSCE, mass spectrometry; CERL, laser spectroscopy	Yes	(Ekaykin et al., 2014)
NVFL-1	77.11	95.072	3775	1711-1944	31	10	CERL, laser spectroscopy	No	this work
NVFL-3	76.405	102.167	3528	1978-2009	34	10	CERL, laser spectroscopy	No	this work
PV-10	72.805	79.934	2800	1976-2009	103	2	CERL, laser spectroscopy	No	this work
200 km	68.25	94.083	1990	1640-1987	271	NA	No	Yes	(Ekaykin et al., 2000)

NA = not applicable

5 **Table 2: Correlation matrix between individual surface air temperature records from meteorological stations in the Princess Elisabeth Land.**

	Casey	Mirny	Davis	Mawson	Vostok
Casey	1	0.82	0.60	0.53	0.54
Mirny		1	0.86	0.77	0.67
Davis			1	0.86	0.58
Mawson				1	0.62
Vostok					1

All the correlation coefficients are statistically significant with 95 % confidence level.




Figure 1: The Princess Elisabeth Land sector of East Antarctica. Blue iso-contours display the spatial pattern of surface snow $\delta^{18}\text{O}$ (Vladimirova et al., in preparation). The light blue contour shows the shoreline of subglacial Lake Vostok. Yellow dots mark the location of individual records used here. Stars depict the location of former or present research stations.




Figure 2: δD records from 6 individual series used in this study.




Figure 3: Normalized and low-pass filtered individual records (with a cut-off for variations on timescales shorter than 27 years), displayed using the same colours as in Figure 2.

The thick grey line is the stacked record (PEL2016). The dashed grey lines show the less robust marginal parts of the stack.

- 5 Vertical dashed lines mark reference horizons that contain the nss-SO_4^{2-} peaks pointing out Tambora (1815) and Pinatubo (1991) volcanic eruptions, respectively deposited until 1816 and 1993 in Antarctica.




Figure 4: Climatic variability in the Southern Hemisphere in 1958-2015.

a – Composite temperature anomaly in the Princess Elisabeth Land (based on records from Mirny, Davis and Vostok). The red shading displays ± 1 standard error of mean.

5 **b** – Antarctic Oscillation Index from NOAA (solid line), and BAS (dashed line). See text for details.

c – Interdecadal Pacific Oscillation Index.

d – Indian Ocean Dipole Index.

Thick lines are low-pass filtered (with a cut-off for variations on timescales shorter than <27 years).




Figure 5: Antarctic climatic variability over the past 350 years.

a – Number of individual records in the stacked isotopic record;

b – Temperature anomaly relative to 1980-2009, based on Princess Elisabeth Land meteorological records (red) and reconstructed from the stacked isotopic record (PEL2016 – magenta). Shading is ± 1 standard error of mean. Dashed lines denote less robust marginal parts of the PEL2016 record.

c – Low-pass filtered values of the IOD index.

d – Antarctic temperature anomaly from (Schneider et al., 2006).

e – Normalized and low-pass filtered stacked isotopic record for East Antarctica (data from PAGES 2k Consortium, 2013).




Figure 6: Normalized (relative to period 1952-1981) and low-pass filtered records of snow accumulation rate at sites “200 km” (purple), “105 km” (magenta) and Vostok (orange).