

## Response to the reviewers

We would like to thank the editor and both reviewers for their support, comments and suggestions that helped clarify some aspects of our manuscript. We respond to all comments and questions raised by the reviewers. The reviewers' comments are in *italic* and our answers in **bold**.

### **Reviewer #1:**

#### Recommendation

*This paper investigates how changes in orbital parameters and ice sheets during the last glacial-interglacial cycle impact Arabian Sea productivity through changes in the monsoon intensity and spatial pattern. This is an important topic, since productivity estimates from Arabian Sea cores are usually interpreted as proxies of the monsoon intensity. Using numerical simulations with a earth system model, this study shows that the relationship between monsoon intensity and productivity is non trivial, because spatial shifts in the monsoon jet axis influence both wind stress and its curl, which both control the influence of the coastal upwelling. The paper topic is interesting, and the analyses are scientifically sound. I however suggest a couple of additional diagnostics, re-arrangement of some results, additional critical discussion of the model, which should improve the clarity of the paper and make it more convincing (see general comments below).*

**We thank reviewer#1 for his very constructive comments on our work.**

#### General comments

- 1. I feel that some diagnostics on the intensity of the upwelling (eg SST anomaly in the coastal box relative to the Indian Ocean average and/or vertical velocities and/or depth of the thermocline . and nutricline-) would be helpful to relate the changes in nutrients with changes in the upwelling intensity. A bit more validation of the upwelling characteristics in the present-day simulation also would not harm.*

**We add an evaluation of the SST anomalies in boreal summer on a new panel on figure 3.**

- 2. There is no discussion of how the biases of the model for the current-day climate (e.g. its underestimated productivity) may influence the overall results of the study.*

**We started to give some explanation on the effects of the model biases on page 15 and on page 6 but we can develop the discussion on this subject.**

**We have added a new paragraph on the model limitations to page 15 L10-21. We need to keep in mind that the model's coarse resolution does not allow for a very precise representation of the region dynamics. This may have altered the relative weight of the processes related to wind stress and wind stress curl and can explain why the astronomical signal is weak in the productivity changes. Arabian Sea productivity in the CTRL simulation shows quite large differences with data, especially for the high coastal productivity extension in the western Arabian Sea (Fig. 3a). This can result from the model coarse resolution which prevents the representation of meso-scale processes such as eddies. These fine-scale processes are shown to be of importance for the coupling between biology and physics (Resplandy et al., 2011). Moreover, these meso-scale processes contribute strongly to the export of nutrients offshore and thus can explain why our high productivity area is more restricted to the coast than in the observations. From our set of simulations we cannot assess the effect of the underestimation of productivity on our results. Our simulations may underestimate productivity levels and variations but since most of the main physical processes are represented we can draw conclusions on the link between these processes and productivity and their changes through time. A way to overcome these limitations and to quantify their effects would be to work with several other models and to analyse the coupling between biology and physics in those different models.**

- 3. The abstract could be improved (suggestions below). It may also be beneficial to show and discuss Fig. 14 much earlier in the paper, in order to describe the relative effects of alongshore stress and near-shore Ekman pumping in the present-day climate ahead of the past climates discussion.*

**We have taken this suggestion into account for the abstract (see below).**

We have moved figure 14 to the evaluation section and its description in the evaluation section on p6 L26-34. It helps justifying that the mechanisms behind productivity changes in the Arabian Sea are represented in the model even though the amplitude of the changes in productivity and winds are underestimated.

4. I am not sure that it is worth discussing the central Arabian Sea region (Figures 6 and 9). A reason for that is that this region is usually viewed as highly influenced by what happens in the coastal upwelling region. I hence feel that focussing on the coastal box is enough. If you want to keep this central Arabian Sea box, a diagnostic as that of figure 14, which relates the interannual variability of productivity to wind stress (here an indicator of vertical mixing) and wind stress curl would be helpful.

**This region allows us to discuss changes in the monsoon pattern, in particular Fig. 9b but we remove the figures and the reference to it throughout the text.**

5. It may be useful to show maps of the JJAS climatological wind stress and Ekman pumping values for all the simulations, to more visually relate how the changes in low pressures over the continent relate to changes in the monsoon flow.

**It is difficult to distinguish changes between the different climates on such maps. We do not think it would be clear enough. We believe that Figure11 (old figure10), which shows the link between the barycenter position and the wind stress and wind stress curl intensity is sufficient to show the link between changes in the low pressures and changes in the monsoon flow.**

### Detailed comments:

P1, L1: maybe ~~the~~ current-climate Indian monsoon. . . + Ok

P1, L5-10: I think that the abstract could be clarified. Maybe mention explicitly that coastal upwelling is fuelled by a combination of alongshore stress intensity and upward Ekman pumping to the west of the jet axis. There is however strong downward Ekman pumping to the east of the jet axis, so that changes in coastal alongshore stress / curl depend both on the jet intensity and position. You can then relate changes in the intensity and position to the exact position of the low pressure over Tibet, with astronomical parameters having impact mostly on the intensity and changes in ice sheets rather influencing the jet position.

**Ok, we have changed P1, L5-10 to: Locally, productivity is fueled by nutrient supply driven by Ekman dynamics. Upward transport of nutrients is modulated by a combination of alongshore wind stress intensity, which drives coastal upwelling, and by a positive wind stress curl to the west of jet axis resulting in upward Ekman pumping. On the East of the jet axis there is however a strong downward Ekman pumping due to a negative wind stress curl. Consequently, changes in coastal alongshore stress/curl depend both on the jet intensity and position. The jet position is constrained by the Indian summer monsoon pattern, which in turn is influenced by the orbital parameters and the ice sheet cover.**

P1, L18: Another useful reference here is McCreary et al. (2009) (see full reference below). Ok

P1, L24: maybe indicate ~~upward~~ Ekman pumping. It may be interesting to mention offshore downward pumping to the right of the jet axis.

**Ok, we have added upward or downward in front of Ekman pumping throughout the text and in the Figure captions (e.g. Figure 6).**

**We have changed the text such as in P1 L23-24 P2 L1-6:**

**In addition the wind's tendency to turn on itself in the horizontal plane, quantified by the wind stress curl, also drives upward and downward water transport (Marshall and Plumb, 2008). Between the axis of the jet and the western coast, the wind stress is cyclonic and the wind stress curl is positive, it drives a divergent flow that causes upward Ekman pumping (Murtugudde et al., 2007; Barber et al., 2001; Anderson et al., 1992; Findlater, 1969). On the other side of the axis jet however, the wind stress is anticyclonic, the wind stress curl is therefore negative and drives a convergent flow and downward Ekman pumping. Coastal upwelling and upward Ekman pumping are responsible for increased productivity in the western coastal Arabian due to a higher supply of nutrients to the surface layer (Anderson et al., 1992; Anderson and Prell, 1992)."**

P2, L5: Also mention the specific role of eddies, quoting Resplandy et al. (2011).

**We change P2 L11-12 :** “and increase productivity in those regions (Resplandy et al., 2011; Wiggert et al., 2005; Prasanna Kumar et al., 2001; Lee et al., 2000). **At the mesoscale, filaments contribute to the lateral advection of nutrients from the coast to the central Arabian Sea (Resplandy et al., 2011).”**

P2, L9-11: Recent studies indicate that, due to changes in atmospheric stability, an increase in rainfall is not necessarily associated with an increase of the associated circulation (e.g. Held and Soden J. Clim. 2006 in the context of anthropogenic climate change). **Ok. On P2 L14 we have added :** **“However, Held and Soden (2006) indicate that, in the context of anthropogenic climate change, an increase in rainfall is not necessarily associated with an increase of the associated circulation due to changes in atmospheric stability. This result questions the reliability of such an indicator for the monsoon intensity.”**

P2, L24: Be more specific: change of ice sheet heights in which regions? Same comment for ice volume L27. **We have changed it to :** **“changes in ice-sheet height between glacial and interglacial climates”** on L33 which includes both changes in the northern and southern hemisphere and to **“changes in ice volume, especially in the Northern Hemisphere”** on L35

P3, L14 and following: ~~the~~ **LMDZ5A atmospheric general circulation model** (and likewise for the other components) **Ok**

P4, L30-34: A recent study (Keerthi et al. 2016, see below for full ref) shows that, in winter (when the cloud cover over the Arabian Sea is low), there is a good agreement between various satellite datasets for the Northern Arabian sea, but large differences in terms of amplitude. You may hence want to add a cautionary note about uncertainties of the observational estimate. **We do not see in Keerthi et al 2016 a discussion on satellite observations discrepancies concerning chlorophyll estimates.**

P5, L1: vertical mixing is more specifically due to convective overturning in presence of strong southward winds that bring dry, cold continental air. **We have changed P5L8-10 to :** **“In boreal winter, the mechanisms behind productivity changes are different compared to summer ones, because the winds reverse and blow from the north-east to the south-west. The presence of strong southward winds generates a convective overturning which induces vertical mixing and brings nutrients to the surface.”**

P5, L11: typo: ? -> **yes this is a typo**

How is productivity computed from SeaWiifs? **Productivity is computed using the VGPM algorithm on SeaWiifs chlorophyll data (Behrenfeld and Falkowski 1997).**

**We add it in the text P5-L20-21 :** **“during the period 1998-2005 processed with the VGPM algorithm (Behrenfeld and Falkowski, 1997) to obtain monthly productivity”**

P5, L18-19: It is also likely that the absence of eddies in the model solution contributes to a weaker offshore export than in observations (but you note it a bit later on p6). The consequences of this underestimated productivity on your results has to be thoroughly discussed at the end of the paper. **We add it in the discussion of the model biases. See response to general comments #2.**

P5, L25-35: Time series of the mean seasonal cycle of alongshore wind-stress, near-shore wind stress curl and of an indicator of the upwelling (e.g. SST) would allow a more quantitative validation of the model than the existing figures. **We only focus on the boreal summer season and don't feel the need to evaluate the simulation in winter as well. Moreover, the use of such maps allows us to evaluate the distribution patterns of the different variables.**

P6, L10: Is ~~the~~ pathway appropriate in this context? **We change it to :** **“links”**

P6, L23-25: You should refer to fig4 to justify this diagnostic better. **Ok**

I am surprised by the position of the black start on figure 4e: it is on the edge of the region with SLP anomalies < -5 hPa, which is surprising for a barycentre. Is there a gridpoint with a very large negative SLP anomaly? Or maybe I'm colorblind: you should probably highlight the -5 hPa contour on that figure. **To clarify, we change panel e) by only plotting the SLPa-5 contour. The SLP anomalies lower than -5 hPa extend up to 55°N**

and the region on which we compute the barycenter goes up to 60°N so that the barycenter is attracted to the north and its position is at the limit of the SLPa-5 region over the Tibetan Plateau.

*P8, L5-16: I generally agree with the interpretation, but I feel that the explanation takes twists and turns. I would reorganize as follows: the shift of the Tibetan low leads to a poleward shift of the monsoon jet (5c). This leads to weaker alongshore winds in the Somalia and stronger alongshore winds in the Oman upwelling (5c). But this also brings the Ekman downwelling to the right of the jet axis closer to both Oman and Somalia coasts (5e). Both factors (alongshore, and offshore curl) contribute to a Somalia upwelling reduction, while the wind stress curl change seems to overwhelm the increased alongshore winds in the Oman region, leading to overall upwelling reduction and less nutrients (5b). The reduced wind stress also reduces the mixed layer depth in the Southern Arabian Sea (5d) but I am not sure it's so relevant to show this here. **Ok it is clearer this way and we have followed the reviewer's suggestion, except we do not write 'Tibetan Low' but 'SLPa-5 barycenter'.***

*Rather, showing SST (which characterizes the upwelling intensity) and/or vertical velocities in the model would allow to better characterize physical changes. Showing the depth of the nutricline would also be really helpful. **We already show the Ekman pumping on panel f) of this figure. Moreover, SST may not be a good indicator of upwelling when comparing different climate states (Emeis et al 1995).***

**Emeis, K.-C., Anderson, D.M., Doose, H., Kroon, D., Schulz-Bull, D., 1995. Sea-Surface Temperatures and the History of Monsoon Upwelling in the Northwest Arabian Sea during the Last 500,000 Years. Quat. Res. 43, 355-361. doi:10.1006/qres.1995.1041**

*Finally, I wonder why nutrient reduction is larger in the Oman region than in the Somalia region. Is it an effect of changes in biological uptake? **We see two hypotheses that can explain this difference in nutrient changes. The first one is a 1D explanation based on the biological uptake as you suggested. The second one is a 3D hypothesis, which implies changes in the horizontal currents. We would need to compute the nitrate balance in the area in order to identify the different contribution of processes such as advection or diffusion. But this is beyond the scope of this paper.***

*P8, L17-23: This figure is relevant for the coastal box, but is it so relevant for the offshore box, for which a lot of the productivity changes are most likely a consequences of changes in the coastal upwelling, exported offshore by the circulation. For the coastal box, it indicates that the curl change (favouring downwelling) wins over the change in coastal winds (favouring upwelling and enhanced vertical mixing). This would be confirmed by looking at the thermocline (and nutricline) depth and/or to vertical velocities, i.e. looking at physical indicators of the intensity of the upwelling. Another interest of this plot relies on the fact that productivity is decreased for all years: i.e. the change that you see is statistically significant. Finally the last interest of this plot is that it allows to evaluate the respective roles of Ekman pumping and alongshore wind stress for interannual variability: although not very clear with the current choice of color scale for panel a, there is a tendency for a weaker decrease in productivity for strong alongshore stress, as would be expected. We would also expect to see a stronger decrease in productivity for larger negative wind stress curl anomaly. If you confirm this (i.e. by performing a bilinear fit of the productivity on the stress & curl), illustrating the competition between those two effects for interannual variability will strengthen your case for explaining changes between CTL and EH. Nota: I later came to realize that you actually show this as figure 14, but it may be interesting to show it at an earlier stage in the paper.*

**We remove the central area analysis here and add a few words on it to the discussion. We can then increase the panels'size so we can better see the colors and the effect of increasing wind stress on productivity. Concerning physical indicators, we believe Ekman pumping on Figure 6 (old figure 5) and 8 (old figure 7) already sheds light on the links between physical and biological changes.**

**We move Fig. 14 to the evaluation section (it becomes figure 4) so that we can discuss the relative role of wind stress and curl earlier.**

*P9, L9-10: This statement is more confusing than helpful (where exactly in the Central AS?) **We remove the central area of the paper and thus this sentence.***

*P9, L11: Figure 8 would be usefully complemented by scatterplots between some of the variables (and the associated correlations): alongshore wind stress is strongly controlled by DTT (panels a and b); nutrients exert a (weaker) control on productivity (b and c): I would combine these scatterplot with Figure 9a (again, I don't think that figure 9b is so relevant, because wind stress is not really an upwelling driver away from the coast and productivity changes in the central Arabian Sea may not be the result of local processes).*

We remove panel b on figure 9 which concerns the central Arabian Sea.

We think that the scatterplots won't add more information than there is already in figure 9 (old figure 8) since there is no one-to-one relationship between most of the variables. It is quite clear when looking at the boxplots that DTT and wind stress are related and that PP and NO3 are linked as well.

P9, L24-25: Doing a scatterplot of alongshore wind stress vs. coastal wind stress curl would also be helpful to characterize the relations of the two parameters that control the upwelling. I would not say they are entirely independent: it seems that there is a tendency for a negative correlation. **We already have plotted this relationship between stress and curl on Figure 10 (old figure 9) and on figure 4 (old figure 14).**

P10, L15-20: It may be better to start by explaining the links between the low pressure position and the wind stress and curl, and then discussing the consequences on the productivity. To understand better the links between the low pressure position and wind stress intensity / curl, it would be good to add a figure with maps of the JJAS wind stress (vectors) and its curl (colors), with the position of the barycentre indicated.

**Such a figure would be difficult to read since changes are small.**

P10, L23: simply point out that the latitude of the barycentre exerts a strong control on the coastal wind stress amplitude. **Ok**

P10, L28: In general, in this section mention the increase of productivity with the central longitude of the Tibetan low. **We precise the longitude SLPa-5 barycentre to avoid any misunderstanding.**

P11, L1-10: I'm not that familiar with those orbital parameters. Wouldn't it be simpler to directly relate the position of the low pressure to the annual (or JJAS) solar heat flux properties over the region, and then describe more qualitatively how orbital parameters control this value? Or at least remind how these parameters influence insolation and hence the low (e.g. copy the text at lines 30-31 here). **Ok, we will add some precisions on the respective roles of the astronomical parameters on insolation here. P11 L8-11 On figure 12, we plotted the values of the climatic precession (defined as  $e \cdot \sin(w - 180^\circ)$  with  $e$  the eccentricity and  $w$  the precession), which modulates the northern hemisphere insolation, and obliquity, which controls the temperature contrast between the high and low latitudes, relative to the SLPa-5 barycentre position, in order to analyse the relationship between the astronomical parameters and the SLPa-5 barycentre's position."**

P11, L14: **remove only Ok**

P13, L8-9: can you explain this choice? **No simulation of the Early Holocene with the remnant Laurentide ice sheet was available at the time we performed our analysis. It is only available for the previous version of the model IPSL-CM4 (Marzin et al 2013).**

**Marzin, C., Braconnot, P., Kageyama, M., 2013. Relative impacts of insolation changes, meltwater fluxes and ice sheets on African and Asian monsoons during the Holocene. Clim. Dyn. 41, 2267-2286. doi:10.1007/s00382-013-1948-9**

**We add P13 L12 The simulation with this model version was not available at the time of our analyses.**

P13, L17: you could locate this core, e.g. on figure 7. **Ok we add the core position on Figure 2 and Figure 8**

P14, L9: provide a reference in support of this statement. **We can refer to Banakar et al. 2005.**

**Banakar, V.K.K., Oba, T., Chodankar, A.R.R., Kuramoto, T., Yamamoto, M., Minagawa, M., 2005. Monsoon related changes in sea surface productivity and water column denitrification in the Eastern Arabian Sea during the last glacial cycle. Mar. Geol. 219, 99-108. doi:10.1016/j.margeo.2005.05.004**

P14, L11-17: in the model. **Ok. We add it on several lines.**

P15, L9-19: This is nice. I would move this much earlier in the paper, because it allows to explain better the relative roles of wind stress and its curl at interannual timescales before moving to applying these explanations for climates of the past. **We move this figure in the evaluation section.**

P15: a thorough discussion on the possible consequences of the present-day climate biases in the model on the results of this study is needed. **We develop this point of the discussion. See response to general comment #2.**

## Figures:

Fig. 2: also draw the Northern box. **Ok**

Fig 3: left panels of b, c, d have some spurious horizontal lines on them. **This is due to the model resolution. What you see is the model grid cells, thus showing the quite coarse resolution used here.**

Some smoothing or median filter would not harm on the sat. wind stress intensity & curl. **Ok we regrid the observations on a 1 degree grid**

On d, also avoid saturating the colorscale. **Ok.**

Panel b: use the same spacing between vectors for the model and data. **Ok**

Fig 5: it would be useful to materialize the boxes that you use for integrated diagnostics on this figure. **Ok, we add the coastal box on the productivity panel only**

Fig 7: draw the coastal box. **We add Ekman pumping change on Figure 8 (old figure 7) so we won't add the coastal box otherwise it will be unreadable.**

Figure 8: describe the whiskers in the caption. I imagine it is a confidence interval: at what significance level? How is it computed? What are the small circles on this figure (and on figure 12). **We add the following precision to the figure caption: "The boxplots highlight the median value (bold line), the first and the third quartile (lower and higher limits of the box) and the 95% confidence interval of the median (upper and lower horizontal tics). The dots are extreme values that happened during the 100 years of simulation."**

**We also add this text in the legend of Figure 13.**

Figure 10: draw the boundaries of panels b, c, d on panel a. You can re-organize the panels into 2 x 2 to save space. **Ok to reorganize and we add the boundaries on panel a.**

Figure 12: it would be nice to locate the core / box that are used for figure 12 on another plot, e.g. figure 7. **Ok Clarify in the caption which panels are model results. We add "a) observed" and "b) simulated" in the caption.**

Figure 13: I don't find this picture very clear. I would do it upside down: start it by changes in orbital parameters / ice sheet driving changes in the Tibetan plateau low, which impact the intensity / pattern, stress / curl, the upwelling of nutrients, etc. . . Materialize the direction of arrows connecting boxes. **Ok, we reversed the order and add the direction of the arrows.**

## Reviewer #2:

The manuscript *From Monsoon to marine productivity in the Arabian Sea: insights from glacial and interglacial climates* by P. Le Mézo et al. is an interesting model study on the driving forces of ocean productivity in the Arabian Sea. A large number of paleo climate simulations of different warm and cold climates, forced by changes in orbital parameters, greenhouse gases and northern hemisphere ice sheets is used to systematically investigate changes in productivity patterns. Against the common paradigm that there is a straightforward relationship between pure Monsoon strength (wind mixing) and marine productivity, which was often used to infer past Monsoon changes from productivity proxies, the study highlights important aspects of changes in the Monsoon pattern and potentially opposing regional trends, for example in coastal and open ocean areas. According to the model results, wind stress curl, the changes of which are mostly driven by changes in the Monsoon pattern, turns out to be an equally important factor influencing productivity. The study is well designed, straightforwardly carried out and the demonstrated results are robust and clearly illustrated in many figures. The text could benefit from a number of clarifications, however, this does not compromise the overall excellent quality of the study. I suggest publishing after some minor revisions.


We thank reviewer#2 for his thoughtful and useful comments that helped improve the manuscript.

### Major comments

Since many readers are maybe not too familiar with the theory of wind stress curl, a short introduction into the most important mechanisms of upward nutrient transport and their interplay would probably be helpful. **Ok we add a more detailed description of the mechanisms in the introduction. P1 L23-24 P2 L1-6:**

“In addition the wind's tendency to turn on itself in the horizontal plane, quantified by the wind stress curl, also drives upward and downward water transport (Marshall and Plumb, 2008). Between the axis of the jet and the western coast, the wind stress is cyclonic and the wind stress curl is positive, it drives a divergent flow that causes upward Ekman pumping (Murtugudde et al., 2007; Barber et al., 2001; Anderson et al., 1992; Findlater, 1969). On the other side of the axis jet however, the wind stress is anticyclonic, the wind stress curl is therefore negative and drives a convergent flow and downward Ekman pumping. Coastal upwelling and upward Ekman pumping are responsible for increased productivity in the western coastal Arabian due to a higher supply of nutrients to the surface layer (Anderson et al., 1992; Anderson and Prell, 1992).”

Apart from MIS3, the control simulation stands out as the most productive period for the Arabian Sea. Can the results be extrapolated to other ocean areas, for example the prominent eastern boundary upwelling regions (EBUs)? What would be the consequences for the underlying oxygen minimum zone? **We have not looked at other regions. The recently submitted paper Bopp et al (in review) analyzes the changes in oxygen in the Indian Ocean in the LGM. The following 2 figures show the averaged changes in oxygenation between the LGM and the current day climate as simulated by the model and reconstructed in Jaccard and Galbraith 2012, between 0-1500m and 2000-5000m.**



The analysis of other upwelling systems is not the scope of this study.

Bopp, L., Resplandy, L., Untersee, A., Le Mézo, P., Kageyama, M., in review. Ocean (de)oxygenation from the Last Glacial Maximum to the 21 st century : insights from Earth System Models. Philosophical Transactions of the Royal Society.

*What are the model limitations? Does spatial resolution play a role? Since a rather coarse resolution was used here, what are the uncertainties and/or probable effects of parameterizations of sub gridscale processes or why would all these not matter? A first tentative explanation is given on page 15, l. 6-8, but this could be further elaborated. We wrote about some effects of the resolution on p6 but add a more complete paragraph on the model limitations and biases and how they affect our results and interpretation in the discussion section of the paper. P15 L10-21.* We need to keep in mind that the model's coarse resolution does not allow for a very precise representation of the region dynamics. This may have altered the relative weight of the processes related to wind stress and wind stress curl and can explain why the astronomical signal is weak in the productivity changes. **Arabian Sea productivity in the CTRL simulation shows quite large differences with data, especially for the high coastal productivity extension in the western Arabian Sea (Fig.3a). This can results from the model coarse resolution which prevents the representation of meso-scale processes such as eddies. These fine-scale processes are shown to be of importance for the coupling between biology and physics (Resplandy et al., 2011). Moreover, these meso-scale processes contribute strongly to the export of nutrients offshore and thus can explain why our high productivity area is more restricted to the coast than in the observations. From our set of simulations we cannot assess the effect of the underestimation of productivity on our results. Our simulations may underestimate productivity levels and variations but since most of the main physical processes are represented we can draw conclusions on the link between these processes and productivity and their changes through time. A way to overcome these limitations and to quantify their effects would be to work with several other models and to analyse the coupling between biology and physics in those different models.**

*When describing changes in wind stress curl often the words higher or lower are used, which may cause misunderstandings. Although the relative effect of the curl tendency on upward nutrient transport is the same, it needs to be kept in mind that the change maybe associated with a change of sign of the curl or not. I therefore suggest that the use of more or less negative/positive curl is a more correct wording. Please check throughout the text. This is a good suggestion. We have changed it throughout the text.*

## Minor points

*p. 1, l. 6-8: This sentence is confusing. It seems that the combination of increased wind stress and more positive curl could result in a reduction of productivity. However, this is not the case (Fig. 9). I think it is more correct to state that both changes in wind stress and curl need to be taken into account and that under opposing changes one or the other may be the ultimately dominating effect. We change L5-10 to: **Locally, productivity is fueled by nutrient supply driven by Ekman dynamics. Upward transport of nutrients is modulated by a combination of alongshore wind stress intensity, which drives coastal upwelling, and by a positive wind stress curl to the west of jet axis resulting in upward Ekman pumping. On the East of the jet axis there is however a strong downward Ekman pumping due to a negative wind stress curl. Consequently, changes in coastal alongshore stress/curl depend both on the jet intensity and position. The jet position is constrained by the Indian summer monsoon pattern, which in turn is influenced by the orbital parameters and the ice sheet cover.***

*p. 1, l. 12: typo, remove ~~an~~ affects* Ok

*p. 1, l. 21 : replace ~~in~~ India ~~by~~ ~~over~~ India* Ok

*p. 3, l. 15: add ~~semi-labile~~ before ~~dissolved~~ organic carbon* Ok

*p. 3, l. 29: add ~~greenhouse~~ before ~~gas~~* Ok



p. 3, l. 30: add *the Northern Hemisphere* before ice sheets (at least from Fig. 1 this seems to be the case) **We do not change it because there are some small changes in the Southern Hemisphere as well. You can mostly see it on the tip of South America on Figure 1.**

p. 4, l. 8: This sentence is confusing. Do you count MIS3 as glacial or interglacial? If glacial then you should mention that five glacial simulations were carried out. Otherwise, it is strange to see it listed among the glacial climates. Please check. **MIS3 is counted as a glacial climate. It is indeed 5 glacial simulations and not 4. We correct that.**

p. 4, l. 9: remove *of* **Ok**

p. 4, l. 26: a few more information on length of simulation and starting conditions would be nice **You can find information on the forcing parameters in Table 1. We can add information on the length of the simulation.**

p. 4, l. 31: I guess that 68\_N should be 28\_N **Yes that's a typo, we will change it throughout the text as it appears several times.**

p. 4, l. 33: replace *sediments bulk* by *bulk sediment* **Ok**

p. 4, l. 34: replace *changes* by *peaks* and *nes* by *maxima* **Ok**

p. 4, l. 34: Does the sentence starting here refer to the winter period? **Yes**  
If yes, then p. 5, l. 1: add *ashed line in* before *Fig. 2b* **Ok**

p. 5, l. 11: typo: NASA **Ok**

p. 5, l. 16: *Arabian* instead of *Arabic* **Ok**

p. 5, l. 29: what do you mean by reconstructions? Observations? Please explain. **We meant the re-analyses, we have changed this to avoid confusion.**

p. 6, l. 1: move *oarse* before *model* **Ok**

p. 6, l. 6 *model* **Ok**

p. 6, l. 20: in several places of the manuscript I'm wondering about the use of the word *global* Do you really mean global here, i.e. average over ALL longitudes and latitudes? It would be perfectly fine to use the regional annual average here. **In this case we mean the average on all latitudes and longitudes.**

p. 7, l. 1: does total productivity refer to net or gross productivity? I guess net, please clarify. **Yes, it refers to the net productivity, we precise it in the text.**

p. 8, l. 2: *concentrations* **Ok**

p. 8, l. 2: The sentence starting here is difficult to understand. Maybe splitting into several sentences will help. **Ok, we also remove the part on central AS as suggested by reviewer #1 (see general comment #4).**

p. 9, l. 12: remove *than CTRL* **Ok**

p. 9, l. 20: *macro-nutrient* **Ok**

p. 9, l. 21: the part *and, productivity...* does not make sense to me. Maybe there's something missing? **We remove this part since the link between productivity and nutrients is already stated in the previous sentence.**

p. 10, l. 4: reorder sentence to *as function of changes in summer wind stress and curl in the central Arabian Sea.* **It is clearer this way but we remove the central Arabian Sea as suggested by reviewer #1 (see general comment #4) so we also remove this sentence.**

p. 10, l. 21: use of word *global* I guess *overall* or *general* would be more appropriate here. Furthermore, the decrease is southeastward, which could be mentioned. **We use *overall* and add *as it moves southeastward***

p. 10, l. 33: replace *suggests and* by *demonstrates the* **Ok**

p. 11, l. 1: what do you mean by *climatic precession*? I am not familiar with the expression, so please define or use *precession* or *precessional index* instead. (see also next sentence) **We only defined the climatic precession in the figure caption. We clarify that in the text P11 L8-11 *On figure 12, we plotted the values of the climatic precession (defined as  $e \cdot \sin(w - 180^\circ)$  with  $e$  the eccentricity and  $w$  the precession), which modulates the northern hemisphere insolation, and obliquity, which controls the temperature contrast between the high and low latitudes, relative to the SLPa-5 barycentre position, in order to analyse the relationship between the astronomical parameters and the SLPa-5 barycentre's position.***

p. 11, l. 3: the barycenter position changes with both longitude and latitude, please mention **We change the sentence to *Climatic precession influences the SLPa-5 barycenter position both in latitude and longitude***

p. 11, l. 17: add *marine sediment* before *cores* **Ok**

p. 11, l. 23: *climates* **Ok**

p. 11, l. 30: which latitude range are you talking about? Shouldn't a high obliquity (= high summer insolation) reduce the latitudinal temperature gradient? **We talk about the latitudes used to compute the  $\Delta T T$  value (10°N-45°N) and (25°S-10°N). In our simulations a high obliquity, increases the summer insolation in the northern hemisphere high latitudes and reduces it in the low latitudes. This higher summer insolation in the NH warms the continent and increases the tropospheric temperature gradient in boreal summer, which is characteristic of a stronger summer monsoon.**

p. 12, l. 8: replace *solar* by *northern hemisphere summer* **Ok**

p. 12, l. 21: *simulations* **Ok**

p. 12, l. 24: here is a contradiction to what you say later (l. 34). First you say that there is a mismatch to Bassinot et al., later you explain why your model matches with Bassinot et al., please clarify.

**We mean that simulated productivity on the whole coastal area on which we made our analysis is not consistent with Bassinot et al. reconstructions. However, if we only focus on the northern part of the coastal area, our results match Bassinot et al's reconstruction for the mid-Holocene. Still they do not match the early Holocene reconstruction.**

**We have changed the formulation : *However, even if the simulated coastal summer productivity compares quite well to the data Rostek et al. 1997 (except for the LGM, their core is in the southern part of the coastal area), it shows some discrepancies when compared to the reconstructions in Bassinot et al. 2011 (their core is in the northern part of the coastal area).***

p. 13, l. 13: please check use of *globally* **We change it to *largely***

p. 13, l. 17-24: this part belongs to the method section **Ok we move it to the Method section**

p. 13, l. 27: replace *are* by *is* **Ok**

p. 13, l. 31: region up to 68\_N? Please check **Yes same typo as previously. We change it to 28N.**

p. 14, l. 2: I don't think that the difference between productivity and export production in particular is a major problem for model and data comparison. It is rather that there are too many unknowns on what is finally best describing the signals contained in sediment records. As it stands, the statement seems to open a completely new aspect, which demands to be further elaborated. However, since it is not the main focus of the study I

*suggest leaving it out. We have chosen to keep it here but state more clearly that it has to be viewed as a perspective of this work.*

*p. 14, l. 11-17: it seems to me that in this paragraph you write about productivity, but it is actually the export, which is explained. Please check. We wanted to explain primary productivity but it is true that we miss some arguments and that this description better fits the export production. We change it to make it clear.*

*p. 15, l. 3: add ~~or less~~ before ~~nutrients~~* **Ok**

*p. 15, l. 20: replace ~~on~~ by ~~of~~* **Ok**

*Again, this sentence seems to give a hint to problems of model data comparison. I ~~do~~ suggest adding a full paragraph on what this study contributes to a improved model-data comparison or leave it out.*

**We change the sentence to: *Í This study allows us to draw attention to certain points that may affect the reconstruction of past climate and productivity and hence the comparison between model and data.Î***

*p. 15, l. 24: replace ~~should~~ by ~~is~~ expected to* **Ok**

## **Figures:**

*Figure 2, caption: 68\_N? Probably this should be 28\_N? **Yes***

*Figure 8: caption: replace ~~dash~~ by ~~dashed~~* **Ok**

*Figure 14: ~~circles~~+or ~~color~~ of the circles* **Ok**

## NOTE:

We realized that the computing of the barycenter was erroneous: in the submitted version, the barycenter was computed by taking the SLP anomalies relative to the regional mean (20W-150E,30S-60N) and not to the global mean as we intended. We corrected that. This correction made us change figures 5, 11 and 12. The barycenters move slightly north, especially in some glacial climates, but this does not change our conclusions. The new figures are added at the end of this document.

We added a few word in section 3.2.2 p11 L2-7:

**This latitudinal movement of the SLPa-5 barycentre with the monsoon strength seems also to be dependent on the glacial or interglacial state of the simulation. Indeed, in glacial simulations the SLPa-5 barycentre is north of the CTRL barycentre even if the monsoons are less intense (LGM, MIS4D and MIS4M), however the other glacial simulations with higher summer monsoon intensity (MIS3 and MIS4F) have their barycentre located north of these glacial simulations (Fig. 10). Similarly, in inter-glacial climates, the Holocene simulations have a barycentre north of the CTRL one (Fig. 10).**

And we change a sentence on p.12 l.17-19

**The changes in the ITCZ position highlighted by Fleitmann et al. (2007) are consistent with our results, especially with the changes in the position of the SLPa-5 barycentre in latitude given a similar background state: glacial or interglacial (Fig.11).**

instead of ~~The~~ changes in the ITCZ position highlighted by Fleitmann et al. (2007) are consistent with our results, especially with the changes in the position of the SLPa-5 barycentre in latitude (Fig.11).

# List of all relevant changes made in the manuscript:

---

1. We removed all the discussion and figures related to the central Arabian Sea as suggested by the reviewers.
2. We added a more developed paragraph on the model biases in our discussion.
3. We modified some of the figures to add more physical variables (in the evaluation for instance) and we re-organized some of the figure's order (Fig.14 becomes Fig.4)
4. We modified the computation of the SLPa-5 barycenter, which modifies figure 5, 11 and 12.

# From Monsoon to marine productivity in the Arabian Sea : insights from glacial and interglacial climates

Priscilla Le Mézo<sup>1</sup>, Luc Beaufort<sup>2</sup>, Laurent Bopp<sup>1</sup>, Pascale Braconnot<sup>1</sup>, and Masa Kageyama<sup>1</sup>

<sup>1</sup>LSCE/IPSL, UMR 8112 - CEA/CNRS/UVSQ, Centre CEA-Saclay, Orme des Merisiers, 91191 Gif-sur-Yvette, France

<sup>2</sup>CEREGE, UMR 7330, CNRS-IRD-Aix Marseille Université, Av. Louis Philibert, BP80, 13545 Aix en Provence, France

Correspondence to: Priscilla Le Mézo (priscilla.le-mezo@lsce.ipsl.fr)

**Abstract.** The current-climate Indian monsoon is known to boost biological productivity in the Arabian Sea. This paradigm has been extensively used to reconstruct past monsoon variability from paleo-proxies indicative of changes in surface productivity. Here, we test this paradigm by simulating changes in marine primary productivity for 8 contrasted climates from the last glacial-interglacial cycle. We show that there is no straightforward correlation between boreal summer productivity of the ~~western and central~~ Arabian Sea and summer monsoon strength across the different simulated climates. Locally, productivity is fueled by nutrient supply driven by Ekman dynamics ~~depending on both wind stress and~~. Upward transport of nutrients is modulated by a combination of alongshore wind stress intensity, which drives coastal upwelling, and by a positive wind stress curl ~~. In our simulations, a stronger monsoon with intensified wind stress on the Arabian Sea can lead to either increased or reduced productivity depending on the exact ocean surface on which increased wind stress and a positive~~ to the west of jet axis resulting in upward Ekman pumping. On the East of the jet axis there is however a strong downward Ekman pumping due to a negative wind stress curl ~~are acting. The effects of winds onto the ocean are modulated~~. Consequently, changes in coastal alongshore stress/curl depend both on the jet intensity and position. The jet position is constrained by the Indian summer monsoon ~~intensity and pattern (e.g. position of the low-level jet over the Arabian Sea)~~ pattern, which in turn ~~are driven by the orbital~~ is influenced by the astronomical parameters and the ice sheet cover. ~~The orbital~~ The astronomical parameters are indeed shown to impact wind stress intensity in the Arabian Sea through large scale changes in the meridional gradient of upper tropospheric temperature. But both the ~~orbital~~ astronomical parameters and the ice sheets ~~affects~~ affect the pattern of wind stress curl through the position of the sea level depression barycentre over the monsoon region (20°W-150°W, 30°S-60°N). The combined changes in monsoon intensity and pattern lead to some higher glacial productivity during the summer season, in agreement with some paleo-productivity reconstructions.

## 20 1 Introduction

The Arabian Sea biological productivity is influenced by the strong seasonal activity of the atmospheric circulation

(~~Schott and McCreary, 2001; Ivanova et al., 2003; Lee et al., 2000; Luther et al., 1990~~)

(McCreary et al., 2009; Ivanova et al., 2003; Schott and McCreary, 2001; Lee et al., 2000; Luther et al., 1990). During the boreal summer, the Southwest monsoon consists of strong winds blowing from the south-west to the north-east of the Indian

Ocean. These winds result from the rapid heating of the landmass relative to the ocean, which creates a pressure gradient between the southern Indian Ocean high pressure cell and the low pressure cell over the Tibetan Plateau. During this season, heavy precipitation occurs ~~in-over~~ India and south-east Asia. In the Arabian Sea, the alongshore winds off the coast of Somalia focus into a low-level jet, called the Somali Jet and generate a strong coastal upwelling (Anderson et al., 1992; Findlater, 1969). In addition, the ~~positive-curl-of-the-wind-stress-between-wind's tendency to turn on itself in the horizontal plane, quantified by the wind stress curl, also drives upward and downward water transport (Marshall and Plumb, 2008).~~ Between the axis of the jet and the western coast ~~induces Ekman pumping,~~ the wind stress is cyclonic and the wind stress curl is positive, it drives a divergent flow that causes upward Ekman pumping (Murtugudde et al., 2007; Barber et al., 2001; Anderson et al., 1992; Findlater, 1969). ~~These two processes~~ On the other side of the jet axis however, the wind stress is anticyclonic, the wind stress curl is therefore negative and drives a convergent flow and downward Ekman pumping. Coastal upwelling and upard Ekman pumping are responsible for increased productivity in the western coastal Arabian Sea ~~due thanks~~ to a higher supply of nutrients to the surface layer (Anderson et al., 1992; Anderson and Prell, 1992). The upwelled nutrients are advected from the coast to the north and the interior of the sea. ~~So~~ Thus, productivity in the central and northern Arabian Sea also increases during the Southwest Monsoon (Caley et al., 2011; Prasanna Kumar et al., 2001; Keen et al., 1997). Wind stress and mixing of the upper layers, as well as Ekman pumping generated by the positive wind stress curl, also contribute to the supply of nutrients to the surface layers and increase productivity in those regions (~~Resplandy et al., 2011; Wiggert et al., 2005; Prasanna Kumar et al., 2001; Lee et al., 2000).~~ (Wiggert et al., 2005; Prasanna Kumar et al., 2001; Lee et al., 2000). At the mesoscale, filaments contribute to the lateral advection of nutrients from the coast to the central Arabian Sea (Resplandy et al., 2011).

20 Monsoon intensity can be characterised in different ways, depending on the observational scale and on the studied processes. Precipitation is a major indicator of monsoonal changes. For example, the rainfall-based index, defined as the seasonally averaged precipitation over all the Indian subcontinent from July to September, is used to monitor the strength of the monsoon over India (Mooley and Parthasarathy, 1984). However, Held and Soden (2006) indicate that, in the context of anthropogenic climate change, an increase in rainfall is not necessarily associated with an increase of the associated circulation due to changes in atmospheric stability. This result questions the reliability of such an indicator for the monsoon intensity. A second indicator of the monsoon strength is based on the sea level pressure (SLP) that is a large-scale fingerprint of the monsoon. The monsoon strength can be determined by the SLP anomaly gradient between a northern region over the Tibetan Plateau, where the Tibetan Low develops during the monsoon months, and a southern region over the southern Indian Ocean, where the Mascarene high develops. The large-scale changes of SLP impact the local dynamics over the Arabian Sea (Schott and McCreary, 2001).

30 Monsoon intensity can also be related to the strength of the winds over the Arabian Sea and the associated upwelling. The general paradigm is that a stronger summer monsoon generates stronger upwelling that enhances productivity. Based on this paradigm, past monsoon intensities have been reconstructed using proxies of productivity from marine sediment cores (Caley et al., 2011; Ivanova et al., 2003; Clemens and Prell, 2003).

Monsoon reconstructions and modelling studies (Marzin and Braconnot, 2009; Braconnot et al., 2008; Anderson and Prell, 1992; Prell et al., 1992) have shown that insolation variations are the major driver of fluctuations in the summer monsoon

intensity : the monsoon is stronger when the Northern Hemisphere summer insolation is higher (e.g. during the Holocene). Changes in the ~~orbital~~astronomical parameters, such as the precession that is defined as the longitude of the perihelion, or the obliquity that is defined as the angle between the equator and the orbital plane, modify the seasonal cycle of insolation. Along with ~~orbital~~astronomical parameters, changes in ice-sheet height between glacial and interglacial climates also have an impact  
5 on the monsoon intensity (Masson et al., 2000; Emeis et al., 1995; Prell et al., 1992; Anderson et al., 1992).

There has been some concern about the fact that marine proxies for productivity may be influenced by other processes than monsoon intensity, such as changes in ice volume, especially in the Northern Hemisphere, aeolian transport of nutrients or the Atlantic Meridional Overturning Circulation (Ruddiman, 2006; Ziegler et al., 2010; Caley et al., 2011). Moreover, most studies linking monsoon and productivity in the past have focused on the monsoon intensity but, the monsoon pattern, e.g  
10 wind orientation, can also change in time. Sirocko et al. (1991) have shown that summer monsoon mean position was shifted southward during glacial periods. The monsoon pattern affects the position and the orientation of the low-level jet over the Arabian Sea, which modifies the upwelling of nutrients in the Arabian Sea (Anderson and Prell, 1992). Furthermore, Bassinot et al. (2011) showed opposite evolutions of the upwelling behaviour in the western coastal Arabian Sea and the south-western tip of India during the Holocene, which they related to a southward shift of the monsoonal winds.

15 Here, we investigate the relationship between the summer monsoon intensity and the Arabian Sea biological productivity. How do changes in the summer monsoon ~~structure~~pattern along with changes in its intensity impact productivity in the Arabian Sea ? Could the variations in the summer monsoon structure explain higher productivity rates in some glacial climates ? To answer these questions, we test the effects of a range of astronomical parameters and different ice-sheet states on the Arabian Sea productivity.

20 In section 2, we describe the model we use and the experiments we performed, we evaluate the model results for the pre-industrial and detail the analyses we performed. In section 3, we explain the changes in productivity in the Early Holocene and then, look at several glacial and interglacial climates to link productivity changes to local dynamics and boundary conditions. In section 4, we discuss our results in the light of the summer monsoon paradigm and, we perform a simple model-data comparison and discuss the effects of seasonality on productivity. Finally we summarise our results and give some perspectives.

## 25 **2 Model, experiments, evaluation and diagnostics**

### **2.1 The model**

This study uses an Earth System Model (ESM) that explicitly represents the global climate, oceanic circulation and marine productivity. We use the IPSL-CM5A-LR model developed at the Institut Pierre Simon Laplace (IPSL) (Dufresne et al., 2013). This ESM is composed of the LMDZ5A atmospheric general circulation model ~~LMDZ5A~~ (Hourdin et al., 2013) coupled to the  
30 ORCHIDEE land-surface model ~~ORCHIDEE~~ (Krinner et al., 2005) and the ~~ocean model~~-NEMO v3.2 ocean model (Madec, 2011), which includes the OPA9 ocean general circulation model ~~OPA9~~, LIM-2 the sea-ice component ~~LIM-2~~ (Fichefet and Maqueda, 1997) and the ~~biogeochemical model~~-PISCES ~~PISCES~~ biogeochemical model (Aumont and Bopp, 2006). These components are coupled once a day using the OASIS coupler (Valcke, 2012).



We use the low-resolution (LR) version of the model with a regular atmospheric grid of  $96 \times 96$  points horizontally and 39 vertical levels and, an irregular horizontal oceanic grid (ORCA2.0) with  $182 \times 149$  points corresponding to a nominal resolution of  $2^\circ$ , enhanced near the equator and over the Arctic and sub-polar North Atlantic. The ocean vertical grid comprises 31 levels with intervals from 10 meters for the first 150 meters and up to 500 meters for the bottom of the ocean.

5 The PISCES model simulates marine bio-geochemistry and lower trophic levels. PISCES includes two phytoplankton types (nano-phytoplankton and diatoms), two zooplankton size-classes (micro- and meso-zooplankton) and two detritus compartments distinguished by their vertical sinking speed (small and large organic matter particles), a [semi-labile](#) dissolved organic carbon pool, and five nutrients (Fe,  $NO_3$ ,  $NH_4$ , Si, and  $PO_4^{3-}$ ) (Aumont and Bopp, 2006). In PISCES, phytoplankton growth is a function of temperature, light, mixed layer depth and nutrients.

## 10 2.2 Experiments

Here, we exploit 8 simulations of IPSL-CM5A-LR forced by different boundary conditions ([orbital parameters](#), [astronomical parameters](#), [greenhouse](#) gas concentrations and ice sheets cover), to account for different climates throughout the last glacial-interglacial cycle, as detailed in Table 1 and Figure 1.

The reference simulation (CTRL), is a pre-industrial climate with no external forcing such as volcanoes or anthropogenic activities (Dufresne et al., 2013), forced by pre-industrial CMIP5 forcings (Taylor et al., 2012) and the present day ice sheet (0k on Figure 1). A mid-Holocene (MH) simulation, 6 kyr BP ([Kageyama et al., 2013](#)), part of Paleoclimate Modeling Intercomparison Project phase 3 (PMIP3) (Braconnot et al., 2012) and an early-Holocene (EH) simulation, 9.5 kyr BP, are used to study productivity changes in different interglacial climates. The EH simulation trace gas concentrations are the same as for the CTRL simulation whereas  $CH_4$  and  $N_2O$  concentrations are slightly lower for the MH simulation compared to CTRL (Table 1). MH and EH simulations mainly differ in their astronomical parameters, especially in the precession value (Table 1, Fig. 1). Both Holocene simulations are forced by the present-day ice sheet cover (Fig. 1). ~~Four~~ [Five](#) glacial simulations have also been performed ~~for~~ the last glacial maximum (LGM, 21 kyr BP), the [marine](#) isotopic stage 3 (MIS3, 46 kyr BP) and three [marine](#) isotopic stage 4 states : MIS4F (60 kyr BP), MIS4M (66 kyr BP) and MIS4D (72 kyr BP). The LGM, which has also been performed for PMIP3 ([Kageyama et al., 2013](#)), has the largest ice sheet (Fig. 1) (Abe-Ouchi et al., 2015), which modifies the land-sea distribution and topography since the sea-level is reduced by about 120 meters. The LGM run has the lowest greenhouse gas concentrations of this set of 8 simulations (Table 1). Two of the three MIS4 simulations (MIS4F and MIS4D) are described in Woillez et al. (2014). The MIS4 ice-sheets have been prescribed by using the 16 kyr BP ice-sheet, which is the period for which we have an ice sheet reconstruction for the same sea level as during MIS4, i.e. 70 meters lower than today (Ice6g-16k on Fig. 1) (Peltier et al., 2015). This is the most realistic we could do given the available reconstruction at the time of running the MIS4 experiments (Woillez et al., 2014). However, our MIS4 runs are different from the ones described in Woillez et al. (2014) since we added the nutrient inputs from dust, rivers and sediments that are essential to marine productivity. Large changes in precession occur between the three MIS4 simulations (Table 1, Fig. 1). The MIS3 simulation uses the same ice-sheet reconstruction as MIS4 and it has the lowest eccentricity and highest obliquity of all 8 simulations (Table 1, Fig. 1).

In PISCES, three source terms contribute to the input of nutrients in the ocean: atmospheric dust deposition, river input and sediment mobilisation. The change in sea-level in glacial climate simulations modifies the land-sea mask, thus in the LGM, MIS3 and all MIS4 (F,M,D) simulations, the source terms were adjusted so that the ocean receives the same quantity of associated nutrient supply as in CTRL. In these simulations, no attempt was made to account for the dustier glacial states (Bopp et al., 2003). All productivity changes are therefore due to other factors.

Our analyses are performed on 100 years of monthly outputs from the last stable part of each simulation.

### 2.3 Modern evaluation of the summer mean

This study focuses on primary productivity in the Indian Ocean for the last glacial-interglacial cycle as simulated by the IPSL-CM5A-LR coupled model. Figure 2 shows the seasonal cycle of ~~productivity for three~~ observed present-day productivity (data from SeaWIFS 1998-2014 (Lévy et al., 2007)) for two areas in the Arabian Sea: a coastal area in the western Arabian Sea (Fig. 2a, orange area) ~~;~~ an area that covers the central-east Arabian Sea and a northern region (Fig. 2a, grey area) ~~and a northern region (60E-68E, 20N-68N). Productivity~~ black box. Current-day productivity has two periods of bloom: one in summer and one in winter. In ~~both the coastal and central regions~~ the coastal Arabian Sea, the summer season is the most productive period of the year (Fig. 2b) and contributes the most to the ~~sediments bulk~~ bulk sediment composition. In the northern Arabian Sea, both seasons are equally productive in the data (Fig. 2b). In boreal winter, the mechanisms behind productivity ~~changes~~ peaks are different compared to ~~summer ones.~~ The the summer maxima, the winds reverse and ~~nutrients are brought to the surface thanks to~~ blow from the north-east to the south-west. The presence of strong southward winds generates a convective overturning which induces vertical mixing and ~~advected from the central sea to the coast.~~ brings nutrients to the surface. During this period, productivity is high in the north-western Arabian Sea (~~Fig. dashed line in figure~~ 2b). Following these observations, we focus our analyses on the boreal summer season, defined as June-July-August-September (JJAS) to account for the whole summer monsoon, and we will especially analyse the coastal ~~and central areas (orange and grey shadings on fig.~~ Arabian Sea (orange area on figure 2a).

A comparative global evaluation of the marine bio-geochemical component of the ESM has been published in ~~?~~ Séférian et al. (2013). Even if the model poorly represents the deep-ocean circulation, especially in the Southern Ocean, it has a quite good representation of annual wind patterns, wind stress, mixed-layer depth and geostrophic circulation. The model is able to represent the global ocean biological fields such as macro-nutrients, with correlations higher than 0.9, and surface chlorophyll concentration, with a correlation coefficient of 0.42 ~~(?)~~ (Séférian et al., 2013). We focus here on the representation of the physical processes and productivity distributions in the Indian Ocean, especially in the Arabian Sea. We use satellite products from remote sensing by NASA's Sea-viewing Wide-of-view Sensor (SeaWiFS) during the period 1998-2005 ~~for~~ processed with the VGPM algorithm (Behrenfeld and Falkowski, 1997) to obtain monthly productivity (Lévy et al., 2007) ~~and~~ the NOAA Multiple-Satellite 1995-2005 climatological cycle for wind intensity and wind stress (Zhang, 2006) and the ERA-interim re-analysis (1979-2014) for sea surface temperature (SST) (Dee et al., 2011). We compute the observed and ~~modeled~~ modeled wind stress curl intensity from the wind stress data and model output, respectively. We compare the observations to the pre-industrial (CTRL) simulation outputs.

Figure 3a shows that simulated boreal summer productivity integrated over the whole water column is underestimated relative to the reconstructed boreal summer productivity, especially in the regions of upwelling, along the coast of the ~~Arabie~~ Arabian Peninsula and Somalia. The spatial Pearson's correlation coefficient,  $R$ , between the observed and simulated productivity is 0.44. Underestimation of productivity is first caused by an underestimated wind intensity (Fig. 3b), which affects the  
5 extent and intensity of the coastal upwelling and the supply of nutrients to the surface layer. The boreal summer wind patterns, which are characteristic of the boreal summer monsoon system, are better represented than productivity with a correlation of 0.86. Sperber et al. (2013) studied the representation of the Asian summer monsoon in the CMIP5 models, which comprises the IPSL model. They showed that the monsoon was better represented in the CMIP5 models compared to the CMIP3 models, especially the monsoonal winds. We can however note that the alongshore winds in the western Arabian Sea have a more  
10 northerly orientation in the CTRL simulation than in the observations, which can affect the dynamical processes in the region (Fig. 3b).

In the Arabian Sea, summer productivity is affected by the winds through different mechanisms implying the wind stress and the wind stress curl (Anderson et al., 1992). The strong winds along the Arabian coast, called the Somali Jet, generate a positive wind stress which increases Ekman transport off the coast. The water that leaves the coastal area is being replaced  
15 by subsurface water: this is the coastal upwelling. Similarly to the wind intensity, the CTRL simulation wind stress intensity is underestimated compared to the ~~reconstructions-re-analyses~~ : the maximum wind stress intensity is lower and it does not extend far north in the Arabian Sea as the reconstructed wind stress (Fig. 3c). The wind stress orientation is also more zonal in the simulation than in the observations which causes the simulated wind stress to be higher close to the Oman coast relative to the observations (Fig. 3b,c). Figure 3d represents the wind stress curl, computed from the wind stress, in the simulation and  
20 in the observations. The simulated distribution resembles the reconstructed one: on the left hand side of the strong low-level wind jet, between the coast and the maximum wind intensity, the curl in the wind stress is positive and, on the other side of the jet, the wind stress curl is negative. The differences seen in the jet position and width are transmitted to the wind stress and wind stress curl intensity and distribution-, with an overall less positive curl close to the coast and less negative offshore in the simulation. On Figure 3e, we can see that the modeled SST anomalies, relative to the global averaged SST, in boreal  
25 summer, are underestimated in the Arabian Sea, especially close to the Oman coast suggesting a less intense upwelling activity compared to the observations. This is coherent with the underestimated wind stress and wind stress curl intensities, which control the upward Ekman pumping intensity (Fig. 3c,d).

Discrepancies between our pre-industrial simulation and the observations may be ~~attributed to the model coarse~~ due to the coarse model resolution. In Resplandy et al. (2011), a higher resolution version of the model was used to study the effects of  
30 meso-scale dynamics on productivity. They showed that the model is able to reproduce the observed meso-scale dynamics, such as the Great Whirl and filaments that transport nutrients from the coast to the open sea. They highlighted the major role of the eddy-driven transports in the establishment of biological blooms in the Arabian Sea and the model's ability to represent the different physical processes at stake behind productivity blooms in summer and in winter in the region. Nevertheless, even though both the winds and productivity are underestimated in CTRL by the lower resolution version of the model, the  
35 physical mechanisms playing a role in the marine productivity are represented, which is therefore adapted to our study. Figure

4 illustrates the combined effects of wind stress and wind stress curl on productivity in the coastal Arabian Sea (orange area on Figure 2a) at the inter-annual time-scale. It shows that if the summer (JJAS) wind stress and wind stress curl intensities are higher than their summer average, productivity is higher than average in coastal Arabian Sea (upper-right quadrant). It also highlights that the higher the wind stress and the wind stress curl anomalies, the higher the productivity change, and conversely.

5 This is coherent with the fact that both the wind stress and the wind stress curl act to bring nutrient rich waters to the surface and fuel productivity in boreal summer (Anderson et al., 1992; Murtugudde et al., 2007). Figure 4 also shows that a high wind stress curl (resp., wind stress) can compensate a reduced wind stress (resp., wind stress curl) intensity and lead to higher than average productivity (lower-right and upper-left quadrants of figure 4, respectively).

## 2.4 Diagnostics

10 In this section we briefly describe the variables and the methods we use throughout the paper. We are interested in the [pathways links](#) between the large-scale Indian summer monsoon system and the Arabian Sea primary productivity. To characterise the boreal summer monsoon intensity, we use the meridional gradient of upper tropospheric temperature (TT, averaged from 200 to 500 hPa) between a northern region covering India, south-east Asia and the Tibetan Plateau (60°E-120°E, 10°N-45°N) and a southern region over the tropical Indian Ocean (60°E-120°E, 25°S-10°N) (Marzin and Braconnot, 2009; Goswami

15 et al., 2006). This gradient,  $\Delta TT$ , is associated with the temperature land-sea contrast (Marzin and Braconnot, 2009).  $\Delta TT$  is averaged over the boreal summer period (JJAS for June-July-August-September) and the higher its value, the stronger the Indian summer monsoon.

Changes in the monsoon intensity and pattern affect the sea level pressure (SLP) field. We compute the SLP from the model outputs (i.e. model air temperature and pressure, within and between the atmospheric grid levels, and orography) using the

20 extrapolation described in Yessad (2016). We define "SLP anomalies" as the SLP minus the global annual average of SLP. In order to characterise the monsoon pattern, we compute the barycentre of the region defined by an SLP anomaly lower than -5 hPa over the region covering the African, East Asian and Indian monsoons regions of influence (20°W-150°E, 30°S-60°N) and we call "SLPa-5" the region delimited by the -5 hPa contour in SLP anomalies ~~-(Fig. 5c-f)~~. This SLPa-5 barycentre is representative of the balance between the different monsoons as well as of the Indian summer monsoon wind position and

25 direction over the Arabian Sea. A modification of the monsoon pattern, which can have impacts on productivity through the atmospheric forcing onto the ocean circulation, can then be related to movements of the SLPa-5 barycentre. We only focus on the Tibetan Low since the Mascarene High, the region of high SLP in the southern Indian Ocean, barycentre remains quite similar in the different simulations.

Anderson et al. (1992) showed that the wind stress intensity generates coastal upwelling and that the positive wind stress

30 curl is responsible for [upward](#) Ekman pumping offshore. ~~In the central Arabian Sea, the negative wind stress curl generates downwelling.~~ We focus our work on these two wind variables ~~and on two areas in the Arabian Sea: the coastal area in the in the coastal~~ western Arabian Sea, ~~which covers the a~~ region of positive curl, for the CTRL climate, between the axis of the Jet and the coast (Fig.2a, orange area) ~~and the area covering the central-east Arabian Sea (Fig. 2a, grey area).~~

In the following sections of the paper, total primary productivity (TPP) is defined as the sum of nano-phytoplankton and diatoms [total net primary](#) productivity integrated over the whole water column. We also analyse nitrate concentrations in the first 30 meters of the water column, nitrate being the major limiting [nutrients-nutrient](#) in the region and, its supply to the surface layers being mainly driven by atmospheric changes via coastal upwelling and [upward](#) Ekman pumping.

5 We use the CTRL simulation as a reference. All changes are then defined relative to this pre-industrial simulation.

### 3 Simulated paleoproductivity and monsoon changes

In this section, we investigate the changes in summer productivity in past climate simulations with respect to the CTRL simulation, starting with the early Holocene and then generalising to all the climates.

#### 3.1 The early Holocene case

10 The early Holocene (EH) experiences a stronger Indian summer monsoon than the pre-industrial (Fig. 5) therefore, we would expect higher productivity in the Arabian Sea. However, the EH simulation shows lower levels of productivity than in CTRL (Fig. 6). We explain this counter-intuitive result by a change in the monsoon pattern instead of a change in its intensity (Fig. 7).

The early Holocene, which we choose to represent with a snapshot at 9.5 kyr BP, is an interglacial period that mainly differs  
15 from the pre-industrial because of the imposed obliquity ( $24.2306^\circ$  vs  $22.391^\circ$ ) and precession ( $303.032^\circ$  vs  $102.7^\circ$ ) (Table 1). These changes in astronomical parameters cause the boreal summer insolation in the northern hemisphere to be higher than in the pre-industrial climate (Marzin and Braconnot, 2009). In [our simulation](#) EH, the boreal summer (JJAS) northern hemisphere (0-90°N, NH) mean insolation is  $20 \text{ W}\cdot\text{m}^{-2}$  higher than in CTRL (Fig. 5a,b). This change in insolation modifies the upper tropospheric temperature gradient,  $\Delta\text{TT}$ , represented on [Figure-figure 5c,d](#). ~~In~~ in EH,  $\Delta\text{TT}$  is 1 K higher than in CTRL  
20 supporting a stronger monsoon intensity consistent with previous studies (Marzin and Braconnot, 2009; Goswami et al., 2006).

The large-scale spatial pattern of the summer monsoon is also different in EH compared to CTRL. Maps e) and f) on [Figure-figure 5](#) show the SLP anomalies and the location of the SLPa-5 barycentre. The EH depression extends further to the north-west and onto Africa and ~~onto~~ the Arabic peninsula than in CTRL and, the EH minimum SLP anomaly is moved to the north-west compared to the CTRL one (Fig. 5e,f). The SLPa-5 barycentre, which is representative of the balance between the  
25 different monsoons and of the Somali Jet position and direction, moves to the north-west in EH relative to CTRL (Fig. 5f). This suggests a modification of the monsoon structure with potential impacts on productivity through the atmospheric forcing onto the ocean.

Observation-based (Bauer et al., 1991; Prasanna Kumar et al., 2000) and model-based (Murtugudde et al., 2007) studies  
30 have shown that a stronger monsoon, with increased wind strength over the Arabian Sea, leads to higher productivity through intensified supply of nutrients in the photic zone. EH's Indian summer monsoon is enhanced compared to CTRL. One would therefore expect the EH Arabian Sea productivity to be higher in EH than in CTRL. However, our model shows that the EH productivity is reduced in the Arabian Sea (Fig. 6a).

In this region, productivity is mainly nutrient-limited (Koné et al., 2009; Prasanna Kumar et al., 2000) and the levels of surface  $NO_3^-$  concentration concentrations are lower for EH than for CTRL (Fig. 6b). Most of the nutrients are provided to the surface layer via Ekman dynamics: coastal upwelling, upward Ekman pumping and mixing of the upper layers close to the coast and offshore and, mixing and advection processes further offshore

5 ~~in central Arabian Sea (Resplandy et al., 2011; Murtugudde et al., 2007; Prasanna Kumar et al., 2000; Lee et al., 2000; Bauer et al., 1991; Figure 6e shows that wind stress intensity increases close to the Omani coast and decreases offshore. This leads to less mixing in the central Arabian Sea because of a shallower mixed layer (ML) (Fig.6d). The ML is deeper in EH than in CTRL close to the Oman coast~~  
~~(Resplandy et al., 2011; Murtugudde et al., 2007; Prasanna Kumar et al., 2000; Lee et al., 2000; Bauer et al., 1991; McCreary et al., 2009)~~  
10 ~~The shift of the SLPa-5 barycenter (Fig.6d), which suggests a reduction in the upwelling intensity in EH. Wind stress intensity changes along with ML changes and the wind stress distribution of the CTRL simulation (Fig.3c) imply that the maximum wind intensity has moved closer to the Oman coast in the EH. This displacement of the jet reduces the wind stress intensity in the central sea. Moreover, the stronger EH wind stress very close to the coast 5f) leads to a poleward and westward shift of the monsoon jet (Fig. 6c) has a reduced effect on the ocean as it affects a smaller part of the coastal ocean. The reduction in~~  
15 ~~coastal upwelling and mixing then contribute to the decrease in nutrients concentration in the upper layer. This causes weaker alongshore winds in the Somalia upwelling and stronger alongshore winds in the Oman upwelling with visible effects on the mixed layer depth (Fig.6b). The positive wind stress curl on the left hand side of the jet has also a reduced influence onto the ocean (Fig.6e). The negative wind stress curl on the other side of the wind jet, affects a region closer to the coast leading to reduced Ekman pumping 6c,d). But this also brings the Ekman downwelling to the right of the jet axis closer to both Oman~~  
20 ~~and Somalia coasts (Fig.6f). In summary, the EH atmospheric forcing through wind stress and wind stress curl is not able to generate as much coastal upwelling and Ekman pumping as in the CTRL simulation, 6e). Both factors (alongshore stress and offshore curl) contribute to a Somalia upwelling reduction, while the wind stress curl change seems to overwhelm the increased alongshore winds in the Oman region, leading to overall upwelling reduction and less nutrients even though the monsoon is stronger more intense (Fig. 6b,f).~~

25 On ~~Figure figure~~ 7, we represented the 100 summers of the EH simulation minus the seasonal summer mean of the CTRL simulation for the variables TPP, wind stress and wind stress curl in the coastal ~~and central~~ Arabian Sea (~~regions on Figure see~~ Fig. 2a). In the coastal area, even though wind stress is always higher in EH (y-axis), productivity is always lower in EH than in CTRL (Fig. 7a). This highlights the role of the wind stress curl that is always ~~lower less positive~~ in EH than in CTRL (x-axis) : a ~~smaller less positive~~ wind stress curl is responsible for lower productivity levels in the coastal western Arabian Sea. The  
30 wind stress intensity also affects the intensity of the changes in TPP: the stronger the wind stress the smaller the ~~change in TPP~~. ~~Wind reduction of TPP. Increased wind~~ stress can oppose the negative effect of reduced wind stress curl, but not overcome it as it is restricted very close to the coast (Fig. 6g).

~~In the central-east Arabian Sea, productivity is also lower in EH than in CTRL. The wind stress and the wind stress curl intensities are reduced almost every year compared to the CTRL seasonal summer mean (Fig.7b). This causes an increased~~

~~downwelling and less wind mixing of the upper layers, which leads to reduced entrainment of nutrients and less productivity than in CTRL.~~

In summary, in the EH simulation the summer monsoon intensity is stronger than in the CTRL simulation but the productivity in the Arabian Sea is lower. This is caused by a shift in the Somali Jet position, which reduces coastal upwelling and upward Ekman pumping. This shift in the maximum wind intensity position closer to the coast can be inferred from the north-western movement of the SLPa-5 barycentre that translates into a modification of the monsoon pattern (Fig. 5).

### 3.2 Generalisation

In the previous section, we saw that a stronger monsoon in the EH does not imply more productivity in the Arabian Sea and that it is important to consider the spatial movements of the monsoonal winds. We now examine the links between productivity, monsoon intensity and boundary conditions in the remaining set of 6 glacial and inter-glacial simulations.

Figure 8 shows the changes in productivity, in the Arabian Sea, in all the remaining climates compared to the CTRL climate. Similar to the EH results, the MH, LGM, MIS4M and MIS4D coastal productivities are reduced (Fig. 8a-b,d-f). MH and MIS4F coastal productivity are reduced in average but present a dipole-like pattern in the western coastal Arabian Sea, with higher productivity in the north and reduced productivity in the south compared to CTRL (Fig. 8a,d). Coastal productivity in the western Arabian Sea is enhanced in the MIS3 simulation (Fig. 8c). ~~The central Arabian Sea Along with these TPP changes, Ekman pumpind changes are represented on figure 8 in black contour. In the simulations where coastal productivity is higher than CTRL in all glacial simulations except MIS4F while it is reduced in MH (and EH as seen previously)-~~

in CTRL, upward Ekman pumping also increases close to the coast (Fig. 8a,c,d) and inversely (Fig. 8a,b,e,f). Ekman pumping patterns suggest that the wind orientation has changed throughout the different time periods, affecting differently the supply of nutrients and then productivity.

The tropospheric temperature gradient ( $\Delta TT$ ) for each simulation, on Figure-figure 9a, informs that the Indian summer monsoon intensity is stronger ~~than CTRL~~ in MH, EH, MIS3 and MIS4F (i.e. higher  $\Delta TT$  values) and less intense compared to CTRL in LGM, MIS4M and MIS4D. The changes in productivity for the western coastal Arabian Sea are also summarised on figure 9b. By only looking at these two variables,  $\Delta TT$  and TPP, we cannot conclude on a direct link between monsoon intensity and productivity because stronger monsoons compared to CTRL, as characterised by  $\Delta TT$ , do not necessarily imply higher productivity (Fig. 9a,b), in particular for MH and EH.

#### 3.2.1 Productivity and local dynamics

Productivity is nutrient-limited in the region and coastal productivity changes are similar to the changes in the nitrate content of the upper 30 m of the ocean (Fig. 9b,c). When the upper layer receives more nutrients from the subsurface, there is either a stronger upwelling or a higher ~~macro-nutrients~~ macro-nutrient ( $NO_3^-$  and  $PO_4^{3-}$ ) concentration under the mixed layer associated with enhanced entrainment ~~and, productivity is higher than in CTRL.~~ The stronger monsoon intensity, characterised by a higher  $\Delta TT$  value, is associated with higher values of coastal wind stress (Fig. 9a,d). But changes in wind stress curl are independent of the monsoon intensity since it is lower than CTRL in MH, EH and MIS4F and higher-more positive in

all the other glacial simulations (Fig. 9e). Wind stress curl intensity is also ~~higher~~ more positive in all the glacial climates compared to the Holocene. In MH, EH and MIS4F, even though wind stress intensity is stronger and should generate more coastal upwelling, the highly reduced wind stress curl overcomes this positive effect on productivity and induces lower levels of macro-nutrients, which in turn limit productivity (Fig. 9b-e). Conversely, in LGM, MIS4M and MIS4D, even though the  
5 wind stress curl is ~~higher~~ more positive than in CTRL, the lower wind stress intensity seems to prevail and productivity is reduced because of lower concentrations of nutrients (Fig. 9b-e). In MIS3, both the wind stress and the wind stress curl are more ~~intense~~ positive, more nutrients are brought to the surface and productivity increases (Fig. 9b-e).

Figure 10a summarises the links between productivity, wind stress and wind stress curl intensities in the coastal Arabian Sea. It shows that changes in wind stress and wind stress curl are drivers of productivity changes. Both variables modulate the  
10 change in productivity, with higher values associated with higher productivity.

~~In the central Arabian Sea, the enhanced productivity appears to result from more intense wind stress, which deepens the mixed layer and brings more nutrients to the upper layer through wind mixing and advection, consistent with Murtugudde et al. (2007), Wiggert et al. (2005) and Bauer et al. (1991). These results are summarised in Figure 10b, where the summer productivity changes are plotted as function of wind stress and wind stress curl summer changes in the central Arabian Sea. When the wind stress intensity change is higher than in CTRL and wind stress curl not too low, productivity is enhanced (LGM, MIS4m and MIS4d)(Fig. 10b). If the wind stress is high enough it can compensate for a more reduced wind stress curl (MIS3) otherwise, productivity is lower (MIS4f, EH and MH)(Fig. 10b). Surprisingly, the simulations with a stronger summer monsoon do not systematically have stronger wind stress intensity over the central Arabian Sea but they have lower wind stress curl. This is the result of a change in the monsoon pattern, like a shift in the position or the orientation of the maximum wind intensity.~~

### 20 3.2.2 Relation to the large-scale forcing and boundary conditions

In order to understand how changes in the monsoon pattern are linked to the imposed boundary conditions and influence productivity and local monsoonal changes, we use the SLPa-5 barycentre. The position of the barycentre of each simulation is plotted on a map on figure 11a. We also added on ~~Figure~~ figure 11, the mean boreal summer values (color-scales) of productivity, wind stress and wind stress curl as a function of the longitude (x-axis) and the latitude (y-axis) of the SLPa-5  
25 barycentre.

Productivity shows an increasing trend with the longitude of the SLPa-5 barycentre between 70°E and 94°E (Fig.11b). It reaches a maximum of 30 molC.m<sup>-2</sup>.yr<sup>-1</sup> around 94°E. For higher values of longitude, which correspond to simulations where the monsoon intensity is reduced, productivity decreases (Fig. 11b). The higher values of productivity occur in the simulations for which the SLPa-5 barycentre's longitude and latitude have medium values (MH, CTRL, MIS4F and  
30 MIS3)(Fig. 11b). The trends in productivity can be explained by the variations of wind stress (Fig. 11c) and wind stress curl (Fig. 11d) with the SLPa-5 barycentre's position.

Wind stress exhibits ~~a global~~ an overall decrease with the longitude and latitude of the SLPa-5 barycentre as it moves southeastward (Fig. 11c). If we except the CTRL simulation, wind stress is quite constant for the 5 first simulations (i.e. lower value of longitude) and then it decreases strongly for MIS4M, MIS4D and LGM. ~~The wind stress shows a clear separation~~



between simulations in which the monsoon is enhanced and wind stress is higher (EH, MH, MIS4F and MIS3) and those in which monsoon is less intense and wind stress is reduced (i.e. LGM, MIS4D, MIS4M and CTRL). The latitude of the barycentre exerts a strong control on the coastal wind stress amplitude (Fig. 11c). Wind stress curl shows an increasing trend with longitude for the 5 simulations having lower values of longitude and then the wind stress curl becomes quite constant for MIS4M, MIS4D and LGM (Fig. 11d). The wind stress curl has also a tendency to decrease with the SLPa-5 barycentre's latitude (Fig. 11d).

The increase of productivity with longitude is mostly due to an increase in wind stress curl intensity and ~~then~~ the reduction of productivity with higher longitude is caused by a strong reduction in wind stress while the wind stress curl remains constant (Fig. 11b-d). These plots also show that all the glacial simulations have higher values of SLPa-5 barycentre longitude compared to CTRL. This ~~suggests~~ demonstrates a major role of the ice sheet cover over the longitudinal position of the SLPa-5 barycentre. The simulations having a stronger monsoon intensity than CTRL have the highest value of SLPa-5 barycentre latitude, which suggests an influence of the ~~orbital~~ astronomical parameters on the latitudinal position of the SLPa-5 barycentre. This latitudinal movement of the SLPa-5 barycentre with the monsoon strength seems also to be dependent on the glacial or interglacial state of the simulation. Indeed, in glacial simulations the SLPa-5 barycentre is north of the CTRL barycentre even if the monsoons are less intense (LGM, MIS4D and MIS4M), however the other glacial simulations with higher summer monsoon intensity (MIS3 and MIS4F) have their barycentre located north of these glacial simulations (Fig. 11). Similarly, in inter-glacial climates, the Holocene simulations have a barycentre north of the CTRL one (Fig. 11).

On figure 12, we plotted the values of the climatic precession ~~and obliquity~~ (defined as  $e \times \sin(\omega - 180^\circ)$  with  $e$  the eccentricity and  $\omega$  the precession), which modulates the northern hemisphere insolation, and obliquity, which controls the temperature contrast between the high and low latitudes, relative to the SLPa-5 barycentre position, in order to analyse the relationship between the ~~orbital~~ astronomical parameters and the SLPa-5 barycentre's position. Climatic precession influences the SLPa-5 barycentre position ~~in longitude~~ both in longitude and latitude: when the climatic precession is high, the barycentre tends to move to the south-east (Fig. 12a). Obliquity modulates the latitudinal changes of the SLPa-5 barycentre: high obliquities are associated with a SLPa-5 barycentre farther north (Fig. 12b). MIS3 has a climatic precession value similar to CTRL and a much higher obliquity than CTRL, so the changes in MIS3 winds and productivity related to insolation are mostly obliquity-driven (Fig. 12). Inversely, MIS4F has a similar obliquity as CTRL and a smaller climatic precession which implies that the changes in monsoon intensity in MIS4F are related to precession (Fig. 12). The Holocene simulations are influenced by both obliquity and precession while, the LGM, MIS4M and MIS4D seem to reflect a stronger link with the obliquity signal than with the climatic precession (Fig. 12).

## 4 Discussion

### 4.1 The summer monsoon paradigm

In the simulations, the general paradigm stating that a stronger summer monsoon intensity induces a stronger upwelling and therefore increases marine productivity, is not always verified. Our results show that the **only**-characterisation of the summer monsoon intensity is probably insufficient to assess past productivity changes and reciprocally (Fig. 9).

Our results for the summer productivity are consistent with the reconstructed productivity of Rostek et al. (1997). In their study, they analyse two marine sediment cores in the Arabian Sea: one in the south-east (5°04' N - 73°52' E) and one in the upwelling region close to the Oman coast (13°42' N - 53°15' E). They show that paleo-productivity in the south-eastern core was higher in glacial stages than in interglacial stages, which they interpret as the fingerprint of a stronger winter monsoon. In the other core, the productivity signal is more complex and they could find some glacial stages (e.g. stage 2) with high productivity, some interglacial stages with low productivity (e.g. stage 1) and high productivity during stage 3. Similarly, in the simulations, in the central Arabian Sea, glacial productivity is higher than interglacial productivity (except for MIS4F) (Fig. 40b 8). In the western coastal Arabian Sea, the simulated MIS3 productivity is higher than CTRL while the other climates' productivity is lower than CTRL. Hints on the sources of discrepancies between our results and Rostek et al. (1997) results and their interpretation of the productivity changes are given later in this section.

We explain the simulations' summer productivity changes by analysing the variations in the wind forcing (Fig. 9). Given the productivity changes in the different simulations, the summer monsoon intensity only is not able to explain the changes in productivity and therefore, we also investigate the changes the monsoon pattern (Fig. 11).

The large-scale definition of the monsoon intensity, via  $\Delta TT$ , is mainly driven by **orbital-astronomical** changes. The simulations with a strong summer monsoon either have a high obliquity, which enhances the temperature contrast between low and high latitudes in summer (e.g. MIS3) or a small climatic precession that intensifies the summer insolation (MIS4F), or both (MH and EH) (Fig. 42) (?) 12) (Prell and Kutzbach, 1987). Simulations with a weak summer monsoon all have a small obliquity forcing (Fig. 12b). The local wind stress that affects productivity is tightly coupled to the monsoon intensity (Fig. 9a,b) and is associated with a SLPa-5 barycentre movement to the North (Fig. 11c). This simulated latitudinal movement of the SLPa-5 barycentre, according to the monsoon intensity, is consistent with the study of Anderson and Prell (1992). In this study, the authors show that with a stronger monsoon the Somali Jet is moved further North, which would indeed translate into a northward movement of the SLP barycentre (and inversely). In Fleitmann et al. (2007), the authors investigate the variations in the precipitation records of stalagmites located in Oman and Yemen. They link the changes in precipitation to the position and structure of the ITCZ, which affects the tropical climate. They explain that during the early-Holocene, a northward movement of the mean latitudinal position of the summer ITCZ is responsible for the decrease in precipitation. Throughout the Holocene, they show that the ITCZ shifted southward concomitantly with a decrease in the monsoon precipitation, induced by the reduction of **solar-northern hemisphere summer** insolation (Fleitmann et al., 2007). The changes in the ITCZ position highlighted by Fleitmann et al. (2007) are consistent with our results, especially with the changes in the position of the SLPa-5

barycentre in latitude ~~given a similar background glacial or interglacial state~~ given a similar glacial or interglacial background state (Fig. 11).

Marine productivity is not only influenced by the wind stress but also by the wind stress curl (Anderson et al., 1992) and the latter is also strongly influenced, in our simulations, by the glacial or interglacial state of the climate (Fig. 11). The glacial-  
5 interglacial distribution of the wind stress curl is associated with a longitudinal movement of the SLPa-5 barycentre : the SLPa-5 barycentre is moved to the East in glacial climates and to the West in interglacial climates (Fig. 11d). Pausata et al. (2011) ~~analyse~~, who analysed the effects of different LGM boundary conditions on the atmospheric circulation ~~and~~, indeed found that ice sheet topography is responsible for changes in many features of the SLP field, e.g. position of lows and highs and their variability. ~~Consequently, changes in the ice-sheet cover between our simulations can indeed be responsible for the~~  
10 ~~longitudinal movement of the SLPa-5 barycentre~~. Ivanova et al. (2003) also evoke an eastward shift in the low-level jet position in summer as a possible mechanism to explain some productivity changes in the eastern Arabian Sea. Furthermore, we showed that climatic precession can also act to move the SLPa-5 barycentre eastward (Fig. 12a) and therefore affect wind stress curl and productivity.

15 We find a valid physical explanation for our simulations' productivity changes through the effects of the simulations' boundary conditions (~~orbital~~ astronomical parameters and ice sheets) on the monsoon intensity and pattern. However, even if the simulated summer productivity compares quite well to the data in Rostek et al. (1997) in the southeastern Arabian Sea (except for ~~the LGM MIS4F~~), they ~~do not match other reconstructions such as~~ are more discrepancies with their second core in the western coastal Arabian Sea (core in the southern part of the our coastal area), and also when compared to the reconstructions  
20 in Bassinot et al. (2011) (e.g. the Holocene productivity core in the northern part of our the coastal area) -(Fig. 10). These differences can arise from several sources, the first one being linked to the area on which we computed our averages (Fig. 2a). Indeed, in Rostek et al. (1997) the core in the upwelling region is taken in the southern part of our coastal area (Fig.2a) ~~whereas~~ and the core close to the Oman coast in Bassinot et al. (2011) is located in the northern part of our coastal area (Fig. 2a). Bassinot et al. (2011) reconstructed productivity is high in the early-Holocene and gradually decreases throughout the Holocene whereas, in  
25 our simulations, the Holocene productivity is low compared to the pre-industrial productivity. A closer look at the productivity changes in the MH simulation on figure 8a can reconcile our simulations and this reconstructed productivity. In the simulated MH climate, the northern part of the coastal area exhibits a positive productivity change whereas, the southern part of the coastal area is characterised by a negative productivity change compared to CTRL ~~-(Fig. 8a)~~. Therefore, our results are coherent with the results of Bassinot et al. (2011) for the mid-Holocene since their core is located in the northern part of our  
30 coastal area where we simulate higher MH productivity than CTRL. However, we do not observe this dipole-like pattern in the EH productivity (Fig. 6a) and consequently, the EH simulation does not agree with this reconstruction. In the EH simulation, we do not have the remnant Laurentide ice sheet that is supposed to be present at this time-period (Licciardi et al., 1998). The simulation with this model version was not available at the time of our analyses. The addition of a remnant ice-sheet over Europe and North America in the EH has been shown, in Marzin et al. (2013), to induce a southward shift of the Inter-Tropical  
35 Convergence Zone (ITCZ) and a strengthening of the Indian monsoon. A southward shift of the jets would modify the large

scale pattern of the SLP and therefore, the wind stress and wind stress curl effects on the Arabian Sea. Based on our findings, the addition of this residual ice sheet would move the SLPa-5 barycentre to the south-east, which could increase the wind stress curl and therefore productivity (Fig. 11).

Another source of mismatch between our simulations and data resides in the fact that we looked at productivity and not at the export production that will eventually reach the bottom of the ocean. We also focused ~~on~~ on the boreal summer season while it is often advanced that the winter monsoon is responsible for higher productivity in glacial climates compared to interglacial climates, e.g. for the south-eastern core productivity in Rostek et al. (1997). It is likely that the boreal winter productivity may also affect the overall recorded signal in the sediments. In the next section, we especially discuss the effect of seasonality on productivity.

## 10 4.2 Seasonality

Here, we investigate new paleo-productivity reconstructions for the Arabian Sea and we compare them to our simulations. The simulations ~~globally~~ largely agree with the reconstructions, with glacial productivities higher than Holocene productivities in north-western Arabian Sea, even during boreal summer.

We use data from the sediment core MD04-2873 located at 23°32N-63°~~50W~~ 50E in northern Arabian Sea on the Murray Ridge (Böning and Bard, 2009) ~~-(Figures. 2a and 8, yellow circle).~~ This core is well dated by C14 dates from 50 kyr to present and has a marked Toba Ash layer (74 kyr BP, (Storey et al., 2012)) giving a significant robust stratigraphic marker. The coccolithophores are well preserved and abundant at this location. Their assemblages are used to reconstruct paleo-productivity by using a transfer function that has been designed for the Indian Ocean including the Arabian Sea (Beaufort et al., 1997). Samples have been prepared by settling onto cover-slips (Beaufort et al., 2014) every 10 cm for stratigraphic intervals covering 2000 years above and below each time period simulated by the model. In average 6 samples were studied by time intervals. Coccolithophore analysis has been automatically generated by a software, SYRACO, that has been trained to recognise coccolithophores (Beaufort and Dollfus, 2004). Figure 13a shows the resulting paleo-productivity for 7 of the 8 time periods we previously analysed. This reconstruction indicates that glacial productivity is higher than Holocene productivity in this core. At the core location, the effect of the boreal winter monsoon on productivity ~~are~~ is known to be strong (Lévy et al., 2007). Consequently, the stronger winter monsoons during glacial time periods are often used to explain how glacial productivity can be higher than interglacial productivity (e.g. Rostek et al. (1997); Banakar et al. (2005)).

On figure 13, we also plotted the box-plots of simulated annual and summer (JJAS) productivity and export production at 100 m for the same time periods in the northern Arabian Sea (60°E-68°E, 20°~~N-68N~~ N-28°N). Our simulated annual and boreal summer productivity show smaller differences between glacial and interglacial climates than the reconstructions (Fig. 13a-c). The export production shows a clearer separation between the glacial and interglacial climates in both the summer and annual plots (Fig. 13d,e). The differences between productivity and export production (~~fig-Fig.~~ 13b-e) highlight that water column processes modify the recorded signal, which add difficulties when comparing model to data.

In all simulated climates, the mean boreal summer productivity is lower than the mean annual productivity (~~Fig-Figs.~~ 8 and 13). This indicates that, in this region, in the simulations, the mean boreal winter productivity is higher than the mean boreal

summer productivity. Indeed, in the simulations, in the region of the core, boreal winter productivity accounts for more than 40% of the annual productivity and boreal summer productivity accounts for more than 20% (not shown). The hypothesis, stating that glacial productivity is higher than interglacial productivity because of a stronger winter monsoon, could explain the variations in this core ~~-(e.g. Banakar et al. (2005))~~. However, the observed present-day seasonal cycle of productivity in the northern Arabian Sea shows equal contributions of the winter and summer seasons to the annual productivity (Fig. 2b), suggesting that the simulations may underestimate the boreal summer contribution to productivity compared to the boreal winter contribution.

Interestingly, the annual and boreal summer productivity plots look alike, with higher glacial than interglacial productivity in the model (Fig. 13). Since the summer monsoon is able to affect the north-western Arabian Sea, as seen on figure 2a 2b, it contributes to the recorded signal in the sediment (e.g. Caley et al. (2011)). The boreal summer monsoon effect on the recorded signal is then non-negligible and, we see that, even during the boreal summer season, the simulations show higher glacial productivity than interglacial productivity (Fig. 10 and 13), especially ~~in~~ as well as in the central Arabian Sea (Fig. 8). Consequently, the boreal winter productivity is not the sole contributor to the higher glacial productivity signal compared to the interglacial productivity in the model, even in the northern Arabian Sea.

## 5 Summary and perspectives

We use the coupled IPSL-CM5A-LR model to study the Arabian Sea paleo-productivity in 8 different climates of the past. We focus on the processes behind the boreal summer productivity changes in the coastal western ~~and central-eastern~~ Arabian Sea. We show that a stronger Indian summer monsoon, which is mostly driven by higher NH insolation, does not necessarily enhance the Arabian Sea productivity, and conversely.

We show that glacial climates can be more productive, in boreal summer, in the Arabian Sea (~~coastal and/or central~~), compared to the pre-industrial (Fig. 8 and 10). Even more, the glacial climates are more productive than the early Holocene, which was supposed to be the most productive period in the region (Figs. 8 and 13). We found that the paradigm between monsoon intensity and productivity is valid for MIS3, ~~both in the coastal and central~~ in the western coastal sea: a stronger monsoon leads to more productivity. The paradigm is also valid in the coastal Arabian Sea for the LGM, MIS4M and MIS4D simulations: a reduced monsoon intensity leads to a reduction in productivity. However it is not the case for MH, EH and MIS4F simulations for which a stronger summer monsoon is associated with reduced productivity, ~~both in the coastal and central Arabian Sea. Moreover, the LGM, MIS4M and MIS4D simulations that have less intense summer monsoons have higher productivity levels in the central Arabian Sea.~~

Our analyses highlight the importance of considering the monsoon pattern, especially the position of the maximum wind intensity over the Arabian Sea. The mechanisms behind productivity changes are summarised on figure 14. We examine the monsoon pattern through the SLP barycentre position of the depression covering the summer monsoons regions (SLPa-5 barycentre). The SLPa-5 barycentre is moved to the East in glacial climates and far North in climates where monsoon is enhanced (Fig. 11), which highlights the influence of the ice sheet cover (Pausata et al., 2011) and of the orbital-astronomical

parameters (Anderson and Prell, 1992). The monsoon pattern affects the wind stress and wind stress curl efficiency to bring more or less nutrients to the surface layers. A change in the pattern can reduce or increase the area on which the winds are effective. This study also highlights the combined effects of wind stress and wind stress curl related processes on productivity. Neither wind intensity nor wind stress curl alone can explain productivity changes.

5 We need to keep in mind that the model's coarse resolution does not allow for a very precise representation of the region dynamics. This may have altered the relative weight of the processes related to wind stress and wind stress curl and can explain why the ~~orbital-astronomical~~ signal is weak in the productivity changes. Arabian Sea productivity in the CTRL simulation shows quite large differences with observations, especially for the high coastal productivity extension in the western Arabian Sea (Fig. 3a). This can result from the model coarse resolution which prevents the representation of meso-scale processes such as eddies. These fine-scale processes are shown to be of importance for the coupling between biology and physics (Resplandy et al., 2011). Moreover, these meso-scale processes contribute strongly to the export of nutrients offshore and thus can explain why our high productivity area is more restricted to the coast than in the observations. From our set of simulations we cannot assess the effect of the underestimation of productivity on our results. Our simulations may underestimate productivity levels and variations but since most of the main physical processes are represented we can draw

10 conclusions on the link between these processes and productivity and their changes through time. A way to overcome these limitations and to quantify their effects would be to work with several other models and to analyse the coupling between biology and physics in those different models.

15

We demonstrated that both changes in wind stress and wind stress curl can affect productivity at the time-scales of thousands of years (Fig. 10). The same effects of wind stress and wind stress curl changes on productivity can be found at the inter-annual time-scale (Fig. 4). ~~Figure 4 illustrates the combined effects of wind stress and wind stress curl on productivity in the coastal Arabian Sea at the inter-annual time-scale. It shows that if the summer (JJAS) wind stress and wind stress curl intensities are higher than their summer average, productivity is higher than average in coastal Arabian Sea (upper-right quadrant). It also highlights that the higher the wind stress and the wind stress curl anomalies, the higher the productivity change, and conversely. Figure 4 also shows that a high wind stress curl (resp., wind stress) can compensate a reduced wind stress (resp., wind stress curl) intensity and lead to higher than average productivity (lower-right and upper-left quadrants of figure 4, respectively).~~ The relationship between changes in stress or curl and productivity is similar to the one we found for the glacial-interglacial climate changes (Figs. 4 and 10). It could be interesting to further investigate these relationships by looking at high resolution models and re-analyses.

20

25

This study allows us to draw attention ~~on to~~ certain points that may affect the reconstruction of past climate ~~climate~~ and productivity as well as the comparison between model and data. In addition, in regards of projected changes in the monsoon intensity and structure, these results can add some constraints on future productivity changes in the region. In chapter 14 of the 2013 IPCC report (Christensen et al., 2013), it has been shown, through the use of climate projections, that the future Indian summer monsoon ~~should is expected to~~ strengthen in regards of precipitation but become less intense in regards of the monsoon flow. Moreover, Sandeep and Ajayamohan (2014) have shown that the projected low-level jet over the Arabian Sea will shift north because of global warming. A northward shift of the low-level jet is consistent with an increased monsoon intensity in

30

35

our simulations. Then, if a stronger summer monsoon calls for increased productivity, a northward shift of the Somali Jet can either lead to reduced productivity, as in the Holocene and MIS4F simulations, or to an increased productivity as in the MIS3 simulation, depending on the degree of the shift and on the wind stress curl change.

*Acknowledgements.* Priscilla Le Mézo ~~is~~was funded by a grant from the Initiative D'EXcellence (IDEX) Paris-Saclay. This work was supported by the French ANR Project ELPASO (No.2010 BLANC 608 01), the computing time was provided by GENCI (Grand Equipement National de Calcul Intensif) and the simulations were performed on Curie at TGCC (CEA, France).

## References

- Abe-Ouchi, A., Saito, F., Kageyama, M., Braconnot, P., Harrison, S. P., Lambeck, K., Otto-Bliesner, B. L., Peltier, W. R., Tarasov, L., Peterschmitt, J.-Y., and Takahashi, K.: Ice-sheet configuration in the CMIP5/PMIP3 Last Glacial Maximum experiments, *Geoscientific Model Development Discussions*, 8, 4293–4336, doi:10.5194/gmdd-8-4293-2015, <http://www.geosci-model-dev-discuss.net/8/4293/2015/>, 2015.
- Anderson, D. M. and Prell, W. L.: The structure of the southwest monsoon winds over the Arabian Sea during the late Quaternary: Observations, simulations, and marine geologic evidence, *Journal of Geophysical Research*, 97, 15 481, doi:10.1029/92JC01428, <http://doi.wiley.com/10.1029/92JC01428>, 1992.
- Anderson, D. M., Brock, J. C., and Prell, W. L.: Physical upwelling processes, upper ocean environment and the sediment record of the southwest monsoon, *Geological Society, London, Special Publications*, 64, 121–129, doi:10.1144/GSL.SP.1992.064.01.08, <http://sp.lyellcollection.org/cgi/doi/10.1144/GSL.SP.1992.064.01.08>, 1992.
- Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global Biogeochemical Cycles*, 20, doi:10.1029/2005GB002591, <http://doi.wiley.com/10.1029/2005GB002591>, 2006.
- Banakar, V. K., Oba, T., Chodankar, A. R., Kuramoto, T., Yamamoto, M., and Minagawa, M.: Monsoon related changes in sea surface productivity and water column denitrification in the Eastern Arabian Sea during the last glacial cycle, *Marine Geology*, 219, 99–108, doi:10.1016/j.margeo.2005.05.004, <http://linkinghub.elsevier.com/retrieve/pii/S0025322705001544>, 2005.
- Barber, R. T., Marra, J., Bidigare, R. C., Codispoti, L. a., Halpern, D., Johnson, Z., Latasa, M., Goericke, R., and Smith, S. L.: Primary productivity and its regulation in the Arabian Sea during 1995, *Deep Sea Research Part II: Topical Studies in Oceanography*, 48, 1127–1172, doi:10.1016/S0967-0645(00)00134-X, <http://linkinghub.elsevier.com/retrieve/pii/S096706450000134X>, 2001.
- Bassinot, F. C., Marzin, C., Braconnot, P., Marti, O., Mathien-Blard, E., Lombard, F., and Bopp, L.: Holocene evolution of summer winds and marine productivity in the tropical Indian Ocean in response to insolation forcing: data-model comparison, *Climate of the Past*, 7, 815–829, doi:10.5194/cp-7-815-2011, <http://www.clim-past.net/7/815/2011/>, 2011.
- Bauer, S., Hitchcock, G. L., and Olson, D. B.: Influence of monsoonally-forced Ekman dynamics upon surface layer depth and plankton biomass distribution in the Arabian Sea, *Deep Sea Research Part A. Oceanographic Research Papers*, 38, 531–553, doi:10.1016/0198-0149(91)90062-K, <http://linkinghub.elsevier.com/retrieve/pii/019801499190062K>, 1991.
- Beaufort, L. and Dollfus, D.: Automatic recognition of coccoliths by dynamical neural networks, *Marine Micropaleontology*, 51, 57–73, doi:10.1016/j.marmicro.2003.09.003, 2004.
- Beaufort, L., Lancelot, Camberlin, Cayre, Vincent, Bassinot, and Labeyrie: Insolation cycles as a major control of equatorial indian ocean primary production, *Science (New York, N.Y.)*, 278, 1451–4, <http://www.ncbi.nlm.nih.gov/pubmed/9367955>, 1997.
- Beaufort, L., Barbarin, N., and Gally, Y.: Optical measurements to determine the thickness of calcite crystals and the mass of thin carbonate particles such as coccoliths., *Nature protocols*, 9, 633–42, doi:10.1038/nprot.2014.028, <http://www.ncbi.nlm.nih.gov/pubmed/24556786>, 2014.
- Behrenfeld, M. J. and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concentration, *Limnology and Oceanography*, 42, 1–20, doi:10.4319/lo.1997.42.1.0001, 1997.
- Böning, P. and Bard, E.: Millennial/centennial-scale thermocline ventilation changes in the Indian Ocean as reflected by aragonite preservation and geochemical variations in Arabian Sea sediments, *Geochimica et Cosmochimica Acta*, 73, 6771–6788, doi:10.1016/j.gca.2009.08.028, <http://linkinghub.elsevier.com/retrieve/pii/S0016703709005365>, 2009.



- Bopp, L., Kohfeld, K. E., Le Quéré, C., and Aumont, O.: Dust impact on marine biota and atmospheric CO<sub>2</sub> during glacial periods, *Paleoceanography*, 18, n/a–n/a, doi:10.1029/2002PA000810, <http://doi.wiley.com/10.1029/2002PA000810>, 2003.
- Braconnot, P., Marzin, C., Grégoire, L., Mosquet, E., and Marti, O.: Monsoon response to changes in Earth's orbital parameters: comparisons between simulations of the Eemian and of the Holocene, *Climate of the Past Discussions*, 4, 459–493, doi:10.5194/cpd-4-459-2008, <http://hal.archives-ouvertes.fr/hal-00330744/http://www.clim-past-discuss.net/4/459/2008/>, 2008.
- Braconnot, P., Harrison, S. P., Kageyama, M., Bartlein, P. J., Masson-Delmotte, V., Abe-Ouchi, A., Otto-Bliesner, B., and Zhao, Y.: Evaluation of climate models using palaeoclimatic data, *Nature Climate Change*, 2, 417–424, doi:10.1038/nclimate1456, <http://www.nature.com/nclimate/journal/vaop/ncurrent/full/nclimate1456.html><http://www.nature.com/doi/finder/10.1038/nclimate1456>, 2012.
- Caley, T., Malaizé, B., Zaragosi, S., Rossignol, L., Bourget, J., Eynaud, F., Martinez, P., Giraudeau, J., Charlier, K., and Ellouzi-Zimmermann, N.: New Arabian Sea records help decipher orbital timing of Indo-Asian monsoon, *Earth and Planetary Science Letters*, 308, 433–444, doi:10.1016/j.epsl.2011.06.019, <http://dx.doi.org/10.1016/j.epsl.2011.06.019><http://linkinghub.elsevier.com/retrieve/pii/S0012821X11003785>, 2011.
- Christensen, J. H., Kanikicharla, K. K., Marshall, G., and Turner, J.: *Climate phenomena and their relevance for future regional climate change*, 2013.
- Clemens, S. C. and Prell, W. L.: A 350,000 year summer-monsoon multi-proxy stack from the Owen Ridge, Northern Arabian Sea, *Marine Geology*, 201, 35–51, doi:10.1016/S0025-3227(03)00207-X, 2003.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., and Others: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Quarterly Journal of the royal meteorological society*, 137, 553–597, 2011.
- Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J.-P., Ethé, C., Fairhead, L., Fichefet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D. A., Hourdin, F., Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A., Lefebvre, M.-P., Lefebvre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, *Climate Dynamics*, 40, 2123–2165, doi:10.1007/s00382-012-1636-1, <http://link.springer.com/10.1007/s00382-012-1636-1>, 2013.
- Emeis, K.-C., Anderson, D. M., Doose, H., Kroon, D., and Schulz-Bull, D.: Sea-Surface Temperatures and the History of Monsoon Upwelling in the Northwest Arabian Sea during the Last 500,000 Years, *Quaternary Research*, 43, 355–361, doi:10.1006/qres.1995.1041, <http://linkinghub.elsevier.com/retrieve/pii/S0033589485710411>, 1995.
- Fichefet, T. and Maqueda, M. A. M.: Sensitivity of a global sea ice model to the treatment of ice thermodynamics and dynamics, *Journal of Geophysical Research: Oceans*, 102, 12 609–12 646, doi:10.1029/97JC00480, <http://doi.wiley.com/10.1029/97JC00480>, 1997.
- Findlater, J.: A major low-level air current near the Indian Ocean during the northern summer, *Quarterly Journal of the Royal Meteorological Society*, 95, 362–380, doi:10.1002/qj.49709540409, <http://doi.wiley.com/10.1002/qj.49709540409>, 1969.
- Fleitmann, D., Burns, S. J., Mangini, A., Mudelsee, M., Kramers, J., Villa, I., Neff, U., Al-Subbary, A. A., Buettner, A., Hippler, D., and Matter, A.: Holocene ITCZ and Indian monsoon dynamics recorded in stalagmites from Oman and Yemen (Socotra), *Quaternary Science Reviews*, 26, 170–188, doi:10.1016/j.quascirev.2006.04.012, <http://linkinghub.elsevier.com/retrieve/pii/S0277379106002265>, 2007.


- Goswami, B. N., Madhusoodanan, M. S., Neema, C. P., and Sengupta, D.: A physical mechanism for North Atlantic SST influence on the Indian summer monsoon, *Geophysical Research Letters*, 33, L02 706, doi:10.1029/2005GL024803, <http://doi.wiley.com/10.1029/2005GL024803>, 2006.
- Held, I. M. and Soden, B. J.: Robust responses of the hydrological cycle to global warming, *Journal of Climate*, 19, 5686–5699, doi:10.1175/JCLI3990.1, 2006.
- Hourdin, F., Foujols, M.-A., Codron, F., Guemas, V., Dufresne, J.-L., Bony, S., Denvil, S., Guez, L., Lott, F., Ghattas, J., Braconnot, P., Marti, O., Meurdesoif, Y., and Bopp, L.: Impact of the LMDZ atmospheric grid configuration on the climate and sensitivity of the IPSL-CM5A coupled model, *Climate Dynamics*, 40, 2167–2192, doi:10.1007/s00382-012-1411-3, <http://link.springer.com/10.1007/s00382-012-1411-3>, 2013.
- 10 Ivanova, E., Schiebel, R., Singh, A. D., Schmiedl, G., Niebler, H.-S., and Hemleben, C.: Primary production in the Arabian Sea during the last 135 000 years, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 197, 61–82, doi:10.1016/S0031-0182(03)00386-9, <http://linkinghub.elsevier.com/retrieve/pii/S0031018203003869>, 2003.
- Kageyama, M., Braconnot, P., Bopp, L., Caubel, A., Foujols, M.-A., Guilyardi, E., Khodri, M., Lloyd, J., Lombard, F., Mariotti, V., Marti, O., Roy, T., and Woillez, M.-N.: Mid-Holocene and Last Glacial Maximum climate simulations with the IPSL model—part I: comparing IPSL\_CM5A to IPSL\_CM4, *Climate Dynamics*, 40, 2447–2468, doi:10.1007/s00382-012-1488-8, <http://link.springer.com/10.1007/s00382-012-1488-8>, 2013.
- 15 Keen, T. R., Kindle, J. C., and Young, D. K.: The interaction of southwest monsoon upwelling, advection and primary production in the northwest Arabian Sea, *Journal of Marine Systems*, 13, 61–82, doi:10.1016/S0924-7963(97)00003-1, 1997.
- Koné, V., Aumont, O., Lévy, M., and Resplandy, L.: Physical and biogeochemical controls of the phytoplankton seasonal cycle in the Indian Ocean: A modeling study, *Ocean Biogeochemical*, pp. 147–166, doi:10.1029/2008GM000700, [http://www.lodyc.jussieu.fr/~marina/PUBLI/Kone\[\\_\]GMS\[\\_\]2009.pdf](http://www.lodyc.jussieu.fr/~marina/PUBLI/Kone[_]GMS[_]2009.pdf)<http://doi.wiley.com/10.1029/2008GM000700><http://onlinelibrary.wiley.com/doi/10.1029/2008GM000700/summary>, 2009.
- Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I. C.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, *Global Biogeochemical Cycles*, 19, n/a—n/a, doi:10.1029/2003GB002199, <http://doi.wiley.com/10.1029/2003GB002199>, 2005.
- 25 Lee, C. M., Jones, B. H., Brink, K. H., and Fischer, A. S.: The upper-ocean response to monsoonal forcing in the Arabian Sea: Seasonal and spatial variability, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 47, 1177–1226, doi:10.1016/S0967-0645(99)00141-1, <http://linkinghub.elsevier.com/retrieve/pii/S0967064599001411>, 2000.
- Lévy, M., Shankar, D., André, J.-M., Shenoi, S. S. C., Durand, F., and de Boyer Montégut, C.: Basin-wide seasonal evolution of the Indian Ocean's phytoplankton blooms, *Journal of Geophysical Research*, 112, C12 014, doi:10.1029/2007JC004090, <http://doi.wiley.com/10.1029/2007JC004090>, 2007.
- 30 Licciardi, J. M., Clark, P. U., Jenson, J. W., and MacAyeal, D. R.: Deglaciation of soft-bedded Laurentide Ice Sheet, *Quaternary Science Reviews*, 17, 427–448, doi:10.1016/S0277-3791(97)00044-9, 1998.
- Luther, M. E., O'Brien, J. J., and Prell, W. L.: Variability in upwelling fields in the northwestern Indian Ocean 1. Model experiments for the past 18,000 years, *Paleoceanography*, 5, 433–445, doi:10.1029/PA005i003p00433, <http://doi.wiley.com/10.1029/PA005i003p00433>, 1990.
- 35 Madec, G.: NEMO\_book\_v3\_3, Manual, pp. 1–332, <http://www.nemo-ocean.eu/About-NEMO/Reference-manuals>, 2011.

- Marshall, J. and Plumb, R. A.: Atmosphere, Ocean and Climate Dynamics: An Introductory Text, Elsevier Academic Press, <http://www.elsevierdirect.com/companion.jsp?ISBN=9780125586917>{% }5Cn<http://www.amazon.jp/dp/0125586914>, 2008.
- Marzin, C. and Braconnot, P.: Variations of Indian and African monsoons induced by insolation changes at 6 and 9.5 kyr BP, *Climate Dynamics*, 33, 215–231, doi:10.1007/s00382-009-0538-3, <http://link.springer.com/10.1007/s00382-009-0538-3>, 2009.
- 5 Marzin, C., Braconnot, P., and Kageyama, M.: Relative impacts of insolation changes, meltwater fluxes and ice sheets on African and Asian monsoons during the Holocene, *Climate Dynamics*, 41, 2267–2286, doi:10.1007/s00382-013-1948-9, <http://link.springer.com/10.1007/s00382-013-1948-9>, 2013.
- Masson, V., Braconnot, P., Jouzel, J., de Noblet, N., Cheddadi, R., and Marchal, O.: Simulation of intense monsoons under glacial conditions, *Geophysical Research Letters*, 27, 1747–1750, doi:10.1029/1999GL006070, <http://doi.wiley.com/10.1029/1999GL006070>, 2000.
- 10 McCreary, J. P., Murtugudde, R., Vialard, J., Vinayachandran, P. N., Wiggert, J. D., Hood, R. R., Shankar, D., and Shetye, S.: Biophysical processes in the Indian Ocean, in: *Indian Ocean Biogeochemical Processes and Ecological Variability*, pp. 9–32, American Geophysical Union, Washington, D.C., doi:10.1029/2008GM000768, <http://doi.wiley.com/10.1029/2008GM000768>, 2009.
- Mooley, D. A. and Parthasarathy, B.: Fluctuations in all-India summer monsoon rainfall during 1871–1978, *Climatic Change*, 6, 287–301, 1984.
- 15 Murtugudde, R., Seager, R., and Thoppil, P.: Arabian Sea response to monsoon variations, *Paleoceanography*, 22, doi:10.1029/2007PA001467, <http://doi.wiley.com/10.1029/2007PA001467>, 2007.
- Pausata, F. S. R., Li, C., Wettstein, J. J., Kageyama, M., and Nisancioglu, K. H.: The key role of topography in altering North Atlantic atmospheric circulation during the last glacial period, *Climate of the Past*, 7, 1089–1101, doi:10.5194/cp-7-1089-2011, <http://www.clim-past.net/7/1089/2011/>, 2011.
- 20 Peltier, W. R., Argus, D. F., and Drummond, R.: Space geodesy constrains ice age terminal deglaciation: The global ICE-6G\_C (VM5a) model, *Journal of Geophysical Research: Solid Earth*, 120, 450–487, doi:10.1002/2014JB011176, <http://doi.wiley.com/10.1002/2014JB011176>, 2015.
- Prasanna Kumar, S., Madhupratap, M., Dileep Kumar, M., Gauns, M., Muraleedharan, P. M., Sarma, V. V. S. S., and De Souza, S. N.: Physical control of primary productivity on a seasonal scale in central and eastern Arabian Sea, *Proceedings of the Indian Academy of Sciences, Earth and Planetary Sciences*, 109, 433–441, doi:10.1007/BF02708331, 2000.
- 25 Prasanna Kumar, S., Madhupratap, M., Dileep Kumar, M., Muraleedharan, P. M., De Souza, S. N., Gauns, M., and Sarma, V. V. S. S.: High biological productivity in the central Arabian Sea during the summer monsoon driven by Ekman pumping and lateral advection, *Current Science*, 81, 1633–1638, 2001.
- Prell, W. L. and Kutzbach, J. E.: Monsoon variability over the past 150,000 years, *Journal of Geophysical Research*, 92, 8411, doi:10.1029/JD092iD07p08411, <http://doi.wiley.com/10.1029/JD092iD07p08411>, 1987.
- 30 Prell, W. L., Murray, D. W., Clemens, S. C., and Anderson, D. M.: Evolution and variability of the Indian Ocean summer monsoon: Evidence from the western Arabian sea drilling program, in: *Geophysical Monograph Series*, vol. 70, pp. 447–469, American Geophysical Union, Washington, D.C., doi:10.1029/GM070p0447, <http://www.agu.org/books/gm/v070/GM070p0447/GM070p0447.shtml><http://doi.wiley.com/10.1029/GM070p0447>, 1992.
- 35 Resplandy, L., Lévy, M., Madec, G., Pous, S., Aumont, O., and Kumar, D.: Contribution of mesoscale processes to nutrient budgets in the Arabian Sea, *Journal of Geophysical Research*, 116, C11 007, doi:10.1029/2011JC007006, <http://doi.wiley.com/10.1029/2011JC007006>, 2011.


- Rostek, F., Bard, E., Beaufort, L., Sonzogni, C., and Ganssen, G.: Sea surface temperature and productivity records for the past 240 kyr in the Arabian Sea, *Deep Sea Research Part II: Topical Studies in Oceanography*, 44, 1461–1480, doi:10.1016/S0967-0645(97)00008-8, <http://linkinghub.elsevier.com/retrieve/pii/S0967064597000088>, 1997.
- Ruddiman, W. F.: What is the timing of orbital-scale monsoon changes?, *Quaternary Science Reviews*, 25, 657–658, doi:10.1016/j.quascirev.2006.02.004, 2006.
- Sandeep, S. and Ajayamohan, R. S.: Poleward shift in Indian summer monsoon low level jetstream under global warming, *Climate Dynamics*, 45, 337–351, doi:10.1007/s00382-014-2261-y, <http://dx.doi.org/10.1007/s00382-014-2261-y>, 2014.
- Schott, F. a. and McCreary, J. P.: The monsoon circulation of the Indian Ocean, *Progress in Oceanography*, 51, 1–123, doi:10.1016/S0079-6611(01)00083-0, <http://linkinghub.elsevier.com/retrieve/pii/S0079661101000830>, 2001.
- 10 Séférian, R., Bopp, L., Gehlen, M., Orr, J. C., Ethé, C., Cadule, P., Aumont, O., Salas y Mélia, D., Voldoire, A., and Madec, G.: Skill assessment of three earth system models with common marine biogeochemistry, *Climate Dynamics*, 40, 2549–2573, doi:10.1007/s00382-012-1362-8, <http://link.springer.com/10.1007/s00382-012-1362-8>, 2013.
- Sirocko, F., Sarnthein, M., Lange, H., and Erlenkeuser, H.: Atmospheric summer circulation and coastal upwelling in the Arabian Sea during the Holocene and the last glaciation, *Quaternary Research*, 36, 72–93, doi:10.1016/0033-5894(91)90018-Z, <http://linkinghub.elsevier.com/retrieve/pii/003358949190018Z>, 1991.
- 15 Sperber, K. R., Annamalai, H., Kang, I.-S., Kitoh, A., Moise, A., Turner, A., Wang, B., and Zhou, T.: The Asian summer monsoon: an intercomparison of CMIP5 vs. CMIP3 simulations of the late 20th century, *Climate Dynamics*, 41, 2711–2744, doi:10.1007/s00382-012-1607-6, <http://link.springer.com/10.1007/s00382-012-1607-6>, 2013.
- Storey, M., Roberts, R. G., and Saidin, M.: Astronomically calibrated  $^{40}\text{Ar}/^{39}\text{Ar}$  age for the Toba supereruption and global synchronization of late Quaternary records, *Proceedings of the National Academy of Sciences*, 109, 18684–18688, doi:10.1073/pnas.1208178109, <http://www.pnas.org/content/109/46/18684.abstract>{%}5Cn<http://www.pnas.org/content/109/46/18684.full.pdf><http://www.pnas.org/cgi/doi/10.1073/pnas.1208178109>, 2012.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, *Bulletin of the American Meteorological Society*, 93, 485–498, doi:10.1175/BAMS-D-11-00094.1, <http://journals.ametsoc.org/doi/abs/10.1175/BAMS-D-11-00094.1>, 2012.
- 25 Valcke, S.: The OASIS3 coupler: a European climate modelling community software, *Geoscientific Model Development Discussions*, 5, 2139–2178, doi:10.5194/gmdd-5-2139-2012, <http://www.geosci-model-dev-discuss.net/5/2139/2012/>, 2012.
- Wiggert, J. D., Hood, R. R., Banse, K., and Kindle, J. C.: Monsoon-driven biogeochemical processes in the Arabian Sea, *Progress in Oceanography*, 65, 176–213, doi:10.1016/j.pocean.2005.03.008, 2005.
- Willez, M.-N., Levavasseur, G., Daniau, A.-L., Kageyama, M., Urrego, D. H., Sánchez-Goñi, M.-F., and Hanquiez, V.: Impact of precession on the climate, vegetation and fire activity in southern Africa during MIS4, *Climate of the Past*, 10, 1165–1182, doi:10.5194/cp-10-1165-2014, <http://www.clim-past.net/10/1165/2014/>, 2014.
- 30 Yessad, K.: Full-pos in the cycle 43 of APEGE/IFS, Tech. rep., <https://www.cnrm-game-meteo.fr/gmapdoc/IMG/pdf/ykfp03.pdf>, 2016.
- Zhang, H.-M.: Blended and gridded high resolution global sea surface wind speed and climatology from multiple satellites: 1987–present, in: American Meteorological Society 2006 Meeting, 2006.
- 35 Ziegler, M., Lourens, L. J., Tuenter, E., Hilgen, F., Reichert, G. J., and Weber, N.: Precession phasing offset between Indian summer monsoon and Arabian Sea productivity linked to changes in Atlantic overturning circulation, *Paleoceanography*, 25, 1–16, doi:10.1029/2009PA001884, 2010.

**Table 1.** Boundary conditions for the 8 simulations studied in this work. Precession is defined as the longitude of the perihelion, relative to the moving vernal equinox, minus  $180^\circ$ . Ice sheets are represented on figure 1. "Pmip3" ice-sheet stands for the PMIP3 ice sheet reconstruction (Abe-Ouchi et al., 2015), ICE6g-16k stands for the ICE6G reconstruction at 16 kyr BP (Peltier et al., 2015)


Simulation name	Interglacial climates			Glacial climates				
	CTRL	MH	EH	LGM	MIS3	MIS4F	MIS4M	MIS4D
<b>Time (kyr BP)</b>	0	6	9.5	21	46	60	66	72
<b>Ice sheets</b>	present	present	present	pmip3	Ice6g-16k	Ice6g-16k	Ice6g-16k	Ice6g-16k
<b>Sea level difference vs CTRL (m)</b>	0	0	0	-120	-70	-70	-70	-70
<b>Eccentricity</b>	0.016715	0.018682	0.0193553	0.018994	0.0138427	0.018469	0.021311	0.024345
<b>Obliquity (<math>^\circ</math>)</b>	22.391	24.105	24.2306	22.949	24.3548	23.2329	22.493	22.391
<b>Precession (<math>\omega - 180^\circ</math>)</b>	102.7	0.87	303.032	114.42	101.337	266.65	174.82	80.09
<b><math>CO_2</math> (ppm)</b>	284	280	284	185	205	200	195	230
<b><math>N_2O</math> (ppm)</b>	275	270	275	200	260	230	217	230
<b><math>CH_4</math> (ppb)</b>	791	650	791	350	500	426	450	450




**Figure 1.** a) The three ice sheet covers used as boundary conditions. The 0k ice sheet is used in the CTRL, MH and EH simulations, the pmp3 ice sheet (Abe-Ouchi et al., 2015) is used in the LGM simulation and the ICE6g-16k ice sheet (Peltier et al., 2015) is used in the MIS3 and all the MIS4 (F, M and D) simulations and b) the different values of the other forcing parameters for all the simulations.



**Figure 2.** a) Coastal (orange shading) and central-east-northern (grey shading/black dotted contour) Arabian Sea areas, position of core MD04-2873 (yellow circle) and, b) Seasonal-seasonal cycles of current-day productivity in the coastal (bold orange line) -central-eastern (simple line) and northern (60E-68E, 20N-68N, black dashed line) Arabian Sea. Productivity data comes from the SeaWiFS satellite data for the period 1998-2005 (Lévy et al., 2007).




**Figure 3.** Modelled and observed seasonal (JJAS) patterns of a) productivity ( $\text{molC.m}^{-2}.\text{yr}^{-1}$ ), b) surface wind intensity ( $\text{m.s}^{-1}$ ), c) surface wind stress intensity ( $10^{-3}\text{N.m}^{-2}$ ) and d) surface wind stress curl intensity ( $10^{-7}\text{N.m}^{-3}$ ) and e) seas surface temperature (SST) ( $^{\circ}\text{C}$ ). We used SeaWiFS data in 1998-2005 for productivity (Lévy et al., 2007) and the NOAA Multiple-Satellite 1995-2005 product (Zhang, 2006) for the climatology of surface wind, wind stress and wind stress curl intensity and the ERA-interim reanalysis 1979-2014 for SST (Dee et al., 2011).



**Figure 4.** Anomalies of total primary productivity (TPP,  $\text{molC.m}^{-2}\text{.yr}^{-1}$ , represented by the color-scale) integrated over the whole water column as a function of wind stress anomalies ( $10^{-3}\text{N.m}^{-2}$ , y-axis) and wind stress curl anomalies ( $10^{-7}\text{N.m}^{-3}$ , x-axis) in the coastal western Arabian Sea (see Fig. 2a) for the CTRL simulation. Anomalies are computed as the difference between each summer average (JJAS) and the seasonal summer mean of the corresponding variable in the CTRL simulation. The color of the circles represents the value of the change in TPP compared to CTRL. The color-scale, x-axis and y-axis ranges are the same as those in Figure 7 and Figure 10.







**Figure 5. a,b** : Seasonal cycle of the insolation at the top of the atmosphere ( $\text{W.m}^{-2}$ ) in a) the CTRL simulation and b) the difference between EH and CTRL. The Northern Hemisphere ( $0-90^\circ\text{N}$ ) mean summer (JJAS) insolation,  $\text{NH}_{mean}(\text{JJAS})$ , in CTRL and the difference between EH and CTRL,  $\Delta\text{NH}_{mean}(\text{JJAS})$ , is given under panel a) and b), respectively.

**c,d** : Upper tropospheric temperature (TT) averaged between 200 hPa and 500 hPa in c) CTRL and d) the difference between EH and CTRL.  $\Delta\text{TT}$  value, under panel c), is the TT gradient between a northern region ( $60^\circ\text{E}-120^\circ\text{E}$ ;  $10^\circ\text{N}-45^\circ\text{N}$ ) and a southern region ( $60^\circ\text{E}-120^\circ\text{E}$ ;  $25^\circ\text{S}-10^\circ\text{S}$ ) (black boxes on the maps).  $\Delta(\Delta\text{TT})$ , under panel d), is the difference of TT gradients between EH and CTRL.


**e,f** : Boreal summer (JJAS) sea level pressure (SLP) anomaly (from the annual mean) lower than -5 hPa for e) CTRL and f) EH-CTRL. The SLPa-5 barycentre (i.e. barycentre of the SLP anomalies lower than -5 hPa over the region  $20^\circ\text{W}-150^\circ\text{E}$ ;  $30^\circ\text{S}-60^\circ\text{N}$ ) is represented by a black star for CTRL on e) and f) and by a red star for EH on f).




**Figure 6.** Boreal summer mean differences between EH and CTRL for a) total primary productivity (TPP,  $\text{molC.m}^{-2}.\text{yr}^{-1}$ ) integrated over the whole water column, b)  $\text{NO}_3^-$  concentration ( $\text{mmolC.m}^{-2}$ ) in the first 30 m of the water column, c) wind stress intensity ( $\text{N.m}^{-2}$ ) and direction (arrows), d) mixed layer depth (MLD, m), e) wind stress curl intensity ( $10^{-7}\text{N.m}^{-3}$ ) and f) Ekman pumping ( $\text{m.yr}^{-1}$ , positive upward). The black contour on each panel represents the coastal area on which we averaged the variables throughout the text. Coastal areas for atmospheric and oceanic variables differs slightly because of the different model grids.




**Figure 7.** Seasonal (JJAS) anomalies of total primary productivity (TPP,  $\text{molC}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$ ), integrated over the whole water column, as a function of wind stress anomalies ( $10^{-3}\text{N}\cdot\text{m}^{-2}$ ) and wind stress curl anomalies ( $10^{-7}\text{N}\cdot\text{m}^{-3}$ ) in ~~a) the coastal western Arabian Sea and b) the central-east Arabian Sea~~ (see Fig. 2a). Anomalies are computed as the difference between each yearly summer average in the EH simulation and the seasonal summer mean of the CTRL simulation. The color of the circles represents the value of the change in TPP between EH and CTRL. The color-scale, x-axis and y-axis ranges are the same as those in Figure ~~10~~4 and Figure ~~4~~10.




**Figure 8.** Seasonal (JJAS) productivity changes ( $\text{molC.m}^{-2}.\text{yr}^{-1}$ , colorscale) and Ekman pumping (black contour every  $20\text{m.yr}^{-1}$ ) changes compared to the CTRL simulation for a) MH, b) LGM, c) MIS3, d) MIS4F, e) MIS4M and f) MIS4D simulation in the Indian Ocean. The yellow circle indicates the position of core MD04-2873 used later in this paper.



**Figure 9.** Box-plots of summer (JJAS) a)  $\Delta\text{TT}$  (K) values, coastal Arabian Sea b) productivity ( $\text{molC.m}^{-2}.\text{yr}^{-1}$ ) integrated over the whole water column, c) nitrate concentration ( $\text{mmolC.m}^{-2}$ ) in the first 30 meters, d) wind stress intensity ( $10^{-3}\text{N.m}^{-2}$ ) and e) wind stress curl intensity ( $10^{-7}\text{N.m}^{-3}$ ), for all 8 simulations. Dash-Dashed grey line indicates the CTRL value for each variable. The black simple line on e) panel indicates the zero value. The boxplots highlight the median value (bold line), the first and the third quartile (lower and higher limits of the box) and the 95% confidence interval of the median (upper and lower horizontal ticks). The dots are extreme values that happened during the 100 years of simulation.




**Figure 10.** ~~Seasonal~~ Coastal Arabian Sea seasonal (JJAS) productivity changes ( $\text{molC.m}^{-2}\text{.yr}^{-1}$ ) related to wind stress intensity ( $10^{-3}\text{N.m}^{-2}$ , y-axis) and wind stress curl intensity ( $10^{-7}\text{N.m}^{-3}$ , x-axis) changes compared to the CTRL simulation, ~~for a) coastal and b) central Arabian Sea.~~ The color scale, x-axis and y-axis ranges are the same as those in Figure 4 and Figure 7.




**Figure 11.** Seasonal (JJAS) coastal productivity ( $\text{molC}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$ ), wind stress intensity ( $10^{-3}\text{N}\cdot\text{m}^{-2}$ , y-axis) and wind stress curl intensity ( $10^{-7}\text{N}\cdot\text{m}^{-3}$ , x-axis) as a function of the SLP barycentre a-c) longitude and d-f) latitude. Errors bars give standard deviation of the 100 summers. [The dotted black box on panel a\) represents the region on which we zoomed for panels b, c and d.](#)


**a) Climatic precession**



**b) Obliquity**




**Figure 12.** Mean a) climatic precession ( $e \cdot \sin(\omega - 180^\circ)$  where  $e$  is the eccentricity and  $\omega - 180^\circ$  is the precession) and b) obliquity values as a function of the longitude and latitude of the SLPa-5 barycentre for the 8 climate simulations. Errors bars give standard deviation of the SLPa-5 position over the 100 summers of each simulation.



**Figure 13.** Box-plots of a) yearly reconstructed productivity from core MD04-2873 in the north-western Arabian Sea and, b) annual total primary productivity (TPP), c) summer total primary productivity, d) annual export production (EPC) at 100 m and e) summer export production at 100 m, in the northern Arabian Sea (60°E-68°E, 20°N-68°N) for the Holocene and glacial time periods. The boxplots highlight the median value (bold line), the first and the third quartile (lower and higher limits of the box) and the 95% confidence interval of the median (upper and lower horizontal ties). The dots are extreme values that happened during the 100 years of simulation





**Figure 14.** Identified seasonal (JJAS) processes behind productivity changes in glacial and inter-glacial climates. Bold lines highlight the major pathways.