

1                   **THE LAST GLACIAL TERMINATION ON THE EASTERN FLANK OF THE CENTRAL**  
2   **PATAGONIAN ANDES (47°S)**

3  
4 William I. Henríquez<sup>1,2</sup>, Rodrigo Villa-Martínez<sup>3</sup>, Isabel Vilanova<sup>4</sup>, Ricardo De Pol-Holz<sup>3</sup>, and  
5   Patricio I. Moreno<sup>2,\*</sup>

6  
7   <sup>1</sup>Victoria University of Wellington, Wellington, New Zealand

8   <sup>2</sup>Instituto de Ecología y Biodiversidad, Departamento de Ciencias Ecológicas, Universidad  
9   de Chile, Casilla 653, Santiago, Chile

10   <sup>3</sup>GAIA-Antártica, Universidad de Magallanes, Avda. Bulnes 01855, Punta Arenas, Chile

11   <sup>4</sup>Museo Argentino de Ciencias Naturales Bernardino Rivadavia, Avda. Angel Gallardo 470,  
12   Buenos Aires, Argentina.

13   \*Corresponding author: [pimoreno@uchile.cl](mailto:pimoreno@uchile.cl)

14

15 ABSTRACT

16

17 Few studies have examined in detail the sequence of events during the last glacial  
18 termination (T1) in the core sector of the Patagonian Ice Sheet (PIS), the largest ice mass  
19 in the southern hemisphere outside Antarctica. Here we report results from Lago Edita  
20 (47°8'S, 72°25'W, 570 m.a.s.l.), a small closed-basin lake located in a valley overridden by  
21 eastward-flowing Andean glaciers during the Last Glacial Maximum (LGM). The Lago Edita  
22 record shows glaciolacustrine sedimentation until 19,400 yr BP and a mosaic of cold-  
23 resistant, hygrophilous conifers and rainforest trees, along with alpine herbs between  
24 19,400-11,000 yr BP. Increases in arboreal pollen at 13,200 and 11,000 yr BP led to the  
25 establishment of forests near Lago Edita between 9000-10,000 yr BP. Our data suggest  
26 that the PIS retreated at least ~90 km from its LGM limit between ~21,000-19,400 yr BP  
27 and that scattered, low-density populations of cold-resistant hygrophilous conifers,  
28 rainforest trees, high Andean and steppe herbs thrived east of the Andes during the LGM  
29 and T1, implying high precipitation and southern westerly wind (SWW) intensity at 47°S.  
30 We interpret large-magnitude increases in arboreal vegetation as treeline-rise episodes  
31 driven by warming pulses at 13,200 and 11,000 yr BP coupled with a decline in SWW  
32 influence at ~11,000 yr BP, judging from the disappearance of cold-resistant hygrophilous  
33 trees and herbs. We propose that the PIS imposed a regional cooling signal along its  
34 eastern, downwind margin through T1 that lasted until the separation of the North and  
35 South Patagonian icefields along the Andes. We posit that the withdrawal of glacial and  
36 associated glaciolacustrine environments through T1 provided a route for the dispersal of  
37 hygrophilous trees and herbs from the eastern flank of the central Patagonian Andes,  
38 contributing to the afforestation of the western Andean slopes and pacific coasts of  
39 central Patagonia during T1.

40

## 41 INTRODUCTION

42

43 The Patagonian ice sheet (PIS) was the largest ice mass in the southern hemisphere  
44 outside Antarctica during the last glacial maximum (LGM). Outlet lobes from the PIS  
45 flowed westward into the Pacific coast south of 43°S and eastward toward the extra-  
46 Andean Patagonian plains, blanketing a broad range of environments and climatic zones  
47 across and along the Andes. Land biota from formerly ice-free sectors underwent local  
48 extinction or migrated toward the periphery of the advancing PIS during the last glaciation  
49 until its culmination during the LGM. The PIS then underwent rapid recession and thinning  
50 through the last glacial termination (termination 1= T1: between ~18,000-11,000 yr BP)  
51 toward the Andes as illustrated by stratigraphic, geomorphic and radiocarbon-based  
52 chronologies from northwestern Patagonia (39°-43°S) (Denton et al., 1999; Moreno et al.,  
53 2015). These data, along with the Canal de la Puntilla-Huelmo pollen record (~41°S)  
54 (Moreno et al., 2015) (Figure 1), indicate abandonment from the LGM margins in the  
55 lowlands at 17,800 yr BP, abrupt arboreal expansion, and accelerated retreat that exposed  
56 Andean cirques located above 800 m.a.s.l. within 1000 years or less in response to abrupt  
57 warming. Similarly, glaciers from Cordillera Darwin (54°-55°S), the southernmost icefield  
58 in South America, underwent rapid recession from their LGM moraines located in central  
59 and northern Tierra del Fuego prior to 17,500 yr BP, and led to ice-free conditions by  
60 16,800 yr BP near the modern ice fronts (Hall et al., 2013). Sea surface temperature  
61 records from the SE Pacific (Caniupán et al., 2011) are consistent with these terrestrial  
62 records, however, their timing, structure, magnitude and rate of change may be  
63 overprinted by the vicinity of former ice margins and shifts in marine reservoir ages  
64 (Caniupán et al., 2011; Siani et al., 2013).

65 In contrast, very few studies have been conducted in the Andean sector of central-  
66 west Patagonia (45°-48°S) about the timing of glacial advances near the end of the LGM as  
67 well as the structure/ chronology of glacial retreat and climate changes during T1. Recent  
68 chronologies include cosmogenic radionuclides of terminal moraines of the Río Blanco and  
69 recessional moraines deposited by the Lago Cochrane ice lobe (LCIL) (Boex et al., 2013;

70 Hein et al., 2010) (Figure 1), and optically stimulated luminescence dating of  
71 glaciolacustrine beds associated with Glacial Lake Cochrane (GLC) (47°S) (Glasser et al.,  
72 2016). These studies reported ages between 29,000-19,000 yr BP for the final LGM  
73 advance and drainage of GLC toward the Pacific between 13,000-8000 yr BP caused by  
74 breakup of the North and South Patagonian Icefields during the final stages of T1 (Turner  
75 et al., 2005). Palynological interpretations from the Lago Shaman (44°26'S, 71°11'W, 919  
76 m.a.s.l.) and Mallín Pollux (45°41'S, 71°50'W, 640 m.a.s.l.) sites (de Porrás et al., 2012;  
77 Markgraf et al., 2007), located east of the Andes between 44°S and 45°S respectively  
78 (Figure 1), indicate predominance of cold and dry conditions during T1 and negative  
79 anomalies in southern westerly wind (SWW) influence. The validity and regional  
80 applicability of these stratigraphic, chronologic and palynologic interpretations, however,  
81 awaits replication by detailed stratigraphic/geomorphic data from sensitive sites  
82 constrained by precise chronologies.

83 Paleoclimate simulations (Bromwich et al., 2005; Bromwich et al., 2004) and  
84 stratigraphic studies (Kaufman et al., 2004) in the periphery of the Laurentide Ice Sheet in  
85 North America, have detected that large ice sheets exerted important impacts on the  
86 thermal structure and atmospheric circulation at regional, continental and zonal scale  
87 from the LGM to the early Holocene. This aspect has remained largely unexplored in the  
88 PIS region, and might be a factor of importance for understanding the dynamics of the  
89 SWW and climatic/biogeographic heterogeneities through T1 at regional scale. Progress in  
90 this field requires understanding the deglacial chronology of the PIS and a suite of  
91 sensitive paleoclimate sites across and along the residual ice masses through the last  
92 transition from extreme glacial to extreme interglacial conditions.

93 In this study we report high-resolution pollen and macroscopic charcoal records  
94 from sediment cores we collected from Lago Edita (47°8'S, 72°25'W, ~570 m.a.s.l.), a  
95 medium-sized closed-basin lake (radius ~250 m) located in Valle Chacabuco ~16 km  
96 northeast of the Cochrane township, east of the central Patagonian Andes (Figure 1). The  
97 relevant source area for pollen from lakes of this size is about 600-800 m from the lake's  
98 edge, according to numerical simulations using patchy vegetation landscapes (Sugita,

99 1994). Stratigraphic and chronologic results from Valle Chacabuco are important for  
100 elucidating the timing and rates of deglaciation in this core region of the PIS because this  
101 valley is located approximately two thirds (90 km) upstream from the LGM moraines  
102 deposited by LCIL east of Lago Cochrane relative to the modern ice fronts, and its  
103 elevation spans the highest levels of GLC during T1. The Lago Edita data allow assessment  
104 of vegetation, fire-regime and climate changes during the last global transition from  
105 extreme glacial to extreme interglacial conditions in central-west Patagonia. The aim of  
106 this paper is to contribute toward: (1) the development of a recessional chronology of the  
107 LCIL and (2) regressive phases of GLC, (3) documenting the composition and geographic  
108 shifts of the glacial and deglacial vegetation, (4) understanding the tempo and mode of  
109 vegetation and climate changes during T1 and the early Holocene, (5) constraining the  
110 regional climatic influence of the PIS during T1 in terrestrial environments, and (6)  
111 identifying possible dispersal routes of tree taxa characteristic of modern evergreen  
112 forests in central-west Patagonia during T1.

113

#### 114 Study Area

115

116 Central Chilean Patagonia, i.e. the Aysén region (43°45'S-47°45'S), includes  
117 numerous channels, fjords, islands, and archipelagos along the Pacific side, attesting for  
118 tectonic subsidence of Cordillera de la Costa and intense glacial erosion during the  
119 Quaternary. The central sector features an intricate relief associated to the Patagonian  
120 Andes with summits surpassing 3000 m.a.s.l., deep valleys, lakes of glacial origin, and  
121 active volcanoes such as Hudson, Macá, Cay, Mentolat and Melimoyu (Stern, 2004). The  
122 Andes harbors numerous glaciers and the North Patagonian Icefield (Figure 1), which  
123 acted as the source for multiple outlet glacier lobes that coalesced with glaciers from the  
124 South Patagonian Icefield and formed the PIS during Quaternary glaciations, blocked the  
125 drainage toward the Pacific funneling large volumes of glacial meltwater toward the  
126 Atlantic (Turner et al., 2005). Farther to the east the landscape transitions into the back-  
127 arc extra-Andean plains and plateaus.

128 Patagonia is ideal for studying the paleoclimate evolution of the southern mid-  
129 latitudes including past changes in the SWW because it is the sole continental landmass  
130 that intersects the low and mid-elevation zonal atmospheric flow south of 47°S.  
131 Orographic rains associated to storms embedded in the SWW enhance local precipitation  
132 by the ascent of moisture-laden air masses along the western Andean slopes, giving way  
133 to subsidence and acceleration of moisture-deprived winds along the eastern Andean  
134 slopes (Garreaud et al., 2013). This process accounts for a steep precipitation gradient  
135 across the Andes, illustrated by the annual precipitation measured in the coastal township  
136 of Puerto Aysén (2414 mm/year) and the inland Balmaceda (555 mm/year)  
137 (<http://explorador.cr2.cl/>), localities separated by ~80 km along a west-to-east axis. The  
138 town of Cochrane, located ~15 km south of our study site features annual precipitation of  
139 680 mm/year and mean annual temperature of 7.8 °C (Figure 1).

140 Weather station and reanalysis data along western Patagonia show positive  
141 correlations between zonal wind speed and local precipitation, a relationship that extends  
142 to sectors adjacent to the eastern slopes of the Andes (Garreaud et al., 2013; Moreno et  
143 al., 2014). Therefore, changes in local precipitation in the Aysén region are good  
144 diagnostics for atmospheric circulation changes associated with the frequency/intensity of  
145 storms embedded in the SWW over a large portion of the southeast Pacific. This  
146 relationship can be applied to paleoclimate records from central Chilean Patagonia for  
147 inferring the behavior of the SWW on the basis of past changes in precipitation or  
148 hydrologic balance.

149 The steep precipitation gradient, in conjunction with adiabatic cooling and enhanced  
150 continentality toward the east, influences the distribution and composition of the  
151 vegetation, inducing altitudinal, latitudinal and longitudinal zonation of plant communities  
152 throughout the Patagonian Andes. Physiognomic and floristic studies (Gajardo, 1994;  
153 Luebert and Pliscoff, 2006; Pisano, 1997; Schmithüsen, 1956) have recognized five units or  
154 communities which we characterize succinctly in the following sentences: 1) Magellanic  
155 Moorland: this unit occurs in maritime sectors with high precipitation (3000-4000  
156 mm/year and low seasonality) along the islands, fjords and channels, it is dominated by

157 cushion-forming plants such as *Donatia fascicularis*, *Astelia pumila* and *Tetroncium*  
158 *magallanicum*. Also present are the hygrophilous cold-resistant trees *Nothofagus*  
159 *betuloides* and the conifers *Pilgerodendron uviferum*, *Lepidothamnus fonkii* and  
160 *Podocarpus nubigena*. 2) Evergreen rainforest: present in humid, temperate (1500 -3000  
161 mm/year; <600 m.a.s.l.) sectors of Aysén, this unit is characterized by the trees  
162 *Nothofagus nitida*, *N. betuloides*, *Drimys winteri*, along with *P. uviferum* in waterlogged  
163 environments. 3) Winter deciduous forests: located in cooler and/or drier sectors with  
164 higher seasonality (400-1000 mm/year; 500-1180 m.a.s.l.). The dominant tree is  
165 *Nothofagus pumilio*, which intermingles with *N. betuloides* in western sites and the  
166 Patagonian steppe eastward. In the latter *N. pumilio* forms monospecific stands and  
167 presents a species-poor understory. A study of the spatial and temporal variation in *N.*  
168 *pumilio* growth at treeline along its latitudinal range (35°40'S-55°S) in the Chilean Andes  
169 (Lara et al., 2005) showed that temperature has a spatially larger control on tree growth  
170 than precipitation, and that this influence is particularly significant in the temperate Andes  
171 (> 40°S). These results suggest that low temperatures are the main limiting factor for the  
172 occurrence of woodlands and forests at high elevations in the Andes, considering that  
173 precipitation increases with elevation at any given latitude (Lara et al., 2005). The modern  
174 treeline near Cochrane lies between 800-1180 m.a.s.l. 4) Patagonian steppe: occurs in  
175 substantially drier (<500 mm/year) lowland areas with heightened continentality. This unit  
176 is dominated by herbs of the families Poaceae (*Festuca*, *Deschampsia*, *Stipa*, *Hordeum*,  
177 *Rytidosperma*, *Bromus*, *Elymus*), Rubiaceae (*Galium*), and shrubs of families Apiaceae  
178 (*Mulinum*), Rosaceae (*Acaena*), Fabaceae (*Adesmia*) and Rhamnaceae (*Discaria*). 5) High  
179 Andean Desert: occurs in the wind-swept montane environments above the treeline  
180 (>1000 m.a.s.l.) and is represented by herbs of the families Poaceae (*Poa*, *Festuca*),  
181 Asteraceae (*Nassauvia*, *Senecio*, *Perezia*), Berberidaceae (*Berberis*), Brassicaceae  
182 (*Cardamine*), Santalaceae (*Nanodea*), Rubiaceae (*Oreopulus*) Apiaceae (*Bolax*), Ericaceae  
183 (*Gaultheria*, *Empetrum*), along with *Gunnera magellanica* and *Valeriana*, with occasional  
184 patches of *Nothofagus antarctica*.

185

186 MATERIALS AND METHODS

187

188 We collected overlapping sediment cores over the deepest sector of Lago Edita (8 m  
189 water depth) from an anchored coring rig equipped with 10-cm diameter aluminum casing  
190 tube, using a 5-cm diameter Wright piston corer and a 7.5-cm diameter sediment-water  
191 interface piston corer with a transparent plastic chamber. We characterized the  
192 stratigraphy through visual descriptions, digital X radiographs to identify stratigraphic  
193 structures and loss-on-ignition to quantify the amount of organic, carbonate and  
194 siliciclastic components in the sediments (Heiri et al., 2001).

195 The chronology of the record is constrained by AMS radiocarbon dates on bulk  
196 sediment and chronostratigraphic correlation of the H1 tephra from Volcán Hudson (Stern  
197 et al., 2016). The radiocarbon dates were calibrated to calendar years before present (yr  
198 BP) using the CALIB 7.0 program. We developed a Bayesian age model using the Bacon  
199 package for R (Blaauw and Christen, 2011) to assign interpolated ages and confidence  
200 intervals for each level analyzed.

201 We processed and analyzed continuous/contiguous sediment samples (2 cm<sup>3</sup>) for  
202 pollen and fossil charcoal. The samples were processed using a standard procedure that  
203 includes 10% KOH, sieving with a 120 µm mesh, 46% HF and acetolysis (Faegri and Iversen,  
204 1989). We added exotic *Lycopodium* spores tablets to calculate concentration  
205 (particles\*cm<sup>-3</sup>) and accumulation rates of pollen and microscopic charcoal (particles\*cm<sup>-2</sup>  
206 \*years<sup>-1</sup>) from each level. We counted between 200-300 pollen grains produced by trees,  
207 shrubs and herbs (terrestrial pollen) for each palynological sample and calculated the  
208 percent abundance of each terrestrial taxon relative to this sum. The percentage of  
209 aquatic plants was calculated in reference to the total pollen sum (terrestrial + aquatic  
210 pollen) and the percentage of ferns from the total pollen and spores sum. Zonation of the  
211 pollen record was aided by a stratigraphically constrained cluster analysis on all terrestrial  
212 pollen taxa having ≥2%, after recalculating sums and percentages.

213 We identified the palynomorphs based on a modern reference collection housed at  
214 the laboratory of Quaternary Paleoecology of Universidad de Chile, along with published



215 descriptions and keys (Heusser, 1971). In most cases the identification was done at family  
216 or genus level, in some cases to the species level (*Podocarpus nubigena*, *Drimys winteri*,  
217 *Gunnera magellanica*, *Lycopodium magellanicum*). The palynomorph *Nothofagus dombeyi*  
218 type includes the species *N. antarctica*, *N. pumilio*, *N. betuloides* and *N. dombeyi*, the  
219 morphotype *Fitzroya/Pilgerodendron* includes the cupressaceous conifers *Fitzroya*  
220 *cupressoides* and *Pilgerodendron uviferum*. We calculated running means of selected  
221 pollen taxa using a triangular weighing function of values along 7 adjacent levels.

222 We tallied microscopic (<120 µm) and macroscopic (>106 µm) charcoal particles to  
223 document regional and local fire events, respectively. Microscopic particles were counted  
224 from each pollen slide, while macroscopic charcoal was counted from 2-cm<sup>3</sup> sediment  
225 samples obtained from 1-cm thick and continuous-contiguous sections. The samples were  
226 prepared using a standard procedure which involves deflocculation in 10% KOH, careful  
227 sieving through 106 and 212 µm-diameter meshes to avoid rupture of individual particles,  
228 followed by visual inspection on a ZEISS KL 1500 LCD stereoscope at 10x magnification.  
229 These results were analyzed by a time-series analysis to detect local fire events using the  
230 CharAnalysis software (Higuera et al., 2009), interpolating samples at regular time interval  
231 based in the median time resolution of the record. We deconvoluted the CHAR signal into  
232 a peaks and background component using a lowess robust to outliers smoothing with a  
233 100-yr window width. We calculated locally defined thresholds to identify statistically  
234 significant charcoal peaks or local fires events (99<sup>th</sup> percentile of a Gaussian distribution).

235

## 236 RESULTS

237

238 The sediment stratigraphy (Figure 2) reveals a basal unit of blue-gray mud between  
239 1726-819 cm, horizontally laminated for the most part, in some sectors massive and  
240 sandier with small amounts of granule and gravel immersed in a clayey matrix (segment  
241 PC0902AT9). These inorganic clays are overlain by organic silt between 819-678 cm and  
242 organic-rich lake mud (gytjja) in the topmost 678 cm. We found laminated authigenic  
243 carbonates between 794-759 and 394-389 cm (range: 5-20%), for the remainder of the

244 record carbonate values are negligible or null (<5%). The record includes 2 tephras  
245 between 630-628 and 661-643 cm, which exhibit sharp horizontal contacts with the over  
246 and underlying mud and, consequently, we interpret them as aerial fallout deposits from  
247 explosive events originated from Volcán Hudson (H1 tephra) and from Volcán Mentolat  
248 (M1 tephra) based on geochemical data, respectively (Stern et al., 2016).

249 The radiocarbon results show an approximately linear increase of age with depth  
250 between 19,000-9000 yr BP (Figure 3) which, in conjunction with the sediment  
251 stratigraphy, suggests undisturbed in-situ pelagic deposition of lake mud and tephras in  
252 the Lago Edita basin. This study focuses on the interval between 19,000-9000 yr BP (Figure  
253 2, Table 1), and consists of 155 contiguous palynological and macroscopic charcoal levels  
254 with a median time step of 65 years between analyzed samples.

255

#### 256 Pollen stratigraphy

257

258 We divided the record in 6 zones to facilitate its description and discussion, based  
259 on conspicuous changes in the pollen stratigraphy and a stratigraphically constrained  
260 cluster analysis (Figure 4). The following section describes each pollen zone indicating the  
261 stratigraphic and chronologic range, and the mean abundance of major taxa in  
262 parenthesis.

263 Zone Edita-1 (795-780 cm; 19,000-18,100 yr BP) is co-dominated by Poaceae (33%)  
264 and *Empetrum* (32%). This zone starts with a gradual increase in *Empetrum*, attaining its  
265 maximum abundance (~53%) at the end of this zone. Asteraceae subfamily Asteroideae  
266 (7%), *Acaena* (4%), Caryophyllaceae (3%) and Cyperaceae (9%) decrease, while Poaceae  
267 shows fluctuations in its abundance between 2-16 % over the entire interval. Other herbs  
268 and shrubs such as Ericaceae (3%), *Phacelia* (~2%), *Valeriana* (1%), *Gunnera magellanica*  
269 (~2%), Apiaceae (<1%), and Asteraceae subfamily Cichorioideae (<1%) remain relatively  
270 steady. The arboreal taxa *N. dombeyi* type (10%), *Fitzroya/Pilgerodendron* (2%), *P.*  
271 *nubigena* (<1%) and *D. winteri* (<1%) are present in low abundance, as well as the ferns *L.*  
272 *magellanicum* (~1%) and *Blechnum* type (5%) and the green-microalgae *Pediastrum* (2%).

273 Zone Edita-2 (780-758 cm; 18,100-16,800 yr BP) begins with a decline in *Empetrum*  
274 (30%) and an increase in Poaceae (34%) followed by its decrease until the end of this zone.  
275 *N. dombeyi* type (15%), Caryophyllaceae (5%) and Asteraceae subfamily Asteroideae (5%)  
276 show a rising trend during this zone, while other arboreal taxa (*Fitzroya/Pilgerodendron*  
277 (3%), *P. nubigena* (<1%) and *D. winteri* (<1%) and most of the herbs maintain similar  
278 abundance to the previous zone. *L. magellanicum* (2%) and *Pediastrum* (4%) rise slightly,  
279 along with high variability in Cyperaceae (7%).

280 Zone Edita-3 (758-701 cm; 16,800-13,200 yr BP) is characterized by a sharp rise in  
281 Poaceae (45%) and declining trend in *Empetrum* (15%). The conifer *P. nubigena* (2%) starts  
282 a sustained increase, while *N. dombeyi* type (13%) and *Fitzroya/Pilgerodendron* (3%)  
283 remain relatively invariant. *D. winteri* (<1%) and *Misodendrum* (<1%), a mistletoe that  
284 grows on *Nothofagus* species, appear in low abundance in an intermittent manner.  
285 *Pediastrum* (30%) shows a rapid increase until 15,600 yr BP, followed by considerable  
286 variations in its abundance until the end of this zone (between 19% and 55%). *L.*  
287 *magellanicum* (3%) shows a steady increase, while *Blechnum* type (6%) remains invariant  
288 and Cyperaceae (7%) exhibits large fluctuations superimposed upon a declining trend.

289 Zone Edita-4 (701-681 cm; 13,200-11,600 yr BP) starts with an increase in *N.*  
290 *dombeyi* type (29%) and a minor rise in *Misodendrum* (1%). *P. nubigena* (5%) starts this  
291 zone with variability and stabilizes toward the end of this zone, concurrent with  
292 *Fitzroya/Pilgerodendron* (3%) and traces of *D. winteri* (<1%). Poaceae (38%) shows a  
293 steady decrease, while *Empetrum* (6%) continues with a declining trend that started  
294 during the previous zone. Asteraceae subfamily Asteroideae (5%) and Caryophyllaceae  
295 (2%) decrease, *L. magellanicum* (3%), Cyperaceae (4%) and *Pediastrum* (24%) decline  
296 gradually with considerable fluctuations, while *Blechnum*- type (11%) shows modest  
297 increases.

298 Zone Edita-5 (681-674 cm; 11,600-11,100 yr BP) shows marked declines in *N.*  
299 *dombeyi* type (27%) and Poaceae (33%) in concert with a conspicuous increase in the  
300 conifers *Fitzroya/Pilgerodendron* (12%) and *P. nubigena* (9%) that reach their peak  
301 abundance in the record. The abundance of herbs and shrubs decreases or remains

302 steady, with the exception of an ephemeral increase in *Phacelia* (3%). *Blechnum* type  
303 (39%) shows a remarkable increase to its peak abundance in the entire record, while *L.*  
304 *magellanicum* (3%), Cyperaceae (8%) and *Pediastrum* (17%) rise slightly.

305 Zone Edita-6 (674-640 cm; 11,100-8940 yr BP) is characterized by an abrupt increase  
306 in *N. dombeyi* type (62%) and *Misodendrum* (2%), along with conspicuous decline in  
307 *Fitzroya/Pilgerodendron* (2%) and *P. nubigena* (2%) at the beginning of this zone. Poaceae  
308 (26%) shows a downward trend over this period, while other herbs and shrubs  
309 (*Empetrum*, Ericaceae, Caryophyllaceae, Asteraceae subfamily Asteroideae, *Acaena*,  
310 *Phacelia*, *Valeriana*, *Gunnera magellanica*, Apiaceae and Asteraceae subf. Cichorioideae)  
311 show their lowest abundance in the record. *Blechnum* type (7%) drops sharply, followed  
312 by a gradual decline in concert with *L. magellanicum* (1%). Cyperaceae (7%) and  
313 *Pediastrum* (6%) show initial declines followed by increases toward the end of this zone.

314

315 Charcoal stratigraphy

316

317 The record from Lago Edita shows absence of macroscopic charcoal particles  
318 between 19,000-14,300 yr BP followed by an increase in charcoal accumulation rate  
319 (CHAR) that led to a variable plateau between 12,000-13,200 yr BP, a 1000-year long  
320 decline, and a sustained increase led to peak abundance at 9700 yr BP. Charcoal values  
321 then declined rapidly to intermediate levels by 9000 yr BP. We note a close  
322 correspondence between the *Nothofagus* abundance (%) and the CHAR suggesting that  
323 charcoal production was highly dependent upon quantity and spatial continuity of coarse  
324 woody fuels in the landscape (Figure 5).

325 Time-series analysis of the macroscopic charcoal record revealed 11 statistically  
326 significant peaks we interpret as local fire events within the Lago Edita watershed (Figure  
327 5). The temporal structure of these events indicates a sequence of millennial-scale peaks  
328 in fire frequency with maxima at 9600, 10,900, 12,000, 13,100, and 14,100 yr BP. We  
329 observe a steady increase in the fire frequency maxima from 14,100 to 10,900 yr BP  
330 (Figure 5).

331

332 DISCUSSION

333 Paleovegetation

334

335         Given the size of Lago Edita (radius ~250 m) its pollen record is adequate to reflect  
336 local vegetation within 600-800 m from the lake's edge. An extra-local component is also  
337 present considering that species of the genus *Nothofagus* also produce large quantities of  
338 pollen grains susceptible to long-distance transport (Heusser, 1989). These attributes  
339 suggest that the Lago Edita fossil pollen record might be a good sensor of the vegetation  
340 located on the western end of Valle Chacabuco and the Lago Cochrane basin. The record  
341 (Figures 4, 6) documents dominance of herbs and shrubs (chiefly Poaceae, *Empetrum*,  
342 Asteraceae, accompanied by Caryophyllaceae, *Acaena*, Ericaceae, *Phacelia*, *Valeriana*, and  
343 Apiaceae in lower abundance) found above the modern treeline and the Patagonian  
344 steppe between 19,000 and 11,000 yr BP, followed by increasing *Nothofagus* we interpret  
345 as the establishment of scrubland (13,200-11,000 yr BP), woodland (11,000-10,500 yr BP)  
346 and forest (10,500-9000 yr BP). Within the interval dominated by non-arboreal taxa we  
347 distinguish an initial phase with abundant *Empetrum* between 19,000-16,800 yr BP,  
348 followed by diversification of the herbaceous assemblage and preeminence of Poaceae  
349 during the interval 16,800-11,000 yr BP (Figures 4, 6). This change is contemporaneous  
350 with a sustained rise of *P. nubigena* and the mistletoe *Misodendrum* coeval with  
351 conspicuous increases in *Lycopodium magellanicum* and the green microalga *Pediastrum*.  
352 We emphasize the continuous presence of the arboreal *Nothofagus* and  
353 *Fitzroya/Pilgerodendron* in low but constant abundance (~15% and ~3%, respectively)  
354 between 19,000-13,000 yr BP, along with traces (<3%) of hygrophilous trees (*Podocarpus*  
355 *nubigena*, *Drimys winteri*) and herbs (*Gunnera magellanica*, *Lycopodium magellanicum*)  
356 accounting, in sum, for a persistent ~25% of the pre-13,200 yr BP pollen record (Figures 4,  
357 6).

358         The mixed palynological assemblage between ~19,400-11,000 yr BP has no modern  
359 analogues in the regional vegetation (Luebert and Plischoff, 2006; Mancini, 2002). Possible

360 explanations for its development involve: (a) downslope migration of High Andean  
361 vegetation driven by snowline and treeline lowering associated with intense glaciation in  
362 the region, coupled with (b) the occurrence of scattered, low-density populations of  
363 hygrophilous trees and herbs along the eastern margin of the PIS during the LGM and T1.  
364 We rule out the alternative explanation that pollen grains and spores of hygrophilous  
365 trees and herbs in Lago Edita represent an advected signal through the Andes from ice-  
366 free humid Pacific sectors harboring these species because: (i) no empirical basis is  
367 currently available for ice-free conditions and occurrence of cold-resistant hygrophilous  
368 taxa along the western Andean slopes or the Pacific coast of central Patagonia during the  
369 LGM; in fact, the oldest minimum limiting dates for ice-free conditions in records from  
370 Taitao Peninsula and the Chonos archipelago yielded ages of  $14,335 \pm 140$  and  $13,560 \pm 125$   
371  $^{14}\text{C}$  yr BP (median age probability [MAP]: 17,458 and 16,345 yr BP), respectively (Haberle  
372 and Bennett, 2004; Lumley and Switsur, 1993); (ii) the appearance of  
373 *Fitzroya/Pilgerodendron* and *Podocarpus nubigena* at  $\sim 15,000$  and  $\sim 14,000$  yr BP,  
374 respectively, occurred 4000-5000 years later in coastal Pacific sites relative to the Lago  
375 Edita record (Figure 7); (iii) background levels of *Nothofagus* between 15-20% in Lago  
376 Edita predate the appearance and expansion of this taxon in coastal Pacific sites and, once  
377 realized, its abundance in Lago Edita did not follow the trend and magnitude observed in  
378 western sites, as expected if the palynological signal in Lago Edita was attributed to long-  
379 distance transport from that source (Figure 7).

380 Previous palynological studies from bogs located east of the central Patagonian  
381 Andes (de Porras et al., 2012; Markgraf et al., 2007) (Mallín Lago Shaman and Mallín  
382 Pollux, Figure 1) interpreted dry conditions prior to  $\sim 12,000$  yr BP, based on the premise  
383 that low abundance of arboreal taxa and preeminence of herbs and shrubs were indicative  
384 of Patagonian Steppe communities. The glacial-to-interglacial vegetation change in those  
385 studies was interpreted as a westward shift of the forest-steppe boundary brought by  
386 lower-than-present SWW influence at  $44^\circ$ - $46^\circ\text{S}$ , followed by a rise in temperature and  
387 precipitation at the end of the last glaciation. In contrast, the Lago Augusta site (located in  
388 Valle Chacabuco  $\sim 7$  km northeast of Lago Edita) (Figure 1) shows a pollen assemblage

389 prior to 15,600 yr BP dominated by high Andean herbs and shrubs, along with taxa  
390 characteristic of hyperhumid environments along the Pacific coasts of central Patagonia  
391 (*Nothofagus*, *Fitzroya/Pilgerodendron*, *Podocarpus nubigena*, *Saxegothaea conspicua*,  
392 *Drimys winteri*, *Dysopsis glechomoides* and the ferns *Blechnum*, Hymenophyllaceae,  
393 *Cystopteris*) (Villa-Martinez et al., 2012). It appears then that floristic elements of modern  
394 Patagonian forests were present in low abundance and in a discontinuous manner along  
395 the eastern flank of the PIS between 44°-47°S. The data from Lago Edita shown in this  
396 paper, along with the results from Lago Augusta, suggest that Valle Chacabuco harbored  
397 cryptic refugia (Bennett and Provan, 2008) of rainforest trees and herbs during the  
398 interval 19,000-11,000 yr BP, therefore the interpretation of lower-than-present  
399 precipitation of SWW origin in previous studies (de Porrás et al., 2012; Markgraf et al.,  
400 2007), is not applicable to the Valle Chacabuco area over this time interval. Plant  
401 colonization of Valle Chacabuco must have started from the LGM limits located east of  
402 Lago Cochrane and followed the shrinking ice masses to the west, once the newly  
403 deglaciated sectors were devoid of glaciolacustrine influence through T1.

404 Declines and virtual disappearance of the cold-resistant hygrophilous trees  
405 *Fitzroya/Pilgerodendron*, *Podocarpus nubigena* and the herbs *Gunnera magellanica* and  
406 *Lycopodium magellanicum* took place at ~11,000 yr BP in the Lago Edita record (Figures 4,  
407 6), in response to a sudden decline in precipitation. These changes were  
408 contemporaneous with a sustained rise in *Nothofagus*, decreases in all other shrubs and  
409 herbs, and a major increase in macroscopic charcoal (Figure 5), signaling an increment in  
410 arboreal cover, higher spatial continuity of coarse fuels and forest fires. We interpret this  
411 arboreal increase and fire-regime shift as driven by warming which might have triggered a  
412 treeline rise and favored the spread/densification of woody species and coarse fuels  
413 (Figures 4, 5, 6). *Nothofagus* forests (~70% abundance) established near Lago Edita  
414 between 10,000-9000 yr BP.

415

416 Glacial recession in Valle Chacabuco and the Lago Cochrane basin

417

418 Stratigraphic and chronologic results from Lago Edita are key for deciphering the evolution  
419 of Valle Chacabuco and for constraining the timing and rates of deglaciation in this core  
420 region of the PIS. Previous studies (Hein et al., 2010) indicate that Valle Chacabuco was  
421 overridden by the Lago Cochrane ice lobe (LCIL) during the LGM and deposited the Río  
422 Blanco moraines ~90 km downstream from Lago Edita, distal to the eastern end of Lago  
423 Cochrane in Argentina (Argentinian name: Lago Pueyrredón). Cosmogenic radionuclide  
424 dating of three main moraine limits by Hein et al. (2010) yielded cosmogenic  $^{10}\text{Be}$   
425 exposure ages, recently recalculated by Kaplan et al. (2011) at ~21,100, ~25,100, and  
426 ~28,700 yr BP. This was followed by glacial recession starting at 19,600±800 yr BP,  
427 formation of Glacial Lake Cochrane (GLC), stabilization and deposition of the Lago  
428 Columna and Lago Posada moraines at 17,600±900 yr BP, ~55 km upstream from the Río  
429 Blanco moraines (Hein et al., 2010; Kaplan et al., 2011) (Figure 1). Further glacial recession  
430 led to the westward expansion and lowering of GLC until the LCIL stabilized and deposited  
431 moraines in Lago Esmeralda between 13,600-12,800 yr BP ~60 km upstream from the  
432 Lago Columna and Lago Posada moraines (Turner et al., 2005). Recession from this  
433 position led to sudden drainage of GLC toward the Pacific Ocean via Río Baker, once the  
434 continuity between the North and South Patagonian icefields was breached by glacial  
435 recession and thinning. These data suggest that Valle Chacabuco may have been ice-free  
436 and devoid of glaciolacustrine influence after ~17,600 yr BP. More recently Boex et al.  
437 (2013) reported a cosmogenic radio nuclide-based reconstruction of vertical profile  
438 changes of the LCIL through the LGM and T1 that reveals deposition of (i) the Sierra  
439 Colorado lower limit by 28,980±1206 yr BP which can be traced to the Río Blanco  
440 moraines, (ii) the highest summits of Cerro Oportus and Lago Columna moraines by  
441 18,966±1917 yr BP, and (iii) the María Elena moraine by 17,088±1542 yr BP. According to  
442 these data Valle Chacabuco may have been ice-free after ~17,000 yr BP.

443 Lago Edita is a closed-basin lake located ~11 km east of the Cerro Tamango summit  
444 along the ridge that defines the southern edge of the Valle Chacabuco watershed (Figure  
445 1). Lacustrine sedimentation in Lago Edita started when ice-free conditions developed in  
446 Valle Chacabuco, as the LCIL snout retreated eastward to a yet unknown position. The



447 Lago Edita cores show 9 meters of blue-gray clays with millimeter-scale laminations,  
448 interrupted by sporadic intervals of massive pebbly mud appreciable in x radiographs and  
449 the LOI<sub>550</sub> record as increases in the inorganic density data (Figure 2). We also found  
450 exposed glaciolacustrine beds and discontinuous fragments of lake terraces in the vicinity  
451 of Lago Edita, attesting for a large lake that flooded Valle Chacabuco in its entirety.  
452 Differential GPS measurements of 570 m.a.s.l. for the Lago Edita surface and 591 m.a.s.l.  
453 for a well-preserved terrace fragment located ~150 m directly south of Lago Edita, provide  
454 minimum-elevation constraints for GLC during this stage. The Lago Augusta site (Villa-  
455 Martínez et al., 2012), located ~7 km northeast of Lago Edita on the Valle Chacabuco floor  
456 at 444 m.a.s.l. (Figure 1), shows 8 meters of basal glaciolacustrine mud (Figure 2) lending  
457 support to our interpretation.

458         Glaciolacustrine sedimentation persisted in Lago Edita and Lago Augusta until the  
459 surface elevation of GLC dropped below 570 and 444 m.a.s.l., respectively, and the closed-  
460 basin lakes developed. The chronology for this event is constrained by statistically  
461 identical AMS dates of 16,250±90 and 16,020±50 <sup>14</sup>C yr BP (UCIAMS-133418 and CAMS-  
462 144454, respectively) (Table 1) from the same level in the basal portion of the organic  
463 sediments in the Lago Edita record; this estimate approaches the timing for the cessation  
464 of glaciolacustrine influence in Lago Augusta, radiocarbon-dated at 16,445±45 <sup>14</sup>C yr BP  
465 (CAMS-144600) (Table 1). Because we observe approximately the same age for the  
466 transition from glaciolacustrine to organic-rich mud in both stratigraphies, we interpret  
467 the weighted mean age of those three dates (16,254±63 <sup>14</sup>C yr BP, MAP: 19,426 yr BP, two  
468 different laboratories) as a minimum-limiting age for ice-free conditions and nearly  
469 synchronous glaciolacustrine regression from elevations 591 and 444 m.a.s.l. in Valle  
470 Chacabuco. We acknowledge that Villa-Martínez et al. (2012) excluded the age of date  
471 CAMS-144600 from the age model of the Lago Augusta palynological record because it  
472 was anomalously old in the context of other radiocarbon dates higher up in the core.

473         Comparison of the radiocarbon-dated stratigraphy from Lago Edita record with the  
474 exposure-age-dated glacial geomorphology from Lago Cochrane/Pueyrredón, Valle  
475 Chacabuco and surrounding mountains reveals the following:

- 476 • The geochronology for the innermost (third) belt of Río Blanco moraines (~21,100  
477 yr BP) (Hein et al., 2010; Kaplan et al., 2011), glacial deposits on the highest  
478 summits of Cerro Oportus and the Lago Columna moraines (18,966±1917 yr BP)  
479 (Boex et al., 2013) are compatible (within error) with the onset of organic  
480 sedimentation in Lago Edita and Lago Augusta at 19,426 yr BP in Valle Chacabuco.  
481 If correct, this implies ~90 km recession of the LCIL from its LGM limit within ~1500  
482 years.
- 483 • Hein et al. (2010)'s dates for the "final LGM limit", Lago Columna and Lago Posada  
484 moraines should be considered as minimum-limiting ages, as well as Boex et al.  
485 (2013)'s chronology for the María Elena moraine. This is because cosmogenic radio  
486 nuclide ages for these landforms postdate the onset of organic sedimentation in  
487 Lago Edita and Lago Augusta, despite being morphostratigraphically distal (older)  
488 than Valle Chacabuco.
- 489 • As shown in Figure 1, Lago Edita is located along a saddle that establishes the  
490 southern limit of the Río Chacabuco catchment and the northern limit of the Lago  
491 Cochrane basin. According to Hein et al. (2010) the drainage divide on the eastern  
492 end of Lago Cochrane/Pueyrredón basin is located at 475 m.a.s.l., therefore the  
493 presence of this perched glacial lake with a surface elevation of 591 m.a.s.l.  
494 requires ice dams located in the Valle Chacabuco and the Lago Cochrane basin.  
495 This suggests that both valleys remained partially ice covered and that enough  
496 glacier thinning and recession early during T1 enabled the development of a  
497 topographically constrained glacial lake that covered Valle Chacabuco up to the  
498 aforementioned saddle.
- 499 • The high stand of GLC at 591 m.a.s.l. lasted for less than 1500 years during the  
500 LGM and was followed by a nearly instantaneous lake-level lowering of at least  
501 ~150 m at ~19,400 yr BP in Valle Chacabuco. The abrupt large-magnitude drainage  
502 event of this "predecessor lake" was recently recognized by Bourgois et al. (2016),  
503 but its chronology, hydrographic and climatic implications have not been  
504 addressed in the Quaternary literature.

505

506 Biogeographic and paleoclimatic implications

507 The persistence of scattered, low-density populations of rainforest trees and herbs  
508 east of the Andes during the LGM and T1 (Figures 4, 6) implies that precipitation delivered  
509 by the SWW must have been substantially higher than at present (680 mm/year measured  
510 in the Cochrane meteorological station). Because local precipitation in western Patagonian  
511 is positively and significantly correlated with low-level zonal winds (Garreaud et al., 2013;  
512 Moreno et al., 2010; Moreno et al., 2014), we propose that the SWW influence at 47°S  
513 was stronger than present between 19,000-11,000 yr BP, in particular between 16,800-  
514 11,000 yr BP. Subsequent increases in arboreal vegetation, chiefly *Nothofagus*, at 13,200  
515 and 11,000 yr BP led to the establishment of forests near Lago Edita between 10,000-9000  
516 yr BP (Figures 4, 6). We interpret these increases as treeline-rise episodes driven by  
517 warming pulses coupled with a decline in SWW strength at 47°S, as suggested by the  
518 disappearance of cold-resistant hygrophilous trees and herbs at ~11,000 yr BP. We  
519 speculate that the warm pulse and decline in SWW influence at ~11,000 yr BP might  
520 account for the abandonment of early Holocene glacier margins in multiple valleys in  
521 central Patagonia (Glasser et al., 2012)

522 Four salient aspects of the Lago Edita record are relevant for deciphering the pattern  
523 and rates of climate change and dispersal routes of the vegetation in Central Patagonia  
524 (47°S) during the last glacial termination (T1):

525 1- Absence of stratigraphically discernable indications of deglacial warming  
526 between 19,400-13,200 yr BP, in contrast to northwestern Patagonian records  
527 (the Canal de la Puntilla and Huelmo sites, Figure 1) (Moreno et al., 2015) which  
528 show that 75-80% of the glacial-interglacial temperature recovery was  
529 accomplished between 17,800-16,800 yr BP (Figure 8). The record from Lago  
530 Stibnite (46°26'S, 74°25'W), located in central-west Patagonia upwind from the  
531 PIS and Lago Edita (Figure 1), shows a rapid increase in arboreal pollen from ~2%  
532 to >80% in less than 1000 years starting at 16,200 yr BP (Figure 8). We posit that

533 cold glacial conditions lingered along the periphery of the shrinking PIS during  
534 T1, affecting adjacent downwind sectors such as Valle Chacabuco. According to  
535 Turner et al. (2005) the LCIL stabilized and deposited moraines in Lago  
536 Esmeralda, located ~10 km upstream along the glacier flowline and ~240 m  
537 lower in elevation than Lago Edita, between 13,600-12,800 yr BP. We propose  
538 that the climatic barrier for arboreal expansion vanished in downwind sectors  
539 such as Valle Chacabuco once glacial recession from the Lago Esmeralda (Figure  
540 1) margin breached the continuity of the North and South Patagonian icefields  
541 along the Andes. Thus, we propose that regional cooling induced by the PIS  
542 along its eastern margin through T1 accounts for the delayed warming in Valle  
543 Chacabuco relative to records located in western and northwestern sectors  
544 (Figure 8).

545 2- Cold and wet conditions prevailed between 19,400-16,800 yr BP, followed by an  
546 increase in precipitation at 16,800 yr BP. The latter event is contemporaneous  
547 with the onset of a lake-level rise in Lago Lepu  (43 S, central-east Isla Grande  
548 de Chilo ) (Figure 8), which Pesce & Moreno (2014) interpreted as a northward  
549 shift of the SWW as they recovered from a prominent southward shift from  
550 latitudes ~41 -43 S (Figure 8) following the onset of T1 (Moreno et al., 2015).

551 3- Significant ice recession (~90 km) from the eastern LGM margin of the Lago  
552 Cochrane Ice lobe (LCIL) was accomplished between ~21,000-19,400 yr BP, at  
553 times when northwestern Patagonian piedmont glacier lobes experienced  
554 moderate recession during the Varas interstade (Denton et al., 1999; Moreno et  
555 al., 2015) (Figure 8). In contrast to the LCIL, northwestern Patagonian piedmont  
556 glacier lobes readvanced to their youngest glacial maximum position during a  
557 cold episode between 19,300-17,800 yr BP that featured stronger SWW  
558 influence at 41 -43 S (Moreno et al., 2015) (Figure 8). One explanation for this  
559 latitudinal difference might be that northward-shifted SWW between 19,300-  
560 17,800 yr BP fueled glacier growth in northwestern Patagonia while reducing the

561 delivery of moisture to central Patagonia, causing the LCIL to continue the  
562 recession it had started during the Varas interstade.

563 4- A mosaic of cold-resistant and hygrophilous trees and herbs, currently found  
564 along the humid western slopes of the Andes of central Chilean Patagonia, and  
565 cold-resistant shrubs and herbs common to high-Andean and Patagonian steppe  
566 communities developed along the eastern margin of the PIS during the LGM and  
567 T1 (Figures 4, 6). We posit that glacial withdrawal and drainage of GLC through  
568 T1 provided a route for the westward dispersal of hygrophilous trees and herbs,  
569 contributing to the forestation of the newly deglaciated sectors of central-west  
570 Patagonia.

571 We conclude that warm pulses at 13,200 and 11,000 yr BP and a decline in SWW  
572 influence at 47°S starting at 11,000 yr BP brought T1 to an end in central-west Patagonia.  
573 The earliest of these events overlaps in timing with the culmination of Patagonian (Garcia  
574 et al., 2012; Moreno et al., 2009; Strelin et al., 2011; Strelin and Malagnino, 2000) and  
575 New Zealand glacier advances during the Antarctic Cold Reversal. Our data suggest that  
576 the subsequent warm pulse, which was accompanied by a decline in SWW strength at  
577 11,000 yr BP (Moreno et al., 2010; Moreno et al., 2012), was the decisive event that led to  
578 the end of T1 in the study area.

579

#### 580 ACKNOWLEDGEMENTS

581 This study was funded by Fondecyt #1080485, 1121141, ICM grants P05-002 and  
582 NC120066, and a CONICYT M.Sc. Scholarship to W.I.H. We thank E.A. Sagredo, O.H. Pesce,  
583 E. Simi, and I. Jara for assistance during field work, K.D. Bennett and S. Haberle for sharing  
584 published palynological data. We thank C. Saucedo from Agencia de Conservación  
585 Patagónica for permission to work and collect samples in Hacienda Valle Chacabuco  
586 (Parque Patagonia).

587

588

589 FIGURE AND TABLE CAPTIONS

590 Table 1. Radiocarbon dates from the Lago Edita core. The radiocarbon dates were  
591 calibrated to calendar years before present using the CALIB 7.0 program.

592

593 Figure 1. Sketch map of the study area showing the location of central-west Patagonia, the  
594 position of Valle Chacabuco relative to the Río Blanco, María Elena, Lago Columna (LC) and  
595 Lago Posada (LP) ice limits east Lago of Cochrane, and the North Patagonian icefield and  
596 Peninsula Taitao to the west. We also included Sierra Colorado, Lago Esmeralda and Cerro  
597 Oportus for reference. The lower portion of the figure shows a detail on the Cerro  
598 Tamango area and the portion of Valle Chacabuco where Lago Edita and Lago Augusta are  
599 located. Also shown are palynological sites discussed in the main text (Canal de la Puntilla,  
600 Huelmo, Mallín Lago Shaman, Mallín Pollux, Lago Stibnite, Lago Augusta).

601

602 Figure 2. Stratigraphic column, radiocarbon dates and loss-on-ignition data from the Lago  
603 Edita record. The labels on the right indicate the identity and stratigraphic span (dashed  
604 horizontal lines) of each core segment.

605

606 Figure 3. Age model of the Lago Edita record, the blue zones represent the probability  
607 distribution of the calibrated radiocarbon dates, the grey zone represents the calculated  
608 confidence interval of the Bayesian age model.

609

610 Figure 4. Percentage pollen diagrams from the Lago Edita core. The labels on the right  
611 indicate the identity and stratigraphic span (dashed horizontal lines) of each pollen  
612 assemblage zone. The black dots indicate presence of *Drimys winteri* pollen grains,  
613 normally <2%.

614

615 Figure 5. Macroscopic charcoal record from the Lago Edita core and results of  
616 CharAnalysis: blue line: background component, red line: locally defined threshold,

617 triangles: statistically significant charcoal peaks, magnitude: residual abundance that  
618 supersedes the threshold.

619

620 Figure 6. Selected palynomorph abundance of the Lago Edita record shown in the time  
621 scale domain. The red lines correspond to weighted running means of seven adjacent  
622 samples with a triangular filter. The taxa shown in the left panel are characteristic of  
623 humid environments currently found in sectors adjacent to the Pacific coast and/or the  
624 Andean treeline in the study area. The taxon *Nothofagus dombeyi* type, which includes  
625 multiple species with contrasting climatic tolerances, is also found in (relatively) humid  
626 sectors east of the Andes. The herbs and shrubs shown in the right panel are either  
627 cosmopolitan or present in the Patagonian Steppe and sectors located at or above the  
628 Andean treeline in central-west Patagonia.

629

630 Figure 7. Comparison of selected tree pollen recorded in Lago Fácil, Lago Oprasa, Lago  
631 Stibnite (Lumley and Switsur, 1993) and Lago Edita. The red line corresponds to a  
632 weighted running mean in each record of seven adjacent samples with a triangular filter.  
633 The lower panels show the curves from all sites expressed in a common percent scale  
634 (Lago Fácil: purple line, Lago Oprasa: blue line, Lago Stibnite: black line, and Lago Edita:  
635 red line).

636

637 Figure 8. Comparison of the percent sum of arboreal pollen (AP) in records from Lago  
638 Edita, Lago Stibnite (Lumley and Switsur, 1993) and the spliced Canal de la Puntilla-  
639 Huelmo time series (Moreno et al., 2015), as proxies for local rise in treeline driven by  
640 deglacial warming. These data are compared against the  $\delta$  Deuterium record from the  
641 Antarctic Epica Dome Concordia (EDC) ice core (Stenni et al., 2010), and hydrologic  
642 estimates from northwestern Patagonia. The latter consist of the percent abundance of  
643 Magellanic Moorland species found in the spliced Canal de la Puntilla-Huelmo record  
644 (Moreno et al., 2015), indicative of a hyperhumid regime, and the percent abundance of  
645 the littoral macrophyte *Isoetes savatieri* from Lago Lepué (Pesce and Moreno, 2014),

646 indicative of low lake level (LL) during the earliest stages of T1 and the early Holocene  
647 (9000-11,000 yr BP). The vertical dashed lines constrain the timing of the early Holocene  
648 SWW minimum at 41°-43°S (9000-11,000 yr BP) (Fletcher and Moreno, 2011), a low-  
649 precipitation phase during the early termination at 41°-43°S (16,800-17,800 yr BP)  
650 associated with a southward shift of the SWW (Pesce and Moreno, 2014), the final LGM  
651 advance of piedmont glacier lobes (17,800-19,300 yr BP) and the final portion of the Varas  
652 interstade (19,300-21,000 yr BP) in the Chilean Lake District (Denton et al., 1999; Moreno  
653 et al., 2015). The dashed green horizontal lines indicate the mean AP of each pollen record  
654 prior to their increases during T1 (Lago Edita: 17%, Lago Stibnite:2%, spliced Canal de la  
655 Puntilla-Huelmo: 31%). The ascending oblique arrow represents a northward shift of the  
656 SWW, the descending arrow a southward shift of the SWW at the beginning of T1.  
657  
658



659 REFERENCES CITED

- 660 Bennett, K.D., Provan, J., 2008. What do we mean by 'refugia'? *Quaternary Science Reviews* 27,  
661 2449-2455.
- 662 Blaauw, M., Christen, J.A., 2011. Flexible Paleoclimate Age-Depth Models Using an Autoregressive  
663 Gamma Process. *Bayesian Analysis* 6, 457-474.
- 664 Boex, J., Fogwill, C., Harrison, S., Glasser, N.F., Hein, A., Schnabel, C., Xu, S., 2013. Rapid thinning of  
665 the late Pleistocene Patagonian Ice Sheet followed migration of the Southern Westerlies.  
666 *Scientific reports* 3.
- 667 Bourgeois, J., Cisternas, M.E., Braucher, R., Bourlès, D., Frutos, J., 2016. Geomorphic Records along  
668 the General Carrera (Chile)–Buenos Aires (Argentina) Glacial Lake (46°–48°S), Climate  
669 Inferences, and Glacial Rebound for the Past 7–9 ka. *The Journal of Geology* 124, 27-53.
- 670 Bromwich, D.H., Toracinta, E.R., Oglesby, R.J., Fastook, J.L., Hughes, T.J., 2005. LGM Summer  
671 Climate on the Southern Margin of the Laurentide Ice Sheet: Wet or Dry? *Journal of Climate*  
672 18, 3317-3338.
- 673 Bromwich, D.H., Toracinta, E.R., Wei, H., Oglesby, R.J., Fastook, J.L., Hughes, T.J., 2004. Polar MM5  
674 Simulations of the Winter Climate of the Laurentide Ice Sheet at the LGM. *Journal of Climate*  
675 17, 3415-3433.
- 676 Caniupán, M., Lamy, F., Lange, C.B., Kaiser, J., Arz, H., Kilian, R., Baeza Urrea, O., Aracena, C.,  
677 Hebbeln, D., Kissel, C., Laj, C., Mollenhauer, G., Tiedemann, R., 2011. Millennial-scale sea  
678 surface temperature and Patagonian Ice Sheet changes off southernmost Chile (53°S) over  
679 the past ~60 kyr. *Paleoceanography* 26, n/a-n/a.
- 680 de Porras, M.E., Maldonado, A., Abarzúa, A.M., Cárdenas, M.L., Francois, J.P., Martel-Cea, A.,  
681 Stern, C.R., Méndez, C., Reyes, O., 2012. Postglacial vegetation, fire and climate dynamics at  
682 Central Chilean Patagonia (Lake Shaman, 44°S). *Quaternary Science Reviews* 50, 71-85.
- 683 Denton, G.H., Lowell, T.V., Heusser, C.J., Schluchter, C., Andersen, B.G., Heusser, L.E., Moreno, P.I.,  
684 Marchant, D.R., 1999. Geomorphology, stratigraphy, and radiocarbon chronology of  
685 Llanquihue drift in the area of the southern Lake District, Seno Reloncavi, and Isla Grande de  
686 Chiloe, Chile. *Geografiska Annaler Series a-Physical Geography* 81A, 167-229.
- 687 Faegri, K., Iversen, J., 1989. *Textbook of pollen analysis*. John Wiley & Sons.
- 688 Fletcher, M.S., Moreno, P.I., 2011. Zonally symmetric changes in the strength and position of the  
689 Southern Westerlies drove atmospheric CO<sub>2</sub> variations over the past 14 k.y. *Geology* 39,  
690 419-422.
- 691 Gajardo, R., 1994. *La Vegetación Natural de Chile. Clasificación y Distribución Geográfica*. Editorial  
692 Universitaria, Santiago, Chile.
- 693 Garcia, J.L., Kaplan, M.R., Hall, B.L., Schaefer, J.M., Vega, R.M., Schwartz, R., Finkel, R., 2012.  
694 Glacier expansion in southern Patagonia throughout the Antarctic cold reversal. *Geology* 40,  
695 859-862.
- 696 Garreaud, R., Lopez, P., Minvielle, M., Rojas, M., 2013. Large-Scale Control on the Patagonian  
697 Climate. *Journal of Climate* 26, 215-230.
- 698 Glasser, N.F., Harrison, S., Schnabel, C., Fabel, D., Jansson, K.N., 2012. Younger Dryas and early  
699 Holocene age glacier advances in Patagonia. *Quaternary Science Reviews* 58, 7-17.
- 700 Glasser, N.F., Jansson, K.N., Duller, G.A.T., Singarayer, J., Holloway, M., Harrison, S., 2016. Glacial  
701 lake drainage in Patagonia (13-8 kyr) and response of the adjacent Pacific Ocean. *Scientific*  
702 *Reports* 6, 21064.
- 703 Haberle, S.G., Bennett, K.D., 2004. Postglacial formation and dynamics of North Patagonian  
704 Rainforest in the Chonos Archipelago, Southern Chile. *Quaternary Science Reviews* 23, 2433-  
705 2452.

706 Hall, B.L., Porter, C.T., Denton, G.H., Lowell, T.V., Bromley, G.R.M., 2013. Extensive recession of  
707 Cordillera Darwin glaciers in southernmost South America during Heinrich Stadial 1.  
708 Quaternary Science Reviews 62, 49-55.

709 Hein, A.S., Hulton, N.R.J., Dunai, T.J., Sugden, D.E., Kaplan, M.R., Xu, S., 2010. The chronology of  
710 the Last Glacial Maximum and deglacial events in central Argentine Patagonia. Quaternary  
711 Science Reviews 29, 1212-1227.

712 Heiri, O., Lotter, A.F., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and  
713 carbonate content in sediments: reproducibility and comparability of results. Journal of  
714 Paleolimnology 25, 101-110.

715 Heusser, C.J., 1971. Pollen and Spores from Chile. University of Arizona Press, Tucson.

716 Heusser, C.J., 1989. Late Quaternary Vegetation and Climate of Southern Tierra-Del-Fuego.  
717 Quaternary Research 31, 396-406.

718 Higuera, P.E., Brubaker, L.B., Anderson, P.M., Hu, F.S., Brown, T.A., 2009. Vegetation mediated the  
719 impacts of postglacial climate change on fire regimes in the south-central Brooks Range,  
720 Alaska. Ecological Monographs 79, 201-219.

721 Kaplan, M.R., Strelin, J.A., Schaefer, J.M., Denton, G.H., Finkel, R.C., Schwartz, R., Putnam, A.E.,  
722 Vandergoes, M.J., Goehring, B.M., Travis, S.G., 2011. In-situ cosmogenic  $^{10}\text{Be}$  production  
723 rate at Lago Argentino, Patagonia: Implications for late-glacial climate chronology. Earth and  
724 Planetary Science Letters 309, 21-32.

725 Kaufman, D.S., Ager, T.A., Anderson, N.J., Anderson, P.M., Andrews, J.T., Bartlein, P.J., Brubaker,  
726 L.B., Coats, L.L., Cwynar, L.C., Duvall, M.L., Dyke, A.S., Edwards, M.E., Eisner, W.R., Gajewski,  
727 K., Geirsdottir, A., Hu, F.S., Jennings, A.E., Kaplan, M.R., Kerwin, M.N., Lozhkin, A.V.,  
728 MacDonald, G.M., Miller, G.H., Mock, C.J., Oswald, W.W., Otto-Bliesner, B.L., Porinchu, D.F.,  
729 Ruhland, K., Smol, J.P., Steig, E.J., Wolfe, B.B., 2004. Holocene thermal maximum in the  
730 western Arctic (0-180 degrees W). Quaternary Science Reviews 23, 529-560.

731 Lara, A., Villalba, R., Wolodarsky-Franke, A., Aravena, J.C., Luckman, B.H., Cuq, E., 2005. Spatial and  
732 temporal variation in *Nothofagus pumilio* growth at tree line along its latitudinal range (35  
733 degrees 40 ' -55 degrees S) in the Chilean Andes. Journal of Biogeography 32, 879-893.

734 Luebert, F., Plischoff, P., 2006. Sinopsis Bioclimática y Vegetacional de Chile. Editorial Universitaria,  
735 Santiago, Chile.

736 Lumley, S., Switsur, R., 1993. Late Quaternary chronology of the Taitao Peninsula, Southern Chile. .  
737 Journal of Quaternary Science 8, 161-165.

738 Mancini, M.V., 2002. Vegetation and climate during the holocene in Southwest Patagonia,  
739 Argentina. Review of Palaeobotany and Palynology 122, 101-115.

740 Markgraf, V., Whitlock, C., Haberle, S., 2007. Vegetation and fire history during the last 18,000 cal  
741 yr BP in Southern Patagonia: Mallin Pollux, Coyhaique, Province Aisen (45 degrees 41 ' 30 ''  
742 S, 71 degrees 50 ' 30 '' W, 640 m elevation). Palaeogeography Palaeoclimatology  
743 Palaeoecology 254, 492-507.

744 Moreno, P.I., Denton, G.H., Moreno, H., Lowell, T.V., Putnam, A.E., Kaplan, M.R., 2015.  
745 Radiocarbon chronology of the last glacial maximum and its termination in northwestern  
746 Patagonia. Quaternary Science Reviews 122, 233-249.

747 Moreno, P.I., Francois, J.P., Villa-Martínez, R., Moy, C.M., 2010. Covariability of the Southern  
748 Westerlies and atmospheric CO<sub>2</sub> during the Holocene. Geology 39, 727-730.

749 Moreno, P.I., Kaplan, M.R., Francois, J.P., Villa-Martinez, R., Moy, C.M., Stern, C.R., Kubik, P.W.,  
750 2009. Renewed glacial activity during the Antarctic cold reversal and persistence of cold  
751 conditions until 11.5 ka in southwestern Patagonia. Geology 37, 375-378.

752 Moreno, P.I., Vilanova, I., Villa-Martínez, R., Garreaud, R.D., Rojas, M., De Pol-Holz, R., 2014.  
753 Southern Annular Mode-like changes in southwestern Patagonia at centennial timescales  
754 over the last three millennia. *Nat Commun* 5.

755 Moreno, P.I., Villa-Martinez, R., Cardenas, M.L., Sagredo, E.A., 2012. Deglacial changes of the  
756 southern margin of the southern westerly winds revealed by terrestrial records from SW  
757 Patagonia (52 degrees S). *Quaternary Science Reviews* 41, 1-21.

758 Pesce, O.H., Moreno, P.I., 2014. Vegetation, fire and climate change in central-east Isla Grande de  
759 Chiloé (43°S) since the Last Glacial Maximum, northwestern Patagonia. *Quaternary Science*  
760 *Reviews* 90, 143-157.

761 Pisano, E., 1997. Los bosques de Patagonia Austral y Tierra del Fuego chilenas. *Anales del Instituto*  
762 *de la Patagonia, Serie Ciencias Naturales (Chile)* 25, 9:19.

763 Schmithüsen, J., 1956. Die raumliche Ordnung der chilenischen Vegetation. *Bonner Geographische*  
764 *Abhandlungen* 17, 1-86.

765 Siani, G., Michel, E., De Pol-Holz, R., DeVries, T., Lamy, F., Carel, M., Isguder, G., Dewilde, F.,  
766 Lourantou, A., 2013. Carbon isotope records reveal precise timing of enhanced Southern  
767 Ocean upwelling during the last deglaciation. *Nat Commun* 4.

768 Stenni, B., Masson-Delmotte, V., Selmo, E., Oerter, H., Meyer, H., Rothlisberger, R., Jouzel, J.,  
769 Cattani, O., Falourd, S., Fischer, H., Hoffmann, G., Iacumin, P., Johnsen, S.J., Minster, B.,  
770 Udisti, R., 2010. The deuterium excess records of EPICA Dome C and Dronning Maud Land  
771 ice cores (East Antarctica). *Quaternary Science Reviews* 29, 146-159.

772 Stern, C., 2004. Active Andean volcanism: its geologic and tectonic setting. *Revista Geológica de*  
773 *Chile* 31, 161-206.

774 Stern, C.R., Moreno, P.I., Henriquez, W.I., Villa-Martinez, R., Sagredo, E., Aravena, J.C., De Pol-Holz,  
775 R., 2016. Holocene tephrochronology around Cochrane (~47°S), southern Chile. *Andean*  
776 *Geology* 43, 1-19.

777 Strelin, J.A., Denton, G.H., Vandergoes, M.J., Ninnemann, U.S., Putnam, A.E., 2011. Radiocarbon  
778 chronology of the late-glacial Puerto Bandera moraines, Southern Patagonian Icefield,  
779 Argentina. *Quaternary Science Reviews* 30, 2551-2569.

780 Strelin, J.A., Malagnino, E.C., 2000. Late-Glacial History of Lago Argentino, Argentina, and Age of  
781 the Puerto Bandera Moraines. *Quaternary Research* 54, 339-347.

782 Sugita, S., 1994. Pollen Representation of Vegetation in Quaternary Sediments: Theory and  
783 Method in Patchy Vegetation. *Journal of Ecology* 82, 881-897.

784 Turner, K.J., Fogwill, C.J., McCulloch, R.D., Sugden, D.E., 2005. Deglaciation of the eastern flank of  
785 the North Patagonian Icefield and associated continental-scale lake diversions. *Geografiska*  
786 *Annaler: Series A, Physical Geography* 87, 363-374.

787 Villa-Martinez, R., Moreno, P.I., Valenzuela, M.A., 2012. Deglacial and postglacial vegetation  
788 changes on the eastern slopes of the central Patagonian Andes (47 degrees S). *Quaternary*  
789 *Science Reviews* 32, 86-99.

790

## 791 Table 1


Laboratory code	Core	Material	Length (cm)	$^{14}\text{C}$ yr BP $\pm 1\sigma$	Median probability (cal yr BP)	2 $\sigma$ range (cal BP)
UCIAMS-133501	PC0902AT7	Bulk	660-661	8935 $\pm$ 25	10,029	9794-10,177
UCIAMS-133416	PC0902AT8	Bulk	705-706	11,350 $\pm$ 60	13,229	13,109-13,350
UCIAMS-133417	PC0902AT8	Bulk	757-758	13,740 $\pm$ 70	16,863	16,684-17,055
UCIAMS-133418	PC0902AT8	Bulk	795-796	16,250 $\pm$ 90	19,414	18,934-19,779
CAMS-144454	PC0902BT8	Bulk	795-796	16,020 $\pm$ 50	19,164	18,922-19,408

792

793

794 Figure 1


795



796

797


798 Figure 2



799


800

801 Figure 3




802  
803

804 Figure 4




805



806  
807




808 Figure 5




809

810 Figure 6



811


812 Figure 7



813

814

815 Figure 8



816