

8 November 2017

Dr. Nathalie Combourieu-Nebout, Editor Climate of the Past  
Resubmission of Manuscript (Ref. No. cp-2017-104)

Dear Nathalie,

Thank so much for considering our manuscript entitle “*Holocene aridification trend and human impact interrupted by millennial- and centennial-scale climate fluctuations from a new sedimentary record from Padul (Sierra Nevada, southern Iberian Peninsula)*”, authored by María J. Ramos-Román, Gonzalo Jiménez-Moreno, Jon Camuera, Antonio García-Alix, R. Scott Anderson, Francisco J. Jiménez-Espejo, José S. Carrión, for publication in Climate of the Past. Here we are submitting a major revision of our previously submitted manuscript (cp-2017-104).

We have gone in detail over all the very constructive suggestions that the three reviewers made, which improved our manuscript to a much stronger paper. This revised manuscript complements point by point responses to the general suggestions and comments of the referees.

The most important issue that reviewers had was disentangling human vs. climate as the main forcing shaping the environment in the area. We made significant changes to the manuscript in this direction adding more discussion about trying to separate human induced changes in the environment. In this respect we added a new Figure 8, which clarifies and separates as much as we can evidences of natural vs. anthropic signals. In addition, the title was revised as some of the reviewers requested to include the importance of human activity in the area, notably for the last ~1550 yr. We have also made some general editorial changes thorough the manuscript for improve the readability.

We hope you and the reviewers find merit in this work, and look forward to your comments.

Yours truly,

María J. Ramos-Román  
Departamento de Estratigrafía y Paleontología  
Universidad de Granada, Spain

## **Author's response to reviewers:**

Firstly, we would like to thank to the three reviewers comments and suggestion, which improved the manuscript, and for recommending this study for publication in *Climate of the Past*.

Our manuscript has been reviewed by three referees. One of them, Graciela Gil-Romera was satisfied with our response to her comments in the discussion process. However, here we added again all the answers to her questions so we show all the changes we have made in this new version of the manuscript.

The referee suggestions and comments are displayed in black, and our answers are in blue. We indicate the lines in which we made the modifications according to the marked-up manuscript version (see below; in **red** text that was deleted, in **blue** text that was modified and in **green** text that was moved). In the case that we do not follow the reviewer suggestions we discussed the reasons.

## **Author's response to Graciela Gil-Romera:**

### **General**

I find that the manuscript led by Ramos-Roman is a useful contribution to understanding the recent palaeoenvironments of an otherwise, poorly studied region of Southern Europe. The study presents a multiproxy analysis of Late Holocene change from the well known record of Padul. The main objective of the paper is to distinguish climate from human action driving landscape dynamics. The age model is coherent and well built, and the subsequent time series analyses performed on geochemistry and pollen is sound and within the accuracy of the age-depth model. I attach an edited version of the pdf with some, minor, comments and then I have a couple of wider suggestions:

### **General suggestions for the review**

1. Despite the main question posed in this work is essential to the paleosciences (human vs climate driven changes) quite often it's impossible to tell which is the main driving factor as they superimpose. This is also kind of patent in the study led Ramos-Roman and cols. where despite all proxies is difficult to detangle these effects. I would probably include the "human" factor in the title as it includes a large part of the discussion.
2. Despite authors present several proxies and connect with other terrestrial and marine records they lack charcoal as a proxy of fire occurrence. Considering the sampling has been done continuously, adding charcoal as a proxy may illustrate postfire responses of vegetation that are now been attributed to climate or human activities indirectly. Likewise

1 it may help understanding the human-climate dialectic. Please do have a look to my  
2 comments in the attached documents (minor corrections and comments). Let me know if  
3 the document can't be accessed for any reason. I'd be pleased in seeing this study published  
4 in Climate of the Past.

5 **Author's response to general suggestions:**

- 6 1. We agree with your comment about the importance of human impact on the environments  
7 during the Late Holocene. In this record human impact doesn't seem to be very strong until  
8 the last 1.5 cal ka BP and that is the reason why we mostly focused on understanding the  
9 climate influence on the environments during the Late Holocene in the southern Iberian  
10 Peninsula. Anyhow, we agree in that human impact is also important and should appear in  
11 the title so we changed it for "Holocene aridification trend and human impact interrupted  
12 by millennial- and centennial-scale climate fluctuations from a new sedimentary record  
13 from Padul (Sierra Nevada, southern Iberian Peninsula)".
- 14 2. The suggestion of including the charcoal analysis from this sediment record to this  
15 manuscript is a great idea but this is the Masters thesis study that Cole Webster is carrying  
16 out at present under supervision of R. Scott Anderson at NAU, Arizona, USA. So even if  
17 we understand how important is to have a fire-proxy record to compare with the vegetation  
18 and sedimentation dynamics we do not have these data yet.

19  
20 **Minor comments insert in the pdf**

- 21 1. line 63: Gil-Romera et al., 2014 is not highlighting any aridity trend at that time.

22  
23 Ok, we deleted that one and inserted a correct reference ([line 69](#)).

- 24  
25 2. line 80: quote which ones.

26  
27 Thanks. We moved the references to the correct place ([line 89](#)).

- 28 3. line 98: it'd help to have a vegetation profile here summarising the communities that you  
29 explain below, so you don't have to go in such detail with taxa that you can't actually find later  
30 on in your pollen spectra.

31 Ok, this is a good idea. We summarized the vegetation to the principal species present in this  
32 vegetation belts, related with our pollen results.

33 [Modifications in lines 137 to 154.](#)

- 34  
35 4. line 216: can you please state how many samples did you analyze in the end? are these 115?

36 The total number of samples analyzed was 103. This number does not correspond to the 115  
37 cm-long sediment record studied because there was some compaction. Thanks for noticing, we  
38 also clarified this in the manuscript.

[Modifications in lines 266-267.](#)

5. line 234: Please detail average sedimentation rate for core Padul 15-05

Ok, we detailed the average sedimentation rate.

[Modifications in line 287.](#)

6. line 238: as far as I can see you have not done linear interpolation. Indeed you explain that you did smooth splines in Clam?? Please correct inconsistency

Yes, we used a smooth spline age model. However, the sedimentation rates were calculated by lineal interpolation between the radiocarbon dates. We agree in that they have to be consistent so we modified this SAR in Figure 2 and we now show the average SAR between radiocarbon dates using the modeled ages.

[Modifications in lines 289-290.](#)

“Six distinct sediment accumulation rate (SAR) intervals can be differentiated between 0 and 122.96 cm between radiocarbon dates in the studied core”

7. line 409-415: I see the aridification argument as a more likely explanation than a real summer cooling due to reduced insolation. If anything, at these latitudes, that would help the forest reducing evapotranspiration. In other northern mountain systems as the alps and the reduced summer insolation does not seem to affect the forest during the Mid-late Holocene (Perez et al., 2013, Leunda et al., 2017) so I would see as more feasible the second aridification trend due to the westerlies moving northwards than the reduced growing season due to cooling.

Yes, we agree and also think that aridification was the main trigger of this vegetation change but we cannot underestimate the effect that summer insolation (perhaps very little) might have caused in the vegetation with a reduction of the growing season.

[We try to clarify this point over the manuscript in lines 502-508.](#)

## **Author’s response to anonymous reviewer 2:**

### **General**

This manuscript is quite interesting, bringing lights on a period not yet explored in this notorious site and worthy to be published. Nevertheless I suspect the authors to partly over-interpret their datasets in order to demonstrate that they confirm a model of late-Holocene climate changes identified in other western Mediterranean sites (reinforcement syndrom). Here are my remarks along the text:



1 Conclusion: A further study on the micro charcoal could trace the rhythms in aridification (or  
2 land-use). I invite the authors to clarify some elements of their demonstration.

3  
4 Thanks for your suggestions. In this study we found a linear increasing trend in deforestation  
5 very similar to the decrease in insolation and we do not find clear evidences (with any of the  
6 studied proxies) of humans impacting the area until the last ca. 1500 years so we cannot really  
7 assume that humans were also contributing to that decrease in forest species in the area.  
8 Therefore we related that deforestation mostly to climate. In addition, there seems to be a  
9 correlation of our record with global climate variability, which also supports our suggestion.  
10 In the new version of the manuscript we inserted a new figure 8 (see below in the new  
11 manuscript version) trying to clarify and separate natural vs. anthropic signals. Considering the  
12 importance of human activity in the area, notably in the top part of the sedimentary record, we  
13 changed the title for “Holocene aridification trend and human impact interrupted by millennial-  
14 and centennial-scale climate fluctuations from a new sedimentary record from Padul (Sierra  
15 Nevada, southern Iberian Peninsula)”.

16 The suggestion of including the charcoal analysis from this sediment record to this manuscript  
17 is a great idea but this is the Masters thesis study that Cole Webster is carrying out at present  
18 under supervision of R. Scott Anderson at NAU, Arizona, USA. So even if we understand  
19 how important is to have a fire-proxy record to compare with the vegetation and sedimentation  
20 dynamics we do not have these data yet.

## 21 **Comments**

22 1/ Study site: the information on the geological context (lines: 127-133) could be (at least  
23 partly) presented before the short description of the regional vegetation.

24 Ok, this is a good idea. We moved the part of the specific vegetation description from Padul  
25 right after the geological setting so it reads better.

26 Text moved to lines 164-173.

27 The description of the “vegetation in the Padul area” (line 194) seems to concern potential  
28 vegetation as, looking on “google earth” I observe that the surrounding of the site is on large  
29 almost totally cultivated: this agricultural activity (and maybe its historical) could be described.  
30 Concerning the vegetation on and around the lake, the floristic list, is a description of different  
31 assemblages possible? One of them could constitute a modern analogue to the pollen spectra  
32 in PAZ 3 and 4a? *Study site:*

33 Ok, we believe that the reviewer refers to line 154 - about the “vegetation in the Padul area”.  
34 The Padul area was cultivated since historical times as we indicated in the text (line 776-777).  
35 We agree with the reviewer that this information could be better presented and we added some  
36 more information about the agricultural activities in the Padul area at present. The species  
37 cultivated in the past might have changed through time but there are not historical records  
38 available for a deep review and our pollen data could provide with a clearer image of what

happened in the past. See modifications in section of “human activities” concerning Cichorioideae and other nitrophilous taxa that could have possibly related with livestock or agricultural activities.

Modifications about information for agricultural activities [in lines 213-215](#). Modifications about human activities [in lines 752-760 and 773-777](#).

A paper by Pons and Reille (1988) mentions a peat exploitation: It would be of interest to know the connection between this exploitation and its potential impact on the lake beside the coring point.

We do not quite understand the question. The Padul-15-05 core was drilled nearby but outside the peat exploitation area, and thus in a different location than the previous cores (e.g. Pons and Reille 1988; Ortiz et al., 2004) that were taken directly from the peat mine. This is the reason why the first (Late Holocene) part of the sediment record was not previously studied as it was lacking from that area.

[This information is given in lines 88-92.](#)

2/ Line 218: Lycopodium tablets are mentioned here but the changes in pollen concentrations not discussed later? It would be of interest to know if the transition from unit 1 to Unit 2 is marked by a major change in pollen concentrations?

Yes, thank you for the suggestions. We inserted the pollen concentration in the text and pollen diagram (Fig. 4). As you indicated a decrease in pollen concentration occurred in the transition from Units 1 and 2, which could be explained by the decrease in organic matter from a shallow water lake to an ephemeral environment.

[Modification in lines 330-331.](#)

[See modifications in figure 4.](#)

3/ Line 220: It would be of interest to mention the state of preservation of the pollen grains as far as some authors (Bottema) suggested that a high amount of the very resistant grains of Cichorioideae could be due to a differential destruction of pollen grains.

That is a good question. We also firstly thought the idea of a major degradation in the pollen during this arid period. However, other proxies such as the increase in other nitrophilous plants and other NPP as *Tilletia* (covarying with Cichorioideae) are indicative of livestock or agricultural activities. In addition, a decrease in this taxon happened since ca. 1500 CE when an increase in clastic material occurred, not supporting a preservation issue.

To clarify this we [modified the text between lines 610-618](#).

4/ Line 269 (fig. 4). It would be of interest to recall the lithology beside the pollen zones (even if the scale refers to ages).

1    Ok, this is a good idea. We added the lithology to the pollen diagram.

2    See modifications in figure 4.

3    5/ Line 288: this increase in Mediterranean forest types is far to be evident!

4    Thank you, we modified this line.

5    Modifications in lines 350-351.

6    6/ lake level, lines 322-331: At a time when human impact is evident together with a high  
7    minerogenic influx it is difficult to decide if it is a lake level lowering or simply an infilling.

8    What we are seeing in this record is a progressive trend in somerization of the lake environment  
9    and that is the reason we believe it is mostly due to lake level lowering. In the discussion we  
10   explain the environmental evolution of the shallow water lake to an ephemeral lake and then  
11   to an emerged environment with a maximum in clastic and MS values in the last ca. 400 cal yr  
12   BP.

13   Modification in lines 388-400.

14   7/ discussion p.9,

15   Line 58-360: it could simply mean that *Botryococcus* greatly suffers in ephemeral lakes

16   Yes, for this reason we calculated two different correlations of *Botryococcus* with temperature  
17   record from Bond et al (2001): one for the total record and another for only the first (oldest)  
18   part, during the deeper shallow water lake phase, where we appreciated a higher correlation.

19   To clarify this we modified the text in line 565.

20   Line 373: clastic input could simply be linked with increasing land use in the catchment  
21   leading to more active erosion!

22   Yes, we agree with the reviewer, but in this sentence we are talking about the interpretation of  
23   variation in detritics in lake environments without any human disturbances – we specified this  
24   in the text. Later on, in the discussion, we try disentangling the natural vs. human-induced  
25   causes for enhanced erosion, which is sometimes difficult and that is why we give some  
26   possible explanations (natural vs human induced).

27   Modifications is lines 462-466.

28

29   Line 377: it would be of interest to connect this evidence with local historical and  
30   archaeological data on the land use around the site across time.

31   That would be great, however there is a lack of published historical records from the area for a  
32   deep review.

Line 492: this increase in *Botryococcus* starts quite at least 500 years before the beginning of IRHP: the argument is faint.

The increase in *Botryococcus* started around ca. 3.1 cal ka BP with an increasing trend towards a higher water lake reaching a *Botryococcus* maximum during the IRHP.

Line 506: Cichorioideae! First try to identify the ecosystem (or the disturbance) which generates this incredible amount of Cichorioideae!

We agree that some further discussion must be added about this matter. See answer to question 1 and 3.

Line 603: I am not a statistician but I am afraid by this suggestion of periodicity bases on honest but rather low pollen and algae counts (260 terrestrial).

We are aware of the limited amount of terrestrial pollen counted, however the data show clear cyclical patterns that are not random and are statistically sound (above confidence levels). These results are confirmed by replication of the same patterns through statistical treatments in the TOC%. Therefore we are quite confident that these results are not an artifact.

Line 618: human impact; my feeling is that a part of the information provided here could be included in the initial presentation of the site.

Yes, we agree and included some Human impact information in the initial presentation of the site (see answer to questions 1).

## Author's responses to Laura Sadori

### General

I found the new data from Padul record, presented by María J. Ramos-Román and colleagues, quite important and necessary to improve our knowledge of the environmental history of the last 5000 years.

The chronology is very well assessed and the sediment and pollen data are consistent.

I was very surprised by the fact that such a multidisciplinary dataset is not used to disentangle between the two main drivers of deforestation in the Mediterranean region: human impact and climate forcing. The authors in fact start the interpretation with pollen, embracing the climate party, but they HAVE TO DEMONSTRATE THIS with clear data, and we have to admit that pollen alone is not enough. I suggest re-writing of interpretation and discussion under this light.

I just noticed that this is also the main concern of the other reviewer, I totally agree with her. This is in fact a never- ending dilemma of Holocene palaeoecology: is it possible to separate the effects due to climate change and human impact in the pollen records of the last millennia?

(see for example the discussion in Marignani et al., 2017. Plant Biosystems). I want to add that my personal opinion is that climate is the most important factor in shaping the present landscape, but it is just an opinion if it is not clearly supported by data! Sometimes, in my personal experience, charcoal counting together with concentration data seemed to be resolute (Sadori et al., 2004. The Holocene), but most times it is just the use of independent sediment and geochemical proxies (Giraudi et al., 2011, The Holocene; Morellon et al., 2016. Quaternary Science Reviews; Sadori et al., 2016 Quaternary Science Reviews) that can disentangle drivers, solving the "dilemma". You have good data from your own record that can be used in this sense! I found that the comparisons with other sites are too many and not always meaningful, so that the reader gets lost. The references are mostly up to date, but the discussion present in the pollen community about the cause of the deforestation (aridification vs. increased land use) is completely ignored. It should be included. I do think that the paper absolutely deserves to be published, and I recommend publication in *Climate of the Past*, but just after the above mentioned issues will be assessed. Please have a look also at the file with my comments.

Thank you. We agree with your comment about the importance of disentangling human impact and climate forcing in this study. In our opinion the most important process shaping the environment in the area was climate as we do not have clear evidences of human activities in this record until the last 1500 cal yr BP. Humans left their footprint in the area since then, with indications of cultivars and livestock but still when we expect to see a significant forest reduction the deforestation trend continues to be linear and even opposite to what expected with slight increases in forest during the MCA. In this new version of the manuscript we inserted a new figure 8 (see below in the new manuscript version) trying to clarify and separate natural vs. anthropic signals. We hope that with this new figure together with some modifications and reorganizations of the text the reader gets a better idea of what we found in this record. Considering the importance of human activity in the area, notably for the last 1500 years, we modified the title for "Holocene aridification trend and human impact interrupted by millennial- and centennial-scale climate fluctuations from a new sedimentary record from Padul (Sierra Nevada, southern Iberian Peninsula)".

## Comments

1/ Line 22: What is the source?

We try to clarify this later on in the manuscript:

[Modifications in lines 502-508.](#)

2/ Line 68: see also...

Thank you for the reference.

3/ Line 107-125: I would recommend the addition of species' authors

We appreciate this comment but we had to reorganize this part following one of the reviewer's comment to make it more synthetic and succinct and we believe adding the species' authors would make it too long.

4/ Line 154-170: same as above (vegetation)

See response to previous comment above.

5/ Line 236: here you have to quote the dates obtained for the published interval, that is 8. In the chapter on chronology assessment you have explained what you have used.

Thank you for the suggestion - we added this information in the manuscript.

Modification in lines 237-239.

6/ Line 269-270: As this work is a part of the complex study you are carrying out on this core, I suggest to number zones from the top to the bottom. In this way in the following you will be able to continue numbering...I also suggest to name the zone PA-1 (PA-1a, PA-1b for subzones). Can you imagine a future quotation of your paper with "zone Padul-15-05-1"? Isn't simpler and more effective PA-1?

It is a good idea. Thank you for the suggestion. We changed the names of the pollen zones throughout the manuscript and figures.

7/ Line 296: better to use Thecamoebians undiff.

Thank you, we agree it is more appropriate. We will modify this throughout the manuscript.

8/ Line 341-342: this is true in absence of human pressure...human impact is overlapping to climate signals and the two are quite difficult to disentangle! To distinguish between them, especially in the last 5000 years, palynology needs to be confirmed by other evidences. This should be emphasized and it would explain the multi-proxy approach you used.

We agree with your comment. However, in this part of the text we are only trying to introduce the different proxies and how can they usually help us. Anyhow, we added some more information about how we tried to discriminate human impact in the area using the palynological proxies. We added the following modification to that part of the text:

Modifications in lines 412-414, 436-447 and 462-466.

9/ Line 346-347: It's not enough! A strong land-use can produce the same deforestation effect!

See answer to question 8.

10/ Line 350: in the central Mediterranean the increase of Mediterranean vegetation occurring in the second half of the Holocene is taken as either the evidence of aridification (from mesophilous to sclerophyllous vegetation) or as the result of increased grazing and cultivation

Yes, we agree and we worked on adding some more information and a new figure (Fig. 8) to clarify this question. See also modifications above and below where we try to clarify this matter. As you will see, this new figure 8 shows that human impact in the area does not seem to be evident until the last 1500 years and the Mediterranean forest dynamics point to a regional climatic signal of aridification that is also affected by millennial-scale climate variability rather than human impact.

11/ Line 352: they could either be the evidence of increased water salinity (*Salicornia*, *Sarcocornia* species) or of increased human impact (many *Chenopodium* species...) ...

We agree with your suggestion and we added this to the manuscript. See modifications in answer to question 8.

12/ Line 362-367: These are the proxies you need to confirm your hypothesis based on pollen!

We clarify this in the modifications inserted related with question 8.

13/ Line 367: *Cannabaceae-Urtycaceae* type

Thank you. We corrected it.

14/ Line 380: See *Mercuri*

Thank you for the suggestions. We added this information to the manuscript. See modifications answering questions 8 and 23.

15/ Line 385-387: This interpretation is confirmed by other proxies in the pollen records from a very different areas such as the coastal ones of Butrint lake, Albania and of the Tiber delta, Italy

Thank you for your suggestions. We added these references to the manuscript. See previous modifications inserted answering question 8.

16/ Line 396-399: Again, a deforestation can produce the same effects... you need to emphasize climate proxies first, and then interpret the pollen record.

Ok, we worked on this – see previous comments and new Fig. 8 where we believe this is clarified.

[Modifications in lines 488-493.](#)

17/ Line 401-403: you see... the two factors were probably overlapping. Who was the leading and starting one?

In my opinion (and in yours, I guess) it was climate, but this has to be demonstrated by independent data... that you, by the way, have.



1 In our opinion the most important process shaping the environment in the area was climate and  
2 we start recording human activities in this area in the last 1500 cal yr BP. We tried to clarify  
3 this matter in section 5.4 “human activity”.

4 18/ Line 408: could be probably  
5 We believe so.

6 19/ Line 418: what about grazing in such environments?

7 Yes, a signal of the presence of livestock was described in these alpine environments (i.e.,  
8 Sporormiella) but in those studies there is not a clear signal of grazing affecting the local  
9 vegetation.

10  
11 20/ Line 436-438: the ages appear to be quite different. Again, local human presence can mask  
12 climatic driving.

13 Yes, we agree in that there are age differences between sites but in this section we are talking  
14 about aridity pulses between 4.2 and 3.0 ka so those dates fall within those ages.

15 21/ Line 433: the date is not calibrated!

16 Thank you. We corrected that error.

17 22/ Line 446: this is the best global relation you can find! The age correspondance is quite  
18 good!

19 Yes, we agree. Thanks!

20 23/ Line 496: it's quite long! In other areas independent proxies found also humid phases.  
21 You cannot explain every environmental change ONLY with climate change!

22 Cichorioideae have also other meaning, not just climate! Just think at pasturelands! See:

23 We agree and changed the title and discussion in this section. We also tried to concentrate  
24 everything concerning human impact in the “human activity” section. However, we still believe  
25 these changes are mostly produced by natural climate change and its variability. Reviewer#2  
26 also made some questions about Cichorioideae. We have made some modifications in the  
27 manuscript in reference to this matter.

28 Modifications in lines 610-618 and in human activities section (5.4).  
29

30 24/ Line 622: you have taken them as an evidence of climate change, see line 497. Again, you  
31 should use pollen data to cross check with sediment data. In that way you could be able to  
32 distinguish about the two driving forces, humans and climate



1 Thank you for your suggestions. Yes, we agree in that we can not use the same taxa for  
2 explaining different processes and tried to be more consistent. Cichoroideae is now only  
3 discussed in the human activity section and only explained as a human activity proxy.  
4

5 25/ Line 660-662: again, pollen can have a dual interpretation!

6 Yes, but in this record we strongly believe, and hope it is more clear in this new version of  
7 the manuscript, climate is the main factor controlling these changes until ca. 1500 years ago.  
8  
9

**Holocene aridification trend and human impact interrupted by millennial- and centennial-scale climate fluctuations from a new sedimentary record from Padul (Sierra Nevada, southern Iberian Peninsula)**

María J. Ramos-Román<sup>1</sup>, Gonzalo Jiménez-Moreno<sup>1</sup>, Jon Camuera<sup>1</sup>, Antonio García-Alix<sup>1</sup>, R. Scott Anderson<sup>2</sup>, Francisco J. Jiménez-Espejo<sup>3</sup>, José S. Carrión<sup>4</sup>

<sup>1</sup> Departamento de Estratigrafía y Paleontología, Universidad de Granada, Spain

<sup>2</sup> School of Earth Sciences and Environmental Sustainability, Northern Arizona University, USA.

<sup>3</sup> Department of Biogeochemistry, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Japan.

<sup>4</sup> Departamento de Biología Vegetal, Facultad de Biología, Universidad de Murcia, Murcia, Spain.

*Correspondence to:* María J. Ramos-Román (mjrr@ugr.es)

**Abstract.** Holocene centennial-scale paleoenvironmental variability has been described in a multiproxy analysis (i.e. lithology, geochemistry, macrofossil and microfossil analyses) of a paleoecological record from the Padul basin in Sierra Nevada, southern Iberian Peninsula. This sequence covers a relevant time interval hitherto unreported in the studies of the Padul sedimentary sequence. The ~~~ea-~~4700 yr-long record has preserved proxies of climate variability, with vegetation, lake levels and sedimentological change during the Holocene in one of the most unique and southernmost peat bogs from Europe. The progressive Middle and Late Holocene trend toward arid conditions identified by numerous authors in the western Mediterranean region, mostly related to a decrease in summer insolation, is also documented in this record, being here also superimposed by centennial-scale variability in humidity. In turn, this record shows centennial-scale climate oscillations in temperature that correlate with well-known climatic events during the Late Holocene in the western Mediterranean region, synchronous with variability in solar and atmospheric dynamics. The multiproxy Padul record first shows a transition from a relatively humid Middle Holocene in the western Mediterranean region to more aridity from ~4700 to ~2800 cal yr BP. A relatively warm and humid period occurred between ~2600 to ~1600 cal yr BP, coinciding with persistent negative NAO conditions and the historic Iberian-Roman Humid Period. Enhanced arid conditions, co-occurring with overall positive NAO conditions and increasing solar activity, are observed between ~1550 to ~450 cal yr BP (~400 to ~1400 CE) and colder and warmer conditions happened during the Dark Ages and Medieval Climate Anomaly, respectively. Slightly wetter conditions took place during the end of the MCA and the first part of the Little Ice Age, which could be related to a change towards negative NAO conditions and minima in solar activity. ~~Evidences of higher human impact in the Padul peat bog area are observed in the last ~1550 cal yr BP.~~ Time series analysis performed from local (*Botryococcus* and TOC) and regional ~~signals~~ (Mediterranean forest) signals helped us determining the relationship between southern Iberian climate evolution, atmospheric, oceanic dynamics and solar activity. Our multiproxy record shows little evidence of human impact in the area until ~1550 cal yr BP, when evidence of agriculture and livestock grazing occurs. Therefore climate is the main forcing mechanism controlling environmental change in the area until relatively recently.

43

44 **Keywords:** Holocene, Padul, peat bog, North Atlantic Oscillation, atmospheric dynamics,  
45 southern Iberian Peninsula, Sierra Nevada, western Mediterranean.

## 46 1 Introduction

47 The Mediterranean area is situated in a sensitive region between temperate and subtropical  
48 climates making it an important place to study the connections between atmospheric and oceanic  
49 dynamics and environmental changes. Climate in the western Mediterranean and the southern  
50 Iberian Peninsula is influenced by several atmospheric and oceanic dynamics (Alpert et al.,  
51 2006), including the North Atlantic Oscillation (NAO) one of the principal atmospheric  
52 phenomenon controlling climate in the area (Hurrell, 1995; Moreno et al., 2005). Recent NAO  
53 reconstructions in the western Mediterranean ~~region~~ relate negative and positive NAO conditions  
54 with an increase and decrease, respectively, in winter (effective) precipitation, ~~respectively~~  
55 (Olsen et al., 2012; Trouet et al., 2009). Numerous paleoenvironmental studies in the western  
56 Mediterranean have detected a link at millennial- and centennial-scales between the oscillations  
57 of paleoclimate proxies ~~studied in~~ from sedimentary records with solar variability and  
58 atmospheric (i.e., NAO) and/or ocean dynamics during the Holocene (Fletcher et al., 2013;  
59 Moreno et al., 2012; Rodrigo-Gámiz et al., 2014). Very few montane and low altitude lake  
60 records in southern Iberia document centennial-scale climate change [see, for example Zoñar  
61 Lake (Martín-Puertas et al., 2008)], with most terrestrial records in the western Mediterranean  
62 region evidencing only millennial-scale cyclical changes. Therefore, higher-resolution decadal-  
63 scale analyses are ~~thus necessary in order~~ to analyze the link between solar activity, atmospheric  
64 and oceanographic systems with terrestrial environment in this area at shorter (i.e., centennial)  
65 time scales.

66  
67 Sediments from lakes, peat bogs and marine records from the western Mediterranean have  
68 documented an aridification trend during the Late Holocene (Carrión et al., 2010; Gil-Romera et  
69 al., 2010; Jalut et al., 2009). This trend, however, was superimposed by shorter-term climate  
70 variability, as shown by several recent studies from the region, ~~as well as human pressure~~  
71 (Carrión, 2002; Fletcher et al., 2013; Jiménez-Moreno et al., 2013a; Martín-Puertas et al., 2008;  
72 Ramos-Román et al., 2016). This relationship between climate variability, culture evolution and  
73 human impact during the Late Holocene has also been the subject of recent paleoenvironmental  
74 studies (Carrión et al., 2007; Lillios et al., 2016; López-Sáez et al., 2014; Magny, 2004).  
75 ~~However, it is still unclear what has been the main forcing driving environmental change (i.e.,~~  
76 ~~deforestation) in this area during this time: was it climate or humans?~~ However, it is still unclear  
77 whether climate or human activities have been the main forcing driving environmental change  
78 (i.e., deforestation) in this area during this time.

79  
80 Within the western Mediterranean ~~region~~, Sierra Nevada is the highest and southernmost  
81 mountain range in the Iberian Peninsula and thus presents a critical area for paleoenvironmental  
82 studies. Most high-resolution studies there have come from high elevation sites. The well-known  
83 Padul peat bog site is located at the western foot of the Sierra Nevada (Fig. 1) and bears one of  
84 the longest continental records in southern Europe, with a sedimentary sequence of ~~ca.~~ ~100 m  
85 thick that could represent the last 1 Ma (Ortiz et al., 2004). Several research studies, including  
86 radiocarbon dating, geochemistry and pollen analyses, have been carried out on previous cores

from Padul, and have documented glacial/interglacial cycles during the Pleistocene and up until the Middle Holocene. However, the Late Holocene section of the Padul sedimentary sequence has never been [effectively](#) retrieved and studied ([Florschütz et al., 1971](#); [Ortiz et al., 2004](#); [Pons and Reille, 1988](#)). This was due to the location of these previous corings within [a current](#) the peat mine ~~exploitation-setting~~[operation](#), where the upper (and non- productive) part of the sedimentary sequence was missing (~~Florschütz et al., 1971; Ortiz et al., 2004; Pons and Reille, 1988~~).

Here we present a new record from the Padul peat bog basin: Padul-15-05, a 42.64 m-long sediment core that, for the first time, contains a continuous record of the Late Holocene (Fig. 2). A high-resolution multi-proxy analysis of the upper 1.15 m, the past ~4700 cal yr BP, has allowed us to determine a complete paleoenvironmental and paleoclimatic record at centennial- and millennial-scales. To accomplish that, we reconstructed changes in the Padul peat bog vegetation, sedimentation, climate and human impact during the Holocene throughout the interpretation of the lithology, palynology and geochemistry.

Specifically, the main objective of this paper is to determine environmental variability and climate evolution in the southern Iberian Peninsula and the western Mediterranean region and their linkages to northern hemisphere climate and solar variability during the [latter](#) Holocene. In order to do this, we compared our results with other paleoclimate records from [the regional areas](#) and solar activity from the northern hemisphere for the past ~~~ea.~~[4.7](#)[700](#) cal ~~ka-yr~~ BP (Bond et al., 2001; Laskar et al., 2004; Sicre et al., 2016; Steinhilber et al., 2009).

## 2 Study site

### 2.1 Regional setting: Sierra Nevada climate and vegetation

Sierra Nevada is a W-E aligned mountain range located in Andalusia (southern Spain). Climate in this area is Mediterranean, with cool and humid winters and hot/warm summer drought. In the Sierra Nevada, the mean annual temperature at approximately 2500 m asl is 4.5 °C and the mean annual precipitation is 700 mm/yr (Oliva et al., 2009). Sierra Nevada is strongly influenced by thermal and precipitation variations due to the altitudinal gradient (from ca. 700 to more than 3400 m), which control plant taxa distribution in different bioclimatic vegetation belts due to the variability in thermotypes and ombrotypes (Valle Tendero, 2004). ~~According to the climatophilous series classification, Sierra Nevada is divided in four different vegetation belts (Fig. 1). The erioromediterranean vegetation belt, occurring above ~2800 m, is characterized principally by Festuca clementei, Hormatophylla purpurea, Erigeron frigidus, Saxifraga nevadensis, Viola crassiuscula, and Linaria glacialis. The oromediterranean belt, between ~1900 to 2800 m, bears Pinus sylvestris, P. nigra, Juniperus communis subsp. hemisphaerica, J. sabina var. humilis, J. communis subsp. nana, Genista versicolor, Cytisus oromediterraneus, Hormatophylla spinosa, Prunus prostrata, Deschampsia iberica and Astragalus sempervirens subsp. nevadensis as the most representative species. The supramediterranean belt, from approximately 1400 to 1900 m of elevation, principally includes Quercus pyrenaica, Q. faginea, Q. rotundifolia, Acer opalus subsp. granatense, Fraxinus angustifolia, Sorbus torminalis, Adenocarpus decorticans, Helleborus foetidus, Daphne gnidium, Clematis flammula, Cistus laurifolius, Berberis hispanicus, Festuca scariosa, Thymus serpylloides subsp. gadoriensis, Helichrysum italicum subsp. serotinum, Santolina rosmarinifolia subsp. canescens and Artemisia~~

~~*glutinosa*. The mesomediterranean vegetation belt occurs between ~600 and 1400 m of elevation and is characterized by *Quercus rotundifolia*, *Retama sphaerocarpa*, *Paeonia coriacea*, *Juniperus oxycedrus*, *Rubia peregrina*, *Asparagus acutifolius*, *D. gnidium*, *Ulex parviflorus*, *Genista umbellata*, *Cistus albidus* and *C. laurifolius* (Al Aallali et al., 1998; Valle, 2003). The human impact over this area, especially important during the last millennium, affected the natural vegetation distribution through fire, deforestation, cultivation. (i.e., *Olea*) and subsequent reforestation (mostly *Pinus*) (Anderson et al., 2011).~~ According to the climatophilous series classification, Sierra Nevada is divided in four different vegetation belts (Fig. 1). The crioromediterranean vegetation belt, occurring above ~2800 m, is characterized by tundra vegetation and principally composed by species of Poaceae, Asteraceae, Brassicaceae, Gentianaceae, Scrophulariaceae and Plantaginaceae between other herbs, with a number of endemic plants (e.g. *Erigeron frigidus*, *Saxifraga nevadensis*, *Viola crassiuscula*, *Plantago nivalis*). The oromediterranean belt, between ~ 1900 to ~2800 m, is principally made up of *Pinus sylvestris*, *P. nigra* and *Juniperus* spp. and other shrubs such as species of Fabaceae, Cistaceae and Brassicaceae. The supramediterranean belt, from ~ 1400 to 1900 m of elevation, bears principally *Quercus pyrenaica*, *Q. faginea* and *Q. rotundifolia* and *Acer opalus* ssp. *granatense* with other trees and shrubs, including members of the Fabaceae, Thymelaeaceae, Cistaceae and *Artemisia* sp. being the most important. The mesomediterranean vegetation belt occurs between ~600 and 1400 m of elevation and is principally characterized by *Quercus rotundifolia*, some shrubs, herbs and plants as *Juniperus* sp., and some species of Fabaceae, Cistaceae and Liliaceae with others (Al Aallali et al., 1998; Valle, 2003). The human impact over this area, especially important during the last millennium, affected the natural vegetation distribution through fire, deforestation, cultivation (i.e., *Olea*) and subsequent reforestation (mostly *Pinus*) (Anderson et al., 2011).

## 2.2 Padul peat bog

The Padul basin is situated at approximately 725 m of elevation in the southeastern part of the Granada Basin, at the foothill of the southwestern Sierra Nevada, Andalucía, Spain (Fig. 1). This is one of the most seismically active areas in the southern Iberian Peninsula with numerous faults in NW-SE direction, with the Padul fault being one of these active normal faults (Alfaro et al., 2001). It is a small extensional basin approximately 12 km long and covering an area of approximately 45 km<sup>2</sup>, which is bounded by the Padul normal fault. The sedimentary in-filling of the basin consists of Neogene and Quaternary deposits; Upper Miocene conglomerates, calcarenites and marls, and Pliocene and Quaternary alluvial sediments, lacustrine and peat bog deposits (Sanz de Galdeano et al., 1998; Delgado et al., 2002; Domingo et al., 1983). Vegetation in the Padul area is dominated by *Q. rotundifolia* (and in less amounts *Q. faginea*), which is normally accompanied by *Pistacia terebinthus*. Shrub species in the area include *Juniperus oxycedrus*, *Crataegus monogyna*, *Daphne gnidium* and *Ruscus aculeatus*. Creepers such as *Lonicera implexa*, *Rubia peregrina*, *Hedera helix*, *Asparagus acutifolius* also occur in this area and some herbs, such as *Paeonia broteroi*. *Quercus coccifera* also occurs in ~~erests~~ ridges and very sunny rocky outcrops. *Retama sphaerocarpa* and *Genista cinerea* subsp. *speciosa* and the *Thymo-Stipetum tenacissime* series also occur in sunny areas and in more xeric soils. Nitrophilous communities occur in soils disrupted by livestock, pathways or open forest, normally related with anthropization (Valle, 2003).



The Padul peat bog is an-endorheic area with a surface of approximately 4 km<sup>2</sup> placed in the Padul basin that contains a sedimentary sequence mostly characterized mostly by peat accumulation. The basin fill is asymmetric, with thicker sedimentary and peat infill to the northeast (~100 m thick; Domingo-García et al., 1983; Florschütz et al., 1971; Nestares and Torres, 1997) and progressively becoming disappearing-thinner to the southwest (Alfaro et al., 2001). The main source area of the allochthonous sediments in the Padul peat-bog is the Sierra Nevada, which is characterized at higher elevations by Paleozoic siliceous metamorphic rocks (mostly mica-schists and quartzites) from the Nevado-Filabride complex and, at lower elevations and acting as bedrock, by Triassic dolomites, limestones and phyllites from the Alpujárride Complex (Sanz de Galdeano et al. 1998). Geochemistry in the Padul sediments is influenced by detritic materials also primarily from coming from the neighboring mountains, mainly the Sierra Nevada mountain range (Ortiz et al., 2004). In addition, groundwater inputs into the Padul basin are the main reason why there is a wetland in this area. These inputs come from the Triassic carbonates aquifers (N and S edge to the basin), the out flow of the Granada Basin (W edge to the basin) and the conglomerate aquifer to the east edge (Castillo Martín et al., 1984; Ortiz et al., 2004). The main water output is by evaporation and evapotranspiration, water wells and by canals (“madres”) that drain the water to the Dúrcal river to the southeast (Castillo Martín et al., 1984). Climate in the Padul area is characterized by a mean annual temperature of 14.4 °C and a mean annual precipitation of 445 mm (<http://www.aemet.es/>).

~~Vegetation in the Padul area is dominated by *Q. rotundifolia* (and in less amounts *Q. faginea*), which is normally accompanied by *Pistacia terebinthus*. Shrub species in the area include *Juniperus oxycedrus*, *Crataegus monogyna*, *Daphne gnidium* and *Ruscus aculeatus*. Creepers such as *Lonicera implexa*, *Rubia peregrina*, *Hedera helix*, *Asparagus acutifolius* also occur in this area and some herbs, such as *Paeonia broteroi*. *Quercus coccifera* also occurs in crests and very sunny rocky outcrops. *Retama sphaerocarpa* and *Genista cinerea* subsp. *speciosa* and the *Thymo-Stipetum tenacissimum* series also occur in sunny areas and in more xeric soils. Nitrophilous communities occur in soils disrupted by livestock, pathways or open forest, normally related with anthropization (Valle, 2003).~~

The Padul-15-05 drilling site was located around 50 m south of the present-day Padul lake shore area. This basin area is presently subjected to seasonal water level fluctuations and is principally dominated by *Phragmites australis* (Poaceae). The lake environment is dominated by aquatic and wetland communities with *Chara vulgaris*, *Myriophyllum spicatum*, *Potamogeton pectinatus*, *Potamogeton coloratus*, *Phragmites australis*, *Typha domingensis*, *Apium nodiflorum*, *Juncus subnodulosus*, *J. bufonius*, *Carex hispida*, *Juncus bufonius* and *Ranunculus muricatus*, between among others (Pérez Raya and López Nieto, 1991). Some sparse riparian trees occur in the northern lake shore, such as *Populus alba*, *Populus nigra*, *Salix* sp., *Ulmus minor* and *Tamarix*. At present *Phragmites australis* is the most abundant plant bordering the lake. Surrounding this area are cultivated crops with cereals, such as *Triticum* spp., as well as *Prunus dulcis* and *Olea europea*.

### 3 Material and methods

Two sediment cores, Padul-13-01 (37°00'40''N; 3°36'13''W) and Padul-15-05 (37°00'39.77''N; 3°36'14.06''W) with a length of 58.7 cm and 42.64 m, respectively, were collected between 2013 and 2015 from the peat bog (Fig. 1). The cores were taken using a Rolatec RL-48-L drilling

machine equipped with a hydraulic piston corer from the Scientific Instrumentation Centre of the University of Granada (UGR). The sediment cores were wrapped in film, put in core boxes, transported to UGR and stored in a dark cool room at  $\pm 4^{\circ}\text{C}$ .

### 3.1 Age-depth model (AMS radiocarbon dating)

The core chronology was constrained using fourteen AMS radiocarbon dates from plant remains and organic bulk samples taken throughout from the cores (Table 1). In addition, one sample with gastropods was also measured-submitted for AMS radiocarbon analysis, although it was rejected due to important reservoir effect, which-that provided a very old date. Thirteen of these samples came from Padul-15-05 and-with one from the nearby Padul-13-01 (Table 1). We were able to use this date from Padul-13-01 core as there is a very significant correlation between the upper part of Padul-15-05 and Padul-13-01 cores, shown by identical lithological and geochemical changes (Supplementary information 1; Figure S1). The age model for the upper ~3 m until-minus the upper 21 cm from the surface was built using the R-code package ‘Clam 2.2’ (Blaauw, 2010) employing the calibration curve IntCal 13 (Reimer et al., 2013), a 95 % of confidence range, a smooth spline (type 4) with a 0.20 smoothing value and 1000 iterations (Fig. 2). The chronology of the uppermost 21 cm of the record was built using a linear interpolation between the last radiocarbon date and the top of the record (Present; 2015 CE). Even though the length of the Padul-15-05 core is ~43 m, the studied interval in the work presented here is the uppermost 115 cm of the record that are constrained by seven AMS radiocarbon dates (Fig. 2). The studied interval in the present work are the uppermost 115 cm of the record that are constrained by six AMS radiocarbon dates (Fig. 2).

### 3.2 Lithology, MS, XRF and TOC

~~The length for the Padul-15-05 core is ~43 m. In this study, we focus in the uppermost ~115 cm from that core. Padul-15-05 core~~ was split longitudinally and was described in the laboratory with respect to lithology and color (Fig. 3).

Magnetic susceptibility (MS) was measured with a Bartington MS3 operating with a MS2E sensor. MS measurements (in SI units) were obtained directly from the core surface every 0.5 cm (Fig. 3).

Elemental geochemical composition was measured in an X-Ray fluorescence (XRF) Avaatech core scanner® at the University of Barcelona (Spain). A total of thirty-three chemical elements were measured in the XRF core scanner at 10 mm of spatial resolution, using 10 s count time, 10 kV X-ray voltage and a X-ray current of 650  $\mu\text{A}$  for lighter elements and 35 s count time, 30 kV X-ray voltage, X-ray current of 1700  $\mu\text{A}$  for heavier elements. Thirty-three chemical elements were measured but only the most representative with a major number of counts were considered (Si, K, Ca, Ti, Fe, Zr, Br and Sr). Results for each element are expressed as intensities in counts per second (cps) and normalized (norm.) for the total sum in cps in every measure (Fig. 3).

Total organic carbon (TOC) was analyzed every 2 or 3 cm throughout the core. Samples were previously decalcified with 1:1 HCl in order to eliminate the carbonate fraction. The percentage of organic Carbon (OC %) was measured in an Elemental Analyzer Thermo Scientific Flash 2000 model from the Scientific Instrumentation Centre of the ~~University of Granada~~UGR (Spain). Percentage of TOC per gram of sediment was calculated from the percentage of organic

carbon (OC %) yielded by the elemental analyzer, and recalculated by the weight of the sample prior to decalcification (Fig. 3).

### 3.3 Pollen and NPP

Samples for pollen analysis (1-3 cm<sup>3</sup>) were taken every 1 cm throughout the core, with a total of 103 samples analyzes. Pollen extraction methods followed a modified Faegri and Iversen (1989) methodology. Processing included the addition of *Lycopodium* spores for calculation of pollen concentration. Sediment was treated with NaOH, HCl, HF and the residue was sieved at 250  $\mu$ m previous to an acetolysis solution. Counting was performed using a transmitted light microscope at 400 magnifications to an average pollen count of ca. 260 terrestrial pollen grains. Fossil pollen was identified using published keys (Beug, 2004) and modern reference collections at University of Granada (Spain). Pollen counts were transformed to pollen percentages based on the terrestrial pollen sum, excluding aquatics. The palynological zonation was executed by cluster analysis using twelve ~~different~~-primary pollen taxa- *Olea*, *Pinus*-~~undifferentiated~~, deciduous *Quercus*, evergreen *Quercus*, *Pistacia*, Ericaceae, *Artemisia*, Asteroideae, Cichorioideae, Amaranthaceae and Poaceae (Grimm, 1987) (Fig. 4). Non-pollen palynomorphs (NPP) include fungal and algal spores, and thecamoebians (testate amoebae). The NPP percentages were calculated and represented with respect to the terrestrial pollen sum (Fig. 4). Furthermore, some pollen taxa were grouped, according to present-day ecological bases, into Mediterranean forest and xerophytes (Fig. 4). The Mediterranean forest taxa is composed of *Quercus* total, *Olea*, *Phillyrea* and *Pistacia*. The xerophyte group includes *Artemisia*, *Ephedra*, and Amaranthaceae.

## 4 Results

### 4.1 Chronology and sedimentation rates

The age-model of the ~~studied upper 115 cm of~~ Padul-15-05 core (Fig. 2) shows ~~that the top 115 cm, with~~ an average sedimentation rate (SAR) of 0.058 cm/yr, ~~continuously cover approximately the over~~ last ca. 4700 cal yr BP, being the age constrained by ~~fourteen~~-seven AMS <sup>14</sup>C dates (Table 1). However, SARs of individual core segments vary from 0.01 to 0.11 cm/yr (Fig. 2). Five distinct sediment accumulation rate (SAR) intervals can be differentiated between 0 and 122.96 cm based on the linear interpolation between radiocarbon dates in the studied core (Fig. 2).

### 4.2 Lithology, MS, XRF and TOC

The ~~lithology~~-stratigraphy of the upper ca. 115 cm of the Padul-15-05 sediment core was ~~mainly~~-deduced ~~from~~-primarily by visual inspection. However, our visual inspections were support together with the by comparison with the element geochemical composition (XRF), ~~and the correlation of these data, and~~ the MS of the split cores, and. ~~In addition, this information was complemented with the~~ TOC (Fig. 3) to determine shifts in sediment facies. The lithology for this sedimentary sequence consists in clays with variable carbonates, siliciclastics and organic content (Fig. 3). We also used a :  
A-Linear r (Pearson) correlation ~~was to~~ calculated ~~for~~-relationship for the XRF data, The correlation for the inorganic geochemical elements ~~show us~~determined two different groups of



elements that covary (Table 2): Group 1) Si, K, Ti, Fe and Zr with a high positive correlation between them; Group 2) Ca, Br and Sr have negative correlation with Group 1. ~~The lithology for this sedimentary sequence consists in clays with variable carbonates, siliciclastics and organic content (Fig. 3).~~ Based on this, the sequence is subdivided in two principal sedimentary units. The lower ~87 cm of the record is designated to ~~bottom of the record corresponds with~~ Unit 1, characterized principally by relatively lower ~~er~~ values of MS and higher values of Ca. The top upper ~28 cm of the sequence ~~can be subdivided~~ is designated to ~~in~~ Unit 2, in which the mineralogical composition is lower in Ca with higher values of MS in correlation with mostly siliciclastics elements (Si, K, Ti, Fe and Zr).

Within these two units, four different facies can be identified by visual inspection and by the elemental geochemical composition and TOC of the sediments. *Facies 1*; ~~between~~ (115-110 cm depth, ~4700 to 4650 cal yr BP; and 89-80 cm depth ~4300 to 4000 cal yr BP) are characterized by dark brown organic clays that bear charophytes and macroscopic plant remains. They also have depicted relative higher values of TOC values (Fig. 3). *Facies 2* (110-89 cm depth ~4650 to 4300 cal yr BP; and 80-42 cm depth, ~4000 to 1600 cal yr BP) is compose of brown clays, with the occurrence of gastropods and charophytes. ~~These~~ This facies is ~~are~~ also characterized with decreasing by lower TOC values. *Facies 3*; ~~between~~ (42-28 cm depth, ~~(ca.~~ 1600 to 450 cal yr BP) is ~~are~~ characterized by grayish brown clays with the occurrence of gastropods, and lower values of TOC, ~~the and an~~ increasing trend in MS and in siliciclastic elements. *Facies 4*; ~~between~~ (28-0 cm, ~~ca.~~ 450 cal yr BP to Present); are is made up of light grayish brown clays and features a strong increase in siliciclastic linked to a strong increase in MS.

### 4.3 Pollen and NPP

~~A total of seventy-two~~ Several terrestrial and aquatic pollen taxa were identified but only the most representative taxa are here plotted in a the summary pollen diagram (Fig. 4). Selected NPP percentages are also displayed in Figure 4. Four pollen zones (PA) ~~Fig. 4~~ were visually identified with the help of a cluster analysis using the program CONISS (Grimm, 1987). Pollen concentration was higher during Unit 1 with a decreasing trend in the transition to Unit 2 and a later increase during the pollen subzone PA-4b (Fig. 4). Pollen zones are described below:

#### 4.3.1 Zone ~~PA-1~~ adul-15-05-1 [~4720 to 3400 cal yr BP/ ~2800 to 1450 BCE (115-65 cm)]

Zone 1 is characterized by the abundance of Mediterranean forest species reaching up to ca. 70 %. Another important taxon in this zone is *Pinus*, with average values around 18 %. Herbs are largely represented by Poaceae, averaging around 10 %, and reaching up to ca. 25 %. This pollen zone is subdivided into subzones PA-1a, PA-1b and PA-1c (Fig. 4). The principal characteristic that differentiate differentiating subzone PA-1a to from subzone PA-1b (boundary at ca. 4650 cal yr BP/ca. 2700 BCE) is the decrease in Poaceae ~~from an average value of ca. 18 to 10 %~~, the increase in *Pinus* ~~from ca. 7 to 18 %~~, and the appearance of cf. *Vitis*. The subsequent decrease in Mediterranean forest pollen to average values around 40 %, the increase in *Pinus* to average values around 25 % and a progressive increase in Ericaceae ~~with average occurrences from ca. to ~6 to 11 %~~, allow to distinguish seem subzones PA-subzones-1b and PA-1c (boundary at ca. 3950 cal yr BP).

#### 4.3.2. Zone ~~PA-2~~ adul-15-05-2 PA-2 [~3400 to 1550 cal yr BP/~1450 BCE to 400 CE (65-41 cm)]

The main features of this zone are the increase in Ericaceae up to ~~~ea-~~16 % and in deciduous *Quercus*, reaching values ~~~around-~~20 %. Therefore, the Mediterranean forest component progressively ~~decreased-~~declined to values around 34 %. Some herbs, such as Cichorioideae, became more abundant reaching average ~~occurrences around~~percentages of ~7 %. This pollen zone can be subdivided in subzones PA-2a and PA-2b with a boundary at ~2850 cal yr BP (~900 BCE). The principal characteristics that differentiate these subzones is marked by the ~~increase in Mediterranean forest types and the~~ increasing trend in deciduous *Quercus* and Ericaceae. The increase in *Botryococcus*, which averages ~~ea-~~4 to 9 %. Also notable is ~~and~~ the expansion of *Mougeotia* and *Zygnema* types ~~are also noticeable~~.

#### 4.3.3 Zone ~~Padul-15-05-3~~PA-3 [~1550 to ~~4050~~ cal yr BP/~400 CE to ~~15500~~ CE (41-29 cm)]

This zone is distinguished by the ~~maximum-depletion~~continuing decline of Mediterranean forest elements. Cichorioideae reached average values of about 40 %, and is paralleled by- ~~The-the~~ decrease in Ericaceae ~~is also significant-~~. A ~~decrease-~~decline in *Botryococcus* and other algal remains is also observed in this zone, although there is an increase in ~~other~~total -Thecamoebians from average ~~values of~~<1 % to 10 %. This pollen zone is subdivided in subzones PA-3a and PA-3b at ~1000 cal yr BP (~950 CE). The main features that differentiate these subzones are the increase in *Olea* from subzone PA-3a to PA-3b from average values of ~~~ea-~~1 to 5 %. The increasing trend in Poaceae is also a feature in this subzone, as well as the slight increase in Asteroideae at the top. Significant changes are documented in NPP percentages in this subzone with the ~~expansion-~~increase of some fungal remain such as *Tilletia* and *Glomus* type. Furthermore, a decrease in *Botryococcus* and the near disappearance of other algal remains such as *Mougeotia* occurred.

#### 4.3.4 Zone ~~Padul-15-05-4~~PA-4 [~last ~~4050~~ cal yr BP/ ~ ~~15500~~ CE to Present (29-0 cm)]

The main feature in this zone is the significant increase in *Pinus*, reaching maximum values (~~ea. of~~ ~32 %), an increase in Poaceae (~~ea. to~~ ~40 %) and the decrease in Cichorioideae (~~from~~ ~~~average occurrences of ea-~~44 to 16 %). Other important changes are the nearly total disappearance of some shrubs such as *Pistacia* and a decreasing trend in Ericaceae, as well as ~~an~~ ~~important decrease~~ further decline in Mediterranean forest pollen. An increase in wetland pollen taxa, mostly *Typha*, also occurred. A significant increase in xerophytes, ~~with the expansion of mostly~~ Amaranthaceae to ~14 %, is also observed in this period. Other herbs such as *Plantago*, Polygonaceae and Convolvulaceae show moderate increases. ~~This zone~~PA-4 is subdivided into subzones PA-4a and PA-4b (Fig. 4). The top of the record (PA-4b), which corresponds with approximately the last ~1830 CE to Present120 yr, is ~~characterized by the~~differentiated from subzone PA-4a (from ~400 – 120 cal yr BP) by, ~~the main characteristic that differentiate subzone 4b from the previous 4a is a~~ ~~decrease-~~decline in some herbs such as Cichorioideae. However, an increase in ~~some other~~ xerophytic herbs such as Amaranthaceae ~~-occurred and~~ Poaceae occurred. The increase in *Plantago* is also significant during this period. ~~A PA-4b also~~ has a noteworthy increase in *Pinus* (from ~~~an average of ea-~~14 to 27 %) and a slight increase in *Olea* and evergreen *Quercus* are also characteristic of this subzone. With respect to NPPs, ~~there is an increase in~~ thecamoebians such as *Arcella* type and in the largely coprophilous sordariaceous (Sordariales) group spores also increase. This zone also documents the decrease in fresh-water algal spores, in *Botryococcus* concomitant with *Mougeotia* and *Zygnema* type.

#### 4.4 Estimated lake level reconstruction

Different local proxies from the Padul-15-05 record [Si, Ca, TOC, MS, ~~h~~Hygrophytes (~~made-up~~ ~~of~~ Cyperaceae and *Typha*), ~~—~~ Poaceae and ~~a~~Algae (including *Botryococcus*, *Zygnema* types and *Mougeotia*) groups] have been depicted in order to understand the relationship between lithological, geochemical, and palynological variability and the water lake level oscillations. Sediments with higher values of TOC (more algae and hygrophytes) ~~and~~ rich in Ca (related with the occurrence of shells and charophytes remains) most likely characterized a shallow water environment (Unit 1). The continuous decline in *Botryococcus*, the disappearance of charophytes and the progressively increase in detritics (increase in MS and Si values) could be associated with shallower and even ephemeral lake environment (transition from Unit 1 to Unit 2; ~41 to 28 cm). The absence of aquatic ~~shells~~remains, almost disappearance of *Botryococcus* and decreasing Ca and a lower TOC and/or a higher input of clastic material (higher MS and Si values) into the lake, could be related with lake level lowering, and ~~a shallower wetland environment~~even emerged conditions (increase in Poaceae; Unit 2) (Fig. 5).

#### 4.5 Spectral analysis

Spectral analysis was performed on selected pollen and NPP time series (Mediterranean forest and *Botryococcus*), as well as TOC in order to identify millennial- and centennial-scale periodicities ~~in the Padul-15-05 record~~. The mean sampling resolution for pollen and NPP is ~~~ea.~~ 50 yr and for geochemical data is ~~~ea.~~ 80 yr. Statistically significant cycles, above the 90, 95 and 99 % of confident levels, were found around 800, 680, 300, 240, 200, 170 (Fig. 7).

#### 5 Discussion

~~Different~~ Numerous proxies have been used in this study to interpret the paleoenvironmental and hydrodynamic changes recorded in the Padul ~~peat bog~~ sedimentary record during the last 4700 cal yr BP. Palynological analysis (pollen and NPP) is commonly used as a proxy for vegetation and climate change, and lake level variations, as well as ~~and~~ human impact and land uses (e.g. Faegri and Iversen, 1990; van Geel et al., 1983). Disentangling natural vs. anthropogenic effects on the environment in the last millennia is sometimes challenging but can be persuaded using a multi-proxy approach (Roberts et al., 2011; Sadori et al., 2011). In this study, we used the variations between Mediterranean forest taxa, xerophytes and algal communities for paleoclimatic variability and the occurrence of nitrophilous and ruderal plant communities and some NPPs for identifying human influence in the study area ~~(Fig. 4)~~. Variations in arboreal pollen (AP, including Mediterranean tree species) have previously been used in ~~the~~ previous Sierra Nevada records as a proxy for humidity changes (Jiménez-Moreno and Anderson, 2012; Ramos-Román et al., 2016). The ~~abundance-increase or decrease of the~~ in Mediterranean forest species has been used as a proxy for climate change in other studies in the western Mediterranean region, with ~~higher~~ greater forest development generally meaning higher humidity (Fletcher et al., 2013; Fletcher and Sánchez-Goñi, 2008). On the other hand, increases in xerophyte pollen taxa (i.e., *Artemisia*, *Ephedra*, *Amaranthaceae*) have been used as an indication of aridity in this area (Anderson et al., 2011; Carrión et al., 2007).

The chlorophyceae alga *Botryococcus* sp. has been ~~described~~ used as an indicator of freshwater environments, in relatively productive fens, temporary pools, ponds or lakes (Guy-Ohlson,

1992). The high visual and statistical correlation between *Botryococcus* from Padul-15-05 and North Atlantic temperature estimations (Bond et al., 2001;  $r = -0.63$ ;  $p < 0.0001$ ; between ca. 4700 to 1500 cal ka-yr BP – the decreasing and very low *Botryococcus* occurrence in the last 1500 cal yr BP makes this correlation moderate:  $r = -0.48$ ;  $p < 0.0001$  between 4700 and -65 cal yr BP) seems to show that in this case *Botryococcus* is driven by temperature change and would reflect variations in lake productivity (increasing with warmer water temperatures).

Human impact can be investigated using several palynomorphs. Nitrophilous and ruderal pollen taxa, such as *Convolvulus*, *Plantago lanceolata* type, Urticaceae type and *Polygonum avicularis* type, are often proxies for human impact (Riera et al., 2004), and abundant Amaranthaceae has also been used as well (Sadori et al., 2003). Some species of Cichorioideae have been described as nitrophilous taxa (Abel-Schaad and López-Sáez, 2013) and as grazing indicators (Florenzano et al., 2015; Mercuri et al., 2006; Sadori et al., 2016). At the same time, NPP taxa such as some coprophilous fungi, Sordariales and thecamoebians are also used as indicators of anthropization and land use (Carrión et al., 2007; Ejarque et al., 2015; van Geel et al., 1989; Riera et al., 2006). *Tilletia* a grass-parasitizing fungi has been described as an indicator of grass cultivation in other Iberian records (Carrión et al., 2001a). In this study we follow the example of others (van Geel et al., 1989; Morellón et al., 2016; Sadori et al., 2016) who used the NPP soil mycorrhizal fungus *Glomus* sp. as a proxy for erosive activity.

~~In addition to~~ the palynological analysis, variations in the lithology, geochemistry and macrofossil remains (gastropod shells and charophytes) from the Padul-15-05 core helped us reconstruct the estimated lake level and the local environment changes in the Padul ~~peat bog area~~ and their relationship with regional climate (Figs. 5). Several previous studies on Late Holocene lake records from the Iberian Peninsula show that lithological changes can be used as a proxy for lake level reconstruction (Martín-Puertas et al., 2011; Morellón et al., 2009; Riera et al., 2004). For example, carbonate sediments formed by biogenic remains of gastropods and charophytes are indicative of shallow lake waters (Riera et al., 2004). Furthermore, van Geel et al. (1983), described occurrences of *Mougeotia* and *Zygnema* type (Zygnemataceae) as typical of shallow water environments. The increase in organic matter accumulation deduced by TOC (and Br) could be considered as characteristic of high productivity (Kalugin et al., 2007) in these shallow water environments. On the other hand, increases in clastic input in lake sediments have been interpreted as due to lowering of lake level and more influence of terrestrial-fluvial deposition in a very shallow/ephemeral lake (Martín-Puertas et al., 2008). Carrión (2002) related the increase in some fungal species and Asteraceae as indicators of seasonal desiccation stages in lakes. Nevertheless, in natural environments with potential interactions with human activities the increase in clastic deposition related with other indications of soil erosion (e.g. *Glomus* sp.) may be assigned to intensification in land use (Morellón et al., 2016; Sadori et al., 2016). ~~We used the variations between those proxies to estimate water level (Fig. 5):~~

~~Nitrophilous and ruderal pollen taxa (*Convolvulus*, *Plantago lanceolata* type, Urticaceae type and/or *Polygonum avicularis* type) are also very useful as proxies for human impact (Riera et al., 2004). Some species of Cichorioideae have also been described in different studies from the Iberian Peninsula as nitrophilous taxa (Abel-Schaad and López-Sáez, 2013). At the same time, NPP taxa such as some coprophilous fungi, Sordariales and thecamoebians are also used as indicators of anthropization and land use (Carrión et al., 2007; Ejarque et al., 2015; van Geel et al., 1989; Riera et al., 2006). *Tilletia* a grass-parasitizing fungi has been described as an indicator~~



of grass cultivation in other Iberian records (Carrión et al., 2001b). In this study we also used the NPP mycorrhizal fungus *Glomus* sp. as a proxy for erosive activity. This interpretation comes from a study from van Geel et al. (1989), who correlated erosive events with elevated percentages of *Glomus* cf. *fasciculatum*.

## 5.1 Late Holocene aridification trend

~~This study shows that a~~Our work confirms the progressive aridification trend that occurred during at least the last ~~ca.~~ 4700 cal yr BP in the southern Iberian Peninsula, as shown here. ~~The increase in aridity is shown in the Padul 15-05 core by the a~~ progressive decrease in Mediterranean forest component and the increase in herbs (Figs. 4 and 67). ~~Our lake level interpretations from our record agree with the pollen data, and showing~~ an overall decrease during the Late Holocene, from a shallow water table containing relatively abundant organic matter (high TOC, indicating higher productivity), gastropods and charophytes (high Ca values) to a low-productive ephemeral/emerged environment (high clastic input and MS and decrease in Ca) (Fig. 5). This natural progressive ~~aridification~~ confirmed by the decrease in Mediterranean forest taxa and increase in siliciclastics pointing to a change towards ephemeral (even emerged) environments became more prominent since ~~the last ca.~~ about 1550 cal yr BP and then enhanced again ~~in the last~~ since 300 ~~ca.~~ 400 cal yr BP to Present. A clear increase in human land use is also observed during the last ca. 1550 cal yr BP (see below), including abundant Glomus from erosion, which shows that humans were at least partially responsible for this sedimentary change. Furthermore, the increase in some proxies indicating human land use during this last period suggests that humans were more active in this area since then. ~~These results are supported by~~ A suite of proxies previous studies supports our conclusions regarding the ~~from the Mediterranean area using different proxies documenting an~~ aridification trend since the Middle Holocene (Carrión, 2002; Carrión et al., 2010; Fletcher et al., 2013; Fletcher and Sánchez-Goñi, 2008; Jiménez-Espejo et al., 2014; Jiménez-Moreno et al., 2015). In the western Mediterranean region ~~the~~ is decline in forest development during the Middle and Late Holocene is related with a decrease in summer insolation (Fletcher et al., 2013; Jiménez-Moreno and Anderson, 2012), which may have decreased, winter rainfall as a consequence of a northward shift of the westerlies - a long-term enhanced positive NAO trend - which induced drier conditions in this area since 6000 cal yr BP (Magny et al., 2012). Furthermore, the decrease in summer insolation would produce a progressive cooling, with a reduction in the length of the growing season as well as a decrease in the sea-surface temperature (Marchal et al., 2002), generating a decrease in the land-sea contrast that would be reflected in a reduction of the wind system and a reduced precipitation gradient from sea to shore during the fall-winter season. This would produce a progressive cooling, with a reduction in the length of the growing season as well as a decrease in the sea-surface temperature (Marchal et al., 2002), generating a decrease in the land-sea contrast that would be reflected in a reduction of the wind system and a reduced precipitation gradient from sea to shore during the fall-winter season. Also, a reorganization of the general atmospheric circulation with a northward shift of the westerlies - a long-term enhanced positive NAO trend - has been interpreted, inducing drier conditions in this area since 6000 cal yr BP (Magny et al., 2012). The aridification trend can clearly be seen in the nearby alpine records from the Sierra Nevada, where there was little influence by human activity (Anderson et al., 2011; Jiménez-Moreno et al., 2013a; Jiménez-Moreno and Anderson, 2012; Ramos-Román et al., 2016).

## 5.2 Millennial- and centennial-scale climate variability in the Padul peat bog during the Late Holocene

The multi-proxy paleoclimate record from Padul-15-05 shows an overall aridification trend. However, this trend seems to be modulated by millennial- and centennial-scale climatic variability.

### 5.2.1 Aridity pulses around 4200 (4500, 4300 and 4000 cal yr BP) and around 3000 cal yr BP (3300 and 2800 cal yr BP)

Marked aridity pulses are registered in the Padul-15-05 record around 4200 and 3000 cal yr BP (Unit 1; ~~Pollen-zones~~ [PA-1](#) and [PA-2a](#); Figs. 6 and 7). These arid pulses are mostly evidenced in this record by declines in Mediterranean forest taxa, as well as lake level drops and/or cooling evidenced by a decrease in organic component as TOC and the decrease in *Botryococcus* algae. However, a discrepancy between the local and regional occurs between 3000-2800 cal yr BP, with an increase in the estimated lake level and a decrease in the Mediterranean forest during the late Bronze Age until the early Iron Age (Figs. 6 and 7). The disagreement could be due to deforestation by humans during a very active period of mining in the area observed as a peak in lead pollution in the alpine records from Sierra Nevada (García-Alix et al., 2013). The aridity pulses agree regionally with recent studies carried out at higher elevation in the Sierra Nevada, a decrease in AP percentage in Borreguil de la Caldera record around 4000-3500 cal yr BP (Ramos-Román et al., 2016), high percentage of non-arboreal pollen around 3400 cal ka BP in Zoñar lake [Southern Córdoba Natural Reserve; (Martín-Puertas et al., 2008)], and lake desiccation at ca. 4100 and 2900 cal yr BP in Lake Siles (Carrión et al., 2007). Jalut et al. (2009) compared paleoclimatic records from different lakes in the western Mediterranean region and also suggested a dry phase between 4300 to 3400 cal yr BP, synchronous with this aridification phase. Furthermore, in the eastern Mediterranean basin other pollen studies show a decrease in arboreal pollen concentration toward more open landscapes around ~~3.7-ka-cal BP~~ [4 cal ka BP](#) (Magri, 1999).

Significant climatic changes also occurred in the Northern Hemisphere at those times and polar cooling and tropical aridity are observed at ca. 4200-3800 and 3500-2500 cal yr BP; (Mayewski et al., 2004), cold events in the North Atlantic [cold event 3 and 2; (Bond et al., 2001)], decrease in solar irradiance (Steinhilber et al., 2009) and humidity decreases in the eastern Mediterranean area at 4200 cal yr BP (Bar-Matthews et al., 2003) that could be related with global scale climate variability (Fig. 6). These generally dry phases between 4.5 and 2.8 in Padul-15-05 are generally in agreement with persistent positive NAO conditions during this time (Olsen et al., 2012).

The high-resolution Padul-15-05 record shows that climatic crises such as [the essentially global event at the one occurred at ca. 4200 cal yr BP, which seems to be recorded worldwide](#) (Booth et al., 2005), are ~~the product of the sum of more than one single~~ [actually multiple events in climatic event \(i.e., ca. 4500, 4300, 4000 cal yr BP\)](#) and thus are affected by climatic variability at centennial-scales [\(i.e., ca. 4500, 4300, 4000 cal yr BP\)](#).

### 5.2.2 Iberian-Roman Humid Period (~2600 to 1600 cal yr BP)

High relative humidity is recorded in the Padul-15-05 record between ca. 2600 and 1600 cal yr BP, synchronous with the well-known Iberian-Roman Humid Period (IRHP; between 2600 and

1600 cal yr BP; (Martín-Puertas et al., 2009). This is interpreted in our record due to an increase in the Mediterranean forest species at that time (Unit 1; ~~PA-ollen-Zone-2-b~~; Figs. 6 and 7). In addition, there is a simultaneous increase in *Botryococcus* algae, which is probably related to higher productivity during warmer conditions and relatively higher water level. Evidence of a wetter climate around this period has also been shown in ~~other regional records and~~ several alpine records from Sierra Nevada. For example, in the Laguna de la Mula core (Jiménez-Moreno et al. (2013) ~~studying a sediment record from the Laguna de la Mula, showed~~ an increase in deciduous *Quercus* ~~is~~ correlated with the maximum in algae between 2500 to 1850 cal yr BP, also evidencing the most humid period of the Late Holocene. A geochemical study from the Laguna de Río Seco (also in Sierra Nevada) also evidenced humid conditions around 2200 cal yr BP by the decrease in Saharan dust input and the increase in detritic sedimentation into the lake suggesting higher rainfall (Jiménez-Espejo et al., 2014). In addition, Ramos-Román et al. (2016) showed an increase in AP in the Borreguil de la Caldera record around 2200 cal yr BP, suggesting an increase in humidity at that time.

Other records from the Iberian Peninsula also show this pattern to wetter conditions during the IRHP. For example, high lake levels are recorded in Zoñar Lake in southern Spain between 2460 to 1600 cal yr BP, only interrupted by a relatively arid pulse between 2140 and 1800 cal yr BP (Martín-Puertas et al., 2009). An increase in rainfall is described in the central region of the Iberian Peninsula in a study from the Tablas de Daimiel National Park between 2100 and 1680 cal yr BP (Gil García et al., 2007). Deeper lake levels at around 2650 to 1580 cal yr BP, also interrupted by an short arid event at ca. 2125-1790 cal yr BP, were observed to the north, in the Iberian Range (Currás et al., 2012). The fact that the Padul-15-05 record also shows a relatively arid-cold event between 2150-2050 cal yr BP, just in the middle of this relative humid-warm period, seems to point to a common feature of centennial-scale climatic variability in many western Mediterranean and North Atlantic records (Fig. 6). Humid climate conditions at around 2500 cal yr BP are also interpreted in previous studies from lake level reconstructions from Central Europe (Magny, 2004). Increases in temperate deciduous forest are also observed in marine records from the Alboran Sea around 2600 to 2300 cal yr BP, also pointing to high relative humidity (Combourieu Nebout et al., 2009; Fletcher et al., 2007). Overall humid conditions between 2600 and 1600 cal yr BP seem to agree with predominant negative NAO reconstructions at that time, which would translate into greater winter (and thus more effective) precipitation in the area triggering ~~more~~ greater development of forest species in the area.

Generally warm conditions are interpreted between 1900 and 1700 cal yr BP in the Mediterranean Sea, with high sea surface temperatures (SSTs), and in the North Atlantic area, with the decrease in Drift Ice Index. In addition, persistent positive solar irradiance occurred at that time. The increase in *Botryococcus* algae reaching maxima during the IRHP also seems to point to very productive and perhaps warmer conditions in the Padul peat bog area (Fig. 6).

### 5.2.3 DA and MCA (~~—aridity between—~~1550 cal yr BP ~~and to~~ 600 cal yr BP)

Enhanced aridity occurred right after the IRHP in the Padul peat bog area ~~between ca. 1550 and 600 cal yr BP (ca. 400—1350 CE)~~. This is deduced in the Padul-15-05 record by a significant forest decline, with a prominent decrease in Mediterranean forest elements, an increase in ~~Cichorioideae~~ herbs ~~and the decline in the estimated water level~~ (Unit 1; ~~Pollen-Zone~~ PA-3; Figs. 4 and 7). In addition, our evidence suggests a transition from a shallow lake to a more ephemeral

wetland. ~~A significant change since the end of the IRHP took place in the lake environment, suggesting the transition from a shallow water table to an ephemeral environment.~~ This is ~~deduced~~ suggested by the disappearance of charophytes, a significant decrease in ~~Algae~~ algae component and higher Si and MS and lower TOC values (Unit 1; Figs. 6 and 7). Humans probably also contributed to enhancing erosion in the area during this last ~1550 cal yr BP. The significant change during the transition from Unit 1 to Unit 2 with a decrease in the pollen concentration and the increase in Cichorioideae could be due to enhanced pollen degradation as Cichorioideae have been found to be very resistant to pollen deterioration (Bottema, 1975). However, the occurrence of other pollen taxa (e.g. *Quercus*, *Ericaceae*, *Pinus*, *Poaceae*, *Olea*) showing climatic trends and increasing between ca. 1500-400 cal yr BP and a decrease in Cichorioideae in the last ~400 cal yr BP, when an increase in clastic material occurred, do not entirely support a preservation issue (see section of Human activity; 5.4).

This ~~arid~~ phase could be separated into two different periods. The first period occurred between ~~~ca. 1550 cal yr BP (ca. 400 CE)~~ and ~~ca. 1100 cal yr BP (~400 to ca. 900 CE)~~ and is characterized by a decreasing trend in Mediterranean forest and *Botryococcus* taxa ~~and the increase in Cichorioideae~~. This period corresponds with the Dark Ages [from ca. 500 to 900 CE; (Moreno et al., 2012)]. ~~A visual~~ eCorrelation between the ~~decrease~~ decline in Mediterranean forest, the increase in the Drift Ice Index in the North Atlantic record (cold event 1; Bond et al., 2001), the ~~decrease~~ decline in SSTs in the Mediterranean Sea and maxima in positive NAO reconstructions suggests drier and colder conditions during this time (Fig. 6). Other Mediterranean and central-European records agree with our climate interpretations, for example, a decrease in forest ~~extent is~~ pollen types is shown in a marine record from the Alboran Sea (Fletcher et al., 2013) and a decrease in lake levels is also observed in Central Europe (Magny et al., 2004) pointing to aridity during the DA. Evidences of aridity during the DA have been shown too in the Mediterranean part of the Iberian Peninsula, for instance, cold and arid conditions were suggested in the northern Betic Range by the increase in xerophytic herbs around 1450 and 750 cal yr BP (Carrión et al., 2001b) and in southeastern Spain by a forest decline in lacustrine deposits around 1620 and 1160 cal yr BP (Carrión et al., 2003). Arid and colder conditions during the Dark Ages (around 1680 to 1000 cal yr BP) are also suggested for the central part of the Iberian Peninsula using a multiproxy study of a sediment record from the Tablas de Daimiel Lake (Gil García et al., 2007).

A second period that we could differentiate ~~within this overall arid phase~~ occurred around 1100 to 600 cal yr BP/900 to 1350 CE, during the well-known MCA (900 to 1300 CE after Moreno et al., 2012). During this period the Padul-15-05 record shows a slight increasing trend in the Mediterranean forest taxa with respect to the DA, but the decrease in *Botryococcus* and the ~~higher abundance of~~ increase in herbs still point to overall arid conditions. This change could be related to an increase in temperature, favoring the development of temperate forest species, and would agree with inferred increasing temperatures in the North Atlantic areas, as well as the increase in solar irradiance and the increase in SSTs in the Mediterranean Sea (Fig.7). This hypothesis would agree with the reconstruction of persistent positive NAO and overall warm conditions during the MCA in the western Mediterranean (see synthesis in Moreno et al., 2012). A similar pattern of increasing xerophytic vegetation during the MCA is observed in alpine peat bogs and lakes in the Sierra Nevada (Anderson et al., 2011; Jiménez-Moreno et al., 2013; Ramos-Román et al., 2016) and arid conditions are shown to occur during the MCA in southern



and eastern Iberian Peninsula deduced by increases in salinity and lower lake levels (Corella et al., 2013; Martín-Puertas et al., 2011). However, humid conditions have been reconstructed for the northwestern of the Iberian Peninsula at this time (Lebreiro et al., 2006; Moreno et al., 2012), as well as northern Europe (Martín-Puertas et al., 2008). The different pattern of precipitation between ~~the northwestern Iberian Peninsula~~ / northern Europe and the Mediterranean area is undoubtedly a function of the ~~well-known-NAO dipole in~~-precipitation ~~dipole pattern~~ (Trouet et al., 2009).

#### 5.2.4 The last ~600 cal yr BP: LIA (~600 to 100 cal yr BP/~1350 to-1850 CE) and IE (~100 cal yr BP to Present/~1850 CE-Present)

Two climatically ~~different-distinct~~ periods can be distinguished during the last ca. 600 ~~cal-yr BP~~ years (end of ~~PA-Zone-3b~~ to ~~Zone-PA-4~~; Fig. 4) in the area. However, the climatic signal is more difficult to interpret due to a higher human impact at that time. The first phase around ~~1350-600-1450-500 CE-cal yr BP~~ was characterized ~~by-as~~ increasing relative humidity by the decrease in xerophytes and the increase in Mediterranean forest taxa and *Botryococcus* after a period of decrease during the DA and MCA, corresponding to the LIA. The second phase is characterized here by the decrease in the Mediterranean forest around ~~1700-300-1850-100 CE-cal yr BP~~, pointing to a return to more arid conditions during the last part of the LIA (Figs. 4 and 7). This climatic pattern agrees with an increase in precipitation by the transition from positive to negative NAO mode and from warmer to cooler conditions in the North Atlantic area during the first phase of the LIA and a second phase characterized by cooler (cold event 0; Bond et al., 2001) and drier conditions (Fig. 6). A stronger variability in the SSTs is described in the Mediterranean Sea during the LIA (Fig. 6). Mayewski et al. (2004) described a period of climate variability during the Holocene at this time (600 to 150 cal yr BP) suggesting a polar cooling but more humid in some parts of the tropics. Regionally, (Morellón et al., 2011) also described a phase of more humid conditions between 1530 to 1750 CE (420 to 200 cal yr BP) in a lake sediment record from NE Spain. An alternation between wetter to drier periods during the LIA are also shown in the nearby alpine record from Borreguil de la Caldera in the Sierra Nevada mountain range (Ramos-Román et al., 2016).

The environmental transition from ephemeral ~~to-emerged-conditions~~, observed in the last ca. 1550 cal yr BP (Unit 1; Fig. 5), to emerged conditions ~~intensified~~ in the last ca. ~~300-400~~ cal yr BP. This is shown by the highest MS and Si values the increase in wetland plants and the stronger decrease in Ca and organic components (TOC) in the sediments in the uppermost part of the Padul-15-05 record (Unit 2; Figs. 3 and 6).

### 5.3 Centennial-scale variability

Time series analysis has become important in determining the recurrent periodicity of cyclical oscillations in paleoenvironmental sequences (e.g. Jiménez-Espejo et al., 2014; Ramos-Román et al., 2016; Rodrigo-Gámiz et al., 2014; Fletcher et al., 2013). This analysis also assists in understanding possible relationships between the paleoenvironmental proxy data and the potential triggers of the observed cyclical changes: i.e., solar activity, atmospheric, oceanic dynamics and climate evolution during the Holocene. The cyclostratigraphic analysis on the pollen (Mediterranean forest; regional signal), algae (*Botryococcus*; local signal) and TOC (local signal) times series from the Padul-15-05 record evidence centennial-scale cyclical patterns with periodicities around ~800, 680, 300, 240, 200 and 170 years above the 90 % confidence levels

(Fig. 7).

Previous cyclostratigraphic analysis in Holocene western Mediterranean records suggest cyclical climatic oscillations with periodicities around 1500 and 1750 yr (Fletcher et al., 2013; Jiménez-Espejo et al., 2014; Rodrigo-Gámiz et al., 2014). Other North Atlantic and Mediterranean records also present cyclicities in their paleoclimatic proxies of ca. 1600 yr (Bond et al., 2001; Debret et al., 2007; Rodrigo-Gámiz et al., 2014). However, this cycle is absent from the cyclostratigraphic analysis in the Padul-15-05 record (Fig. 7). In contrast, the spectral analysis performed in the Mediterranean forest time series from Padul peat bog record, pointing to cyclical hydrological changes, shows a significant ~800 yr cycle that could be related to solar variability (Damon and Sonett, 1991) or could be the second harmonic of the ca. ~1600 yr oceanic-related cycle (Debret et al., 2009). A very similar periodicity of ca. 760 yr is detected in the *Pinus* forest taxa, also pointing to humidity variability, from the alpine Sierra Nevada site of Borreguil de la Caldera and seems to show that this is a common feature of cyclical paleoclimatic oscillation in the area.

A significant ~680 cycle is shown in the *Botryococcus* time series most likely suggesting recurrent centennial-scale changes in temperature (productivity) and water availability. A similar cycle is shown in the *Artemisia* signal in an alpine record from Sierra Nevada (Ramos-Román et al., 2016). This cycle around ~650 yr is also observed in a marine record from the Alboran Sea, and was interpreted as the secondary harmonic of the 1300 yr cycle that those authors related with cyclic thermohaline circulation and sea surface temperature changes (Rodrigo-Gámiz et al., 2014).

A statistically significant ~300 yr cycle is shown in the Mediterranean forest taxa and TOC from the Padul-15-05 record suggesting shorter-scale variability in water availability. This cycle is also observed in the cyclical *Pinus* pollen data from Borreguil de la Caldera at higher elevations in the Sierra Nevada (Ramos-Román et al., 2016). This cycle could be principally related to NAO variability as observed by Olsen et al. (2012), which follows variations in humidity observed in the Padul-15-05 record. NAO variability also regulates modern precipitation in the area.

The *Botryococcus* and TOC time series shows variability with a periodicity around ~240, 200 and 164 yrs. Sonnet et al. (1984) described a significant cycle in solar activity around ~208 yr (Suess solar cycle), which could have triggered our ~200 cyclicity. The observed ~240 yr periodicity in the Padul-15-05 record could be either related to variations in solar activity or due to the mixed effect of the solar together with the ~300 yr NAO-interpreted cycle and could point to a solar origin of the centennial-scale NAO variations as suggested by previously published research (Lukianova and Alekseev, 2004; Zanchettin et al., 2008). Finally, a significant ~170 yr cycle has been observed in both the Mediterranean forest taxa and *Botryococcus* times series from the Padul-15-05 record. A similar cycle (between 168-174 yr) was also described in the alpine pollen record from Borreguil de la Caldera in Sierra Nevada (Ramos-Román et al., 2016), which shows that it is a significant cyclical pattern in climate, probably precipitation, in the area. This cycle could be related to the previously described ~170 yr cycle in the NAO index (Olsen et al., 2012), which would agree with the hypothesis of the NAO controlling millennial- and centennial-scale environmental variability during the Late Holocene in the area (García-Alix et

al., 2017; Ramos-Román et al., 2016).

#### 5.4 Human activity

Humans probably had an impact in the area since Prehistoric times, however, the Padul-15-05 multiproxy record shows a more significant human impact during the last ca. 1550 cal yr BP, which intensified in the last ~~ca. 500 years~~ cal-yr BP (since 1450 CE to Present). This is deduced by, a significant increase in nitrophilous plant taxa such as Cichorioideae, Convolvulaceae, Polygonaceae and *Plantago* and the increase in some NPP such as *Tilletia*, coprophilous fungi and thecamoebians (Unit 2; ~~PA-ollen-Zone~~ 4; Fig. 4). Most of these pollen taxa and NPPs are described in other southern Iberian paleoenvironmental records as indicators of land uses, for instance, *Tilletia* and covarying nitrophilous plants ~~Cichorioideae and Convolvulaceae~~ have been described as indicators of farming (e.g. Carrión et al., 2001a). Thecamoebians also show a similar trend and have also been detected in other areas being related to nutrient enrichment as consequences of livestock (Fig. 8). The stronger increase in Cichorioideae have also been described as indicators of animal grazing in areas subjected to intense use of the territory (Mercuri et al., 2006). Interestingly, these taxa ~~being-began~~ to decline around ca. 400 cal yr BP (~1450 CE), coinciding with the higher increase in detritic material into the basin. We could then interpreted this increase in Cichorioideae as greater in livestock activity in the surroundings of the lake during this period, which is supported by the increase in these other proxies related with animal husbandry.

Climatically, this event coincides with the start of persistent negative NAO conditions in the area (Trouet et al., 2009), which could have further triggered more rainfall and more detritic input into the basin. (Bellin et al., 2011) in a study from the Betic Cordillera (southern Iberian Peninsula) demonstrate that soil erosion increase in years with higher rainfall and this could be intensified by human impact. Nevertheless, in a study in the southeastern part of the Iberian Peninsula (Bellin et al., 2013) suggested that major soil erosion could have occurred by the abandonment of agricultural activities in the mountain areas as well as the abandonment of irrigated terrace systems during the Christian Reconquest. Enhanced soil erosion at this time is also supported by the increase in *Glomus* type (Fig. 4).

An important change in the sedimentation in the environment is observed during the last ca. ~~300~~ 400 cal yr BP marked by the stronger increase in MS and Si values. This higher increase in detritics occurred during an increase in other plants related with human and land uses such as Polygonaceae, Amaranthaceae, Convolvulaceae, Plantago, Apiaceae and Cannabaceae-Urticaceae type (Land Use Plants; Fig. 8). This was probably related to drainage canals in ~~with~~ the Padul peat bog ~~water-drainage-by-humans-using-canals~~ in the late XVIII century for cultivation purposes (Villegas Molina, 1967). The increase in wetland vegetation and higher values of Poaceae could be due to cultivation of cereals or by an increase in the population of *Phragmites australis* (also a Poaceae), very abundant in the Padul peat bog margins at present due to the increase in drained land surface.

The uppermost part (last ca. 100 cal yr BP) of the pollen record from Padul-15-05 shows an increasing trend in some arboreal taxa at that time, including Mediterranean forest, *Olea* and *Pinus* (Fig. 4). This change is most likely of human origin and generated by the increase in *Olea* cultivation in the last two centuries, also observed in many records from higher elevation sites from Sierra Nevada, and *Pinus* and other Mediterranean species reforestation in the 20<sup>th</sup> century

(Anderson et al., 2011; Jiménez-Moreno and Anderson, 2012; Jiménez-Moreno et al., 2013; Ramos-Román et al., 2016).

## 6 Conclusions

Our multiproxy (~~i.e. lithology, geochemistry, paleontology~~) analysis from the Padul-15-05 sequence has provided a detailed climate reconstruction for the last 4700 ca yr BP for the Padul peat bog area and the western Mediterranean. This study, supported by the comparison with other Mediterranean and North Atlantic records suggests a link between vegetation, atmospheric dynamics and insolation and solar activity during the Late Holocene ~~in this area~~. A climatic aridification trend occurred during the Late Holocene in the Sierra Nevada and the western Mediterranean ~~area~~, probably linked with ~~the an~~ orbital-scale ~~decreasing~~ declining trend in summer insolation. This long-term trend is modulated by centennial-scale climate variability as shown by the pollen (Mediterranean forest taxa), algae (*Botryococcus*) and sedimentary and geochemical data in the Padul record. These events ~~are in~~ can be correlated ~~ion~~ with regional and global scale climate variability. ~~and~~ Cold and arid pulses identified in this study around the 4200 and 3000 cal yr BP ~~that are identified in this study seem to be~~ synchronous with cold events recorded in the North Atlantic and decreases in precipitation in the Mediterranean area, probably linked to persistent positive NAO mode. Moreover, one of the most important humid and warmer periods during the Late Holocene in the Padul area coincides in time with the well-known IRHP, characterized by warm and humid conditions in the Mediterranean and North Atlantic regions and overall negative NAO conditions. A drastic ~~decrease~~ decline in Mediterranean forest taxa, trending towards an open landscape, and pointing to colder conditions with ~~and~~ enhanced aridity, occurred in two steps (DA and end of the LIA) during the last ~~ca.~~ 1550 cal yr BP. However, this trend was slightly superimposed by a more arid but warmer event coinciding with the MCA and a cold but wetter event during the first part of the LIA. Besides natural climatic and environmental variability, strong evidences exists for ~~there seems to be~~ intense human activities in the area during the last the last ~~ca.~~ 1550 ~~cal yr BP~~ years. This suggests that the natural aridification trend during the Late Holocene, which produced a progressive decrease in the Mediterranean forest taxa in the Padul area, could have been intensified by human activities, notably in the last centuries. ~~This suggests that the natural aridification trend during the Late Holocene in the western Mediterranean region could have been intensified due to the higher human activity in this area.~~

Furthermore, time series analyses done in the Padul-15-05 record show centennial-scale changes in the environment and climate that are coincident with the periodicities observed in solar, oceanic and NAO reconstructions and could show a close cause-and-effect linkage between them.

## Acknowledgements

This work was supported by the project P11-RNM-7332 funded by Consejería de Economía, Innovación, Ciencia y Empleo de la Junta de Andalucía, the project CGL2013-47038-R funded by Ministerio de Economía y Competitividad of Spain and fondo Europeo de desarrollo regional FEDER and the research group RNM0190 (Junta de Andalucía). M. J. R.-R. acknowledges the PhD funding provided by Consejería de Economía, Innovación, Ciencia y Empleo de la Junta de Andalucía (P11-RNM-7332). J.C. acknowledges the PhD funding provided by Ministerio de Economía y Competitividad (CGL2013-47038-R). A.G.-A. was also supported by a Ramón y Cajal Fellowship RYC-2015-18966 of the Spanish Government (Ministerio de Economía y Competitividad). Javier Jaimez (CIC-UGR) is thanked for graciously helping with the coring, the drilling equipment and logistics. [We also would like to thanks to the editor \(Nathalie Combourieu-Nebout\) and Graciela Gil-Romera, Laura Sadori and an anonymous reviewer for their comments and suggestions which improved the manuscript.](#)

## 856    **References**

- 857    Abel-Schaad, D. and López-Sáez, J. A.: Vegetation changes in relation to fire history and human  
858    activities at the Peña Negra mire (Bejar Range, Iberian Central Mountain System, Spain) during  
859    the past 4,000 years, *Veg. Hist. Archaeobotany*, 22(3), 199–214, doi:10.1007/s00334-012-0368-  
860    9, 2013.
- 861    Alfaro, P., Galinod-Zaldievar, J., Jabaloy, A., López-Garrido, A. C. and Sanz de Galdeano, C.:  
862    Evidence for the activity and paleoseismicity of the Padul fault (Betic Cordillera, Southern Spain)  
863    [Evidencias de actividad y paleosismicidad de la falla de Padul (Cordillera Bética, sur de  
864    España)], *Acta Geol. Hisp.*, 36(3–4), 283–297, 2001.
- 865    Alpert, P., Baldi, M., Ilani, R., Krichak, S., Price, C., Rodó, X., Saaroni, H., Ziv, B., Kishcha, P.,  
866    Barkan, J., Mariotti, A. and Xoplaki, E.: Chapter 2 Relations between climate variability in the  
867    Mediterranean region and the tropics: ENSO, South Asian and African monsoons, hurricanes  
868    and Saharan dust, *Dev. Earth Environ. Sci.*, 4(C), 149–177, doi:10.1016/S1571-9197(06)80005-4,  
869    2006.
- 870    Anderson, R. S., Jiménez-Moreno, G., Carrión, J. S. and Pérez-Martínez, C.: Postglacial history of  
871    alpine vegetation, fire, and climate from Laguna de Río Seco, Sierra Nevada, southern Spain,  
872    *Quat. Sci. Rev.*, 30(13–14), 1615–1629, doi:https://doi.org/10.1016/j.quascirev.2011.03.005,  
873    2011.
- 874    Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A. and Hawkesworth, C. J.: Sea–land  
875    oxygen isotopic relationships from planktonic foraminifera and speleothems in the Eastern  
876    Mediterranean region and their implication for paleorainfall during interglacial intervals,  
877    *Geochim. Cosmochim. Acta*, 67(17), 3181–3199, doi:https://doi.org/10.1016/S0016-  
878    7037(02)01031-1, 2003.
- 879    Bellin, N., Vanacker, V., van Wesemael, B., Solé-Benet, A. and Bakker, M. M.: Natural and  
880    anthropogenic controls on soil erosion in the internal betic Cordillera (southeast Spain), *Catena*,  
881    87(2), 190–200, doi:10.1016/j.catena.2011.05.022, 2011.
- 882    Bellin, N., Vanacker, V. and De Baets, S.: Anthropogenic and climatic impact on Holocene  
883    sediment dynamics in SE Spain: A review, *Quat. Int.*, 308–309, 112–129,  
884    doi:10.1016/j.quaint.2013.03.015, 2013.
- 885    Beug, H.-J.: *Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete*, Fisch.  
886    Stuttg., 61, 2004.
- 887    Blaauw, M.: Methods and code for “classical” age-modelling of radiocarbon sequences, *Quat.*  
888    *Geochronol.*, 5(5), 512–518, doi:https://doi.org/10.1016/j.quageo.2010.01.002, 2010.
- 889    Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M. N., Showers, W., Hoffmann, S., Lotti-  
890    Bond, R., Hajdas, I. and Bonani, G.: Persistent Solar Influence on North Atlantic Climate During  
891    the Holocene, *Science*, 294(5549), 2130, doi:10.1126/science.1065680, 2001.

- Booth, R. K., Jackson, S. T., Forman, S. L., Kutzbach, J. E., E. A. Bettis, I., Kreigs, J. and Wright, D. K.: A severe centennial-scale drought in midcontinental North America 4200 years ago and apparent global linkages, *The Holocene*, 15(3), 321–328, doi:10.1191/0959683605hl825ft, 2005.
- Bottema, S.: The interpretation of pollen spectra from prehistoric settlements (with special attention of *Liguliflorae*), *Palaeohistoria*, 17, 17–35, 1975.
- Carrión, J. S.: Patterns and processes of Late Quaternary environmental change in a montane region of southwestern Europe, *Quat. Sci. Rev.*, 21(18–19), 2047–2066, doi:https://doi.org/10.1016/S0277-3791(02)00010-0, 2002.
- Carrión, J. S., Munuera, M., Dupré, M. and Andrade, A.: Abrupt vegetation changes in the Segura Mountains of southern Spain throughout the Holocene, *J. Ecol.*, 89(5), 783–797, doi:10.1046/j.0022-0477.2001.00601.x, 2001b.
- Carrión, J. S., Andrade, A., Bennett, K. D., Navarro, C. and Munuera, M.: Crossing forest thresholds: inertia and collapse in a Holocene sequence from south-central Spain, *The Holocene*, 11(6), 635–653, doi:10.1191/09596830195672, 2001a.
- Carrión, J. S., Fernández, S., Jiménez-Moreno, G., Fauquette, S., Gil-Romera, G., González-Sampériz, P. and Finlayson, C.: The historical origins of aridity and vegetation degradation in southeastern Spain, *J. Arid Environ.*, 74(7), 731–736, doi:https://doi.org/10.1016/j.jaridenv.2008.11.014, 2010.
- Carrión, J. S., Sánchez-Gómez, P., Mota, J. F., Yll, R. and Chaín, C.: Holocene vegetation dynamics, fire and grazing in the Sierra de Gádor, southern Spain, *Holocene*, 13(6), 839–849, doi:10.1191/0959683603hl662rp, 2003.
- Carrión, J. S., Fuentes, N., González-Sampériz, P., Quirante, L. S., Finlayson, J. C., Fernández, S. and Andrade, A.: Holocene environmental change in a montane region of southern Europe with a long history of human settlement, *Quat. Sci. Rev.*, 26(11–12), 1455–1475, doi:https://doi.org/10.1016/j.quascirev.2007.03.013, 2007.
- Combourieu-Nebout, N., Peyron, O., Dormoy, I., Desprat, S., Beaudouin, C., Kotthoff, U. and Marret, F.: Rapid climatic variability in the west Mediterranean during the last 25 000 years from high resolution pollen data, *Clim Past*, 5(3), 503–521, doi:10.5194/cp-5-503-2009, 2009.
- Corella, J. P., Stefanova, V., El Anjoumi, A., Rico, E., Giralt, S., Moreno, A., Plata-Montero, A. and Valero-Garcés, B. L.: A 2500-year multi-proxy reconstruction of climate change and human activities in northern Spain: The Lake Arreo record, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 386, 555–568, doi:10.1016/j.palaeo.2013.06.022, 2013.
- Currás, A., Zamora, L., Reed, J. M., García-Soto, E., Ferrero, S., Armengol, X., Mezquita-Joanes, F., Marqués, M. A., Riera, S. and Julià, R.: Climate change and human impact in central Spain

928 during Roman times: High-resolution multi-proxy analysis of a tufa lake record (Somolinos,  
929 1280m asl), *Catena*, 89(1), 31–53, doi:10.1016/j.catena.2011.09.009, 2012.

930 Damon, P. E. and Sonett, C. P.: Solar and terrestrial components of the atmospheric C-14  
931 variation spectrum. In: Sonett, C.P., Giampapa, M.S., Matthews, M.S. (Eds.), *The Sun in Time*.  
932 University of Arizona Press, Tucson, AZ, USA., 1991.

933 Debret, M., Bout-Roumazelles, V., Grousset, F., Desmet, M., McManus, J. F., Massei, N., Sebag,  
934 D., Petit, J.-R., Copard, Y. and Trentesaux, A.: The origin of the 1500-year climate cycles in  
935 holocene north-atlantic records, *Clim. Past*, 3(4), 569–575, 2007.

936 Debret, M., Sebag, D., Crosta, X., Massei, N., Petit, J.-R., Chapron, E. and Bout-Roumazelles, V.:  
937 Evidence from wavelet analysis for a mid-Holocene transition in global climate forcing, *Quat.*  
938 *Sci. Rev.*, 28(25–26), 2675–2688, doi:https://doi.org/10.1016/j.quascirev.2009.06.005, 2009.

939 Delgado, J., Alfaro, P., Galindo-Zaldivar, J., Jabaloy, A., Lopez Garrido, A. and Sanz de Galdeano,  
940 C.: Structure of the Padul-Nigüelas basin (S Spain) from H/V ratios of ambient noise: application  
941 of the method to study peat and coarse sediments, *Pure Appl. Geophys.*, 159(11), 2733–2749,  
942 2002.

943 Domingo-García, M., Fernández-Rubio, R., Lopez, J. and González, C.: Aportación al  
944 conocimiento de la Neotectónica de la Depresión del Padul (Granada), *Tecniterrae*, 53, 6–16,  
945 1983.

946 Ejarque, A., Anderson, R. S., Simms, A. R. and Gentry, B. J.: Prehistoric fires and the shaping of  
947 colonial transported landscapes in southern California: A paleoenvironmental study at Dune  
948 Pond, Santa Barbara County, *Quat. Sci. Rev.*, 112, 181–196,  
949 doi:https://doi.org/10.1016/j.quascirev.2015.01.017, 2015.

950 Faegri, K. and Iversen, J.: *Textbook of Pollen Analysis*. Wiley, New York., 1989.

951 Fletcher, W. J. and Sánchez-Goñi, M. F.: Orbital- and sub-orbital-scale climate impacts on  
952 vegetation of the western Mediterranean basin over the last 48,000 yr, *Quat. Res.*, 70(3), 451–  
953 464, doi:10.1016/j.yqres.2008.07.002, 2008.

954 Fletcher, W. J., Debret, M. and Sánchez-Goñi, M. F.: Mid-Holocene emergence of a low-  
955 frequency millennial oscillation in western Mediterranean climate: Implications for past  
956 dynamics of the North Atlantic atmospheric westerlies, *The Holocene*, 23(2), 153–166,  
957 doi:10.1177/0959683612460783, 2013.

958 Florenzano, A., Marignani, M., Rosati, L., Fascetti, S. and Mercuri, A. M.: Are Cichorieae an  
959 indicator of open habitats and pastoralism in current and past vegetation studies?, *Plant*  
960 *Biosyst. - Int. J. Deal. Asp. Plant Biol.*, 149(1), 154–165, doi:10.1080/11263504.2014.998311,  
961 2015.



- 962 Florschütz, F., Amor, J. M. and Wijmstra, T. A.: Palynology of a thick quaternary succession in  
963 southern Spain, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 10(4), 233–264,  
964 doi:[http://dx.doi.org/10.1016/0031-0182\(71\)90049-6](http://dx.doi.org/10.1016/0031-0182(71)90049-6), 1971.
- 965 García-Alix, A., Jimenez-Espejo, F. J., Lozano, J. A., Jiménez-Moreno, G., Martinez-Ruiz, F.,  
966 Sanjuán, L. G., Jiménez, G. A., Alfonso, E. G., Ruiz-Puertas, G. and Anderson, R. S.:  
967 Anthropogenic impact and lead pollution throughout the Holocene in Southern Iberia, *Sci. Total*  
968 *Environ.*, 449, 451–460, doi:<https://doi.org/10.1016/j.scitotenv.2013.01.081>, 2013.
- 969 García-Alix, A., Jiménez-Espejo, F. J., Toney, J. L., Jiménez-Moreno, G., Ramos-Román, M. J.,  
970 Anderson, R. S., Ruano, P., Queralt, I., Delgado Huertas, A. and Kuroda, J.: Alpine bogs of  
971 southern Spain show human-induced environmental change superimposed on long-term  
972 natural variations, *Sci. Rep.*, 7(1), 7439, doi:10.1038/s41598-017-07854-w, 2017.
- 973 van Geel, B., Hallewas, D. P. and Pals, J. P.: A late holocene deposit under the Westfriese Zeedijk  
974 near Enkhuizen (Prov. of Noord-Holland, The Netherlands): Palaeoecological and archaeological  
975 aspects, *Rev. Palaeobot. Palynol.*, 38(3), 269–335, doi:[http://dx.doi.org/10.1016/0034-](http://dx.doi.org/10.1016/0034-6667(83)90026-X)  
976 [6667\(83\)90026-X](http://dx.doi.org/10.1016/0034-6667(83)90026-X), 1983.
- 977 van Geel, B., Coope, G. R. and Hammen, T. V. D.: Palaeoecology and stratigraphy of the  
978 lateglacial type section at Usselo (the Netherlands), *Rev. Palaeobot. Palynol.*, 60(1), 25–129,  
979 doi:[http://dx.doi.org/10.1016/0034-6667\(89\)90072-9](http://dx.doi.org/10.1016/0034-6667(89)90072-9), 1989.
- 980 Gil García, M. J., Ruiz Zapata, M. B., Santisteban, J. I., Mediavilla, R., López-Pamo, E. and Dabrio,  
981 C. J.: Late holocene environments in Las Tablas de Daimiel (south central Iberian peninsula,  
982 Spain), *Veg. Hist. Archaeobotany*, 16(4), 241–250, doi:10.1007/s00334-006-0047-9, 2007.
- 983 Gil-Romera, G., Carrión, J. S., Pausas, J. G., Sevilla-Callejo, M., Lamb, H. F., Fernández, S. and  
984 Burjachs, F.: Holocene fire activity and vegetation response in South-Eastern Iberia, *Quat. Sci.*  
985 *Rev.*, 29(9), 1082–1092, doi:10.1016/j.quascirev.2010.01.006, 2010.
- 986 Grimm, E. C.: CONISS: a FORTRAN 77 program for stratigraphically constrained cluster analysis  
987 by the method of incremental sum of squares, *Comput. Geosci.*, 13(1), 13–35,  
988 doi:[http://dx.doi.org/10.1016/0098-3004\(87\)90022-7](http://dx.doi.org/10.1016/0098-3004(87)90022-7), 1987.
- 989 Guy-Ohlson, D.: Botryococcus as an aid in the interpretation of palaeoenvironment and  
990 depositional processes, *Rev. Palaeobot. Palynol.*, 71(1), 1–15,  
991 doi:[http://dx.doi.org/10.1016/0034-6667\(92\)90155-A](http://dx.doi.org/10.1016/0034-6667(92)90155-A), 1992.
- 992 Hurrell, J. W.: Decadal Trends in the North Atlantic Oscillation: Regional Temperatures and  
993 Precipitation, *Science*, 269(5224), 676, doi:10.1126/science.269.5224.676, 1995.
- 994 Jalut, G., Dedoubat, J. J., Fontugne, M. and Otto, T.: Holocene circum-Mediterranean vegetation  
995 changes: Climate forcing and human impact, *Quat. Int.*, 200(1–2), 4–18,  
996 doi:<https://doi.org/10.1016/j.quaint.2008.03.012>, 2009.

- 1p97 Jiménez-Espejo, F. J., García-Alix, A., Jiménez-Moreno, G., Rodrigo-Gámiz, M., Anderson, R. S.,  
 998 Rodríguez-Tovar, F. J., Martínez-Ruiz, F., Giralte, S., Delgado Huertas, A. and Pardo-Igúzquiza, E.:  
 999 Saharan aeolian input and effective humidity variations over western Europe during the  
 1000 Holocene from a high altitude record, *Chem. Geol.*, 374–375, 1–12,  
 1001 doi:10.1016/j.chemgeo.2014.03.001, 2014.
- 1p02 Jiménez-Moreno, G. and Anderson, R. S.: Holocene vegetation and climate change recorded in  
 1003 alpine bog sediments from the Borreguiles de la Virgen, Sierra Nevada, southern Spain, *Quat.*  
 1004 *Res.*, 77(1), 44–53, doi:10.1016/j.yqres.2011.09.006, 2012.
- 1p05 Jiménez-Moreno, G., García-Alix, A., Hernández-Corbalán, M. D., Anderson, R. S. and Delgado-  
 1006 Huertas, A.: Vegetation, fire, climate and human disturbance history in the southwestern  
 1007 Mediterranean area during the late Holocene, *Quat. Res.*, 79(2), 110–122,  
 1008 doi:https://doi.org/10.1016/j.yqres.2012.11.008, 2013.
- 1p09 Jiménez-Moreno, G., Rodríguez-Ramírez, A., Pérez-Asensio, J. N., Carrión, J. S., López-Sáez, J. A.,  
 1010 Villarías-Robles, J. J. R., Celestino-Pérez, S., Cerrillo-Cuenca, E., León, Á. and Contreras, C.:  
 1011 Impact of late-Holocene aridification trend, climate variability and geodynamic control on the  
 1012 environment from a coastal area in SW Spain, *Holocene*, 25(4), 607–617,  
 1013 doi:10.1177/0959683614565955, 2015.
- 1p14 Kalugin, I., Daryin, A., Smolyaninova, L., Andreev, A., Diekmann, B. and Khlystov, O.: 800-yr-long  
 1015 records of annual air temperature and precipitation over southern Siberia inferred from  
 1016 Teletskoye Lake sediments, *Quat. Res.*, 67(3), 400–410,  
 1017 doi:https://doi.org/10.1016/j.yqres.2007.01.007, 2007.
- 1p18 Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M. and Levrard, B.: A long-term  
 1019 numerical solution for the insolation quantities of the Earth, *A&A*, 428(1), 261–285,  
 1020 doi:10.1051/0004-6361:20041335, 2004.
- 1p21 Lebreiro, S. M., Francés, G., Abrantes, F. F. G., Diz, P., Bartels-Jónsdóttir, H. B., Stoyanowski, Z.  
 1022 N., Gil, I. M., Pena, L. D., Rodrigues, T., Jones, P. D., Nombela, M. A., Alejo, I., Briffa, K. R., Harris,  
 1023 I. and Grimalt, J. O.: Climate change and coastal hydrographic response along the Atlantic  
 1024 Iberian margin (Tagus Prodelta and Muros Ría) during the last two millennia, *Holocene*, 16(7),  
 1025 1003–1015, doi:10.1177/0959683606hl990rp, 2006.
- 1p26 Lillios, K. T., Blanco-González, A., Drake, B. L. and López-Sáez, J. A.: Mid-late Holocene climate,  
 1027 demography, and cultural dynamics in Iberia: A multi-proxy approach, *Quat. Sci. Rev.*, 135, 138–  
 1028 153, doi:https://doi.org/10.1016/j.quascirev.2016.01.011, 2016.
- 1p29 López-Sáez, J. A., Abel-Schaad, D., Pérez-Díaz, S., Blanco-González, A., Alba-Sánchez, F., Dorado,  
 1030 M., Ruiz-Zapata, B., Gil-García, M. J., Gómez-González, C. and Franco-Múgica, F.: Vegetation  
 1031 history, climate and human impact in the Spanish Central System over the last 9000 years,  
 1032 *Quat. Int.*, 353, 98–122, doi:https://doi.org/10.1016/j.quaint.2013.06.034, 2014.

- 1033 Lukianova, R. and Alekseev, G.: Long-Term Correlation Between the Nao and Solar Activity, *Sol.*  
1034 *Phys.*, 224(1), 445–454, doi:10.1007/s11207-005-4974-x, 2004.
- 1035 Magny, M.: Holocene climate variability as reflected by mid-European lake-level fluctuations  
1036 and its probable impact on prehistoric human settlements, *Quat. Int.*, 113(1), 65–79,  
1037 doi:https://doi.org/10.1016/S1040-6182(03)00080-6, 2004.
- 1038 Magny, M., Peyron, O., Sadori, L., Ortu, E., Zanchetta, G., Vannière, B. and Tinner, W.:  
1039 Contrasting patterns of precipitation seasonality during the Holocene in the south- and north-  
1040 central Mediterranean, *J. Quat. Sci.*, 27(3), 290–296, doi:10.1002/jqs.1543, 2012.
- 1041 Magri, D.: Late Quaternary vegetation history at Lagaccione near Lago di Bolsena (central Italy),  
1042 *Rev. Palaeobot. Palynol.*, 106(3–4), 171–208, doi:https://doi.org/10.1016/S0034-  
1043 6667(99)00006-8, 1999.
- 1044 Marchal, O., Cacho, I., Stocker, T. F., Grimalt, J. O., Calvo, E., Martrat, B., Shackleton, N.,  
1045 Vautravers, M., Cortijo, E., Van Kreveld, S., Andersson, C., Koç, N., Chapman, M., Sbaifi, L.,  
1046 Duplessy, J.-C., Sarnthein, M., Turon, J.-L., Duprat, J. and Jansen, E.: Apparent long-term cooling  
1047 of the sea surface in the northeast Atlantic and Mediterranean during the Holocene, *Quat. Sci.*  
1048 *Rev.*, 21(4–6), 455–483, doi:10.1016/S0277-3791(01)00105-6, 2002.
- 1049 Martín-Puertas, C., Valero-Garcés, B. L., Mata, M. P., González-Sampériz, P., Bao, R., Moreno, A.  
1050 and Stefanova, V.: Arid and humid phases in southern Spain during the last 4000 years: the  
1051 Zoñar Lake record, Córdoba, *The Holocene*, 18(6), 907–921, doi:10.1177/0959683608093533,  
1052 2008.
- 1053 Martín-Puertas, C., Valero-Garcés, B. L., Brauer, A., Mata, M. P., Delgado-Huertas, A. and Dulski,  
1054 P.: The Iberian-Roman Humid Period (2600-1600 cal yr BP) in the Zoñar Lake varve record  
1055 (Andalucía, southern Spain), *Quat. Res.*, 71(2), 108–120, doi:10.1016/j.yqres.2008.10.004, 2009.
- 1056 Martín-Puertas, C., Valero-Garcés, B. L., Mata, M. P., Moreno, A., Giral, S., Martínez-Ruiz, F.  
1057 and Jiménez-Espejo, F.: Geochemical processes in a Mediterranean Lake: A high-resolution  
1058 study of the last 4,000 years in Zoñar Lake, southern Spain, *J. Paleolimnol.*, 46(3), 405–421,  
1059 doi:10.1007/s10933-009-9373-0, 2011.
- 1060 Mayewski, P. A., Rohling, E. E., Stager, J. C., Karlén, W., Maasch, K. A., Meeker, L. D., Meyerson,  
1061 E. A., Gasse, F., Kreveld, S. van, Holmgren, K., Lee-Thorp, J., Rosqvist, G., Rack, F., Staubwasser,  
1062 M., Schneider, R. R. and Steig, E. J.: Holocene climate variability, *Quat. Res.*, 62(3), 243–255,  
1063 doi:https://doi.org/10.1016/j.yqres.2004.07.001, 2004.
- 1064 Mercuri, A. M., Accorsi, C. A., Mazzanti, M. B., Bosi, G., Cardarelli, A., Labate, D., Marchesini, M.  
1065 and Grandi, G. T.: Economy and environment of Bronze Age settlements – Terramaras – on the  
1066 Po Plain (Northern Italy): first results from the archaeobotanical research at the Terramara di  
1067 Montale, *Veg. Hist. Archaeobotany*, 16(1), 43, doi:10.1007/s00334-006-0034-1, 2006.

- 1068 Morellón, M., Valero-Garcés, B., Vegas-Vilarrúbia, T., González-Sampériz, P., Romero, Ó.,  
1069 Delgado-Huertas, A., Mata, P., Moreno, A., Rico, M. and Corella, J. P.: Lateglacial and Holocene  
1070 palaeohydrology in the western Mediterranean region: The Lake Estanya record (NE Spain),  
1071 Quat. Sci. Rev., 28(25–26), 2582–2599, doi:<https://doi.org/10.1016/j.quascirev.2009.05.014>,  
1072 2009.
- 1073 Morellón, M., Valero-Garcés, B., González-Sampériz, P., Vegas-Vilarrúbia, T., Rubio, E.,  
1074 Rieradevall, M., Delgado-Huertas, A., Mata, P., Romero, Ó., Engstrom, D. R., López-Vicente, M.,  
1075 Navas, A. and Soto, J.: Climate changes and human activities recorded in the sediments of Lake  
1076 Estanya (NE Spain) during the Medieval Warm Period and Little Ice Age, J. Paleolimnol., 46(3),  
1077 423–452, doi:10.1007/s10933-009-9346-3, 2011.
- 1078 Morellón, M., Anselmetti, F. S., Ariztegui, D., Brushulli, B., Sinopoli, G., Wagner, B., Sadori, L.,  
1079 Gilli, A. and Pambuku, A.: Human–climate interactions in the central Mediterranean region  
1080 during the last millennia: The laminated record of Lake Butrint (Albania), Spec. Issue Mediterr.  
1081 Holocene Clim. Environ. Hum. Soc., 136(Supplement C), 134–152,  
1082 doi:10.1016/j.quascirev.2015.10.043, 2016.
- 1083 Moreno, A., Cacho, I., Canals, M., Grimalt, J. O., Sánchez-Goñi, M. F., Shackleton, N. and Sierro,  
1084 F. J.: Links between marine and atmospheric processes oscillating on a millennial time-scale. A  
1085 multi-proxy study of the last 50,000 yr from the Alboran Sea (Western Mediterranean Sea),  
1086 Quat. Sci. Rev., 24(14–15), 1623–1636, doi:<https://doi.org/10.1016/j.quascirev.2004.06.018>,  
1087 2005.
- 1088 Moreno, A., Pérez, A., Frigola, J., Nieto-Moreno, V., Rodrigo-Gámiz, M., Martrat, B., González-  
1089 Sampériz, P., Morellón, M., Martín-Puertas, C., Corella, J. P., Belmonte, Á., Sancho, C., Cacho, I.,  
1090 Herrera, G., Canals, M., Grimalt, J. O., Jiménez-Espejo, F., Martínez-Ruiz, F., Vegas-Vilarrúbia, T.  
1091 and Valero-Garcés, B. L.: The Medieval Climate Anomaly in the Iberian Peninsula reconstructed  
1092 from marine and lake records, Quat. Sci. Rev., 43, 16–32,  
1093 doi:<https://doi.org/10.1016/j.quascirev.2012.04.007>, 2012.
- 1094 Nestares, T. and Torres, T. de: Un nuevo sondeo de investigación paleoambiental del  
1095 Pleistoceno y Holoceno en la turbera del Padul (Granada, Andalucía). Geogaceta 23, 99-102.,  
1096 1997.
- 1097 Olsen, J., Anderson, N. J. and Knudsen, M. F.: Variability of the North Atlantic Oscillation over  
1098 the past 5,200 years, Nat. Geosci, 5(11), 808–812, doi:10.1038/ngeo1589, 2012.
- 1099 Ortiz, J. E., Torres, T., Delgado, A., Julià, R., Lucini, M., Llamas, F. J., Reyes, E., Soler, V. and Valle,  
1100 M.: The palaeoenvironmental and palaeohydrological evolution of Padul Peat Bog (Granada,  
1101 Spain) over one million years, from elemental, isotopic and molecular organic geochemical  
1102 proxies, Org. Geochem., 35(11–12), 1243–1260,  
1103 doi:<https://doi.org/10.1016/j.orggeochem.2004.05.013>, 2004.

- 1104 Paillard, D., Labeyrie, L. and Yiou, P.: Macintosh Program performs time-series analysis, *Eos*  
1105 *Trans. Am. Geophys. Union*, 77(39), 379–379, doi:10.1029/96EO00259, 1996.
- 1106 Pérez Raya, F. and López Nieto, J.: Vegetación acuática y helofítica de la depresión de Padul  
1107 (Granada), *Acta Bot Malacit.*, 16(2), 373–389, 1991.
- 1108 Pons, A. and Reille, M.: The holocene- and upper pleistocene pollen record from Padul  
1109 (Granada, Spain): A new study, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 66(3), 243–263,  
1110 doi:http://dx.doi.org/10.1016/0031-0182(88)90202-7, 1988.
- 1111 Ramos-Román, M. J., Jiménez-Moreno, G., Anderson, R. S., García-Alix, A., Toney, J. L., Jiménez-  
1112 Espejo, F. J. and Carrión, J. S.: Centennial-scale vegetation and North Atlantic Oscillation  
1113 changes during the Late Holocene in the southern Iberia, *Quat. Sci. Rev.*, 143, 84–95,  
1114 doi:https://doi.org/10.1016/j.quascirev.2016.05.007, 2016.
- 1115 Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Ramsey, C. B., Buck, C. E., Cheng,  
1116 H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Haflidason, H., Hajdas, I.,  
1117 Hatté, C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B.,  
1118 Manning, S. W., Niu, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Staff, R. A.,  
1119 Turney, C. S. M. and van der Plicht, J.: IntCal13 and Marine13 Radiocarbon Age Calibration  
1120 Curves 0–50,000 Years cal BP, *Radiocarbon*, 55(4), 1869–1887, doi:10.2458/azu\_js\_rc.55.16947,  
1121 2013.
- 1122 Riera, S., Wansard, G. and Julià, R.: 2000-year environmental history of a karstic lake in the  
1123 Mediterranean Pre-Pyrenees: the Estanya lakes (Spain), *{CATENA}*, 55(3), 293–324,  
1124 doi:https://doi.org/10.1016/S0341-8162(03)00107-3, 2004.
- 1125 Riera, S., López-Sáez, J. A. and Julià, R.: Lake responses to historical land use changes in  
1126 northern Spain: The contribution of non-pollen palynomorphs in a multiproxy study, *Rev.*  
1127 *Palaeobot. Palynol.*, 141(1–2), 127–137, doi:https://doi.org/10.1016/j.revpalbo.2006.03.014,  
1128 2006.
- 1129 Roberts, N., Brayshaw, D., Kuzucuoğlu, C., Perez, R. and Sadori, L.: The mid-Holocene climatic  
1130 transition in the Mediterranean: Causes and consequences, *The Holocene*, 21(1), 3–13,  
1131 doi:10.1177/0959683610388058, 2011.
- 1132 Rodrigo-Gámiz, M., Martínez-Ruiz, F., Rodríguez-Tovar, F. J., Jiménez-Espejo, F. J. and Pardo-  
1133 Igúzquiza, E.: Millennial- to centennial-scale climate periodicities and forcing mechanisms in the  
1134 westernmost Mediterranean for the past 20,000 yr, *Quat. Res.*, 81(1), 78–93,  
1135 doi:https://doi.org/10.1016/j.yqres.2013.10.009, 2014.
- 1136 Sadori, L., Jahns, S. and Peyron, O.: Mid-Holocene vegetation history of the central  
1137 Mediterranean, *The Holocene*, 21(1), 117–129, doi:10.1177/0959683610377530, 2011.
- 1138 Sadori, L., Giraudi, C., Masi, A., Magny, M., Ortu, E., Zanchetta, G. and Izdebski, A.: Climate,  
1139 environment and society in southern Italy during the last 2000 years. A review of the

1140 environmental, historical and archaeological evidence, Spec. Issue Mediterr. Holocene Clim.  
 1141 Environ. Hum. Soc., 136(Supplement C), 173–188, doi:10.1016/j.quascirev.2015.09.020, 2016.

1142 Sanz de Galdeano, C., El Hamdouni, R. and Chacón, J.: Neotectónica de la fosa del Padul y del  
 1143 Valle de Lecrín, Itiner. Geomorfológicos Por Andal. Orient. Publicacions Univ. Barc. Barc., 65–81,  
 1144 1998.

1145 Sicre, M.-A., Jalali, B., Martrat, B., Schmidt, S., Bassetti, M.-A. and Kallel, N.: Sea surface  
 1146 temperature variability in the North Western Mediterranean Sea (Gulf of Lion) during the  
 1147 Common Era, Earth Planet. Sci. Lett., 456, 124–133,  
 1148 doi:http://dx.doi.org/10.1016/j.epsl.2016.09.032, 2016.

1149 Sonett, C. P. and Suess, H. E.: Correlation of bristlecone pine ring widths with atmospheric <sup>14</sup>C  
 1150 variations: a climate-Sun relation, Nature, 307(5947), 141–143, doi:10.1038/307141a0, 1984.

1151 Steinhilber, F., Beer, J. and Fröhlich, C.: Total solar irradiance during the Holocene, Geophys.  
 1152 Res. Lett., 36(19), n/a–n/a, doi:10.1029/2009GL040142, 2009.

1153 Trouet, V., Esper, J., Graham, N. E., Baker, A., Scourse, J. D. and Frank, D. C.: Persistent Positive  
 1154 North Atlantic Oscillation Mode Dominated the Medieval Climate Anomaly, Science, 324(5923),  
 1155 78, doi:10.1126/science.1166349, 2009.


1156 Villegas Molina, F.: Laguna de Padul: Evolución geológico-histórica, Estud. Geográficos, 28(109),  
 1157 561, 1967.

1158 Zanchettin, D., Rubino, A., Traverso, P. and Tomasino, M.: Impact of variations in solar activity  
 1159 on hydrological decadal patterns in northern Italy, J. Geophys. Res. Atmospheres, 113(D12),  
 1160 n/a–n/a, doi:10.1029/2007JD009157, 2008.


1161  
 1162  
 1163  
 1164  
 1165  
 1166  
 1167  
 1168  
 1169  
 1170  
 1171  
 1172  
 1173  
 1174  
 1175




Figures and tables




**Figure 1.** Location of Padul peat bog in Sierra Nevada, southern Iberian Peninsula. Panel on the left is the map of the vegetation belts in the Sierra Nevada (Modified from REDIAM. Map of the vegetation series of Andalucía: [http://laboratoriolediam.cica.es/VisorGenerico/?tipo=WMS&url=http://www.juntadeandalucia.es/medioambiente/mapwms/REDIAM\\_Series\\_Vegetacion\\_Andalucia?](http://laboratoriolediam.cica.es/VisorGenerico/?tipo=WMS&url=http://www.juntadeandalucia.es/medioambiente/mapwms/REDIAM_Series_Vegetacion_Andalucia?)). The inset map is the Google earth image of the Iberian Peninsula in the Mediterranean region. Panel on the right is the Google earth image (<http://www.google.com/earth/index.html>) of Padul peat bog area showing the coring locations.



**Figure 2.** Photo of the Padul-15-05 sediment core with the age-depth model showing the part of the record that was studied here (red rectangle). The sediment accumulation rates (SAR) between individual segments are marked. See the body of the text for the explanation of the age reconstructions.




**Figure 3.** Lithology, facies interpretation with paleontology, magnetic susceptibility (MS), and geochemical (X-ray fluorescence (XRF) and total organic carbon (TOC) data from the Padul-15-05 record. XRF elements are represented normalized by the total counts. (a) Magnetic susceptibility (MS; SI [units](#)). (b) Strontium normalized (Sr; norm.). (c) Bromine norm. (Br; norm.). (d) Calcium normalized. (Ca; norm.). (e) Silica normalized (Si; norm.). (f) Potassium normalized (K; norm.). (g) Titanium normalized (Ti; norm.). (h) Iron normalized (Fe; norm.). (i) Zirconium normalized (Zr; norm.). (j) Total organic carbon (TOC %).



1208


1209 **Figure 4.** Percentages of selected pollen taxa and non-pollen palynomorphs (NPPs) from the  
1210 Padul-15-05 record, ~~represented~~calculated with respect to terrestrial pollen sum. Silhouettes  
1211 show 7-time exaggerations of pollen percentages. Pollen zonation, —pollen concentration  
1212 (grains/cc) and lithology ~~is~~are shown on the right. Tree and shrubs are showing in green, herbs  
1213 and grasses in yellow, aquatics in dark blue, algae in blue, fungi in brown and thecamoebians in  
1214 beige. The Mediterranean forest taxa category is composed of *Quercus* total, *Olea*, *Phillyrea* and  
1215 *Pistacia*. The xerophyte group includes *Artemisia*, *Ephedra*, and Amaranthaceae. PA = Pollen  
1216 zones.

1217




**Figure 5.** Estimated lake level evolution and regional palynological component from the last ca. 4700 yr based on the synthesis of determinate proxies from the Padul-15-05 record: (a) Proxies used to estimate the water table evolution from the Padul-15-05 record (proxies were resampled at 50 yr (lineal interpolation) using Past software [http://palaeo-electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm)). [(a.1) Magnetic Susceptibility (MS) in SI; (a.2) Silica normalized (Si; norm.); (a.3) Calcium normalized (Ca; norm.); (a.4) Bromine normalized (Br; norm.); (a.5) Strontium normalized (Sr; norm.); (a.6) Hygrophytes (%); (a.7) Poaceae (%); (a.8) Algae (%) (a.9) Total organic carbon (TOC %)] (b) Mediterranean forest taxa, with a smoothing of three-point in bold. Pink and blue shading indicates Holocene arid and humid regionally events, respectively. See the body of the text for the explanation of the lake level reconstruction. Mediterranean forest smoothing was made using Analyseries software (Paillard et al., 1996) (Paillard et al., 1996). PA = Pollen Zones; CA = Copper Age; BA = Bronze Age; IA = Iron Age; IRHP = Iberian Roman Humid Period; DA = Dark Ages; MCA = Medieval Climate Anomaly; LIA = Little Ice Age; IE = Industrial Era.





**Figure 6.** Comparison of the last ca. 4700 yr between different pollen taxa from the Padul-15-05 record, summer insolation for the Sierra Nevada latitude, eastern Mediterranean humidity and North Atlantic temperature. (a) *Botryococcus* from the Padul-15-05 record, with a smoothing of three-point in bold (this study). (b) Drift Ice Index (reversed) from the North Atlantic (Bond et al., 2001). (c) Summer insolation calculated for 37° N (Laskar et al., 2004). (d) Mediterranean forest taxa from the Padul-15-05 record, with a smoothing of three-point in bold (this study). (e) Alkenone-SSTs from the Gulf of Lion (Sicre et al., 2016), with a smoothing of four-point in bold. (f) North Atlantic Oscillation (NAO) index from a climate proxy reconstruction from Morocco and Scotland (Trouet et al., 2009). (g) North Atlantic Oscillation (NAO) index (reversed) from a climate proxy reconstruction from Greenland (Olsen et al., 2012). (h) Total solar irradiance reconstruction from cosmogenic radionuclide from a Greenland ice core (Steinhilber et al., 2009), with a smoothing of twenty-one-point in bold. Yellow and blue shading correspond with arid (and cold) and humid (and warm) periods, respectively. Grey dash lines show a tentative correlation between arid and cold conditions and the decrease in the Mediterranean forest and *Botryococcus*. Mediterranean forest, *Botryococcus* and solar irradiance smoothing was made using Analyseries software (Paillard et al., 1996), Alkenone-SSTs smoothing was made using Past software ([http://palaeo-electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm)). A linear r (Pearson) correlation was calculated between *Botryococcus* (detrended) and Drift Ice Index (Bond et al., 2001;  $r = -0.63$ ;  $p < 0.0001$ ; between ca. 4700 to 1500 cal ka BP –  $r = -0.48$ ;  $p < 0.0001$  between 4700 and -65 cal yr BP). Previously, the data were detrended (only in *Botryococcus*), resampled at 70-yr (linear interpolation) in order to obtain equally spaced time series and smoothed to three-point average. CA = Copper Age; BA = Bronze Age; IA = Iron Age; IRHP = Iberian Roman Humid Period; DA = Dark Ages; MCA = Medieval Climate Anomaly; LIA = Little Ice Age; IE = Industrial Era.



**Figure 7.** Spectral analysis of (a) Mediterranean forest taxa and (b) *Botryococcus* (mean sampling space = 47 yr) and (c) TOC (mean sampling space = 78 yr) from the Padul-15-05. The significant periodicities above confident level are shown. Confidence level 90 % (blue line), 95 % (green line), 99 % (green dash line) and AR (1) red noise (red line). Spectral analysis was made with Past software ([http://palaeo-electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm)).




Fig. 8. Comparison of the last ca. 4700 yr between regional climatic proxies and local human activity indicators from the Padul-15-05 record. (a) Mediterranean forest taxa, with a smoothing of three-point in bold. (b) Local human activities indicators [(b.1) Total organic carbon (TOC %), soil erosion indicator; (b.2) Si normalized (Si, norm.), soil erosion indicator; (b.3) Poaceae (%), lake drained and/or cultivars indicator; (b.4) Land Use Plants (%), cultivar indicator; (b.5) Cichorioideae (%), livestock occurrence indicator; (b.6) *Tilletia* (%), farming indicator; (b.7) Sordariales (%), livestock indicator; (b.8) Thecamoebians undiff. (%), livestock indicator]. Degraded yellow to red shading correspond with the time when we have evidence of human shaping the environment since ca. 1550 cal yr BP to Present. Previously to that period there is a lack of clear evidences of human impact in the area. Land use plants is composed by Polygonaceae, Amaranthaceae, Convolvulaceae, *Plantago*, Apiaceae and Cannabaceae-Urticaceae type.

1284 **Table 1.** Age data for Padul-15-05 record. All ages were calibrated using R-code package ‘clam  
 1285 2.2’ employing the calibration curve IntelCal 13 (Reimer et al., 2013) at 95 % of confident range.

1286 \*Sample number assigned at radiocarbon laboratory

Laboratory number	Core	Material	Depth (cm)	Age ( $^{14}\text{C}$ yr BP $\pm 1\sigma$ )	Calibrated age (cal yr BP) 95 % confidence interval	Median age (cal yr BP)
Reference ages			0	2015CE	-65	-65
D-AMS 008531	Padul-13-01	Plant remains	21.67	103 $\pm$ 24	23-264	127
Poz-77568	Padul-15-05	Org. bulk sed.	38.46	1205 $\pm$ 30	1014-1239	1130
BETA-437233	Padul-15-05	Plant remains	46.04	2480 $\pm$ 30	2385-2722	2577
Poz-77569	Padul-15-05	Org. bulk sed.	48.21	2255 $\pm$ 30	2158-2344	2251
BETA-415830	Padul-15-05	Shell	71.36	3910 $\pm$ 30	4248-4421	4343
BETA-437234	Padul-15-05	Plant remains	76.34	3550 $\pm$ 30	3722-3956	3838
BETA-415831	Padul-15-05	Org. bulk sed.	92.94	3960 $\pm$ 30	4297-4519	4431
Poz-74344	Padul-15-05	Plant remains	122.96	4295 $\pm$ 35	4827-4959	4871
BETA-415832	Padul-15-05	Plant remains	150.04	5050 $\pm$ 30	5728-5900	5814
Poz-77571	Padul-15-05	Plant remains	186.08	5530 $\pm$ 40	6281-6402	6341
Poz-74345	Padul-15-05	Plant remains	199.33	6080 $\pm$ 40	6797-7154	6935
BETA-415833	Padul-15-05	Org. bulk sed.	217.36	6270 $\pm$ 30	7162-7262	7212
Poz-77572	Padul-15-05	Org. bulk sed.	238.68	7080 $\pm$ 50	7797-7999	7910
Poz-74347	Padul-15-05	Plant remains	277.24	8290 $\pm$ 40	9138-9426	9293
BETA-415834	Padul-15-05	Plant remains	327.29	8960 $\pm$ 30	9932-10221	10107

1287

1288



**Table 2.** Linear r (Pearson) correlation between geochemical elements from the Padul-15-05 record. Statistical treatment was performed using the Past software ([http://palaeo-electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm)).

	<b>Si</b>	<b>K</b>	<b>Ca</b>	<b>Ti</b>	<b>Fe</b>	<b>Zr</b>	<b>Br</b>	<b>Sr</b>
<b>Si</b>		8.30E-80	2.87E-34	7.47E-60	3.22E-60	5.29E-44	0.001152	7.79E-09
<b>K</b>	0.98612		7.07E-29	6.05E-60	8.20E-68	1.77E-51	0.00030317	5.38E-12
<b>Ca</b>	-0.88096	-0.84453		6.09E-42	5.81E-39	8.10E-34	0.35819	0.26613
<b>Ti</b>	0.96486	0.96501	-0.91794		1.74E-74	1.12E-57	0.074223	8.88E-07
<b>Fe</b>	0.96546	0.97577	-0.90527	0.98224		2.77E-66	0.051072	3.32E-08
<b>Zr</b>	0.92566	0.94789	-0.8783	0.96109	0.97398		0.054274	7.16E-08
<b>Br</b>	-0.31739	-0.3506	-0.091917	-0.17755	-0.19372	-0.19116		4.03E-18
<b>Sr</b>	-0.53347	-0.61629	0.11113	-0.46426	-0.51386	-0.50295	0.72852	