

Response to Reviewers' comments on Manuscript 'Simulation of the Greenland Ice sheet over two glacial-interglacial cycles: Investigating a sub-ice shelf melt parameterisation and relative sea level forcing in an ice sheet-ice shelf model.

5 Please find an attached revised manuscript, with and without track changes. We hope this highlights all the revisions that were made to typesetting and following the Reviewers' suggestions.

Reviewer 2:

10

1. I am still confused about the usage of eustatic sea level (ESL) and sea level equivalent (SLE) acronyms. It seems to me that in the first part of the introduction you use "ESL" for both eustatic sea level and the unit of sea level equivalent, while in the second part you refer to "ESL" as the eustatic sea level. I suggest to be more precise in these definitions.

15

Thank you for this comment. We did feel that we used throughout the manuscript the term 'ice volume equivalent sea level', to represent what the reviewer refers to as eustatic sea level. However, we agree, that within many scientific publications there are many different terms used to define eustatic sea level/global mean sea level/ sea level equivalent. We have therefore revised the term ice-volume equivalent sea level (ESL) to the term global mean sea level (GMSL), as this is the terminology used in the two most recent comprehensive publications which discuss eustatic/global mean sea level; Dutton et al., 2015 and Clark & Tarasov 2014. In the Table 3 information, we have added that this number is calculated from the ice volume. We feel this is necessary as the 'global mean sea level' as calculated from an ice sheet model simulated ice volume is different from that calculated from a Glacial isostatic adjustment model, as the latter includes the deformation of the ocean floor.

20

2. This manuscript has many points in common with Tabone et al., 2017, since both works tackle the same topic from two different approaches and have been published in CPD around the same time. Their model sub-shelf melting parameterisation (SSM), which depends on, allows the GrIS to expand onto the continental shelf in glacial times. Thus, I would not say that there isn't any ice-sheet model which is able to simulate a reliable GrIS glacial extent and ice volume (lines 60-63, pag. 2), since this is in contrast to what is found in Tabone et al., 2017. Also, since their sub-shelf melting (SSM) parameterisation is not based on changes in water depth, as it is in this manuscript, but rather on the PD-LGM ocean temperature anomaly, I suggest to spend a few words about it. It could be included, for example, in the

35

40 description of the SSM schemes used to address the expansion of the GrIS beyond the PD margin (lines 72-83, pag.1-2).

Thank you for mentioning to include your recent publication. We have now added a short note to mention that this publication has resolved the previous mismatch between generating a larger GrIS at glacial maximum (see line 61-63). However, as
45 this publication and methods within were not available to the authors of this study, during our modelling and method development we do not want to include extensive reference to the study as this implies that the information within this study may have influenced the results of this paper.

50 3. The description of the RSL forcing (Section 3.3) is clearer than in the first version of the manuscript. However, there is one point which is not completely crystal to me. If I understood well, you are able to create a "predicted" Δ WD signal by taking Δ GNL and Δ RNL from the GIA model and Δ RL from the ice-sheet model, provided with a ELRA scheme (following Eq. 6). Then you compare this signal to the Δ WD obtained from Eq. 5 to
55 get the "missing" Δ GL component (lines 291-293, pag. 8) and you see that this term is negligible for your purposes. What is not clear to me is how you directly produce Fig. 3f, i.e. the total Δ WD signal from Eq. 5. Maybe you directly know the terms Δ GT and Δ RT. If this is the case, where these total geoid (Δ GT) and solid earth (Δ RT) terms come from?

60 We have revised the text in this section, we hope this make the method clearer.

65

70

Simulation of the Greenland Ice sheet over two glacial-interglacial cycles: Investigating a sub-ice shelf melt parameterisation and relative sea level forcing in an ice sheet-ice shelf model.

Sarah L. Bradley^{1,2}, Thomas J. Reerink¹, Roderik S.W. van de Wal¹, Michiel M. Helsen¹

1 Institute for Marine and Atmospheric research Utrecht, Utrecht University, Utrecht, The Netherlands

2 Department of Geoscience & Remote Sensing, Delft University of Technology, Delft, The Netherlands

Correspondence to: Sarah L. Bradley (d80ngv@gmail.com)

Abstract. Observational evidence, including offshore moraines and sediment cores confirm that at the Last Glacial Maximum (LGM) the Greenland ice sheet (GrIS) expanded to a significantly larger spatial extent than seen at present, grounding into Baffin Bay and out onto the continental shelf break. Given this larger spatial extent and its close proximity to the neighboring Laurentide (LIS) and Innuitian Ice sheet (IIS), it is likely these ice-sheets will have had a strong non-local influence on the spatial and temporal behaviour of the GrIS. Most previous paleo ice-sheet modelling simulations recreated an ice-sheet that either did not extend out onto the continental shelf, or utilized a simplified marine ice parameterisation which did not fully include the effect of ice shelves, or neglected the sensitivity of the GrIS to this non-local bedrock signal from the surrounding ice-sheets.

In this paper, we investigated the evolution of the GrIS over the two most recent glacial-interglacial cycles (240 kyr BP to present-day) using the ice-sheet-ice shelf model, IMAU-ICE. We investigated the solid earth influence of the LIS and IIS via an offline relative sea level (RSL) forcing generated by a Glacial Isostatic Adjustment (GIA) model. This RSL forcing controlled, via changes in the water depth below the ice shelves, the spatial and temporal pattern of sub-ice shelf melting, which was parameterized as a function of changes in the water depth.

In the ensemble of simulations, at the glacial maximums, the GrIS coalesced with the IIS to the north and expanded to the continental shelf break to the south west but remained too restricted to the north east. In terms of the global mean sea level contribution, at the Last Interglacial (LIG) and LGM the ice-sheet added 1.46 m and -2.59 m respectively. This LGM contribution by the GrIS is considerably higher (~1.26 m) than most previous studies whereas the contribution to the LIG highstand is lower (~0.7 m). The spatial and temporal behaviour of the northern margin was highly variable in all simulations, controlled by the sub-ice shelf melting which was dictated by the RSL forcing and the glacial history of the IIS and LIS. In contrast, the southwestern part of the ice-sheet was insensitive to these forcings, with a uniform response in all simulations controlled by the surface air temperature, derived from ice cores.

1. Introduction

There have been many ice-sheet modelling studies of the glacial-interglacial evolution of the Northern hemisphere ice sheets (NHIS) (including the Greenland Ice sheet (GrIS) and/or Laurentide Ice Sheet (LIS) (Charbit et al., 2007; Greve et al., 1999; Helsen et al., 2013; Ritz et al., 1996; Quiquet et al., 2013), in which there was no expansion of the ice-sheet beyond the present-day (PD) coastline during glacial periods. The ice-sheet model in these studies modelled solely the evolution of grounded ice, where the edge of the grounded ice margin was determined by the flotation criterion. However, the wealth of new observational data infers that at glacial maximums the GrIS extended beyond the PD coastline, grounding out onto the continental shelf (Vasskog et al., 2015 and references therein Sect. 2). This shows there is a mismatch between the observed and the modelled extents.

A review publication by Dutton et al., (2015) stated that the exact magnitude and contribution of the various global ice sheets to global mean sea level (GMSL) during the Last Interglacial (LIG, 130-115 kyr BP) is still largely unresolved. From the analysis of far-field sea level records, it is estimated to have reached a peak between 6-9 m above PD. However, the contribution from the GrIS is poorly constrained and its reconstructed spatial extent highly variable (Vasskog et al., 2015). Estimates from ice-sheet modelling based studies of the contribution to the LIG highstand range between 0.6 and 3.5 m (Cuffey and Marshall, 2000; Dutton et al., 2015; Helsen et al., 2013; Robinson et al., 2011; Stone et al., 2013). Also, Clark and Tarasov, (2014) highlight that closing the Last Glacial Maximum (LGM) GMSL budget is becoming increasingly problematic. This is mostly due to the reduction in the estimated contribution from the Antarctic Ice sheet (AIS), derived from both modelling and observational studies. In addition, Glacial Isostatic Adjustment (GIA) modelling studies have estimated the contribution of the GrIS to the LGM GMSL budget to be ~ 5m (Lecavalier et al., 2014), whereas most ice-sheet modelling based studies indicate significantly less, typically < 2.5 m (average < 1m) (Fyke et al., 2011; Letreguilly et al., 1991; Ritz et al., 1996). These lower estimates are possibly caused by restricting the glacial maximum extent to the PD coastline. Consequently, the number of ice-sheet modelling based studies which simulate a sufficiently large GrIS during glacial periods, both in terms of maximum spatial extent and total contribution to the GMSL budget are limited, and as a consequence resolving the GrIS GMSL contribution over the last two glacial-interglacial cycles remains problematic. We note however, that the recent Tabone et al., (2018) study which does address this.

There have been two ice-sheet modelling based approaches to address the expansion of the grounded ice-sheet beyond the PD margin. In the first approach, often referred to as *a marine parameterisation*, ice is permitted to flow and ground beyond the PD coastline to a specified 'critical water depth', regardless of the ice thickness. This critical water depth is either a function of changes in GMSL or constrained by a series of masks reconstructed from observational data sets (Zweck and Huybrechts, 2005). This approach has been adopted in many ice-sheet modelling studies, solving only for grounded ice and reconstructed an extended GrIS during glacial periods (i.e. (Huybrechts, 2002; Lecavalier et al., 2014; Simpson et al., 2009). However, rather than the ice-sheet evolving freely, it is preconditioned to match with the observational data and does not use any physically based principles.

The second approach includes ice-shelf dynamics in combination with a calculation of sub-ice shelf melting (SSM). The sub-ice shelf melt is calculated by a parameterisation which is typically

based on changes in water depth, estimated using a GMSL forcing. This heuristic approach allows the ice-sheet to expand onto the continental shelf but not into the open ocean. There have been a number of publications which applied the second approach using, for instance the GRISLI ice-sheet model (Ritz et al., 2001). For example, Colleoni et al., (2014) parameterised the SSM as a uniform value in relation to changes in water depth to examine the growth of the NHIS during MIS7 and MIS5. During glacial periods, the reconstructed GrIS grounded across the Nares Strait, the Smith Sound and out onto the continental shelf to the NE and SW (see Fig. 8, Colleoni et al., 2014). Although in this reconstruction the ice-sheet retreated from the latter two offshore regions (NE and SW) by the LIG minimum (~ 115 kyr BP), it remain grounded across the Nares Strait, which is contrary to the observational data which are reviewed in Sect. 2.

Implicit in both these approaches is that the changes in paleo water depth surrounding the ice-sheet are driven by the GMSL forcing, generally derived either from a benthic $\delta^{18}\text{O}$ record (Lisiecki and Raymo, 2005) or by inverse forward modelling (Bintanja and van de Wal, 2008). However, sea level variations are in fact not simply GMSL (i.e with no spatial variations), but vary spatially due to numerous processes that dominate over different time scales, with GIA the dominant process on the time scales of this study (Kopp et al., 2015; Rovere et al., 2016).

This study advances the second approach, using the ice sheet-ice shelf model 'IMAU-ICE' (Sect. 3.1). The GrIS will be simulated over two glacial-interglacial cycles (240 kyr BP to PD), focusing on the parameterisations adopted for SSM (Sect. 3.2). Secondly, to investigate the influence of the spatial and temporal variability in sea level (or water depth) on the GrIS evolution, an offline forcing derived from a GIA model (Sect. 3.3) will be included. The first goal is to investigate if a larger than PD GrIS can be simulated for glacial maximum conditions, which is coherent with the observational data (Sect. 2), and thereby addresses the current mismatch between ice-sheet model and GIA based GrIS reconstructions. Secondly, we aim to evaluate the spatial and temporal sensitivity of the GrIS to changes in the SSM and the sea level forcing. Finally, we will address the question of the GrIS contribution to GMSL over the last two glacial-interglacial cycles.

2. Observational data

There have been numerous recent publications which have reviewed the wealth of new observational data that can be used to constrain the spatial and temporal history of the GrIS simulated by ice-sheet models (e.g Funder et al., 2011; Vasskog et al., 2015; Cofaigh et al., 2016). It is not the aim of this study to replicate this information, rather a selection of studies are outlined below which were useful to constrain the ice-sheet model simulations (summarised in Table 1, Fig. 1).

There are currently six Greenland ice core records (Fig. 1, white circles) that contain evidence for LIG age ice, and so were used to constrain the minimum extent that the ice-sheet reached during LIG (Fig. 1). Only simulations where these six sites remained glaciated at the LIG were considered valid. From the NEEM record (Dahl-Jensen et al., 2013) it is inferred that at 122 kyr BP, the surface elevation thinned by 130 ± 300 metres. The other five ice core sites remained ice covered, including Dye-3 (Yau et al., 2016). Additionally, analysis of Sr-Nd-Pb isotope ratios in offshore material

collected from Erik Drift (Colville et al., 2011) infer that the southern margin retreated to a smaller than PD extent, but that the ice-sheet did not undergo complete deglaciation during the LIG.

190 Constraining the offshore extent at the Penultimate Glacial maximum (PGM) or earlier
glaciations is complicated as the older geomorphological evidence (i.e moraines) is overridden by the
subsequent readvances. As a consequence, the preservation of offshore sediments is limited.
Therefore, we assumed that the ice extent during the PGM and the LGM are equal. The aim with all
simulations within the study was to reproduce a spatially expanded grounded ice-sheet which reached
the constraints given below during these two glacial maximums.

195 Offshore geomorphological evidence collected from numerous geophysical surveys indicate
that the ice-sheet grounded out onto the continental shelf (Table 1, Fig. 1), specifically to the shelf
break along the SW, north and central east at the LGM. This evidence includes moraines, grounding
zone wedges (Hogan et al., 2016), large scale glacial lineations (Cofaigh et al., 2004) and glacio-
marine sediments dated and analysed from offshore cores (Jennings et al., 2006). Table 1 provides an
overview of the asynchronous nature of the timing of retreat from this expanded glacial maximum
200 towards the PD margin.

Through the expansion of Petermann and Humboldt Glacier at the NW margin into the Kane
and Hall Basin and the Nares Strait (Fig. 1), the ice-sheet coalesced with the Innuitian Ice sheet (IIS) at
the glacial maximums (LGM or PGM). The grounded ice margin reached south into the north of Baffin
Bay and out along the Arctic coastline to the north. Dating from one of the few sediment cores from the
205 offshore NW margin (Table 1), Jennings et al., (2011) documented that the retreat of the grounded ice
from the Kane and Hall Basin initiated after 10.3 kyr BP. The margin retreated in an ‘unzipping
manner’, first from west (Kane Basin) and later to the east (Hall Basin), driven in part by the retreat of
IIS back onto Ellesmere Island. The final opening of the connection between the Arctic Ocean and
Baffin Bay, via the Nares Strait did not occur until after 9 kyr BP, implying that this region was one of
210 the last regions to deglaciate. The retreat of the grounded ice-sheet across the Nares Strait and back to
the PD margin was a key feature which was used to constrain the simulations, and if ice remained
grounded across this margin at PD, the model simulation was rejected.

215 Along the NE margin, Evans et al., (2009) concluded that the ice-sheet advanced out onto the
middle to outer shelf, covering the Westwind trough (open blue square, Fig. 1). It grounded close to (as
indicated by ice-rafted debris (IRD)) but did not extend as far as the Fram Strait, limited by the
continental shelf break (see Fig. 1). No dated material was recovered so the timing is unresolved.

Progressing further south, the lateral extent and timing are better constrained (Table 1) due to
the greater availability of data. Retreat from the central east outer shelf initiated by ~ 15 kyr BP (blue
star, Fig.1), stabilising on the inner shelf at 10 kyr BP (green star, Fig. 1) and reaching the PD margin
220 by 7.4 kyr BP (red star, Fig. 1) (Cofaigh et al., 2004; Evans et al., 2002). Along the SE margin, the
retreat from the outer to inner shelf is highly asynchronous, retreating from the outer Kangerdlugssuaq
trough at ~ 17.8 kyr BP (dark blue triangle, Fig. 1) (earlier than from the central east), reaching the
inner fjord by 11.8 kyr BP at Kangerdlugssuaq (open blue triangle, Fig. 1) and by 10.8 kyr BP at
Sermilik (red cross, Fig. 1). It is suggested that the timing of retreat across this region is strongly
225 influenced by the warm incursion of the Irminger current (Dyke et al., 2014).

The SW region of Greenland, around Disko Bugt and the Uumannaq trough is one of the more extensively studied regions of the ice-sheet, with a range of observational data confirming that the ice-sheet grounded out onto the continental shelf break (Cofaigh et al., 2013; Jennings et al., 2014; Sheldon et al., 2016; Winsor et al., 2015). The retreat from the outer shelf (cluster of red triangles, Fig. 1) between 19.3-18.6 kyr BP is inferred to have been driven by either a change in sea level and/or the ongoing gradual rise in the boreal summer insolation rather than changes in ocean temperatures. The margin stabilised at the middle shelf near the Hellefiske moraine (open red circle, Fig. 1), retreating at 12.24 kyr BP and reaching the inner shelf by 10.9 kyr BP. The question of whether a change in sea level could initiate such a retreat is just one aspect that the inclusion of a RSL forcing in this study will address. The retreat from the outer shelf edge in the vicinity of the Uumannaq fjord (cluster of green triangles, Fig. 1) was later, after 14.9 kyr BP, reaching the outer Uumannaq trough by 10.8 kyr BP (Sheldon et al., 2016). Against this background of geological evidence, we evaluated our model results as presented in Sect. 4.

3. Method.

3.1. IMAU-ICE: ice sheet-ice shelf model.

As the aim of this study is to simulate the expansion onto and retreat from the continental shelf of the GrIS, it is essential to utilise an ice-sheet model which includes the possibility for ice shelves to ground and thereby the ice-sheet to expand beyond the PD margin. To achieve this, we used a 3D thermomechanical ice sheet-ice shelf model IMAU-ICE (previously known as ANICE) (de Boer et al., 2014). For regions of grounded ice, IMAU-ICE uses the commonly adopted SIA approximation (Hutter, 1983) to simulate ice velocities in combination with a 3D thermodynamical approach. For regions of floating ice, the ice-shelf velocities are approximated using the SSA solution (Macayeal, 1989). The model does not accurately solve for grounding line dynamics, rather the grounding line is defined as the transition between ice-sheet (grounded) and ice-shelf (floating) points using the flotation criterion. The complex marginal topography of Greenland, with narrow troughs with steep gradients, can lead to complications when adopting the usual SSA approach. To address this problem, a 2D one-sided surface height gradient discretisation scheme was included for ice-shelf points neighbouring the grounded line.

At regions within the ice-sheet where the basal temperature reaches pressure melting point, the ice-sheet is allowed to slide using a Weertman-type sliding law, which relates the sliding velocity (v_b), to the basal shear stress (τ_b) such that

$$v_b = A_s \frac{\tau_b^p}{z^q} \quad (1)$$

Where A_s is defined as the sliding coefficient which can be taken as inversely proportional to the bed roughness, z is the reduced normal load and p and q are spatially uniform constants over the ice-sheet domain. As the roughness at the base of ice-sheet is a relatively unknown quantity, a range of sliding coefficients (A_s) were investigated, between 0.04×10^{-10} and $1.8 \times 10^{-10} \text{ m}^8 \text{ N}^{-3} \text{ yr}^{-1}$.

Present-day input fields of the ice thickness, the surface elevation and the bed topography are taken from Bamber et al., (2013) with input climate fields (surface mass balance (SMB), refreezing, surface air temperature (SAT)) adapted from the RACMO2 dataset (van Angelen et al., 2014). All these external datasets are interpolated and projected onto the 20x20 km ice model grid using the mapping software OBLIMAP2.0 (Reerink et al., 2010; Reerink et al., 2016). The adopted OBLIMAP grid projection parameters were $\lambda = 371.5$, $\phi = 71.8$ and $\alpha = 7.15$.

First, IMAU-ICE was ran for 100 kyr, under a constant PD climate (using the input climate fields taken from the RACMO2 dataset), to reach an equilibrium state with the aim of replicating the observed PD configuration (Bamber et al., 2013). The sensitivity to the flow enhancement factor (m_{enh} , varied between 1-5) was investigated for the range of sliding coefficients (A_s). The resultant ice volume varied over the range of m_{enh} and A_s by $\sim 0.12 \times 10^{15} \text{ m}^3$, with all simulations producing an underprediction of the ice thickness across the centre of the ice-sheet and overprediction at the narrow outlet glaciers. Based on this preliminary evaluation, a value of $m_{\text{enh}} = 3.5$ was used in all simulations. Also as simulations with an $A_s = 1.8 \times 10^{-10} \text{ m}^8 \text{ N}^{-3} \text{ yr}^{-1}$ resulted in a significant retreat of the SW margin, the range of sliding coefficients was revised to 0.04×10^{-10} and $1.2 \times 10^{-10} \text{ m}^8 \text{ N}^{-3} \text{ yr}^{-1}$ in subsequent simulations. The output of these simulations was used as the initial conditions for the subsequent simulations of the two glacial-interglacial cycles.

Secondly, each simulation was ran for 240 kyr using a spatially uniform SAT forcing taken from Helsen et al., (2013) (Fig. 2a), combined with a SSM parametrisation (Sect.3.2) and sea level forcing (derived from a GIA model, Sect.3.3), to simulate the GrIS over the two glacial-interglacial cycles. As there is no GrIS SAT record that extends beyond 128 kyr BP, this SAT forcing record was produced by combining the Vostok ice core (Petit et al., 1999) with the GRIP ice core record (Johnsen et al., 2001) using the glacial-index method (Greve, 2005). We note that using a SAT forcing record derived from ice cores will not account for any spatial variability in the SAT during these two glacial-interglacial cycle. The SMB-gradient method (Helsen et al., 2012; Helsen et al., 2013) was applied at each time step to calculate a new SMB field resulting from this SAT forcing. In this approach, first this uniform temperature forcing (Fig. 2a) is converted into a spatially variable climate-driven surface elevation change using an atmosphere lapse rate of -7.4 K km^{-1} . Second, the SMB gradient fields are calculated based on a linear regression between this new surface elevation field and the mean SMB in an area with a radius of 150 km. With this approach, the spatially uniform temperature forcing (Fig. 2a) can be translated in the spatially varying SMB field and ensures that the local mass balance height feedback is captured. The resultant suite of simulations was evaluated using the observational data defined in Sect.2.

3.2. Parameterisation of sub-ice shelf melt (SSM)

As full physical based models including SSM are still under development, we investigated a SSM parameterisation (Fig. 3) primarily based around the assumption that for an increase in paleo water depth (or sea level) there will be a corresponding increase in the amount of SSM. Hence the SSM does not depend on temperature: temperature changes only affect the surface mass balance. In this method, the SSM increases with water depth (W_D) by a power law relation with a constant a , and exponent m .

$$SSM = aW_D^m \quad (2)$$

In order to conveniently fit this power law through two points (SSM1, W_{D1}) and (SSM2, W_{D2}), we solve:

$$305 \quad m = \frac{\ln\left(\frac{SSM2}{SSM1}\right)}{\ln\left(\frac{W_{D2}}{W_{D1}}\right)}; \quad a = \frac{SSM1}{W_{D1}^{2m}}; \quad (3)$$

The range of parameter values for SSM1, SSM2, W_{D1} (water depth1) and W_{D2} (water depth2) are listed in Table 2, with three examples illustrated on Fig. 3.

3.3. Relative Sea Level or Water Depth forcing

310 In this study, the output from a GIA model is incorporated into IMAU-ICE to examine the influence of spatial and temporal variability in the RSL forcing via the SSM parameterisation on the expansion and retreat of the GrIS.

Sea Level (or water depth), $W_D(\theta, \psi, t)$ can be defined as the vertical distance between the equilibrium ocean surface, the geoid $G(\theta, \psi, t)$ and the solid earth surface $R(\theta, \psi, t)$ (bed topography) (Mitrovica and Milne, 2003). A change in the water depth $\Delta W_D(\theta, \psi, t)$ can result from any vertical
315 deformation in these two surfaces, and is defined as:

$$\Delta W_D(\theta, \psi, t) = \Delta G_T(\theta, \psi, t) - \Delta R_T(\theta, \psi, t) \quad (4)$$

Where $G_T(\theta, \psi, t)$ and $R_T(\theta, \psi, t)$ are the vertical perturbations in the geoid and solid earth surface, at θ co-latitude, ψ east-longitude and time t .

In most ice-sheet modelling studies of the GrIS, a spatially uniform, time varying GMSL
320 forcing is used to represent the perturbation in the geoid/ocean surface ($G(\theta, \psi, t)$) and the deformation of the solid earth ($R(\theta, \psi, t)$) is calculated using the elastic lithosphere-relaxing asthenosphere (ELRA) method (Le Meur and Huybrechts, 1996). This method only includes the local changes in the solid earth surface resulting from the deglaciation of the GrIS. In reality, the water depth/sea level signal surrounding the GrIS is highly spatially and temporally variable due to the influence of the
325 neighbouring LIS and IIS. On the time scales of this study the main processes driving this spatial and temporal variability is GIA (Rovere et al., 2016). The variability results from the interplay between the GrIS driven local changes, as is typical for near field regions and the non-local changes driven by the LIS and the IIS (Lecavalier et al., 2014). This is because Greenland sits on the resulting forebulge of the LIS. Ideally, the most complete method of incorporating this complex sea level (water depth) signal
330 would be with a coupled ice-sheet-GIA model, as in de Boer et al., (2014). Instead, a simpler alternative method was adopted in this study by coupling offline the output from a GIA model into IMAU-ICE.

To incorporate the output from the GIA model, first Eq. (4) was decomposed into (a) a local (subscript L) signal, driven by changes in the GrIS ($\Delta G_L, \Delta R_L$) and (b) a non-local signal (subscript NL)
335 driven by the influence of all other ice-sheets, primarily the LIS ($\Delta G_{NL}, \Delta R_{NL}$).

Hence the relationship for the change in water depth is written as:

$$\Delta W_D(\theta, \psi, t) = (\Delta G_L(\theta, \psi, t) + \Delta G_{NL}(\theta, \psi, t)) - (\Delta R_L(\theta, \psi, t) + \Delta R_{NL}(\theta, \psi, t)) \quad (5)$$

In order to solve this relationship, a GIA model was used to calculate the non-local contributions
 340 (ΔG_{NL} (Fig. 4d) and ΔR_{NL} (Fig. 4a)). This model has three key input components: a reconstruction of
 Late Pleistocene ice-sheet history (Peltier, 2004), an Earth model that simulates the solid earth
 deformation due to changes in the surface mass redistribution between the oceans and ice-sheets
 (Peltier, 1974) and a model of sea-level change (Farrell and Clark, 1976). The sea-level model included
 perturbations to the rotation vector (Milne and Mitrovica, 1998; Mitrovica et al., 2001; Mitrovica et al.,
 345 2005), time-dependent shoreline migration and an accurate treatment of sea-level change in areas
 characterised by ablating marine based ice (Kendall et al., 2005; Mitrovica and Milne, 2003).

To run the GIA model over the two glacial-interglacial cycles (240 kyr to PD) to produce the
 non-local signals, an input global ice reconstruction is required which reproduces the spatial and
 temporal history of all global ice sheets, apart from the GrIS during this interval. As a basis for this
 350 reconstruction, the ICE5G global ice model (Peltier, 2004) was adopted, which extends from 122 kyr
 BP to PD; one glacial-interglacial cycle. As the history for two glacial-interglacial cycles was required,
 the ice history over the 122 kyr was duplicated to represent the previous glacial-interglacial cycles (240
 kyr to 122 kyr BP), resulting in an ice-sheet reconstruction from 240 kyr BP to PD. The GrIS
 component was removed from the ICE5G global ice model to produce the final ‘non-local’ input ice
 355 history. The adopted earth model is characterised by a 96 km lithosphere, and an upper and lower
 mantle viscosity of 5×10^{20} Pa s and 1×10^{22} Pa s, respectively. These viscosity parameters fall
 approximately within the middle of the range of values commonly inferred. Using this input ice history
 and earth model the GIA model was ran offline to produce the non-local geoid and deformation fields
 (ΔG_{NL} (Fig. 4d) and ΔR_{NL} (Fig. 4a))

360 As the GIA model is used to produce the non-local components, the ‘local’ fields ($\Delta R_L, \Delta G_L$)
 driven by the GrIS are still required to solve Eq. (5). The local driven changes in the solid earth
 surface, ΔR_L were calculated internally within IMAU-ICE, using the ELRA method (Le Meur and
 Huybrechts, 1996) (Fig. 4b). This local field (ΔR_L) is combined with ΔR_{NL} (from the GIA model) to
 calculate the total deformation of the solid earth surface ΔR_T (Fig. 4c). This is combined with the non-
 365 local geoid signal, G_{NL} (Fig. 4d) to produce the final ΔW_D which is used to force the ice-sheet model at
 each time step (Fig. 4e).

Referring back to Eq. (5), using this method result in the following revised equation for ΔW_D .

$$\Delta W_D(\theta, \psi, t) = (\Delta G_{NL}(\theta, \psi, t)) - (\Delta R_L(\theta, \psi, t) + \Delta R_{NL}(\theta, \psi, t)) \quad (6)$$

370 However, comparing Eq. (6) to Eq. (5), it is evident that the local geoid, ΔG_L is not calculated using
 this approach and can be defined as a missing signal. To calculate this local geoid signal ΔG_L , would
 require solving the sea level equation (as within the GIA model, see de Boer et al., (2014)) resulting
 from these local GrIS driven changes within IMAU-ICE, which is not possible within the adopted
 approach. To estimate the magnitude of this missing ‘local geoid’ signal, the difference between the
 375 total signal, which is calculated using the GIA model (Fig. 4f, derived from Eq.5) and the signal as
 obtained from Eq.6 (Fig. 4e) was calculated. This difference is small (contoured on Fig. 4e), but is a
 shortcoming of the modelling that is accepted given the simplicity of the other components of the
 model. It is noted that this approach neglects for example feedbacks between changes in the sea level

and the marine-based ice and the stabilizing influence this may have on the evolution of the ice shelves
380 (de Boer et al., 2014; Gomez et al., 2010).

As the Fig. 4a illustrates, at the PGM there is a significant non-local deflection in the solid
earth surface. Across Ellesmere Island and the NW Greenland, the LIS and IIS produce significant
subsidence, up to 200 m. Central Greenland and Baffin Bay are elevated by up to 100 m and 30 m
respectively, due to the influence of the forebulge. In contrast, the non-local geoid signal is much
385 smaller, with a range of ~ 40 m (Fig. 4d). Comparing these two signals, it is apparent that the deflection
of the solid earth surface will be the main contributor to driving changes in water depth/sea level in this
study (Fig. 4e and 4f).

4. Results of simulations

There were only nine combinations of SSM1, SSM2, W_{D1} and A_s from the ensemble of simulations
390 that resulted in glacial–interglacial retreat over the two glacial-interglacial cycles (Table 3) and fulfilled
the conditions defined in Sect.2. Two additional simulations are included on Table 3
(*Low A_s _lowSSM1-0.25_deep* and *High A_s _highSS1*) which only resulted in a glacial-interglacial
expansion between the LGM and PD, one glacial-interglacial cycle. The spatial extent at selected time
periods is illustrated for one example simulation on Fig. 5.

395 At the glacial maximums (PGM and LGM) the simulated ice-sheet reached the inferred
observational limits along the northern and eastern margin (Fig. 5a and 5c), however at the SW margin
(see red and green triangles, Fig. 5a and 5c) the ice-sheet remains too restricted, possibly related to a
too strong mass balance height feedback in this region. The average LGM GMSL contribution is -2.59
m, which is still ~50% smaller than estimates from GIA modelling based studies (i.e. Lecavalier et al.,
400 2014). Therefore, closing the LGM GMSL budget remains problematic.

The simulated LIG minimum extent in all nine simulations complied with the spatial limits
inferred from the LIG ice core data, with a thinning at NEEM (~ 250m) and a moderate inland retreat
of the SW margin, but with Dye-3 remaining covered with grounded ice (Fig. 5b). The average LIG
GMSL contribution was 1.46 m (Table 3), which lies within the most recent estimated range of
405 between 0.6-3.5 m (Dutton et al., 2015). At the PD, the SW margin has retreated too far inland (Fig. 5d
and 5e) and there is a pronounced overthickening (up to 500 m) along most of the coastline (Fig. 5e).
Preliminary simulations concluded that increasing the resolution to 10x10 km reduced this misfit, but a
more detailed modelling of outlet glaciers at scales down to kilometres is likely needed to fully resolve
this misfit.

410 5: Forcing mechanisms controlling the spatial and temporal variability during deglaciation.

There is an evident correlation between the temporal variability of SAT forcing and the total ice
volume in all simulations; the periods of maximum ice volume (PGM, LGM) corresponding with the
minimum in SAT and vice versa (Fig. 2). This would imply that the timings of the glacial-interglacial
variations are strongly dependent on the adopted SAT forcing. However, there is a spatially variable
415 response between the NW and SW margins which alludes that the two regions are responding

regionally to a different forcing mechanism or at least a different timing of the same mechanism. Therefore, the interplay between the SAT and RSL forcing and the spatial and temporal variability in these two margins is examined in greater detail for the last 20 kyr BP.

It is evident from Fig. 6 that there is minimal variation in total ice volume and spatial extent
420 between the nine simulations from the LGM (~ 19 kyr BP) to 14.6 kyr BP (Fig. 7a). This corresponds to a period of relatively stable SAT ~ -15°C and minimal variations in the non-local RSL forcing (either the predicted bedrock depth or sea surface height (similar to that illustrated on Fig. 4)) due to only minor changes in the glacial history of the LIS (Peltier, 2004). Following this, there are three time periods (highlighted on Fig. 6) where changes in the ice volume and SAT correlate with a significant
425 retreat/readvance along the SW, SE and to a lesser extent NE margins (Fig. 7), but with the NW margin remaining stable. Between 14.6 kyr BP (Fig. 6) and ~13.9 kyr BP there is a rapid retreat in the grounded SW margin (Fig. 6, Fig. 7a-b) and a fall in ice volume of $\sim 1.0 \times 10^{15} \text{ m}^3$ (~ 0.24 m GMSL). This coincides with a warming (~ 10°C (Fig. 6)) and a strong non-local RSL signal due to a significant retreat of the LIS. As the LIS deglaciated, it produced a non-local subsidence of the bedrock (Fig. 4a)
430 across this margin, increasing the water depth and in turn the SSM. Following this retreat, there is a ~ 1 kyr stillstand in the grounded ice extent (Fig. 7c), during which there is a slow gradual cooling (Fig. 6). From ~12.9 till 11.5 kyr BP (Fig. 6, Fig. 7d) during a period of pronounced cooling (~ 15°C, Fig. 6) the ice-sheet readvances along the SW margin, producing a small increase in total ice volume (largest in simulations with high A_s), with the main period of retreat commencing at 11.5 kyr BP at the onset
435 of the sharp rise in SAT (~ 12°C). This readvance (12.9-11.5 kyr BP) coincides with the ongoing large non-local RSL signal (subsidence) which in turn results in an increase in SSM. This interplay implies that changes in the SSM (driven by the RSL signal) have only a secondary influence of the dynamics of the SW margin. This is emphasised by the minimal variations in the behaviour of the SW margins between the nine final simulations. In Sect. 2, from analysis of observational data, it was inferred that
440 the retreat from this margin may, in part be driven by the changes in RSL forcing. The simulations carried out in this study suggests that this is not the case, with the retreat driven primarily by SAT forcing.

The spatial and temporal behaviour of the NW margin (blue box, Fig. 1) in all nine simulations (Table 3) is highly variable, correlating with changes in the SSM, driven by the non-local
445 and local RSL forcing. There is minimal correlation with the timings of the SAT forcing. In all simulations, the timing of final deglaciation of the NW margin was too late compared to observations (~ 10-9 kyr BP), but the spatial pattern as inferred by Jennings et al., (2011) of a retreat initiated first at the western margin and later to the east is replicated. This is due to the faster ice velocity within the narrow outlet fjords to the west, i.e. Humboldt glacier, which feed into the grounded ice-sheet across
450 the Kane Basin (relative to the eastern grounded margin across the Hall Basin). The initiation of this retreat (which is at the earliest 8.9 kyr BP, Table 3) was controlled in part by the timing of the final deglaciation of the LIS within the ICE5G global ice model (Peltier, 2004) but also by the influence of the IIS which was simulated within IMAU-ICE. In ICE5G, the LIS retreats across Hudson Bay at 10 kyr BP with complete deglaciation of the high Arctic by ~ 8 kyr BP. This drives the onset of the non-
455 local subsidence of the solid earth surface (bedrock) (ΔR_{NL}) across this region (Fig. 4a and 4c), as the

LIS forebulge collapses. It is noted that changes in the choice of earth model and/or the spatial and temporal deglaciation history of the LIS during this final deglaciation interval will of course directly impact on the timing of the GrIS retreat.

460 The non-local influence of the IIS (which develops across Ellesmere Island) also strongly governed the timing of the retreat of the NW margin, which can be seen by comparing the results from two simulations: *HighA_s_lowSSM1-0.25* to *HighA_s_lowSSM1* (see Table 3). It could be assumed given the lower SSM1 (0.25 m/yr c.f 0.5 m/yr) in *HighA_s_lowSSM1-0.25* which results in a lower SSM close to the edge of the grounded ice margin that the onset of the retreat would be later. However, the retreat is in fact 1 kyr earlier. In this simulation (*HighA_s_lowSSM1-0.25*) the IIS is considerably thicker (> 465 1500m), increasing the subsidence of the solid earth surface (bedrock) (due to the increased ice loading), the water depth and in turn producing a higher SSM which drives the earlier deglaciation. This highlights the influence of the IIS on controlling the deglaciation of the GrIS across this region.

The amount of basal sliding (via the choice of A_s) also influences the timing of the onset of the NW margin retreat: with a lower amount of basal sliding generally promoting an earlier retreat 470 (comparing the average A_s (labelled AvA_s) simulations to the High A_s simulations (labelled *HighA_s*) on Table 3). This is examined in detail for two simulations: *AvA_s_lowSSM1-0.5_shallow* and *HighA_s_lowSSM1-0.25* (Fig. 8a and 8b). The retreat is initiated 5 kyr earlier in the simulation with a lower A_s value, *AvA_s_lowSSM1-0.5_shallow*. The earlier onset of the retreat with a lower A_s is due in part to the more restricted and thinner grounded ice-sheet across the NW margin, so there is a smaller 475 volume of ice to retreat (compare the red contours in Fig. 8a and Fig. 8b) and also the different SSM parameters. In *AvA_s_lowSSM1-0.5_shallow* the combination of a higher SSM1 (0.5 m/yr c.f 0.25 m/yr) and a shallower W_{D1} (300m c.f 475m) results in SSM that is higher at all water depths. It is this combination of a higher SSM with the lower A_s which drives the earlier onset of retreat and more restricted glacial maximum extent (Fig. 8a).

480 The SSM at deeper water depths (> W_{D1}), controlled by SSM2, also strongly influences the behaviour of the NW margin via the impact on the PGM to LIG glacial history, i.e the first glacial-interglacial cycle. Fig. 8c and 8d compare the difference in the simulated water depth between two simulations (*AvA_s+A_vSSM1* and *AvA_s+A_vSSM1_redSSM2*) where the SSM2 is reduced by 25 m/yr (from 100 m/yr to 75 m/yr). It could be assumed, given the reduction in SSM at deeper water depth, 485 that the retreat would be later. However, the onset of retreat is 2 kyr earlier (8.9 kyr BP c.f 6.9 kyr BP). This is due to the influence of the PGM to LIG glacial history (first glacial-interglacial cycle) on the dynamics of the LGM to PD retreat (second glacial-interglacial cycle). In the *AvA_s+A_vSSM1_redSSM2* simulation, during the first advance of the ice-sheet, the lower SSM at water depths > 400 m results in a thicker ice-sheet across the Nares Strait and eastern Ellesmere Island (part of the IIS). This increases 490 the bedrock subsidence and the water depth (Fig. 8c) resulting in a higher SSM surrounding the retreated ice margin during the subsequent glacial-interglacial cycle (after the LIG minimum). This higher SSM restricts the maximum spatial extent that the grounded ice margin reaches during the subsequent LGM to PD glacial-interglacial cycle (compare Fig. 8d to Fig. 8a and 8b). Therefore, with a smaller ice extent, surrounded by a region of higher SSM, this induces an earlier onset of retreat.

495 **6. Conclusions**

In this study using the ice sheet-ice shelf model, IMAU-ICE, the evolution of the GrIS over the two most recent glacial- interglacial cycles (240 kyr BP to PD) was investigated. The sensitivity of the spatial and temporal behaviour of the ice-sheet to an offline RSL forcing, generated by a GIA model was incorporated. Through this, the influence of the glacial history of the LIS and IIS was explored.

500 This RSL forcing governed the spatial and temporal pattern of SSM via changes in the water depth below the ice-shelves that developed around the ice-sheet. The SSM was parameterised in relation to the water depth, where for an increase in water depth, the SSM increased. We note that we do not investigate the influence of these two ice sheets (LIS and IIS) on the atmospheric circulation; there was no climate model used within our study.

505 At the LIG minimum, all of the LIG ice cores remain ice covered, with a ~ 250 m thinning at NEEM and an inland retreat of the SW margin, contributing 1.46 m to the LIG highstand; a reduction of ~0.7 m relative to previous studies. At the glacial maximums, the ice-sheet expanded offshore to coalesce with the IIS, reaching the Smith Sound at the north of Baffin Bay and out onto the continental shelf along the SW. However, it is still too restricted to the NE. A LGM GMSL contribution of -2.59 m
510 is considerably higher than most previous studies (~ 1.26 m), but closing the LGM GMSL budget remains problematic.

The temporal response of the SW margin was primarily controlled by the adopted SAT forcing (taken from ice core records). The RSL forcing and the choice of SSM parameterisation were of secondary influence. However, the inclusion of the RSL forcing improved the reconstructed PD
515 GrIS by reducing an under prediction along the SW margin (relative to observations). Conversely, the NW margin, where the ice-sheet coalesced with the IIS, was relatively insensitive to the imposed SAT forcing. Instead the spatial and temporal response was controlled by variations in the resultant SSM patterns that are driven by the variability in the RSL forcing and the glacial history of the LIS and IIS. The combined RSL and temperature changes generate a highly variable temporal response, where
520 optimum parameters were found to be a sliding coefficient A_s in the range of 1.0×10^{-10} to 1.2×10^{-10} $m^8 N^{-3} yr^{-1}$, a relatively low SSM close to the grounded ice margin to allow glacial expansion and a higher SSM at deeper water depths to promote interglacial retreat.

Code availability

The IMAU-ICE model is part of the ICEDYN package. The code used in this study is based on the
525 ICEDYN SVN revision 2515. OBLIMAP is an open source package which is available at <https://github.com/oblimap/oblimap-2.0>

Data availability

Output from all simulations, including the GIA model used for the RSL forcing used within this study are available from the S.L.B upon request.

530 **Author contribution**

Competing interests

The authors declare that they have no conflict of interest

Acknowledgements

S.L.B and T.J.R acknowledge support from the Netherlands Earth System Science Centre (NESSC),
535 which is financially supported by the Ministry of Education, Culture and Science (OCW). MH was
supported by The Netherlands Polar Programme (NPP) of the Earth and Life Sciences division of The
Netherlands Organisation for Scientific Research (NWO/ALW).

References

Bamber, J. L., Griggs, J. A., Hurkmans, R., Dowdeswell, J. A., Gogineni, S. P., Howat, I., Mougnot,
540 J., Paden, J., Palmer, S., Rignot, E., and Steinhage, D.: A new bed elevation dataset for Greenland,
Cryosphere, 7, 499-510, 10.5194/tc-7-499-2013, 2013.

Bintanja, R., and van de Wal, R. S. W.: North American ice-sheet dynamics and the onset of 100,000-
year glacial cycles, *Nature*, 454, 869-872, 10.1038/nature07158, 2008.

Charbit, S., Ritz, C., Philippon, G., Peyaud, V., and Kageyama, M.: Numerical reconstructions of the
545 Northern Hemisphere ice sheets through the last glacial-interglacial cycle, *Climate of the Past*, 3, 15-
37, 2007.

Clark, P. U., and Tarasov, L.: Closing the sea level budget at the Last Glacial Maximum, *Proceedings
of the National Academy of Sciences of the United States of America*, 111, 15861-15862,
10.1073/pnas.1418970111, 2014.

550 Cofaigh, C. O., Dowdeswell, J. A., Jennings, A. E., Hogan, K. A., Kilfeather, A., Hiemstra, J. F.,
Noormets, R., Evans, J., McCarthy, D. J., Andrews, J. T., Lloyd, J. M., and Moros, M.: An extensive
and dynamic ice sheet on the West Greenland shelf during the last glacial cycle, *Geology*, 41, 219-222,
10.1130/g333759.1, 2013.

Cofaigh, C. O., Briner, J. P., Kirchner, N., Lucchi, R. G., Meyer, H., and Kaufman, D. S.:
555 PAST Gateways (Palaeo-Arctic Spatial and Temporal Gateways): Introduction and overview,
Quaternary Science Reviews, 147, 1 - 4, <http://dx.doi.org/10.1016/j.quascirev.2016.07.006>, 2016.

Cofaigh, C. Ó., Dowdeswell, J. A., Evans, J., Kenyon, N. H., Taylor, J., Mienert, J., and Wilken, M.:
Timing and significance of glacially influenced mass-wasting in the submarine channels of the
Greenland Basin, *Marine Geology*, 207, 39-54, <http://dx.doi.org/10.1016/j.margeo.2004.02.009>, 2004.

560 Colleoni, F., Masina, S., Cherchi, A., Navarra, A., Ritz, C., Peyaud, V., and Otto-Bliesner, B.:
Modeling Northern Hemisphere ice-sheet distribution during MIS 5 and MIS 7 glacial inception,
Climate of the Past, 10, 269-291, 10.5194/cp-10-269-2014, 2014.

Colville, E. J., Carlson, A. E., Beard, B. L., Hatfield, R. G., Stoner, J. S., Reyes, A. V., and Ullman, D.
565 J.: Sr-Nd-Pb Isotope Evidence for Ice-Sheet Presence on Southern Greenland During the Last
Interglacial, *Science*, 333, 620, 2011.


Cuffey, K. M., and Marshall, S. J.: Substantial contribution to sea-level rise during the last interglacial
from the Greenland ice sheet, *Nature*, 404, 591-594, 10.1038/35007053, 2000.

- 570 Dahl-Jensen, D., Albert, M. R., Aldahan, A., Azuma, N., Balslev-Clausen, D., Baumgartner, M., Berggren, A. M., Bigler, M., Binder, T., Blunier, T., Bourgeois, J. C., Brook, E. J., Buchardt, S. L., Buizert, C., Capron, E., Chappellaz, J., Chung, J., Clausen, H. B., Cvijanovic, I., Davies, S. M., Ditlevsen, P., Eicher, O., Fischer, H., Fisher, D. A., Fleet, L. G., Gfeller, G., Gkinis, V., Gogineni, S., Goto-Azuma, K., Grinsted, A., Gudlaugsdottir, H., Guillevic, M., Hansen, S. B., Hansson, M., Hirabayashi, M., Hong, S., Hur, S. D., Huybrechts, P., Hvidberg, C. S., Iizuka, Y., Jenk, T., Johnsen, S. J., Jones, T. R., Jouzel, J., Karlsson, N. B., Kawamura, K., Keegan, K., Kettner, E., Kipfstuhl, S.,
- 575 Kjaer, H. A., Koutnik, M., Kuramoto, T., Kohler, P., Laepple, T., Landais, A., Langen, P. L., Larsen, L. B., Leuenberger, D., Leuenberger, M., Leuschen, C., Li, J., Lipenkov, V., Martinerie, P., Maselli, O. J., Masson-Delmotte, V., McConnell, J. R., Miller, H., Mini, O., Miyamoto, A., Montagnat-Rentier, M., Mulvaney, R., Muscheler, R., Orsi, A. J., Paden, J., Panton, C., Pattyn, F., Petit, J. R., Pol, K., Popp, T., Possnert, G., Prie, F., Prokopiou, M., Quiquet, A., Rasmussen, S. O., Raynaud, D., Ren, J.,
- 580 Reutenauer, C., Ritz, C., Rockmann, T., Rosen, J. L., Rubino, M., Rybak, O., Samyn, D., Sapart, C. J., Schilt, A., Schmidt, A. M. Z., Schwander, J., Schubach, S., Seierstad, I., Severinghaus, J. P., Sheldon, S., Simonsen, S. B., Sjolte, J., Solgaard, A. M., Sowers, T., Sperlich, P., Steen-Larsen, H. C., Steffen, K., Steffensen, J. P., Steinhage, D., Stocker, T. F., Stowasser, C., Sturevik, A. S., Sturges, W. T., Sveinbjornsdottir, A., Svensson, A., Tison, J. L., Uetake, J., Vallelonga, P., van de Wal, R. S. W., van der Wel, G., Vaughn, B. H., Vinther, B., Waddington, E., Wegner, A., Weikusat, I., White, J. W. C.,
- 585 Wilhelms, F., Winstrup, M., Witrant, E., Wolff, E. W., Xiao, C., Zheng, J., and Community, N.: Eemian interglacial reconstructed from a Greenland folded ice core, *Nature*, 493, 489-494, 10.1038/nature11789, 2013.
- de Boer, B., Stocchi, P., and van de Wal, R. S. W.: A fully coupled 3-D ice-sheet-sea-level model: algorithm and applications, *Geoscientific Model Development*, 7, 2141-2156, 10.5194/gmd-7-2141-2014, 2014.
- Dutton, A., Carlson, A. E., Long, A. J., Milne, G. A., Clark, P. U., DeConto, R., Horton, B. P., Rahmstorf, S., and Raymo, M. E.: Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, 349, 10, 10.1126/science.aaa4019, 2015.
- 595 Dyke, L. M., Hughes, A. L. C., Murray, T., Hiemstra, J. F., Andresen, C. S., and Rodés, Á.: Evidence for the asynchronous retreat of large outlet glaciers in southeast Greenland at the end of the last glaciation, *Quaternary Science Reviews*, 99, 244-259, <http://dx.doi.org/10.1016/j.quascirev.2014.06.001>, 2014.
- 600 Evans, J., Dowdeswell, J. A., Grobe, H., Niessen, F., Stein, R., Hubberten, H. W., and Whittington, R. J.: Late Quaternary sedimentation in Kejser Franz Joseph Fjord and the continental margin of East Greenland, *Geological Society, London, Special Publications*, 203, 149-179, 2002.
- Evans, J., Cofaigh, C. O., Dowdeswell, J. A., and Wadhams, P.: Marine geophysical evidence for former expansion and flow of the Greenland Ice Sheet across the north-east Greenland continental shelf, *Journal of Quaternary Science*, 24, 279-293, 10.1002/jqs.1231, 2009.
- 605 Farrell, W. E., and Clark, J. A.: Post Glacial Sea-Level, *Geophysical Journal of the Royal Astronomical Society*, 46, 647-667, 1976.
- Funder, S., Kjeldsen, K. K., Kjær, K. H., and Ó Cofaigh, C.: Chapter 50 - The Greenland Ice Sheet During the Past 300,000 Years: A Review, in: *Developments in Quaternary Sciences*, edited by: Jürgen Ehlers, P. L. G. a. P. D. H., Elsevier, 699-713, 2011.
- 610 Fyke, J. G., Weaver, A. J., Pollard, D., Eby, M., Carter, L., and Mackintosh, A.: A new coupled ice sheet/climate model: description and sensitivity to model physics under Eemian, Last Glacial Maximum, late Holocene and modern climate conditions, *Geoscientific Model Development*, 4, 117-136, 10.5194/gmd-4-117-2011, 2011.
- 615 Gomez, N., Mitrovica, J. X., Huybers, P., and Clark, P. U.: Sea level as a stabilizing factor for marine-ice-sheet grounding lines, *Nature Geosci*, 3, 850-853, <http://www.nature.com/ngeo/journal/v3/n12/abs/ngeo1012.html - supplementary-information>, 2010.


- Greve, R., Wyrwoll, K. H., and Eisenhauer, A.: Deglaciation of the Northern Hemisphere at the onset of the Eemian and Holocene, *Annals of Glaciology*, Vol 28, 28, 1-8, 10.3189/172756499781821643, 1999.
- 620 Greve, R.: Relation of measured basal temperatures and the spatial distribution of the geothermal heat flux for the Greenland ice sheet, *Annals of Glaciology*, Vol 42, 2005, 42, 424-432, 10.3189/172756405781812510, 2005.
- Helsen, M. M., van de Wal, R. S. W., van den Broeke, M. R., van de Berg, W. J., and Oerlemans, J.: Coupling of climate models and ice sheet models by surface mass balance gradients: application to the Greenland Ice Sheet, *Cryosphere*, 6, 255-272, 10.5194/tc-6-255-2012, 2012.
- 625 Helsen, M. M., van de Berg, W. J., van de Wal, R. S. W., van den Broeke, M. R., and Oerlemans, J.: Coupled regional climate-ice-sheet simulation shows limited Greenland ice loss during the Eemian, *Climate of the Past*, 9, 1773-1788, 10.5194/cp-9-1773-2013, 2013.
- Hogan, K. A., Cofaigh, C. Ó., Jennings, A. E., Dowdeswell, J. A., and Hiemstra, J. F.: Deglaciation of a major palaeo-ice stream in Disko Trough, West Greenland, *Quaternary Science Reviews*, <http://dx.doi.org/10.1016/j.quascirev.2016.01.018>, 2016.
- 630 Hutter, K.: *Theoretical Glaciology: Material Science of Ice and the Mechanics of Glaciers and Ice Sheets*, Springer, 1983.
- Huybrechts, P.: Sea-level changes at the LGM from ice-dynamic reconstructions of the Greenland and Antarctic ice sheets during the glacial cycles, *Quaternary Science Reviews*, 21, 203-231, 10.1016/S0277-3791(01)00082-8, 2002.
- 635 Jennings, A. E., Hald, M., Smith, M., and Andrews, J. T.: Freshwater forcing from the Greenland Ice Sheet during the Younger Dryas: evidence from southeastern Greenland shelf cores, *Quaternary Science Reviews*, 25, 282-298, 10.1016/j.quascirev.2005.04.006, 2006.
- 640 Jennings, A. E., Sheldon, C., Cronin, T. M., Francus, P., Stoner, J., and Andrews, J.: The Holocene history of the Nares Strait: Transition from Glacial Bay to Arctic-Atlantic Throughflow, *Oceanography*, 24, 26-41, 2011.
- Jennings, A. E., Walton, M. E., Cofaigh, C. O., Kilfeather, A., Andrews, J. T., Ortiz, J. D., De Vernal, A., and Dowdeswell, J. A.: Paleoenvironments during Younger Dryas-Early Holocene retreat of the Greenland Ice Sheet from outer Disko Trough, central west Greenland, *Journal of Quaternary Science*, 29, 27-40, 10.1002/jqs.2652, 2014.
- 645 Johnsen, S. J., Dahl-Jensen, D., Gundestrup, N., Steffensen, J. P., Clausen, H. B., Miller, H., Masson-Delmotte, V., Sveinbjornsdottir, A. E., and White, J.: Oxygen isotope and palaeotemperature records from six Greenland ice-core stations: Camp Century, Dye-3, GRIP, GISP2, Renland and NorthGRIP, *Journal of Quaternary Science*, 16, 299-307, 10.1002/jqs.622, 2001.
- 650 Kendall, R. A., Mitrovica, J. X., and Milne, G. A.: On post-glacial sea level - II. Numerical formulation and comparative results on spherically symmetric models, *Geophysical Journal International*, 161, 679-706, 10.1111/j.1365-246X.2005.02553.x, 2005.
- Kopp, R. E., Hay, C. C., Little, C. M., and Mitrovica, J. X.: Geographic Variability of Sea-Level Change, *Current Climate Change Reports*, 1, 192-204, 10.1007/s40641-015-0015-5, 2015.
- 655 Le Meur, E., and Huybrechts, P.: A comparison of different ways of dealing with isostasy: examples from modelling the Antarctic ice sheet during the last glacial cycle", *Annals of Glaciology*, 23, 1996.
- Lecavalier, B. S., Milne, G. A., Simpson, M. J. R., Wake, L., Huybrechts, P., Tarasov, L., Kjeldsen, K. K., Funder, S., Long, A. J., Woodroffe, S., Dyke, A. S., and Larsen, N. K.: A model of Greenland ice sheet deglaciation constrained by observations of relative sea level and ice extent, *Quaternary Science Reviews*, 102, 54-84, 10.1016/j.quascirev.2014.07.018, 2014.
- 660

- Letreguilly, A., Reeh, N., and Huybrechts, P.: The Greenland ice-sheet through the last glacial interglacial cycle *Global and Planetary Change*, 90, 385-394, 1991.
- 665 Lisiecki, L. E., and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic delta O-18 records, *Paleoceanography*, 20, 17, 10.1029/2004pa001071, 2005.
- Macayeal, D. R.: Large-scale ice flow over a viscous basal sediment - Theory and application to ice stream-B, Antarctica. , *Journal of Geophysical Research-Solid Earth and Planets*, 94, 4071-4087, 10.1029/JB094iB04p04071, 1989.
- 670 Milne, G. A., and Mitrovica, J. X.: Postglacial sea-level change on a rotating Earth, *Geophysical Journal International*, 133, 1-19, 10.1046/j.1365-246X.1998.1331455.x, 1998.
- Mitrovica, J. X., Milne, G. A., and Davis, J. L.: Glacial isostatic adjustment on a rotating earth, *Geophysical Journal International*, 147, 562-578, 10.1046/j.1365-246x.2001.01550.x, 2001.
- Mitrovica, J. X., and Milne, G. A.: On post-glacial sea level: I. General theory, *Geophysical Journal International*, 154, 253-267, 10.1046/j.1365-246X.2003.01942.x, 2003.
- 675 Mitrovica, J. X., Wahr, J., Matsuyama, I., and Paulson, A.: The rotational stability of an ice-age earth, *Geophysical Journal International*, 161, 491-506, 10.1111/j.1365-246X.2005.02609.x, 2005.
- Peltier, W. R.: Impulse response of a maxwell Earth, *Reviews of Geophysics*, 12, 649-669, 10.1029/RG012i004p00649, 1974.
- 680 Peltier, W. R.: Global glacial isostasy and the surface of the ice-age earth: The ice-5G (VM2) model and grace, *Annual Review of Earth and Planetary Sciences*, 32, 111-149, 10.1146/annurev.earth.32.082503.144359, 2004.
- 685 Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J. M., Basile, I., Bender, M., Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M., Lipenkov, V. Y., Lorius, C., Pepin, L., Ritz, C., Saltzman, E., and Stievenard, M.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429-436, 10.1038/20859, 1999.
- Quiquet, A., Ritz, C., Punge, H. J., and Melia, D. S. Y.: Greenland ice sheet contribution to sea level rise during the last interglacial period: a modelling study driven and constrained by ice core data, *Climate of the Past*, 9, 353-366, 10.5194/cp-9-353-2013, 2013.
- 690 Reerink, T. J., Klijhuis, M. A., and van de Wal, R. S. W.: Mapping technique of climate fields between GCM's and ice models, *Geosci. Model Dev.*, 3, 13-41, 10.5194/gmd-3-13-2010, 2010.
- Reerink, T. J., van de Berg, W. J., and van de Wal, R. S. W.: OBLIMAP 2.0: a fast climate model-ice sheet model coupler including online embeddable mapping routines, *Geosci. Model Dev.*, 9, 4111-4132, 10.5194/gmd-9-4111-2016, 2016.
- 695 Ritz, C., Fabre, A., and Letreguilly, A.: Sensitivity of a Greenland ice sheet model to ice flow and ablation parameters: Consequences for the evolution through the last climatic cycle, *Climate Dynamics*, 13, 11-24, 10.1007/s003820050149, 1996.
- Ritz, C., Rommelaere, V., and Dumas, C.: Modeling the evolution of Antarctic ice sheet over the last 420,000 years: Implications for altitude changes in the Vostok region, *Journal of Geophysical Research-Atmospheres*, 106, 31943-31964, 10.1029/2001jd900232, 2001.
- 700 Robinson, A., Calov, R., and Ganopolski, A.: Greenland ice sheet model parameters constrained using simulations of the Eemian Interglacial, *Climate of the Past*, 7, 381-396, 10.5194/cp-7-381-2011, 2011.
- Rovere, A., Stocchi, P., and Vacchi, M.: Eustatic and Relative Sea Level Changes, *Current Climate Change Reports*, 2, 221-231, 10.1007/s40641-016-0045-7, 2016.

- 705 Sheldon, C., Jennings, A., Andrews, J. T., Cofaigh, C. O., Hogan, K., Dowdeswell, J. A., and Seidenkrantz, M. S.: Ice stream retreat following the LGM and onset of the west Greenland current in Uummannaq Trough, west Greenland, *Quaternary Science Reviews*, 147, 27-46, 10.1016/j.quascirev.2016.01.019, 2016.
- 710 Simpson, M. J. R., Milne, G. A., Huybrechts, P., and Long, A. J.: Calibrating a glaciological model of the Greenland ice sheet from the Last Glacial Maximum to present-day using field observations of relative sea level and ice extent, *Quaternary Science Reviews*, 28, 1631-1657, 10.1016/j.quascirev.2009.03.004, 2009.
- Stone, E. J., Lunt, D. J., Annan, J. D., and Hargreaves, J. C.: Quantification of the Greenland ice sheet contribution to Last Interglacial sea level rise, *Climate of the Past*, 9, 621-639, 10.5194/cp-9-621-2013, 2013.
- 715 Tabone, I., Blasco, J., Robinson, A., Alvarez-Solas, J., and Montoya, M.: The sensitivity of the Greenland Ice Sheet to glacial–interglacial oceanic forcing, *Clim. Past*, 14, 455-472, 10.5194/cp-14-455-2018, 2018.
- 720 van Angelen, J. H., van den Broeke, M. R., Wouters, B., and Lenaerts, J. T. M.: Contemporary (1960–2012) Evolution of the Climate and Surface Mass Balance of the Greenland Ice Sheet, *Surveys in Geophysics*, 35, 1155-1174, 10.1007/s10712-013-9261-z, 2014.
- Vasskog, K., Langebroek, P. M., Andrews, J. T., Nilsen, J. E. O., and Nesje, A.: The Greenland Ice Sheet during the last glacial cycle: Current ice loss and contribution to sea-level rise from a palaeoclimatic perspective, *Earth-Science Reviews*, 150, 45-67, 10.1016/j.earscirev.2015.07.006, 2015.
- 725 Winsor, K., Carlson, A. E., Welke, B. M., and Reilly, B.: Early deglacial onset of southwestern Greenland ice-sheet retreat on the continental shelf, *Quaternary Science Reviews*, 128, 117-126, 10.1016/j.quascirev.2015.10.008, 2015.
- Yau, A. M., Bender, M. L., Blunier, T., and Jouzel, J.: Setting a chronology for the basal ice at Dye-3 and GRIP: Implications for the long-term stability of the Greenland Ice Sheet, *Earth and Planetary Science Letters*, 451, 1-9, 10.1016/j.epsl.2016.06.053, 2016.
- 730 Zweck, C., and Huybrechts, P.: Modeling of the northern hemisphere ice sheets during the last glacial cycle and glaciological sensitivity, *Journal of Geophysical Research-Atmospheres*, 110, 24, 10.1029/2004jd005489, 2005.



735 Figure 1: Summary of the place names and regions referred to in the main text and locations of observational data. All information describing the symbols and references for the observational data are listed in Table 1. The red and blue boxes highlight the regions shown on Fig. 7 and Fig. 8 respectively.



740

Figure 2: (a) Surface Air Temperature forcing (SAT, °C) taken from Helsen et al., (2013), with the solid red line a 2 kyr mean. (b) Grounded Ice Volume (10^{15} m^3) from the ensemble of simulations (grey lines) and the nine optimum simulations (see Table 3 for colours). The solid orange line marks the present-day ice volume (Bamber et al., 2013)

745

750






Figure 3: Schematic illustration of three examples of the SSM parameterisation.


755



760 Figure 4: Illustration of the various components used within the calculation of the offline RSL forcing at 135
 765 kyr BP. Panels (a-c) are the predicted (solid earth deformation) bedrock depth, ΔR : (a) non-local
 component, ΔR_{NL} (No GrIS), (b) Local component, ΔR_L (GrIS only) and (c) Total signal, ΔR_T (d) Predicted
 non-Local Geoid signal, ΔG_{NL} (No GrIS). The predicted water depth (W_D) signal is illustrated for: (e)
 $\Delta G_{NL} - \Delta R_T$: Combination of Total Predicted bedrock depth and non-local Geoid. It is this signal which
 is used within all simulations. Contoured is the local signal, ΔG_L which is not included. (f) Total water depth
 signal, ΔW_D . See text for extra details.



770 Figure 5: Simulated extent of the grounded ice-sheet and relative water depth using the parameters
 highlighted on Table 3 ($A_vA_s + A_vSSM1$) at four time periods (a) Penultimate Glacial Maximum (PGM), 135
 kyr BP, (b) LIG minimum, 123 kyr BP, (c) Last Glacial Maximum (LGM), 19 kyr BP and (d) Present-day
 (PD). Panel (e) is the difference in the present-day surface elevation (relative to the observed surface
 elevation (Bamber, et al., 2013), where positive value indicates an overprediction). The black circles mark
 775 the location of the GrIS ice core sites. Observed data constraining the timing of retreat is summarised on
 Table 1. Note that the colour bar extended beyond (+/-) 150. Small floating ice shelves formed at the edge of
 the grounded ice-sheet, but these are not shown.



780


Figure 6: Comparison of the Grounded Ice Volume (10^{15} m^3) from the nine optimum simulations (see Table 3 for colours) and the Surface Air Temperature forcing ($^{\circ}\text{C}$) (black line, Fig. 2a) from 15 to 10 kyr BP. Highlighted are the timings of the retreat-readvance-retreat of the SW margin (red box, Fig.1), the spatial pattern of which is illustrated for $\text{AvA}_s + \text{AvSSM1}$ (light red line) on Fig. 7. Results for the NW are illustrated on Fig. 8.

785

790

795


800



805

Figure 7: Retreat of the grounded ice sheet along the SW margin (region bounded by the red box on Fig.1) for AvAs+AvSSM1 (kyr BP). The red contour marks the edge of the grounded ice sheet at an earlier time

810 step (given in red): (a) 14.9 (20.9); (b) 13.9 (14.9); (c) 12.9 (13.9); 11.9 (12.9); (e) 9.9 (10.9) (f) 8.9 (9.9) and (g) 5.9. There is minimal change between the extent at 5.9 and present-day. Observed data constraining the timing of retreat are summarised on Table 1. Small floating ice shelves formed at the edge of the grounded ice sheet, but these are not shown.



815 Figure 8: Examples of the influence of the choice of SSM parameter and RSL forcing on the spatial and temporal retreat pattern across the NW margin (region bounded by the blue box on Fig.1). Panels (a) and
 820 (b) compare the grounded ice sheet extent between two simulations AvA_s_lowSSM1-0.5_shallow and HighA_s_lowSSM1-0.25 to highlight the impact of the choice of A_s in controlling the onset of the LGM to PD retreat. The red contour marks the edge of the grounded ice sheet at an earlier time step. Panels (c) and (d) illustrate the difference in the simulated water depth and maximum grounded ice sheet extent (shaded grey region) between simulations AvA_s+AvSSM1 and AvA_s+AvSSM1_redSSM2. The main difference between these two simulations is the choice of SSM2: 100 m/yr (AvA_s+AvSSM1) compared to 75 m/yr (AvA_s+AvSSM1_redSSM2)

| Symbol | Name | Region | Location and Material | Latitude | Longitude | Timing deglaciation | Reference |
|--------|--|--------------|---------------------------------------|------------------------|----------------------|---------------------------|--|
| ▲ | HLY03-01-05GC | North West | Core -Hall Basin | 81.62° N | 296.74° W | < 9 kyr BP | Jennings et al., 2011 |
| ▣ | Swath bathymetry & TOPAS acoustic data | North East | Near Westwind Trough | ~ 80° N | ~ 346 °W | Evidence for grounded ice | Evans et al., 2009 |
| ★ | Kejser Franz | Central East | Core- Inner shelf Moraine | 73.45° N | 335° W | before 7.4 kyr BP | Evans et al., 2002 |
| ★ | Fosters Bugt | Central East | Core- Mid shelf Moraine | 73.5° N | 338.5° W | stabilised ~ 10 kyr BP | Evans et al., 2002 |
| ★ | PS2630 | Central East | Core - Outer shelf | 73.15° N | 341.9° W | < 15 kyr BP | Evans et al., 2002 |
| ■ | PS2628 | Central East | Cores - NE Greenland Basin | 73.15° N | 344.75° W | ~ 13 kyr BP | O'Cofaigh et al., 2004 |
| ■ | GC25 | Central East | Cores - NE Greenland Basin | 73.01° N | 348.13° W | ~ 13 kyr BP | O'Cofaigh et al., 2004 |
| ○ | MD99-2371 | South East | Core - Grivel Basin Denmark Strait | 68.09° N | 332.06° W | 14.3-13.8 kyr BP | Jennings et al., 2006 |
| ▲ | Core | South East | Outer Kangerdlugssuaq Trough | 65.96° N | 330.63° W | 17.2 kyr BP | Dyke et al., 2004 |
| △ | MD99-2371 | South East | Inner Kangerdlugssuaq Fjord | 68.43° N | 328.06° W | 11.8 kyr BP | Dyke et al., 2004 |
| X | MD99-2371 | South East | Inner Sermilik Fjord | 65.86° N | 322° W | 10.9 kyr BP | Dyke et al., 2004 |
| ▲ | VC15 | South West | Core - Disko Trough | 67.91° N | 301.27° W | ~ 12.24 kyr BP | Hogan et al., 2016 |
| ▲ | VC35 | South West | Core - Disko Trough | 67.70° N | 300.66° W | 12.35 | Jennings et al., 2014 |
| ▲ | VC20 | South West | Core - Disko Trough | 68.20° N | 302.24° W | 12.2 | Jennings et al., 2014 |
| ▲ | MSM343300 | South West | Core - Disko Bugt | 68.47° N | 306.0° W | 10.9 | Hogan et al., 2106 |
| ○ | Hellefiske moraine | South West | moraine - approx central location | ~ 67.0° N ~ 69.0° N | ~ 304° W ~ 304° W | Evidence for grounded ice | Hogan et al., 2016 Hogan et al., 2016 |
| ▲ | JR175-VC45 | Central West | Core - Uummannaq trough | 70.56° N | 299.63° W | > 14.9 kyr BP | Sheldon et al., 2016 |
| ▲ | JR175- VC43 | Central West | Core - Uummannaq trough | 70.62° N | 300.38° W | 11.4 kyr BP | Sheldon et al., 2016 |
| ▲ | MSM343520 | Central West | Core - Uummannaq trough | 70.82° N | 303.152° W | < 10.8 kyr BP | Sheldon et al., 2016 |

825

Table 1: Summary of a selection of the observational data which were used to constraint the timing and spatial extent of the grounded ice sheet (Fig. 1)

| Parameter | Values |
|--------------------------------|------------------------------|
| A_s ($m^8 N^{-3} yr^{-1}$) | 0.04 - 1.2×10^{-10} |
| SSM1 (m/yr) | 0.25 - 10 |
| SSM2 (m/yr) | 10 - 150 |
| water depth1 (m) | 300 - 600 |
| water depth 2 (m) | 1000 |

830

Table 2: Range of parameters adopted in the sub-ice shelf melting (SSM) parameterisations, illustrated on Fig. 3.

835

| Abbreviated Name | Colour Fig. 6 | $A_e \times 10^{-10}$ ($m^8 N^{-3} yr^{-1}$) | SSM1 (m/yr) | $W_D 1$ (m) | SSM2 (m/yr) | Timing (kyr BP) | | LIG GMSL(m) | LGM GMSL(m) |
|--|------------------|---|----------------|----------------|----------------|-----------------|-----------|----------------|----------------|
| | | | | | | PGM - LIG | LGM - PD | | |
| LowA _e _lowSSM1-0.25_deep | - | 0.1 | 0.25 | 550 | 150 | NONE | 3.9 - 2.9 | 1.62 | -1.90 |
| midA _e _AvSSM1 | black | 0.8 | 1 | 400 | 100 | 129.0 - 128.0 | 0.9 - 0 | 1.55 | -2.52 |
| AvA _e _lowSSM1-0.25_shallow | red | 1 | 0.25 | 300 | 150 | 129.0 - 127.9 | 3.9 - 2.9 | 1.44 | -2.57 |
| AvA _e _lowSSM1-0.5_shallow | dark red | 1 | 0.5 | 300 | 150 | 129.0 - 127.9 | 7.9 - 6.9 | 1.36 | -2.47 |
| AvA _e +AvSSM1 | light red | 1 | 1 | 400 | 100 | 129.0 - 127.9 | 6.9 - 5.9 | 1.42 | -2.61 |
| AvA _e +AvSSM1_redSSM2 | dashed red | 1 | 1 | 400 | 75 | 128.9 - 128.0 | 8.9 - 7.9 | 1.50 | -2.58 |
| HighA _e _lowSSM1-0.25 | blue | 1.2 | 0.25 | 475 | 150 | 128.0 - 126.9 | 2.9 - 1.9 | 1.25 | -2.87 |
| HighA _e _lowSSM1-0.5 | dark blue | 1.2 | 0.5 | 475 | 150 | 127.9 - 126.9 | 1.9 - 0.9 | 1.45 | -2.83 |
| HighA _e _AvSSM1 | light blue | 1.2 | 1 | 400 | 100 | 129.0 - 127.9 | 1.9 - 0.9 | 1.50 | -2.81 |
| HighA _e _AvSSM1_deeper | dashed blue | 1.2 | 1 | 475 | 150 | 127.9 - 126.9 | 1.9 - 0.9 | 1.49 | -2.71 |
| HighA _e _highSSM1 | -- | 1.2 | 5 | 475 | 150 | NONE | 6.9 - 5.9 | 1.48 | -2.59 |

840

Table 3: Set of optimum parameters which resulted in a growth beyond the PD margin during glacial maximums (PGM and LGM) and a retreat by present-day (PD). Note that $W_D 2$ is constant in all simulations, 1000 m. The simulation highlighted in grey is shown on Fig. 5. The timing of the retreat across the Nares Strait for two interglacial is given (Timing (kyr BP)): PGM-LIG and LGM-PD, along with the total global mean sea level (GMSL) contribution from the GrIS only for the Last interglacial (LIG) and the Last Glacial Maximum (LGM). The GMSL was calculated using the simulated ice-volume.

845