A Stalagmite Test of North Atlantic SST and Iberian Hydroclimate Linkages over the Last 2 **Two Glacial Cycles**

3

Rhawn F. Denniston^{a*}, Amanda N. Houts^{a1}, Yemane Asmerom^b, Alan D. Wanamaker, Jr.^c, 4 Jonathon A. Haws^d, Victor J. Polyak^b, Diana L. Thatcher^c, Setsen Altan-Ochir^a, Alyssa C. 5 Borowske^{a2}, Sebastian F.M. Breitenbach^e, Caroline C. Ummenhofer^f, Frederico T. Regala^g, 6 7 Michael M. Benedetti^h, Nuno Bichoⁱ 8

- ^a Department of Geology, Cornell College, Mount Vernon, Iowa 52314 USA 9
- ^b Department of Earth and Planetary Sciences, University of New Mexico, Albuquerque, New 10 11 Mexico 87131 USA
- ^c Department of Geological and Atmospheric Sciences, Iowa State University, Ames, Iowa 12 13 50011 USA
- ^d Department of Anthropology, Louisville University, Louisville, Kentucky 40208 USA 14
- ^e Institute for Geology, Mineralogy, and Geophysics, Ruhr-University Bochum 44801 Germany 15
- 16 ^f Department of Physical Oceanography, Woods Hole Oceanographic Institution, Woods Hole, 17 Massachusetts 02543 USA
- ^g Associação de Estudos Subterrâneos e Defesa do Ambiente, Torres Vedras, Portugal 18
- ^h Department of Geography and Geology, University of North Carolina Wilmington, 19 20 Wilmington, North Carolina 28403 USA
- 21 ¹ Center for Archaeology and Evolution of Human Behaviour, Universidade do Algarve, Faro, 22 Portugal
- 23
- 24 * corresponding author
- 25
- ¹ current address: Department of Earth Sciences, University of New Hampshire, Durham, New 26
- 27 Hampshire 03824 USA
- ² current address: Department of Ecology and Evolutionary Biology, University of Connecticut, 28
- 29 Storrs, Connecticut 06269 USA
- 30
- 31 **Keywords**
- Iberia, hydroclimate, stalagmite, oxygen isotope, carbon isotope, δ^{234} U, pollen, sea surface 32
- 33 temperature

35 Abstract

36 Close coupling of Iberian hydroclimate and North Atlantic sea surface temperature (SST) 37 during recent glacial periods has been identified through the analysis of marine sediment and 38 pollen grains co-deposited on the Portuguese continental margin. While offering precisely 39 correlatable records, these time series have lacked a directly-dated, site-specific record of 40 continental Iberian climate spanning multiple glacial cycles as a point of comparison. Here we present a high-resolution, multi-proxy (growth dynamics and $\delta^{13}C$, $\delta^{18}O$, and $\delta^{234}U$ values) 41 42 composite stalagmite record of hydroclimate from two caves in western Portugal across the 43 majority of the last two glacial cycles (~220 ka). At orbital and millennial scales, stalagmitebased proxies for hydroclimate proxies covaried with SST, with elevated $\delta^{13}C$, $\delta^{18}O$, and $\delta^{234}U$ 44 45 values and/or growth hiatuses indicating reduced effective moisture coincident with periods of 46 lowered SST during major ice-rafted debris events, in agreement with changes in palynological 47 reconstructions of continental climate. While in many cases the Portuguese stalagmite record can 48 be scaled to SST, in some intervals the magnitudes of stalagmite isotopic shifts, and possibly 49 hydroclimate, appear to have been somewhat decoupled from SST.

50

51 **1. Introduction**

52 The Portuguese continental margin is an important location for understanding variations 53 in paleoceanographic conditions over orbital and millennial-scales (Hodell et al., 2013; Voelker 54 and de Abreu, 2011). Here, marine sediments record basin-wide oceanographic signals while co-55 deposited pollen grains track coeval vegetation changes occurring across Iberia. Integrated 56 analysis of these proxies has revealed a close coupling of North Atlantic SST, regional climate, 57 and Iberian ecosystems during the last three glacial cycles, including changes in vegetation 58 dynamics (Sánchez Goñi et al., 2002; Tzedakis et al., 2004; Roucoux et al., 2006; Martrat et al., 59 2007; Naugthon et al., 2007; Sánchez Goñi et al., 2008), atmospheric circulation (Sánchez Goñi 60 et al., 2013), and fire frequency (Daniau et al., 2007). One commonly applied palynological 61 metric is the abundance of temperate tree pollen, which rises during warm and wet conditions 62 associated both with interglacials and Greenland interstadials, concomitant with shifts in Iberian 63 margin SST (Sánchez Goñi et al., 2002; Tzedakis et al., 2004; Combourieu-Nebout et al., 2009; 64 Fletcher et al., 2010; Chabaud et al., 2014). However, the nature of such land-sea connections is partially obscured by the size of catchments from which the pollen are derived, with some 65

reaching into central Iberia and spanning a range of environmental settings subject to varying
climatic influences (Martin-Vide and Lopez-Bustins, 2006; Naughton et al., 2007) (Fig. 1).

68 Testing the links between terrestrial and marine systems benefits from continental climate 69 archives that provide precisely-dated and high resolution rainfall-sensitive time series spanning 70 tens of millennia, but such records remain rare in Iberia, particularly near the west Iberian 71 margin (Fletcher et al., 2010; Moreno et al., 2012; Stoll et al., 2013). Here we present a composite stalagmite record of four proxies for hydroclimate – growth dynamics and $\delta^{13}C$, $\delta^{18}O$, 72 and δ^{234} U values – spanning the majority of the last and penultimate glacial cycles (~220 ka) at 73 74 two cave sites in western Portugal. These time series offer a rare, site-specific continental record 75 capable of examining the coherence of SST controls on Iberian climate and ecosystem dynamics 76 across glacial and interglacial periods. The new record provides a continental perspective of 77 hydroclimate dynamics linked to regional oceanographic conditions.

78

79 2. Samples and Regional Setting

80 2.1 Environmental Setting

81 We report the analysis of five stalagmites (BG41, BG66, BG67, BG611, BG6LR) from 82 Buraca Gloriosa (BG; 39°32'N, 08°47'W; 420 m a.s.l.) and one stalagmite (GCL6) from Gruta 83 do Casal da Lebre (GCL; 39°18'N, 9°16'W; 130 m a.s.l.), two caves in western Portugal (Fig. 1). 84 Environmental conditions in BG and GCL are well suited for speleothem paleoclimate 85 reconstruction (see below). BG and GCL are located within the Meso-Mediterranean bioclimatic 86 zone that dominates much of Iberia (Fig. 1). This region is characterized by strong seasonality, 87 with warm, dry summers and cool, wet winters (Fig. 2) associated with the winter westerlies 88 (Blanco Castro et al., 1997). In contrast, the Atlantic zone, north of the Douro River, is cooler, 89 wetter, and less strongly seasonal. In the Pleistocene, the transition between these zones likely 90 shifted southward with Mediterranean-type vegetation restricted to refugia (Rey Benayas and 91 Scheiner, 2002).

Over interannual scales, the hydroclimate of Iberia is tightly coupled with the winter North Atlantic Oscillation (NAO) (Fig. 3), an atmospheric dipole that strongly influences precipitation across much of western Europe and that more broadly reflects the strength and positioning of the Azores high pressure system, which steers storm tracks contained within the westerlies into or north of Iberia (e.g., Trigo et al, 2002; Paredes et al., 2006; Trouet et al., 2009;

97 Cortesi et al., 2014). The NAO is typically measured as the NAO index, which is calculated 98 using atmospheric pressure differences between Iceland and Lisbon (or the Azores) (Barnston 99 and Livezey, 1987). The nature of the influence of the NAO varies across Iberia, but it is 100 strongly correlated to rainfall in western Portugal (Fig. 3), with a positive NAO index associated 101 with a steeper pressure gradient and elevated Iberian aridity. Iberian precipitation has also been 102 linked to SST in regions ranging from the western North Atlantic to the Iberian margin (Lorenzo 103 et al., 2010) where ocean circulation is dominated by the south-flowing Portugal Current and the 104 near-coastal, north-flowing Iberian Poleward Current, two systems that transport pollen from 105 river mouths along the continental shelf (Fig. 1).

106

107 2.2 Cave Settings

Buraca Gloriosa cave is located near the town of Alvados, 30 km from the Atlantic Ocean, within middle Jurassic limestones of the Estremadura Limestone Massif (Rodrigues and Fonseca, 2010), a topographically distinct region in central Portugal (Fig. 1). The \sim 35 m-long cave is accessed through a single, small (\sim 0.5 m²) entrance at the top of a collapse at the base of a 30 m-high escarpment (Fig. 4). The cave is well decorated although little active growth is occurring today. Vegetation above the cave is primarily shrubs, small trees, and mosses, hosted by a thin (0-10 cm) and highly organic soil layer.

Gruta do Casal da Lebre overlooks the coastal town of Peniche and is hosted by upper Jurassic limestones. The cave is 130 m long and contains a single, one m^2 entrance that opens onto a 7 m vertical shaft (Fig. 4). This entrance has been closed with a solid metal door in recent decades in order limit access to the cave, and this modification likely has reduced air exchange in GCL relative to its original state. Like BG, GCL hosts little active calcite deposition, but contains numerous fossil stalagmites and stalactites. The vegetation over the cave has been replaced in recent decades by stands of eucalyptus that grow in thin (<1-5 cm), clay-rich soils.

122

123 *2.3 Pollen Sources*

Pollen deposited on the west Iberian margin is sourced primarily from vegetation inhabiting the watersheds of the major west-flowing stream systems draining Portugal and Spain, which are (from north to south) the Douro, Tagus, and Sado rivers. The areas encompassed by these streams are large (79,000, 81,000, and 7,650 km², respectively) and span a variety of 128 elevations. The Tagus and Sado are primarily responsible for pollen deposited southwest of 129 Portugal, while the Douro plays an important role in delivering pollen to the more northwesterly 130 sites (Fig. 1). Prevailing wind patterns likely prevent substantial transport of pollen from Iberia 131 to the western Portuguese margin (Naughton et al., 2007). The pollen data presented here were 132 collected in three closely spaced cores from the southwest Iberian margin: MD01-2443: 250-194 133 ka (Roucoux et al, 2006; Tzedakis et al., 2004); MD01-2444: 193-136 ka (Margari et al., 2010; 134 Margari et al., 2014); MD95-2042: 141-1 ka (Sánchez Goñi et al., 2008; Sánchez Goñi et al., 135 2013) (Fig. 1) and are integrated here into a single time series.

136

137 **3. Materials and Methods**

138 *3.1 Environmental Monitoring*

Environmental conditions were measured at both cave sites over a multi-year period, with data recorded in two-hour intervals near the areas where the stalagmites were deposited. Temperature and relative humidity were obtained using HOBO U23 automated sensors while barometric pressure was recorded with HOBO U20L loggers. Drip rates were monitored at BG with Stalagmate acoustic drip counters (Collister and Mattey, 2008).

144

145 *3.2 Uranium-Series Dating*

Stalagmite chronologies were constructed with a total of 69 ²³⁰Th dates obtained at the 146 147 University of New Mexico (Table 1) using the methods of Asmerom et al. (2010). For dating of 148 stalagmite carbonate, powders ranging from 100-200 mg were weighed, dissolved in 15N nitric acid, spiked with a mixed ²²⁹Th-²³³U-²³⁶U tracer, and processed using column chemistry 149 150 methods. U and Th fractions were dissolved in 5 ml of 3% nitric acid and transferred to analysis 151 tubes for measurement on a Thermo Neptune MC-ICP-MS. U and Th solutions were aspirated 152 into the Neptune using a Cetac Aridus II low flow desolvating nebulizer and run as static routines. All isotopes of interest were measured in Faraday cups, except for ²³⁴U and ²³⁰Th, 153 154 which were measured in the secondary electron multiplier (SEM). Gains between the SEM and 155 the Faraday cups were determined using standard solutions of NBL-112 for U and an in-house ²³⁰Th-²²⁹Th standard for Th that was measured after every fifth sample; chemistry blanks reveal 156 157 U and Th blanks below 20 pg. Ages are reported using two standard deviation errors.

For BG stalagmites, corrections were made for unsupported 230 Th using a 230 Th/ 232 Th 158 159 ratio of 13.5 ppm (±50%), a value determined from isotopic analysis of cave dripwater. To 160 obtain this value, 108 ml of dripwater were transferred into six 30 ml Teflon beakers. These 161 beakers were fluxed in 6N HCl for an hour, rinsed, and heated gently on a hotplate until 162 approximately 1-2 ml of fluid remained in each. All solutions were then combined into a single 163 30 ml Teflon beaker, spiked with the same tracer described above (which contains HF), fluxed, 164 and then taken to complete dryness. The resulting precipitate was dissolved with 15N HNO₃, 165 dried down, dissolved again in 7N HNO₃, and processed with the same column chemistry 166 methods used for the stalagmite samples. We lack independent constraints on the initial Th ratio 167 for the GCL stalagmite, and thus apply the default value of 4.4 ppm ($\pm 50\%$). This difference in 168 the initial Th ratio impacts the corrected ages of GCL6 by 0.5-3.0 kyr relative to the value used 169 for BG, and thus does not meaningfully influence our interpretations.

170 Age models were developed via multiple polynomial interpolations between dated 171 intervals using the COPRA age modeling software (Breitenbach et al., 2012) (Fig. 5). Aside from 172 providing age models, COPRA also yields mean modeled stable isotope values and confidence intervals (Supp. Fig. S1). Here we rely primarily here on the original δ^{18} O and δ^{13} C values 173 174 because COPRA-derived median values reflect statistically robust variations, but reduce to some 175 degree the range of isotopic variability. For COPRA, a dummy age was included in the age 176 model for BG41 in order to extrapolate below the hiatus, which is only possible with at least two 177 dated points. The value of this dummy age was based on the assumption that it maintains a 178 stratigraphically correct slope (i.e. higher sections of the stalagmite represent younger material). 179 The dummy age was applied a conservative error, meaning that it was as large as possible 180 without causing stratigraphic inversion with respect to the bounding ages.

181

182 *3.3 Stable Isotope Ratios*

A total of 1,510 stable isotope analyses were performed on calcite samples milled from the central axis of each stalagmite. After milling, powders were weighed (~200 μ g) and transferred to reaction vessels that were flushed with ultra-pure helium. Samples were then digested using >100% H₃PO₄ and equilibrated overnight (~16 hours) at 34°C before being analyzed. Isotopic ratios were measured using a GasBench II with a CombiPal autosampler coupled to a Thermo Finnigan Delta Plus XL mass spectrometer at Iowa State University. A 189 combination of internal and external standards was run after every fifth sample, as well as before 190 and after each batch, in order to ensure reproducibility. Oxygen and carbon isotope ratios are 191 presented in parts per mil (‰) relative to the Vienna Pee Dee Belemnite carbonate standard 192 (VPDB). Average precision for both δ^{13} C and δ^{18} O analyses is better than ±0.1‰ (1 σ).

193 For isotopic analyses of soil organic matter and vegetation collected from above the 194 caves, samples were dried, crushed, and transferred to tin boats. Carbon isotopic ratios were 195 measured using a Thermo Finnegan Delta Plus XL mass spectrometer in continuous flow mode 196 coupled with a Costech Elemental Analyzer. Caffeine (IAEA-600), cellulose (IAEA-CH-3), and 197 acetanilide (laboratory standard) isotopic standards yielded an average analytical uncertainty for 198 carbon of $\pm 0.09\%$ 1 σ (VPDB). Dripwater samples were measured using a Picarro L2130-i 199 Isotopic Liquid Water Analyzer, with autosampler and ChemCorrect software. Each sample was 200 measured six times, with only the last three injections used to determine isotopic values in order 201 to minimize memory effects. Three reference standards (VSMOW, IAEA-OH-2, IAEA-OH-3) 202 were used for regression-based isotopic corrections and to assign the data to the appropriate 203 isotopic scale. Reference standards were measured at least once every five samples. The average analytical uncertainty for δ^{18} O measurements was $\pm 0.1\% 1\sigma$ (VSMOW). 204

205

206 *3.4 Stalagmite Mineralogy and Fabrics*

207 The calcite comprising the BG samples ranges across a variety of fabrics including a 208 faster-growing, white, fibrous form and a slower-growing, dense, clear structure (Fig. 6; Supp. 209 Fig. S2). In some samples, sharp changes between the two forms within the same growth 210 horizons mark intervals of recrystallization during which U/Th ages are highly inconsistent, and 211 these intervals were excluded from our data set. BG6LR, which grew discontinuously over much 212 of the last glacial cycle, suffered from alteration of early and middle Holocene material, which 213 was therefore excluded from this analysis. BG67 is characterized primarily by fibrous calcite that 214 has been recrystallized to clear, dense calcite in a narrow band descending through its core. U/Th 215 dates from the fibrous calcite on the margins of the growth surface reveal open system behavior 216 and thus this portion of BG67 was excluded. Recrystallization is evident in portions of GCL6 217 (particularly just above its base) and BG66 but the consistency of U/Th dates and the trends in 218 stable isotopes suggest that this alteration may have occurred soon after original deposition. We 219 tested whether these altered sections retain reliable paleoclimatic information by analyzing stable

isotopes along partial transects located just outside the zones of recrystallization (Fig. 6). Because stable isotopic values and trends between these transects were consistent (within the analytical errors), we retained these sections in the time series. Growth position changed at numerous times in several of these stalagmites, and our sampling strategy accounted for these changes so as to consistently collect samples for stable isotopic analysis from the top surface (cap) of each stalagmite rather than the margins.

226

227 **4. Results**

228 4.1 Environmental Monitoring

Temperature and relative humidity collected inside both caves document environmental conditions over a multi-year period. Relative humidity remained largely stable at ~100% in both caves. Temperatures, while different at the two sites, exhibited similar seasonal variability that approximates the mean average temperature of the region (14.2 \pm 0.4°C at BG and 16.2 \pm 0.3°C at GCL for August 2012-January 2018) (Fig. 7).

234 Dripwater was collected at BG both over the course of minutes during site visits on four 235 separate occasions (November 2014, October 2015, March 2016, January 2018) and as months-236 long integrated samples. A total of 25 dripwater samples were analyzed for stable isotopic values. Dripwater δ^{18} O values range from -2.4‰ to -4.6‰, with a mean of -3.8±0.8‰ (Supp 237 238 Table 1), although as the timing of site visits varied, this value clearly is impacted by seasonal 239 controls on precipitation (and thus infiltration) oxygen isotope values. Drip rates were measured 240 for much of the period spanning June 2014 to January 2018 (for a total of ~36 months) and 241 exhibit seasonal variations tied to the winter wet and summer dry seasons, as well as individual 242 rain events (Fig. 7).

243

244 *4.2 U-Th Dates and Age Models*

245 ²³⁴U-²³⁰Th dating of BG and GCL stalagmites reveals growth across approximately three 246 quarters of the last 220 ka, with periods of deposition interrupted by numerous hiatuses of 247 varying length, with the longest gaps from 160-147, 97-87, 72-60, 41-36, 32-30, and 17-15 ka 248 (Fig. 5 and 6; Supp. Fig. S3). These features, coupled with repeated changes in growth direction 249 and high ²³²Th abundances in select sections, complicate construction of a chronology in some 250 intervals. Macroscopic petrographic discontinuities suggest the presence of several short-lived hiatuses, but these were included as gaps in the age models only where U/Th dates reveal an identifiable temporal offset. For example, the marine isotope stage (MIS) 6/5e boundary recorded by stalagmite BG67 is marked by both a change in drip position and a sharp transition from dense, clear calcite to a white, fibrous form. Taken together, it is clear that a hiatus of some duration occurred at this time. However, these isotope data are presented as being uninterrupted given the continuity of δ^{18} O values and no U/Th evidence for a long-lived hiatus (Fig. 6).

- 257

258 4.3 Assessing Equilibrium in Speleothem $\delta^{18}O$ and $\delta^{13}C$ Values

259 We used two approaches to assess the fidelity of BG/GCL carbon and oxygen isotopes as 260 records of past environmental variability. First, Hendy Tests, in which stalagmite isotopic ratios 261 must satisfy two criteria in order to be considered as having crystallized near isotopic 262 equilibrium with cave dripwater (Hendy, 1971), were performed for each stalagmite. The first 263 half of the Hendy Test involves analysis of multiple isotopic analyses performed on samples 264 drilled at increasing distance from the central growth axis along the same series of growth layers. The conceptual justification for this approach is that dripwater, and thus speleothem calcite, δ^{18} O 265 values should remain constant down the stalagmite flanks because ¹⁶O preferentially lost to CO₂ 266 267 out-gassing is replenished by CO₂ hydration and hydroxylation reactions. Progressive ¹⁸O 268 enrichment associated with kinetic effects tied to Rayleigh distillation suggests isotopic 269 disequilibrium. No such consistent trends toward elevated oxygen isotopic ratios are found (Fig. 8), and thus the BG/GCL stalagmites appear to satisfy the first criterion of the Hendy Test. 270

271 The second portion of the Hendy Test is based on the degree of covariation of carbon and 272 oxygen isotopic ratios. Oxygen isotopic ratios of speleothem calcite reflect those of infiltrating fluids, which are generally close to the δ^{18} O values of meteoric precipitation, and which, in many 273 274 locations, are linked to climate (air temperature, moisture source, seasonality of precipitation, or 275 rainfall amount (Lachniet, 2009)). Interpreting changes in oxygen isotope composition at 276 BG/GCL during intervals of profound climatic change such as marked the last glacial period is complicated by the multiple factors that influenced δ^{18} O values of precipitation at these sites, 277 278 including shifts in moisture source. The potential exists for rainfall in Iberia to be derived from 279 atmospheric moisture sources that change on synoptic/seasonal scales (Moreno et al., 2014; 280 Gimeno et al., 2010; Gimeno et al., 2012) as well as in response to changing glacial boundary 281 conditions (Florineth and Schlüchter, 2000; Kuhlemann et al., 2008; Luetscher et al., 2016). In addition, strong but opposite correlations exist in modern precipitation between rainwater δ^{18} O values and (i) the regional air temperature (r=+0.8) and (ii) rainfall amount (r=-0.8), both of which are related to the strong seasonality of precipitation associated with Meso-Mediterranean climates (IPMA, 2016).

286 Correlations between carbon and oxygen isotope ratios are presented in Figure 8. Three stalagmites – BG6LR, BG66, and BG67 – show strong correlations between $\delta^{13}C$ and $\delta^{18}O$ 287 $(r^2=0.6)$, while the other three samples lack a strong correlation. If one considers the second 288 289 criterion of the Hendy Test, the nature of equilibrium crystallization in stalagmites BG6LR, 290 BG66, and BG67 would be considered suspect. It must be noted, however, that the reliability of 291 the Hendy Test has been questioned because (1) equilibrium may be maintained in some portions 292 of a stalagmite but not others, (2) growth layers thin progressively down the sides of the 293 stalagmite, making it difficult to restrict samples to the same material, and (3) equilibrium 294 covariation of carbon and oxygen isotope ratios may result as the direct or indirect result of 295 climatic variability (Dorale and Liu, 2009; Lechleitner et al., 2017). We therefore interpret both 296 isotope ratios and their covariation as environmental signals.

297

298 4.4 Hydroclimate Proxies

299 4.4.1 Carbon Isotopes

Interpreting speleothem δ^{13} C variability in a climatic context requires understanding, or 300 at least constraining, the origins of these isotopic shifts. Stalagmite δ^{13} C values reflect two 301 302 primary inputs: CO₂ derived from the atmosphere and/or soil zone and bicarbonate derived from dissolution of bedrock carbonate. Speleothem $\delta^{13}C$ values reflect the type (C₃ vs C₄) and density 303 304 of vegetation over the cave, both of which are impacted by changes in air temperature and/or precipitation. The average δ^{13} C value of biogenic CO₂ in the soil zone is tied to the ratio of 305 plants utilizing the C₃ (average δ^{13} C -26‰) versus C₄ (average δ^{13} C -14‰) photosynthetic 306 pathways (Deines, 1980; von Fischer et al., 2008). Similarly, vegetation density and soil 307 308 respiration rates over the cave impact the relative contribution of atmospheric CO₂ (pre-Industrial δ^{13} C -6‰ to -7‰; Francey et al., 1999) as compared to soil-derived CO₂ (Hellstrom 309 and McCulloch, 2000; Genty et al., 2003). Phanerozoic bedrock δ^{13} C values range from -4% to 310 311 +8‰ (Saltzman and Thomas, 2012), but these values are static and do not contribute to temporal 312 variability in stalagmite carbon isotopic ratios.

Superimposed on these inputs are secondary effects capable of influencing the $\delta^{13}C$ 313 314 values of dripwater in the epikarst or cave. When voids in the bedrock are not fully saturated, CO₂ degassing from infiltrated water may occur in the epikarst. This preferential loss of ¹²CO₂ 315 (that may result in crystallization of calcium carbonate - so-called prior calcite precipitation) 316 enriches the residual solution in ¹³C, a signal that can be transferred into underlying stalagmites 317 318 (Baker et al., 1997). Once the solution enters the cave, equilibrium fractionation between 319 dissolved carbon species may be disrupted owing to issues surrounding CO₂-degassing under 320 low drip rate conditions (Breitenbach et al., 2015) or by disequilibrium processes occurring 321 during carbonate crystallization (Mickler et al., 2004; Fairchild et al., 2006). Importantly, δ^{13} C values reflect local infiltration rather than (pan-)regional atmospheric conditions as in the case of 322 δ^{18} O. This difference between both proxies offers the opportunity to investigate environmental 323 324 changes at different spatial scales.

325 Terrestrial deposits preserving pollen spectra spanning substantial portions of the last 326 glacial cycle from western Iberia are rare (Gómez-Orellana et al., 2008; Fletcher et al., 2010; 327 Moreno et al., 2012), and thus pollen in marine sediments represents a particularly important 328 continental climate record. Pollen samples obtained from the Iberian margin contain small 329 percentages of *Poaceae*, the family including the majority of C₄ plants, demonstrating a 330 persistent and overwhelming majority of C₃ (largely shrub and arboreal) vegetation throughout 331 the last glacial cycle including between Greenland stadials (GS) and interstadials (GI) and across 332 Heinrich stadials (HS) (d'Errico and Sánchez Goñi, 2003; Tzedakis et al., 2004; Desprat et al., 333 2006; Sánchez Goñi et al., 2008; Sánchez Goñi et al., 2013; Margari et al., 2014). In the absence 334 of changes in vegetation type, shifts in the source of carbon found in cave dripwater therefore 335 likely originated with the density of vegetation and/or soil respiration rates (Genty et al., 2003). 336 Reductions in these values are generally associated with decreases in temperature and/or 337 increases in aridity, such as have been inferred from Iberian pollen spectra to have characterized 338 Iberia during GS, HS, and glacial maxima (Sánchez Goñi et al., 2008; Margari et al., 2014). 339 Complementing these effects are increases in the contribution of bedrock carbon, as well as prior 340 calcite precipitation, reflecting a combination of longer residence times of infiltrating solutions 341 and desaturation of voids in the epikarst above the cave, both of which are consistent with more arid climates (Baker et al., 1997; Genty et al., 2003). Thus, we interpret the carbon isotopic 342 values of the BG/GCL record as primarily a local (hydro)climate proxy, with higher δ^{13} C values 343

indicative of a cooler, drier climate. Integrating the GCL6 δ^{13} C record into the BG time series is complicated by the slightly different bedrock δ^{13} C values of the host rocks (Supp. Table 1) and what may have been distinct vegetation types and cave hydrologies at each cave when GCL6 was being deposited (187-160 ka). However, similar δ^{13} C values during their period of overlap (187-185 ka) suggests that the two records can be consolidated (see below).

349 A test of equilibrium crystallization in the modern system can be constructed by 350 comparing modeled stalagmite isotopic values to recently deposited calcite. The carbon isotopic 351 composition of speleothem calcite is the result of a complex series of reactions that have been 352 addressed in a number of studies (Hendy, 1971; Mühlinghaus et al., 2007; Dreybodt, 2008). For 353 δ^{13} C in BG stalagmites, we use the equations of Li et al. (2014), which factor in the two primary 354 sources of carbon – soil CO₂ and bedrock carbonate – the proportion of carbon derived from 355 each source, and temperature-induced fractionation of carbon isotopes between dissolved carbon 356 species:

357

358
$$\delta^{13}C_{\text{calcite}} = f_1 * [\delta^{13}C_{\text{ls}} - (\delta^{13}C_{\text{CO2(g)}} + 9.48 \times 10^3/\text{T} - 23.89)] + \delta^{13}C_{\text{CO2(g)}} + 9.48 \times 10^3/\text{T} + 0.049 \text{T} - 37.72$$

where: f_1 = fraction of bicarbonate from limestone (ls)

T = temperature (°K)

359

360

361 362

363 We assume the most straightforward and simple situation: the system remains closed to soil CO2 after entering the epikarst and bedrock carbonate contributes 50% of carbon to 364 365 dripwater bicarbonate ($f_1=0.5$). We apply the average cave temperature of 14.4°C and the measured δ^{13} C values of BG bedrock and the overlying vegetation/soil of +3±1‰ and -28±1‰. 366 367 respectively. This approach, while certainly overly simplified for the BG cave system, yields modeled stalagmite δ^{13} C values averaging -7.7±1‰, similar to calcite crystallized on two glass 368 369 slides installed at the site of two actively growing stalagmites in the loft area of BG, which vielded δ^{13} C values of -8.4±1.2‰. 370

371

372 *4.4.2 Oxygen Isotopes*

373 The origins of BG/GCL isotopic variability appear more complex for oxygen than for 374 carbon. Like δ^{13} C values, local δ^{18} O minima mark interstadials and interglacials. Analysis of 375 modern precipitation data reveals equally strong, albeit inverse, correlations between precipitation δ^{18} O and both amount (r=-0.8) and air temperature (r=+0.8) effects. likely owing to 376 377 the dominance of cool season precipitation in annual water budgets (IAEA/WMO, 2016) (Fig. 378 2). Based on these relationships, it remains possible that changes in air temperature, overall precipitation, and/or precipitation seasonality could impact the δ^{18} O values of effective moisture. 379 380 That air temperature is likely not a prominent driver of stalagmite oxygen isotopic variability is supported by two observations, however. First, the slopes of the air temperature/ δ^{18} O 381 382 relationships (‰/°C) at the three GNIP stations located closest to BG and GCL (Porto, Vila Real, 383 and Portalegre) are nearly identical (average for the three sites of 0.25±0.03‰/°C) but opposite 384 in sign to the calcite-water temperature dependence of oxygen isotopic fractionation $(-0.2\%)^{\circ}C$ (Kim and O'Neil, 1997) (slopes of precipitation amount/ δ^{18} O are -1.6, -3.5, and -3.7‰/100 385 386 mm/month, respectively). In the simplest sense, therefore, a 1°C increase in mean annual air 387 temperature (and thus also cave temperature) would increase precipitation δ^{18} O values by approximately the same amount that the water temperature effect would lower stalagmite calcite 388 389 δ^{18} O values. In this simplified scenario, the net effect is a stalagmite record that is negligibly 390 influenced by multi-decadal/centennial-scale temperature changes alone. Secondly, the observed shift toward lower stalagmite δ^{18} O values during interstadials and interglacials, periods of 391 392 elevated mean annual temperature, demonstrates that the observed positive correlation between precipitation δ^{18} O and air temperature is not a dominant feature over millennial time scales. For 393 example, the 3.5% decrease in δ^{18} O values between MIS 6 and MIS 5e (136-128 ka) (Fig. 9) can 394 be only partially accounted for by the ~1‰ ice volume-related decrease in North Atlantic surface 395 water δ^{18} O values (Schrag et al., 1996). Other factors such as kinetics associated with humidity 396 397 and wind speed at the point of evaporation (Grootes et al., 1993), temperature and source of 398 atmospheric moisture (Herbert et al., 2001), and cloud evolutionary pathways (Rozanski and 399 Araguás, 1995) need also be considered but cannot account for the entirety of this shift. Because 400 of the narrow continental shelf in central Portugal, the LGM shoreline was located close to the 401 modern shoreline, thereby minimizing continental effects, and the magnitude of the impacts of 402 wind speed and ocean temperature do not appear sufficient to account for the observed stalagmite δ^{18} O variability. Thus, the decrease in stalagmite δ^{18} O between the penultimate glacial 403 and last interglacial suggests that stalagmite oxygen isotope ratios are primarily recording (pan-404

405)regional hydroclimate rather than temperature. The origin of the anomalously low δ^{18} O values 406 during GI 1 (dated here from 14.5-13.9 ka) are unclear (unfortunately no other BG or GCL 407 stalagmite also spans this interval) but reinforce this inverse relationship between mean annual 408 temperature and stalagmite oxygen isotope ratios.

409 Speleothem oxygen isotopic ratios were modeled using the paleotemperature equation of 410 Kim and O'Neil (1997), which requires measurements of water (cave) temperature and dripwater δ^{18} O values. The resulting δ^{18} O model value of -3.1±1.0‰ is nearly identical to the glass plate-411 412 grown calcite value of -3.0±0.6‰. It should be noted, however, that assessing equilibrium 413 crystallization in modern calcite/dripwater pairs at BG is complicated by the low temporal 414 resolution associated with integrated, months-long dripwater samples, variable timing of 415 dripwater collecting trips, and any seasonal biases in calcite crystallization that at present remain 416 poorly constrained.

417 Replication between stalagmites of similar age is arguably the single most reliable 418 method for evaluating the impacts of climate versus secondary influences, including evaporation 419 and kinetic effects (Denniston et al., 1999; Mickler et al., 2004), on stalagmite isotopic ratios 420 (Dorale and Liu, 2009; Denniston et al., 2013). When presented as an integrated data set, the 421 BG/GCL stalagmite carbon and oxygen isotopic time series spans the majority of the last 220 ka 422 (Fig. 9), although stalagmites spanning the same periods of time are restricted to 187-185, 111-423 104, 83-81, 78-73, and 58-53 ka. Because these intervals are short, and because the temporal 424 resolution varies substantially between stalagmites, replication tests based on these intervals are of limited utility. However, within the age uncertainties, $\delta^{18}O$ and $\delta^{13}C$ values and trends are 425 426 similar, suggesting that oxygen and carbon isotopic ratios track environmental, rather than dripspecific, variables. The three exceptions in which coeval samples do not replicate well are: $\delta^{13}C$ 427 values offset by 3‰ from 83-81 ka and by 4‰ from 58-53 ka, and δ^{18} O values offset by 1‰ 428 429 from 111-104 ka (Fig. 9; Supp. Fig. S4).

430

431 4.4.3 δ^{234} U Values

432 δ^{234} U values (calculated as the difference between the age-corrected 234 U/ 238 U ratio of a 433 sample and the secular equilibrium 234 U/ 238 U ratio) of speleothem carbonate have also been used 434 as a proxy for paleoprecipitation (Hellstrom and McCulloch, 2000; Oster et al., 2012; Plagnes et 435 al., 2002; Polyak et al., 2012; Zhou et al., 2005). 234 U exists in the stalagmite crystalline lattice

due to incorporation from cave dripwater and through *in situ* production from decay of ²³⁸U. 436 437 Alpha recoil displaces ²³⁴U from its lattice position, increasing its susceptibility to leaching by infiltrating waters, meaning that ²³⁴U is selectively mobilized relative to ²³⁸U in cave dripwater 438 439 (Chabaux et al., 2003; Oster et al., 2012). The flux of infiltrating fluids is therefore tied to δ^{234} U 440 values of dripwater, and thus stalagmite carbonate, such that decreases in effective precipitation and/or bedrock dissolution rate, both of which are tied to increased aridity, are associated with 441 elevated speleothem δ^{234} U values (Hellstrom and McCulloch, 2000; Plagnes et al., 2002; Polvak 442 443 et al., 2012).

As differences in δ^{234} U values between stalagmites may arise from distinct infiltration 444 pathways (Zhou et al., 2005), complicating the integration of δ^{234} U values from multiple 445 stalagmites into a single, cohesive data set, we restrict our analysis to stalagmite BG6LR, which 446 447 represents the longest individual stalagmite record of the BG/GCL time series. While the number of δ^{234} U measurements is small compared to stable isotopic values, the temporal density of the 448 former is sufficient to demonstrate the utility of δ^{13} C and δ^{18} O values as paleohydroclimate 449 450 proxies (Fig. 9). Decreased precipitation/effective moisture is associated with elevated stalagmite δ^{13} C, δ^{18} O, and δ^{234} U values. The relationships between δ^{13} C and δ^{234} U values in all BG and 451 GCL stalagmites are presented in Supp. Fig. S5. 452

453

454 **5. Environmental Conditions at BG/GCL and Links to Iberian Margin SST**

455 The previously discussed tests for isotopic equilibrium, including the reproducibility of 456 carbon and oxygen isotope ratios between coeval BG and GCL stalagmites, support the notion that their δ^{13} C and δ^{18} O values may be integrated into cohesive time series reflecting 457 paleohydroclimatic conditions and used to assess links between continental climate and SST 458 459 (Fig. 10). Over the last several glacial cycles, oceanographic conditions along the western Iberian 460 margin varied at millennial and orbital time scales in close correlation with Greenland air 461 temperature and North Atlantic conditions and circulation (Roucoux et al., 2005; Daniau et al., 462 2007; Sánchez Goñi et al., 2008; Darfeuil et al., 2016). Abrupt changes in SST reflect a balance 463 between southward expansion of subpolar waters and northward migration of subtropical water 464 masses (de Abreu et al., 2003). During the particularly cold conditions characterizing HS and 465 GS, Iberian margin SST decreased by up to 9°C (to as much as 13°C below present values; de 466 Abreu et al., 2003), with these changes helping to position the arctic or subarctic front at $\sim 39^{\circ}$ N,

467 the same latitude as BG and GCL. These cold surface waters reduced the production and 468 transport of atmospheric moisture to Iberia (Eynaud et al., 2009; Voelker and de Abreu, 2011), 469 and would have thereby influenced the timing of speleothem growth and carbon and oxygen 470 isotopic values in BG and GCL stalagmites. Indeed, the composite BG/GCL record documents 471 coherence, at both orbital and millennial scales, between Portuguese hydroclimate, vegetation, 472 and Iberian margin SST during the last two glacial cycles (Fig. 10 and 11). In an attempt to 473 quantify this covariance, we binned the SST and stalagmite stable isotope data into century-long 474 intervals. The relatively short record of BG41 was not included, and model ages for stalagmites 475 BG66 and GLC6 were increased by 4.0 kyr and 1.3 kyr, respectively, to improve correlation with 476 the SST chronology. The resulting inverse correlation between SST and carbon and oxygen is 477 strong (r=-0.55 and -0.52, respectively; p<0.0001) (Supp. Fig. S6).

478

479 *5.1 Growth Intervals*

The single most fundamental prerequisite to speleothem deposition is infiltration of surface waters, and thus the timing of stalagmite growth can reflect changes in mean hydroclimatic state. Deposition of multiple BG stalagmites was punctuated by hiatuses spanning similar time intervals (although the precise ages of the onset and/or termination of the hiatuses are distinct), a relationship that suggests links to changes in hydroclimate rather than random drip site-specific variability.

486 Hiatuses in some BG samples coincide with HS1, HS3, HS4, and HS6, and pollen spectra 487 independently suggest increased aridity during HS and glacial maxima. Decreases in arboreal 488 pollen abundance and concomitant increases in drought-tolerant vegetation coincide with periods 489 of reduced SST. Vegetation patterns during maximal IRD deposition on the Iberian margin 490 reveal not only dramatically reduced forest cover but also a pronounced expansion of semi-desert 491 plants (e.g. Sánchez Goñi et al., 2000; Roucoux et al., 2005; Naughton et al., 2009). These 492 changes mark the long hiatus between HS7 and HS6 (71-59 ka), which overlaps with the some of the coldest SST of the last 70 ka as reconstructed using $U_{37}^{k'}$ at core MD95-2042 (Darfeuil et al., 493 494 2016) (Fig. 10; Fig. 12). An absence in BG stalagmite deposition from ~160-149 ka occurs at the 495 same time as massive seasonal discharges from the Fleuve Manche river and the coldest 496 continental climates and SST (157-154 ka) of the last 220 ka, as determined from pollen and 497 foraminifera from core MD01-2444 (Margari et al., 2014; Fig. 1).

498 Whether hiatuses in BG speleothem deposition are a result of pronounced reductions in 499 precipitation, an extension of below freezing temperatures that limited infiltration (Vaks et al., 500 2013; Fankhauser et al., 2016), or variations in infiltration pathway/drip position is ambiguous. 501 Pollen transfer functions from MD95-2042 suggest winter temperatures dropped below 0°C 502 during HS and annual precipitation was reduced by up to 50% (from 800 mm to 500-400 mm 503 during HS3, HS4, and HS5) (Sánchez Goñi et al., 2002). Applying this temperature 504 reconstruction to western Portugal is complicated, however, by the broad area across which these 505 pollen grains were sourced. Permafrost reconstructions (Vandenberghe et al., 2014) of Iberia 506 argue against the hypothesis that continuous sub-zero temperatures inhibited infiltration and 507 stalagmite growth. We thus suggest that the hiatuses observed at BG and GCL were driven 508 largely by reductions in precipitation.

509 Other western European cave records also share similar growth histories. For example, 510 stalagmites from Villars Cave, southwestern France (Genty et al., 2003; Genty et al., 2010; 511 Wainer et al., 2011), and from multiple caves in northern Spain (Stoll et al., 2013) (Fig. 1) are 512 also punctuated by hiatuses during HS. For example, at or near HS7, stalagmite hiatuses were 513 formed at Villars Cave (78-76 ka), in northern Spain (~75 k), and BG (80-78 ka). No stalagmite 514 deposition has been identified at BG from 71-60 ka or Villars cave from 67-62 ka, a period that 515 includes HS6. Finally, HS1 is marked by a hiatus in northern Spain (18-15.5 ka) and at BG (17-516 15 ka). While the timing of these hiatuses is not identical, and not all hiatuses at Villars Cave and 517 the Spanish caves are coincident with those at BG, the substantial degree of overlap suggests a 518 common origin. Stoll et al. (2013) noted that stalagmite deposition and/or elevated growth rates 519 in northern Spain stalagmites occurred during periods of high Northern Hemisphere summer 520 insolation or during GI, while hiatuses occurred during periods of low insolation and low SST 521 (<13.7°C). The BG record supports the hypothesis that growth interruptions are related to SST 522 controls on regional atmospheric moisture availability, although the impact of insolation is not 523 clear.

524

525 5.2 BG/GCL Stable Isotopic and $\delta^{234}U$ Variability

526 Stalagmite δ^{13} C and δ^{18} O values covary with changes in SST at orbital time scales. The 527 offset between interglacial and glacial isotopic values averages ~3‰ for δ^{18} O and ~7‰ for δ^{13} C 528 values (Fig. 10). Stalagmite δ^{234} U values also preserve these changes in aridity. Millennial-scale

changes are also recorded in stalagmite carbon isotope ratios, with shifts of 3-7‰ associated 529 with GI/GS transitions, and oxygen isotopic changes of ~1-2‰. The large swing in δ^{18} O values 530 531 during the transition from GI-1 to the Younger Dryas (YD) (~5% from 14.0-13.5 ka) is anomalous. Given that the change in δ^{13} C values at this time (6‰) is consistent with other GI 532 transitions, the hydroclimatic implications of this interval require additional study. Similarly, 533 534 oxygen and carbon isotopic variability is pronounced during the late Holocene portion of the BG 535 record. The origin of this high variability is unclear. Replication of the Holocene portion of this 536 record currently underway will help address this question (Thatcher et al., 2018).

537 Where growth is continuous during HS, the link between stalagmite isotopic variations 538 and SST changes is clearly visible (Fig. 11). Prominent positive carbon isotopic excursions 539 define the YD, HS2, HS5, HS6, and HS8, consistent with diminished concentrations of arboreal 540 pollen in cores from the Iberian margin, and serve to document particularly cold and dry 541 conditions at these times (Sánchez Goñi et al., 2000; Roucoux et al., 2006; Sánchez Goñi et al., 2008). Reduced stalagmite δ^{13} C values mark periods of enhanced effective moisture from 170-542 543 160 and 145-135 ka, tracking peaks in temperate tree pollen and alkenone-based SST. The BG record reveals a pronounced increase in stalagmite δ^{13} C values during the YD, at odds with the 544 545 plateau in SST observed in some Portuguese coastal margin sediments at this time. However, a 546 higher resolution SST record reveals a pronounced drop in SST (Rodrigues et al., 2010), well 547 matched with the BG isotopic profile and the stalagmite record from Villars Cave.

548 Hydroclimatic shifts associated with GS and GI are most clearly expressed during MIS 549 5a and 5b in the BG carbon isotope record (Fig. 11). Other European stalagmite records have 550 identified GI/GS events from the last glacial period (Genty et al., 2003; Spötl et al., 2006; Boch 551 et al., 2011; Moseley et al., 2014) (Fig. 10), but the level of resolution recorded in the BG/GCL 552 time series has not been clearly identified previously in western Iberia. A carbon isotope time 553 series (albeit with low temporal resolution) of a flowstone from southeastern Spain does not 554 present clear evidence of either GI or most HS during the last glacial cycle, although it does 555 contain a clear expression of HS11 (Hodge et al., 2008) (Fig. 1). And while some Iberian lakes 556 and peat bogs document environmental changes concurrent with HS, no single record, including 557 one of the longest - the 50 ka time series from the Fuentillejo maar, south-central Spain -558 contains a consistent signal for all HS (Vegas et al., 2010; Moreno et al., 2012) (Fig. 1). GS/GI 559 oscillations during MIS 3 are not clearly defined in BG stalagmites, likely owing to insufficient temporal resolution, although the BG records do share a resemblance to reconstructed SSTvariability (Fig. 11).

562 Whether the apparent inconsistent linkages between Iberian margin SST and Iberian 563 hydroclimate are due to the limitations of these proxies, region-specific responses to SST 564 variations, or a changing influence of SST on precipitation is unclear. However, other points of 565 divergence between SST and the BG/GCL records exist. For example, some marine cores reveal 566 a prominent spike in forest taxa occurring at the start of interglacials, decreasing thereafter for 567 the next 5-10 kyr (Tzedakis et al, 2004; Desprat et al., 2007) (Fig. 10). This early interglacial 568 peak is a common feature in several time series including the Antarctic δD (Petit et al., 1999) 569 and CH₄ records (Loulergue et al., 2008), and in stalagmite isotopic ratios from the eastern 570 Mediterranean (Bar-Matthews et al., 2003) and southern France (Couchoud et al., 2009) (Fig. 10). The BG/GCL δ^{13} C and δ^{18} O records lack this feature, although the previously discussed 571 issues surrounding the continuity of the MIS6/5e transition may complicate identifying it. 572

Stalagmite δ^{13} C and δ^{18} O values are lower during GI 20-22 (MIS 5a/4; 84-72 ka) than in 573 either the Holocene or MIS 5e (Fig. 10 and 12), and BG6LR δ^{234} U values support this 574 575 observation. This interval is of particular interest given that Atlantic forest pollen, which has 576 been used as a proxy for air temperature, was decoupled from SST across northwestern Iberia 577 during cold events (C18-C20) (Rousseau et al., 2006; Rasmussen et al., 2014). This decoupling 578 is interpreted as reflective of a weakened control of SST on Iberian atmospheric temperature 579 that, in turn, enhanced transport of atmospheric vapor to the high latitudes, amplifying 580 production of ice sheets in the early stages of the last glacial cycle (Sánchez Goñi et al., 2013). 581 This process has also been demonstrated for an earlier interglacial (MIS 19; Sánchez Goñi et al., 2016). Other offsets include (1) the gradual change in BG δ^{13} C and δ^{18} O values across the MIS 582 8/7 boundary, in contrast to the sharp rise in SST at this time; (2) the anomalously large δ^{13} C 583 response to ice rafting event C24 (111-108 ka), and (3) the persistence of low δ^{13} C values as SST 584 585 decreased from 205-187 ka (Fig. 11 and 12).

The mechanism linking SST and Iberian hydroclimate over millennial time scales remains unclear. The NAO exerts a strong control over Iberian precipitation, and previous studies have suggested that GS and GI (Moreno et al., 2002; Sánchez Goñi et al., 2002; Daniau et al., 2007) and HS (Naughton et al., 2009) were characterized by distinct NAO modes. The dynamics of the NAO and Azores High pressure system prior to the historical era are only beginning to be understood (Trouet et al., 2009; Olsen et al., 2012; Wassenburg et al., 2013), and the BG/GCL record cannot address this question independently. However, rainfall variability in eastern Iberia is less closely tied to the NAO than is western Iberia and instead reflects other climatic phenomena including the El Niño-Southern Oscillation (Rodó et al., 1997), helping to produce an east-west precipitation gradient. Additional high-resolution speleothem records from central and eastern Iberia could therefore provide a more robust test of the underlying drivers of millennial-scale hydroclimatic changes during recent glacial periods.

598

599 6. Conclusions

600 The BG/GCL composite speleothem record demonstrates that the hydroclimate and 601 vegetation dynamics in west-central Portugal tracked Iberian margin SST over orbital and 602 millennial scales during the past two glacial cycles. Enhanced aridity characterized HS, as 603 evidenced by elevated carbon and oxygen isotopic ratios and/or hiatuses in stalagmite growth, 604 consistent with other regional stalagmite time series. GI/GS variability expressed in the Iberian 605 margin SST record and in co-deposited pollen spectra is also present in the BG/GCL time series, 606 and is particularly well defined in MIS 5a and 5b. Understanding differences between the 607 structures of the stalagmite and SST records during some time intervals will require development 608 of speleothem records from central and southern Iberia.

609

610 Acknowledgements

611 This work was supported by the Center for Global and Regional Environmental Research

and Cornell College (to R.F.D.), and the U.S. National Science Foundation (grant BCS-1118155

to J.A.H., BCS-1118183 to M.M.B., and AGS-135539 to C.C.U.). Field sampling performed

under the auspices of IGESPAR (to J.A.H.) and Associação de Estudos Subterrâneos e Defesa do

615 Ambiente. Brandon Zinsious and Stephen Rasin contributed to fieldwork at BG, and Zachary

616 LaPointe assisted with radioisotopic analyses; Suzanne Ankerstjerne performed stable isotope

617 measurements. Use of the following data sets is gratefully acknowledged: Global Precipitation

618 Climatology Center data by the German Weather Service (DWD) accessed through

619 <u>http://gpcc.dwd.de;</u> NAO Index Data provided by the Climate Analysis Section, NCAR, Boulder,

620 USA, Hurrell (2003). Updated regularly. Accessed through

621 https://climatedataguide.ucar.edu/climate-data/hurrell-north-atlantic-oscillation-nao-index-pc-

- 622 <u>based</u>. This manuscript benefitted tremendously from discussions with Maria F. Sánchez Goñi,
- 623 David Hodell, and Chronis Tzedakis. We thank four anonymous reviewers and associate editor
- 624 Nathalie Combourieu-Nebout, who together substantially improved this manuscript's scope and
- 625 clarity through detailed and thoughtful assessments. Stable and U-series isotope data are
- available at the NOAA National Centers for Environmental Information website.
- 627

628 References

- de Abreu, L., Shackleton, N.J., Schönfeld, J., Hall, M., and Chapman, M.: Millennial-scale
 oceanic climate variability off the Western Iberian margin during the last two glacial periods,
 Marine Geol, 196, 1-20, 2003.
- Asmerom, Y., Polyak, V., Burns, S.: Variable winter moisture in the southwestern United States
 linked to rapid glacial climate shifts, Nat Geosci, 3, 114-117, 2010.
- Baker, A., Ito, E., Smart, P., McEwan, R.: Elevated and variable values of ¹³C in speleothems in
 a British cave system, Chem Geol, 136, 263-270, 1997.
- Barker, S., Knorr, G., Edwards, R.L., Parrenin, F., Putnam, A.E., Skinner, L.C., Wolff, E.,
 Ziegler, M.: 800,000 years of abrupt climate variability, Science, 334, 347-351, 2011.
- Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A., Hawkesworth, C.J.: Sea-land
 oxygen isotopic relationships from planktonic foraminifera and speleothems in the Eastern
 Mediterranean region and their implication for paleo-rainfall during interglacials intervals,
 Geochim Cosmochim Ac, 67, 3181–3199, 2003.
- Barnston, A.G. and Livezey, R.E.: Classification, seasonality, and persistence of low-frequency
 atmospheric circulation patterns, Mon Weather Rev, 115, 1083–1126, 1987.
- Berger, A. and Loutre, M.F.: Insolation values for the climate of the last 10 million years.
 Quaternary Sci Rev 10, 297-318, 1991.
- 646 Blanco Castro, E., Casado González, M.A., Costa Tenorio, M., Escribano Bombín, R., García
- 647 Antón, M., Génova Fuster, M., Gómez Manzaneque, F., Gómez Manzaneque, A., Moreno
- Sáiz, J.C., Morla Juaristi, C., Regato Pajares, P., Sáiz Ollero, H.: Los bosques ibéricos.
 Planeta, Barcelona, 1997.
- 650 Boch, R., Cheng, H., Spötl, C., Edwards, R.L., Wang, X., Hauselmann, Ph.: NALPS: a precisely
- dated European climate record 120-60 ka, Clim Past, 7, 1049-1072, 2011.

- Breitenbach, S.F.M., Rehfeld, K., Goswami, B., Baldini, J.U.L., Ridley, H.E., Kennett, D.J.,
 Prufer, K.M., Aquino, V.V., Asmerom, Y., Polyak, V.J., Cheng, H., Kurths, J., Marwan, N.:
 Constructing proxy records from age models (COPRA), Clim Past, 8, 1765-1779, 2012.
- Breitenbach, S.F.M., Lechleitner, F.A., Meyer, H., Diengdo, G., Mattey, D., Marwan, N.: Cave
- ventilation and rainfall signals in dripwater in a monsoonal setting a monitoring study from
 NE India, Chem Geol, 402, 111-124, 2015.
- 658 CDG: The Climate Data Guide: Hurrell North Atlantic Oscillation (NAO) Index (PC-based).
 659 Retrieved from <u>https://climatedataguide.ucar.edu/climate-data/hurrell-north-atlantic-</u>
 660 oscillation-nao-index-pc-based, accessed 21 May, 2018.
- Chabaud, L., Sánchez Goñi, M.F., Desprat, S., Rossignol, L.: Land-sea climatic variability in the
 eastern North Atlantic subtropical region over the last 14,200 years: Atmospheric and
 oceanic processes at different timescales Holocene, 24, 787-797, 2014.
- Chabaux, F., Riotte, J., Dequincey, O.: U–Th–Ra fractionations during weathering and river
 transport. Rev Mineral Geochem, 52, 533–576. 2003.
- 666 Collister, C. and Mattey, D.: Controls on water drop volume at speleothem drip sites: An
 667 experimental study. J Hydrol, 358, 259-267, 2008.
- 668 Combourieu-Nebout, N., Peyron, O., Dormoy, I., Desprat, S., Beaudouin, C., Kotthoff, U.,
 669 Marret, F.: Rapid climatic variability in the west Mediterannean during the last 25 000 years
 670 from high resolution pollen data, Clim Past, 5, 503-521, 2009.
- 671 Cortesi, N., Gonzalez-Hidalgo, J.C., Trigo, R.M., and Ramos, A.M.: Weather types and spatial
 672 variability of precipitation in the Iberian Peninsula. International Journal of Climatology, 34,
 673 2661-2677, 2014.
- 674 Couchoud, I, Genty, D., Hoffman, D., Drysdale, R., Blamart, D.: Millennial-scale climate
 675 variability during the Last Interglacial recorded in a speleothem from south-western France.
 676 Quaternary Sci Rev, 28, 3263-3274, 2009.
- Daniau, A.-L., Sánchez Goñi, M.F., Beaufort, L., Laggoun-Defarge, F., Loutre, M.-F., Duprat, J.:
 Dansgaard-Oeschger climatic variability revealed by fire emissions in southwestern Iberia.
 Quaternary Sci Rev, 26, 1369-1383, 2007.
- Darfeuil, S., Ménot, G., Giraud, X., Rostek, F., Tachikawa, K., Garcia, M., Bard, É.: Sea surface
 temperature reconstructions over the last 70 kyr off Portugal: Biomarker data and regional
 modeling, Paleocean, 31, 40–65, 2016.

- Deines, P.: The isotopic composition of reduced organic carbon. Handbook of Environmental
 Isotope Geochemistry, The Terrestrial Environment, Part A (Fritz, P. and Fontes, J., Eds.,
 Elseveier, New York, 331-406, 1980.
- Denniston, R.F., González, L.A., Asmerom, Y., Baker, R.G., Reagan, M.K. Bettis, E.A. III.:
 Evidence for increased cool season moisture during the middle Holocene, Geology, 27, 815818, 1999.
- Denniston, R.F., Wyrwoll, K.-H., Polyak, Brown, J. Asmerom, Y., Wanamaker, A. Jr., LaPointe⁶
 Z., Ellerbroek, R., Barthelmes, M., Cleary, D., Cugley, J., Woods, D., Humphreys, W.: A
 Stalagmite Record of Holocene Indonesian-Australian Summer Monsoon Variability from
 the Australian Tropics. Quaternary Sci Rev 78, 155-168, 2013.
- Desprat et al.: Climatic variability of Marine Isotope Stage 7: direct land-sea-ice correlation from
 a multiproxy analysis of a north-western Iberian margin deep-sea core. Quaternary Sci Rev
 25, 1010-1026, 2006.
- Desprat, S., Sánchez Goñi, M.F., Naughton, F., Turon, J.-L., Duprat, J., Malaizé, B., Cortijo, E.,
 Peypouquet, J.-P.: Climate variability of the last five isotopic interglacials: Direct land-seaice correlation from the multiproxy analysis of North-Western Iberian margin deep-sea cores,
 Editor(s): F. Sirocko, M. Claussen, M.F. Sánchez Goñi, T. Litt *In* Developments in
 Quaternary Sciences, Elsevier, pp. 375-386, 2007.
- Dorale, J.A., and Liu, Z.: Limitations of Hendy Test criteria in judging the paleoclimatic
 suitability of speleothems and the need for replication. J Cave Karst Stud, 71, 73-80, 2009.
- 703 Dreybodt, W.: Evolution of the isotopic composition of carbon and oxygen in a calcite
 704 precipitating H2O-CO2-CaCO3 solution and the related isotopic composition of calcite in
 705 stalagmites. Geochim. Cosmochim. Acta, 72, 4712-4724, 2008.
- 706 Eynaud et al.: Position of the Polar Front along the western Iberian margin during key cold 707 episodes of the last 45 ka. Geochem Geophy Geosys, 10, Q07U05, 708 doi:10.1029/2009GC002398, 2009.
- Fairchild, I.J., Smith, C.L., Baker, A., Fuller, L., Spötl, C., Mattey, D., McDermott, F., E.I.M.F.:
- Modification and preservation of environmental signals in speleothems. Earth Sci Rev 75,
 105-153, 2006.

- Fankhauser, A., McDermott, F., and Fleitmann, D.: Episodic speleothem deposition tracks the
 terrestrial impact of millennial-scale last glacial climate variability in SW Ireland, Quaternary
 Sci Rev, 152, 104-117, 2016.
- von Fischer, J.C., Tieszen, L.L., and Schimel, D.S.: Climate controls on C₃ vs. C₄ productivity in
 North American grasslands from carbon isotope composition of soil organic matter. Global
 Change Bio, 14, 1–15, 2008.
- Fletcher, W.J., Sánchez Goñi, M.F., Allen, J.R.M., Cheddadi, R., Combourieu-Nebout, N.,
 Huntley, B., Lawson, I., Londiex, L., Magri, D., Margari, v., Müller, U.C., Naughton, F.,
 Novenko, E., Roucoux, K., Tzedakis, P.C.: Millennial scale variability during the last glacial
 in vegetation records from Europe. Quaternary Sci Rev, 29, 2839-2864, 2010.
- Florineth, D. and Schlüchter, S. Alpine Evidence for Atmospheric Circulation Patterns in Europe
 during the Last Glacial Maximum. Quaternary Research, 54, 295-308, 2000.
- Francey, R. J., Allison, C. E., Etheridge, D. M., Trudinger, C. M., Enting, I. G., Leuenberger, M.,
 Langenfelds, R. L., Michel, E., and Steele, L. P. A.: 1000-year high precision record of ¹³C in
 atmospheric CO₂. Tellus B: Chemical and Physical Meteorology, 51, 170-193, 1999.
- Genty, D., Blamart, D., Ouahdi, R., Gilmour, M., Baker, A., Jouzel, J., Van-Exter, S.: Precise
 dating of Dansgaard-Oeschger climate oscillations in western Europe from stalagmite data,
 Nature, 421, 833-837, 2003.
- Genty, D., Comboruieu-Nebout, N., Peyron, O., Blamart, D., Wainer, K., Mansuri, F., Ghaleb,
 B., Isabello, L., Dormoy, I., von Grafenstein, U., Bonelli, S., Landais, A., Brauer, A.:
 Isotopic characterization of rapid climatic events during OIS3 and OIS4 in Villars Cave
 stalagmites (SW-France) and correlation with Atlantic and Mediterranean pollen records.
 Quaternary Sci. Rev., 29, 2799-2820, 2010.
- Genty, D., Blamart, D., Ghaleb, B., Plagnes, V., Causse, Ch., Bakalowicz, M., Zouari, K., Chkir,
 N., Hellstrom, J., Wainer, K., and Bourges, F.: Timing and dynamics of the last deglaciation
 from European and North African δ¹³C stalagmite profiles comparison with Chinese and
 South Hemisphere stalagmites. Quaternary Sci Rev 25, 2118-2142, 2006.
- Gimeno, L., Nieto, R., Trigo, R.M., Vicente-Serrano, S.M., and López-Moreno, J.I., Where does
 the Iberian Peninsula moisture comr from? An answer based on a Lagrangian approach. J.
- 741 Hydrometeorology, 11, 421-436, 2010.

- Gimeno, L., Stohl, A., Trigo, R.M., Dominguez, F., Yoshimura, K., Yu., L., Drumond, A.,
 Durán-Quesada, A.M., Nieto, R.: Oceanic and terrestrial sources of continental precipitation.
 Rev Geophy, 50, 1-41, 2012.
- Gómez-Orellana, L., Ramil-Rego, P., & Sobrino, C. M.: The Würm in NW Iberia, a pollen
 record from Area Longa (Galicia). Quaternary Res, 67, 438-452, 2008.
- 747 Grootes, P. M.: *Climate Change in Continental Isotopic Records*, P. K. Swart, K. C. Lohmann, J.
- McKenzie, S. Savin, Eds. (American Geophysical Union, Washington, DC), pp. 37-46,
 1993.
- Hellstrom, J. and McCulloch, M.: Multi-proxy constraints on the climatic significance of trace
 element records from a New Zealand speleothem. Earth Planet Sci Lett, 179, 287-297, 2000.
- Hendy, C.: The isotopic geochemistry of speleothems I. The calculation of the effects of
 different modes of formation on the isotopic composition of speleothems and their
 applicability as palaeoclimatic indicators. Geochimica et Cosmochica Acta 35, 801-824,
 1971.
- Herbert, T.D., Schuffert, J.D., Heusser, L., Lyle, M., Mix, A., Ravelo, A.C., Stott, L.D., and
 Herguera, J.C.: Collapse of the California current during glacial maxima linked to climate
 change on land. Science, 293, 71-76, 2001.
- Hodell, D., Crowhurst, S., Skinner, L., Tzedakis, P.C., Margari, V., Channell, J.E.T., Kamenov,
 G., Maclachlan, S., Rothwell, G.: Response of Iberian margin sediments to orbital and
 suborbital forcing over the past 420 ka, Paleoceanography, 28, 185-199, 2013.
- Hodge, E.J., Richards, D.A., Smart, P.L., Andreo, B., Hoffman, D.L., Mattey, D.P., GonzalesRamon, A.: Effective precipitation in southern Spain (~266 to 46 ka) based on a speleothem
 stable carbon isotope record. Quaternary Res, 69, 447-457, 2008.
- 765 IAEA/WMO: Global Network of Isotopes in Precipitation. The GNIP Database. Accessible at:
 766 http://www.iaea.org/water, 2016.
- 767 IPMA: Accessible at http://www.meteo.pt/en/oclima/clima.normais/015/, 2012.
- Justino, F. and Peltier, W.R.: The glacial North Atlantic Oscillation. Geophysical Research
 Letters, 32, L21803, 2008.
- 770 Kim, S.-T. and O'Neil, J.R.: Equilibrium and nonequilibrium oxygen isotope effects in synthetic
- carbonates: Geochim Cosmochim Ac, 61, 3461-3475, 1997.

- Kuhlemann, J et al.: Regional synthesis of Mediterranean atmospheric circulation during the Last
 Glacial Maximum. Science, 321, 1338–1340, 2008.
- Lachniet, M.S.: Climatic and environmental controls on speleothem oxygen isotope values.
 Quaternary Sci Rev 28, 412-432, 2009.
- Lechleitner, F.A., Breitenbach, S.F.M., Cheng, H., Plessen, B.: Climatic and in-cave influences
 on d18O and d13C in a stalagmite from northeastern India through the last deglaciation. Quat
 Res, 88, 458-471, 2017.
- Li, Z-H., Driese, S.G., Cheng, H.: A multiple cave deposit assessment of suitability of
 speleothem isotopes for reconstructing palaeo-vegetation and palaeo-temperature.
 Sedimentology, 61, 749-766, 2014.
- Lorenzo, M.N., Iglesias, I., Taboada, J.J., Gomez-Gesteira, M.: Relationship between monthly
 rainfall in northwest Iberian Peninsula and North Atlantic sea surface temperature. Int J
 Climatology, 30, 980-990, 2010.
- Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola,
 J.-M., Raynaud, D., Stocker, T.F., Chappellaz, J.: Orbital and millennial-scale features of
 atmospheric CH4 over the past 800,000 years, Nature, 453, 383-386, 2008.
- Luetscher, M., Boch, R., Sodemann, H., Spötl, C., Cheng, H., Edwards, R.L., Frisia, S., Hof, F.,
 and Müller, W.: North Atlantic storm track changes during the Last Glacial Maximum
 recorded by Alpine speleothems. Nature Communications, 6, DOI: 10.1038/ncomms7344,
 2016.
- Margari, V., Skinner, L.C., Tzedakis, P.C., Ganopolski, A., Vautravers, M., and Shackleton,
 N.J.: The nature of millennial-scale climate variability during the past two glacial periods:
 Nature Geoscience, v. 3, p. 127–131, doi:10.1038/ngeo740, 2010.
- Margari, V., Skinner, L.C., Hodell, D.A., Martrat, B., Toucanne, S., Grimalt, J.O., Gibbard, P.L.,
 Lunkka, J.P., Tzedakis, P.C.: Land-ocean changes on orbital and millennial time scales and
 the penultimate glaciation, Geology, 42, 183-186, 2014.
- Martin-Vide, J. and Lopez-Bustins, J-A.: The Western Mediterranean Oscillation and rainfall in
 the Iberian Peninsula. Int J Climatol, 26, 1455-1475, 2006.
- 800 Martrat, B., Grimalt, J.O., Shackleton, N.J., de Abreu, L., Hutterli, M.A., Stocker, T.F.: Four
- climate cycles of recurring deep and surface water destabilizations on the Iberian margin,
 Science, 317, 502-507, 2007.

- McManus, J.F., Oppo, D.W., Cullen, J.L.: A 0.5-Million-Year Record of Millennial-Scale
 Climate Variability in the North Atlantic, Science, 283, 971-975, 1999.
- Mickler, P.J. et al.: Stable isotope variations in modern tropical speleothems: Evaluation
 equilibrium vs. kinetic isotope effects. Geochim Cosmochim Acta. 68, 4381-4393, 2004.
- Moreno, A., Cacho, I., Canals, M., Prins, M.A., Sánchez Goñi, M.F., Grimalt, J.O., Weltje, G.J.
 Saharan dust transport and high-latitude glacial climate variability: the Alboran Sea record.
 Quaternary Res, 58, 318-328, 2002.
- Moreno, A., Gonzalez-Samperiz, P., Morellon, M., Valero-Garces, B.L., Fletcher, W.J.:
 Northern Iberian abrupt climate change dynamics during the last glacial cycle: a view from
 lacustrine sediments. Quaternary Sci Rev, 36, 139-153, 2012.
- 813 Moreno, A., Sancho, C., Bartolumé, M., Oliva-Rucia, B., Delgado-Huertas, A., José, Estrela, M.,
- Corell, D., López-Moreno, J.I., Cacho, I.: Climate controls on rainfall isotopes and their effects on cave drip water and speleothem growth: the case of Molinos cave (Teruel, NE Spain). Clim Dyn, 43, 221-241, 2014.
- Moseley, G.E., Spötl, C., Scensson, A., Cheng, H., Brandstatter, S., Edwards, R.L.: Multispeleothem record reveals tightly coupled climate between central Europe and Greenland
 during Marine Isotope Stage 3. Geology, 42, 1043-1946, 2014.
- Mühlinghaus, C., Scholz, D., and Mangini, A.: Modelling stalagmite growth and d13C as a
 function of drip interval and temperature. Geochim. Cosmochim. Acta, 71, 2780-2790, 2007.
- 822 Naughton, F., Sánchez Goñi, M.F., Desprat, S., Turon, J.-L., Duprat, J., Malaizé, B., Joli, C.,
- 823 Cortijo, E., Drago, T., Freitas, M.C.: Present-day and past (last 25,000 years) marine pollen
 824 signal off western Iberia. Mar Micropaleontol, 62, 91–114, 2007.
- Naughton, F., Sánchez Goñi, M.F., Kageyama, M., Bard, E., Duprat, J., Cortijo, E., Desprat, S.,
 Malaizé, B., Joly, C., Rostek, F., Turon, J.-L.: Wet to dry climatic trend in north-western
 Iberia within Heinrich events. Earth Planet Sc Lett, 284, 329-342, 2009.
- North Greenland Ice Core Project members: High-resolution record of Northern Hemisphere
 climate extending into the last interglacial period, Nature, 431, 147-151, 2004.
- 830 Olsen, J., Anderson, N.J., and Knudsen, M.F.: Variability of the North Atlantic Oscillation over
- the past 5,200 years. Nature Geoscience, 5, 808-812, 2012.

- Oster, J.L., Ibarra, D.L., Harris, C.H., Maher, K.: Influence of eolian deposition and rainfall
 amounts on the U-isotopic composition of soil water and soil minerals. Geochim Cosmochim
 Ac, 88, 146 166, 2012.
- Paredes, D., Trigo, R.M., Garcia-Herrera, R., Franco Trigo, I.: Understanding precipitation
 changes in Iberia in early spring: weather typing and storm-tracking approaches. J
 Hydrometeorol, 7, 101-113, 2006.
- Petit, J.R., Jouzel, J., Raynaud, D., Barkov, N.I., Barnola, J.-M., Basile, I., Bender, M.,
 Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V.M., Legrand, M.,
 Lipenkov, V.Y., Lorius, C., Pepin, L., Ritz, C., Saltzman, E., Stievenard, M.: Climate and
 atmospheric history of the 420,000 years from the Vostok ice core, Antarctica. Nature, 399,
 429-436, 1999.
- Plagnes, V., Causse, C., Genty, D., Paterne, M., Blamart, D.: A discontinuous climatic record
 from 187 to 74 ka from a speleothem of the Clamouse Cave (south of France). Earth Planet
 Sci Lett, 201, 87-103, 2002.
- Polyak, V.J., Asmerom, Y., Burns, S.J., and Lachniet, M.S.: Climatic backdrop to the terminal
 Pleistocene extinction of North American mammals, Geology, 40, 1023-1026, 2012.
- Rasmussen, S.O. et al.: A stratigraphic framework for abrupt climatic changes during the Last
 Glacial period based on three synchronized Greenland ice-core records: redefining and
 extending the INTIMATE event stratigraphy, Quaternary Sci Rev, 106, 14–28, 2014.
- Rey Benayas, J.M. and Scheiner, S.M.: Plant diversity, biogeography and environment in Iberia:
 Patterns and possible causal factors. J Veg Sci, 13, 245-258, 2002.
- Rodó, X., Baert, E., Comin, F.A.: Variations in seasonal rainfall in Southern Europe during the
 present century: relationships with the North Atlantic Oscillation and the El Niño-Southern
 Oscillation. Clim Dynam, 13, 275-284, 1997.
- 856 Rodrigues et al.: The last glacial-interglacial transition (LGIT) in the western mid-latitudes of the
- North Atlantic: Abrupt sea surface temperature change and sea level implications.
 Quaternary Sci Rev, 29, 1853-1862, 2010.
- Roucoux, K.H., de Abreu, L., Shackleton, N.J., Tzedakis, P.C.: The response of NW Iberian
 vegetation to North Atlantic climate oscillations during the last 65kyr, Quaternary Sci Rev,
- 861 24, 1637-1653, 2005.

- Roucoux, K.H., Tzedakis, P.C., de Abreu, L., Shackleton, N.J.: Climate and vegetation changes
 180,000 to 345,000 years ago recorded in a deep-sea core off Portugal. Earth Planet Sci Lett,
 249, 307-325, 2006.
- Rousseau, D.D., Kukla, G., McManus, J.: What is what in the ice and the ocean? Quaternary Sci
 Rev, 25, 2025-2030, 2006.
- K. L. Rosanski, Araguas-Araguas, R. Gonfiantini, in *Climate Change in Continental Isotopic Records*, P. K. Swart, K. C. Lohmann, J. McKenzie, S. Savin, Eds. (American Geophysical
 Union, Washington, DC), pp. 1–36, 1993.
- 870 Saltzman, Matthew & Thomas, E. (2012). Carbon Isotope Stratigraphy. The Geologic Time
 871 Scale, 1, 207-232. 2012.
- Sánchez Goñi, M.F., Turon, J.L., Eynaud, F., Gendreau, S., European climatic response to
 millennial-scale changes in the atmosphere-ocean system during the Last Glacial Period.
 Quaternary Res, 54, 394-403, 2000.
- Sánchez Goñi, M.F., Cacho, I., Turon, J-L., Guiot, J., Sierro, F.J., Peypouquet, J.-P., Grimalt,
 J.O., Shackleton, N.J.: Synchroneity between marine and terrestrial responses to millennial
 scale climatic variability during the last glacial period in the Mediterranean region. Clim
 Dynam, 19, 95-105, 2002.
- Sánchez Goñi, M.F., Landais, A., Fletcher, W.J., Naughton, F., Desprat, S., Duprat, J.:
 Contrasting impacts of Dansgaard-Oeschger events over a western European latitudinal
 transect modulated by orbital precession. Quaternary Sci Rev, 27, 1136-1151, 2008.
- Sánchez Goñi, M.F., Bard, E., Landais, A., Rossignol, L., d'Errico, F.: Air-sea temperature
 decoupling in western Europe during the last interglacial-glacial transition. Nat Geosci, 6,
 884 837-841, 2013.
- Sánchez Goñi, M.F., Rodrigues, T., Hodell, D.A., Polanco-Martinez, J.M., Alonso-Garcia, M.,
 Hernandez-Almeida, I., Desprat, S., Ferretti, P.: Tropically-driven climate shifts in
 southwestern Europe during MIS 19, a low eccentricity interglacial. Geophys Res Abst, 18,
 EGU2016-3940, 2016.
- Schneider, U., Becker, A., Finger, P., Meyer-Christoffer, A., Ziese, M., Rudolf, B.: GPCC's new
 land surface precipitation climatology based on quality-controlled in situ data and its role in
 quantifying the global water cycle. Theoret Appl Climatol, 115, 15-40, 2013.

- Schneider, U., A. Becker, P. Finger, A. Meyer-Christoffer, M. Ziese, and B. Rudolf (2014),
 GPCC's new land surface precipitation climatology based on quality-controlled in situ data
 and its role in quantifying the global water cycle, Theor. Appl. Climatol., 115, 1–15.
- Schrag, D.P, Hampt, G., and Murray, D.W.: Pore fluid constraints on the temperature and
 oxygen isotopic composition of the glacial ocean. Science, 272, 5270, 1930-1932, 1996.
- Spötl, C., Mangini, A., Richards, D.A.: Chronology and paleoenvironment of Marine Isotope
 Stage 3 from two high-elevation speleothems, Austrian Alps. Quaternary Sci Rev, 25, 11271136, 2006.
- Stoll, H.M., Moreno, A., Mendez-Vincente, A., Gonzalez-Lemos, S., Jimenez-Sánchez, M.,
 Dominguez-Cuesta, M.J., Edwards, R.L., Cheng, H., Wang, X.: Paleoclimate and growth
 rates of speleothems in the northwestern Iberian Peninsula over the last two glacial cycles.
 Quaternary Res, 80, 284-290, 2013.
- 904 Thatcher, D.L., Wanamaker, A.D., Jr., Denniston, R.F., Asmerom, Y., Ummenhofer, C.C.,
 905 Polyak, V.J., Hasiuk, F., Haws, J.A., and Gillikin, D.P.: Changes in hydroclimate in Iberia in
 906 the last 1200 years: insights from speleothem records from western Portugal. Geological
 907 Society of America North-Central Meeting Abstracts with Programs, Ames, Iowa, 2018.
- 908 Trigo, R.M., Osborn, T.J., Corte-Real, J.M.: The North Atlantic Oscillation influence on Europe:
 909 climate impacts and associated physical mechanisms, Clim Res, 20, 9-17, 2002.
- 910 Trouet, V., Esper, J., Graham, N.E., Baker, A., Scourse, J.D., Grank, D.C.: Persistent positive
 911 North Atlantic Oscillation mode dominated the Medieval Climate Anomaly. Science, 324,
 912 78-80, 2009.
- 913 Tzedakis, P.C., Roucoux, K.H., de Abreu, L., Shackleton, N.J.: The duration of forest stages in
 914 southern Europe and interglacial climate variability, Science, 306, 2231-2235, 2004.
- 915 Vaks, A., Gutareva, O.S., Breitenbach, S.F.M., Avirmed, E., Mason, A.J., Thomas, A.L., Osinev,
- A.V., Kononov, A.M., Henderson, G.M.: Speleothems reveal 500,000-year history of
 Siberian permafrost. Science, 340, 183-186, 2013.
- Vandenberghe, J., French, H.M., Gorbunov, A., Marchenko, S., Velichko, A.A., Jin, H., Cui,
 Z., Zhang, T., Wan, X.: The Last Permafrost Maximum (LPM) map of the Northern
- Hemisphere: permafrost extent and mean annual air temperatures, 25-17 ka BP, Boreas, 43,
- 921 652–666, 2014.

- 922 Vegas, J., Ruiz-Zapata, B., Ortiz, J.E., Galan, L., Torres, T., Garcia-Cortes, A., Gil-Garcia, M.J.,
 923 Perez-Gonzalez, A., Gallardo-Millan, J.L.: Identification of arid phases during the last 50 cal.
 924 ka BP from the Fuentillejo maar-lacustrine record (Campo de Calatrava Volcanic Field,
- 925 Spain), J Quaternary Sci, 25, 1051-1062, 2010.
- 926 Voelker, A.H. L., Rodrigues, T., Stein, R., Hefter, J., Billups, K., Oppo, D., McManus, J.,
- 927 Grimalt, J.O.: Variations in mid-latitude North Atlantic surface water properties during the
- 928 mid-Brunhes (MIS 9-14) and their implications for thermohaline circulation, Clim Past, 6, p.
 929 531-552, 2010.
- Voelker, A.H.L. and de Abreu, L.: A review of abrupt climate change events in the northeastern
 Atlantic Ocean (Iberian Margin): Latitudinal, Longitudinal, and Vertical Gradients. Abrupt
- 932 Climate Change: Mechanisms, Patterns, and Impacts (Eds. Rashid, H., Polyak, L., and
- Mosley-Thompson, E.), Geophysical Monograph Series 193, 15-37, 2011.
- 934 Wainer, K., Genty, D., Blamart, D., Daëron, M., Bar-Matthews, M., Vonhof, H., Dublyansky,
- 935 Y., Pons-Branchu, E., Thomas, L., van Calsteren, P., Quinif, Y., and Caillon, N.: Speleothem 936 record of the last 180 ka in Villars cave (SW France): Investigation of a large δ^{18} O shift
- between MIS6 and MIS5. Quaternary Sci. Rev., 30, 130-146, 2011.
- Wassenburg, J.A., Immenhauser, A., Richter, D.K., Niedermayr, A., Riechelmann, S., Fietzke,
 J., Scholz, D., Jochum, K.P., Fohlmeister, J., Schröder-Ritzrau, Sabaoui, A., Riechelmann,
- 940 D.F.C., Schneider, L., Esper, J.: Moroccan speleothem and tree ring records suggest a
- 941 variable positive state of the North Atlantic Oscillation during the Medieval Warm Period.
- 942 Earth Planet. Sci. Lett., 375, 291-302, 2013.
- 943 Zhou, J., Lundstrom, C.C., Fouke, B., Panno, S., Hackley, K., and Curry, B. Geochemistry of
- 944 speleothem records from southern Illinois: Development of $\binom{234}{238}U$ as a proxy for
- paleoprecipitation. Chemical Geology, 221, 1-20, 2005.
- 946



951 Figure 1. Average annual precipitation (mm) of the Iberian Peninsula for years AD 1901-952 2009 (GPCC v. 6; Schneider et al., 2013) relative to cave study sites (white stars: GLC = 953 Gruta do Casal da Lebre; BG = Buraca Gloriosa). Rectangle denotes location of northwest 954 Spain cave sites (NWSC) (Moreno et al., 2010; Stoll et al., 2013); FM = Fuentillejo maar (Vegas 955 et al., 2010) and GT = Gitana cave (Hodge et al., 2008); VC = Villars Cave (Genty et al., 2003) 956 located just north of map. Also shown are locations of marine cores discussed in text and GNIP 957 stations at Porto, Vila Real, and Portalegre. Bathymetric contours shown in grey (m). Location of 958 currents after Voelker et al. (2010).



Figure 2. Oxygen isotopic composition of precipitation versus rainfall amount (lefthand panels) and air temperature (righthand panels). Data collected at IAEA/GNIP site in Porto, Portugal (see Fig. 1 for location) for 1988-2004. Oxygen isotope data represent multi-year averages of monthly means. The two other closest GNIP stations in Portugal - Vila Real and Portalegre (see Figure 1) - share similar relationships between precipitation oxygen isotopic composition and air temperature (+0.27%/°C; $r^2=0.76$ and +0.26%/°C; $r^2=0.69$, respectively) to that of Porto (+0.21‰/°C). The relationship between precipitation oxygen isotopic composition and monthly precipitation amount is -3.5%/100mm/month (r²=0.64), -3.7%/100mm/month $(r^2=0.49)$, and -1.6%/100 mm/month $(r^2=0.62)$ for the three sites, respectively. Note that right hand y-axis in upper left panel is inverted in order to illustrate inverse nature of rainfall and precipitation oxygen isotopic composition.



978 Figure 3. Iberian rainfall anomalies associated with the North Atlantic Oscillation. 979 Composites of November-March precipitation anomalies (mm/month) during (a) positive and (b) 980 negative NAO winters for the period 1901-2012. Positive/negative NAO winters were 981 determined using the December-March Hurrell principal component-based NAO index (CDG, 982 2018) as those winters with NAO values in the highest/lowest decile of all winters. The PC-983 based NAO index represents the time series of the leading Empirical Orthogonal Function of 984 SLP anomalies over the Atlantic sector, 20°-80°N, 90°W-40°E. Precipitation anomalies are 985 based on the GPCC precipitation, version 7, at 0.5° spatial resolution (Schneider et al. 2014). 986 Yellow stars denote cave sites in this study: BG = Buraca Gloriosa; GCL = Gruta do Casal da 987 Lebre.



Figure 4. Profile and map views of Buraca Gloriosa (top) and Gruta do Casal da Lebre (bottom).

992 Entrance denoted by arrow (top panel) and filled square (bottom panel). Red stars denote

993 locations of stalagmites used in this study.



995

996 Figure 5. COPRA-derived age models for BG/GCL stalagmites. Black lines represent mean 997 of calculated age models while red lines denote 95% confidence intervals. See Table 1 for 998 specific ages and isotopic ratios. Orange square represents a "dummy age" that was included in 999 order to extrapolate below the hiatus, which is only possible with at least two dated points. The 900 bottom of BG611 was based on linear extrapolation through dated intervals. Distances for BG66 1001 were measured relative to topmost section of interval for which stable isotopes were obtained, 1002 and not relative to the cap of the stalagmite (see Figure 6).

- 1003
- 1004



Figure 6. BG/GCL stalagmites and U/Th ages. Red lines denote stable isotope sampling
transects. Blue and white scale bars (cm) define differential enlargement of each stalagmite.
Black arrows represent intervals excluded from this study due to evidence of open system
behavior. Sections without arrows or transect lines are older than the interval examined in this
study. The impact of recrystallization in stalagmite cores was assessed by parallel sampling
transects (parallel red lines on BG66 and GCL6) and demonstrated consistent stable isotopic
values and trends (Supp. Fig. S7).



Figure 7. Temperature and relative humidity variations from (top) Buraca Gloriosa and (middle)
GCL. Drip rate from Buraca Gloriosa and precipitation variability (bottom) from Monte Real,
Portugal (35 km from BG). Temperature sensor in GCL was changed in November 2014 and the
sensitivity of the new instrument varies slightly from the original.



1025 **Figure 8.** Hendy Tests of BG/GCL stalagmites. Top: Covariance plots of carbon and oxygen 1026 isotopic ratios. Correlation coefficients (r^2 values) are listed for each plot. High positive 1027 correlations have been identified as an indicator of non-equilibrium crystallization. Bottom: 1028 Oxygen (blue) and carbon (green) isotopic variations along the same growth layers with distance 1029 (listed in the upper left corner of each panel) from the stalagmite central growth axis. Progressive 1030 increases in δ^{18} O values have been interpreted to reflect disequilibrium crystallization. 1031 Limitations of the Hendy Tests are discussed in text.



1036 **Figure 9. BG/GCL stalagmite isotopic time series**. Carbon (top) and oxygen (bottom) isotopes, 1037 with each stalagmite presented in a different color. δ^{234} U values (yellow circles) for BG6LR are 1038 plotted against carbon isotope ratios (plots showing the δ^{234} U and δ^{13} C values of the other 1039 stalagmites are presented in the Supplemental Material). U/Th ages (with 2 s.d. errors) are also 1040 shown. The "?" at the MIS 6/5e transition denotes uncertainties associated with the continuity of 1041 this interval.

- 1042
- 1043
- 1044





1047 Figure 10. Comparison of Portuguese stalagmite hydroclimate proxies with regional and 1048 global climate records from the last two glacial cycles (A) Ice-rafted debris abundance from 1049 North Atlantic ODP Site 980 (McManus et al., 1999 using Hulu cave time scale as presented in 1050 Barker et al., 2011); (B) composite BG/GCL stalagmite carbon isotopic time series with NH summer insolation (Berger an Loutre, 1991); (C) Carbon isotopic time series from Villars Cave, 1051 southern France (Genty et al., 2003; Genty et al., 2006); (D) Alkenone-based Iberian margin SST 1052 1053 reconstruction (core MD01-2443; Martrat et al., 2007); (E) Temperate forest pollen abundance 1054 from three closely spaced cores (MD01-2443: 250-194 ka (Roucoux et al, 2006; Tzedakis et al., 1055 2004); MD01-2444: 194-136 ka (Margari et al., 2010; Margari et al., 2014); MD95-2042: 136-1 1056 ka (Sánchez Goñi et al., 2008; Sánchez Goñi et al., 2013)); (F) NGRIP (0-122 ka) (North 1057 Greenland Ice Core Project members, 2004) and synthetic Greenland oxygen isotopic record 1058 (Barker et al., 2011) and (G) marine isotope stages.





Figure 11. Iberian margin SST (red) versus stalagmite carbon (black; left column) and
 oxygen (blue; right column) isotopes. Numbers denote select GI events using stratigraphic
 nomenclature of Rasmussen et al. (2014).





1067

Figure 12. BG/GCL stalagmite carbon isotopic time series and Iberian margin SST. Light blue vertical rectangles denote North Atlantic cold events (some of which are labeled). Several interruptions in stalagmite growth coincide, within the errors of the stalagmite chronologies, with periods of depressed SST. Question mark at MIS6/5e transition denotes visible hiatus not resolvable by U/Th dates.

	Distance to Top (mm)	²³⁸ U (ng∕g)	²³² Th (pg/g)	δ ²³⁴ Uª (corr'd)	Error ^b	²³⁰ Th/ ²³⁸ U (activity)		²³⁰ Th/ ²³² Th (ppm)		Uncorrected		Corrected	F
Stalagmite							Error		Error	Age	Error (yr)	Age	Error
										(vr BP) ^c		(vr BP) ^d	(yr)
BG41	67	148	2,892	524.7	2.2	0.779	0.0023	657.7	18.6	82,926	389	82,553	538
BG41	41	293	4,635	522.8	2.2	0.742	0.0030	773.8	8.5	77,026	463	76,724	486
BG41	21	217	1,858	566.6	3.1	0.748	0.0039	1,440.0	40.6	74,906	567	74,746	588
BG41	9	271	2,088	610.8	9.8	0.764	0.0073	1,635.6	22.3	74,392	1,135	74,253	1,142
BG66	266	85	6,980	698.6	9.3	1.283	0.0057	256.5	1.8	223,637	3,252	219,220	3,829
BG66	236	123	4,742	520.6	4.1	1.169	0.0030	500.0	4.0	217,460	1,752	216,719	1,780
BG66	218	101	3,132	532.4	3.1	1.174	0.0015	623.6	4.6	214,835	1,052	213,011	1,379
BG66	207	68	4,057	429.2	3.0 3.1	1.110	0.0025	644.2	7.4	210,091	1,560	211,971	2,470
BG66	194	00	4 3 3 6	370 /	3.1	1.149	0.0015	386.6	7.4	204 768	1,175	200,230	2 063
BG66	154	104	2 1 9 3	443 5	2.6	1 1 0 0	0.0025	864.2	11.9	198 297	930	196 990	1 1 2 8
BG66	86	104	2.661	345.4	2.4	1.041	0.0016	672.5	8.4	197.507	994	195,798	1,298
BG66	54	76	995	564.2	6.2	1.159	0.0057	1,453.3	64.3	189,936	2,538	189,182	2,549
BG67	88	320	2,153	617.8	2.9	1.095	0.0043	2,689.7	51.5	146,174	1,146	145,802	1,158
BG67	79	195	2,799	485.8	2.3	1.014	0.0022	1,164.3	18.2	144,037	695	143,171	814
BG67	66	250	4,187	610.3	4.9	1.072	0.0046	1,057.4	12.7	139,735	1,279	138,803	1,350
BG67	44	162	4,858	484.7	2.4	0.969	0.0023	531.9	5.3	129,620	608	127,800	1,087
BG67	2	216	5,542	401.5	2.6	0.837	0.0039	538.0	5.1	107,150	843	105,501	1,168
BG611	173	119	11,744	202.6	3.5	0.801	0.0041	133.9	0.8	126,291	1,253	118,714	3,908
BG611	160	110	16,828	230.9	4.6	0.792	0.0044	01.2	0.7	118,672	1,277	109,828	4,469
BG611	23	313	552	231.3	1.4	0.762	0.0043	5 168 2	3537	59 726	345	59 608	350
BG611	12	248	2 233	356.2	1.4	0.535	0.0021	1 002 4	25.4	57 908	296	57 115	310
BG611	2	250	4.109	376.7	1.8	0.533	0.0021	535.0	5.9	54,959	284	53.887	604
BG6LR	1,623	72	133	175.0	1.5	0.631	0.0015	5,665.9	1,162	86,532	342	86,392	350
BG6LR	1,593	98	140	165.3	1.4	0.618	0.0014	7,166.0	1,764	84,748	318	84,639	324
BG6LR	1,574	74	905	156.6	1.6	0.615	0.0016	824.8	25.3	84,848	360	83,894	596
BG6LR	1478	159	26	249.2	1.8	0.645	0.0021	63,745.2	114,070	82,068	428	82,056	428
BG6LR	1464	166	1,138	246.8	1.5	0.641	0.0009	1,542.3	35.8	81,475	214	80,983	325
BG6LR	1442	162	77	185.4	1.4	0.634	0.6339	21,885.5	13,015	81,442	396	81,407	396
BG6LR	1375	112	220	202.9	1.5	0.602	0.6016	5,064.2	652.0	77,823	234	77,677	246
BG6LR	1324	120	1,908	150.2	1.4	0.566	0.5660	383.8 1 212 0	15.3	77,213	330	75,946	/ I Z 515
BG6LR	1276	105	353	167.8	2.0	0.564	0.5637	2 766 4	298.1	73 512	425	73 254	444
BG6LR	1246	83	1.232	168.7	1.4	0.561	0.5613	625.8	14.2	72,957	369	71.819	675
BG6LR	1179	62	1,114	252.0	2.6	0.507	0.5071	464.4	15.9	57,877	465	56,584	792
BG6LR	1174	77	2,544	196.0	2.2	0.474	0.4736	235.4	3.8	55,882	375	53,375	1,299
BG6LR	1166	5	367	187.1	2.6	0.482	0.4821	100.4	1.6	57,644	524	51,517	3,066
BG6LR	1153	81	3,460	190.7	2.2	0.433	0.4331	167.2	2.3	49,960	367	46,707	1,654
BG6LR	1141	52	1,159	242.6	2.8	0.359	0.3591	266.4	10.4	37,626	449	36,016	918
BG6LR	1138	55	750	239.5	1.8	0.352	0.3518	426.3	33.1	36,815	381	35,830	625
BG6LR	1101	71	283	235.2	2.0	0.323	0.3234	1,344.2	198.7	33,449	272	33,161	310
BG6LR	1093	101	472	262.1	2.1	0.327	0.3269	802.0	73.4	33,052	331	32,575	409
BGGLR	1077	85	1 034	280.0	1.0	0.290	0.2099	384.9	15 5	27,675	178	26,431	463
BG6LR	1046	56	705	238.2	2.2	0.260	0.2603	339.0	19.7	25 911	265	24 993	531
BG6LR	1026	123	2.093	304.1	1.9	0.262	0.2617	253.3	8.5	24.612	206	23.438	621
BG6LR	1025	123	493	296.4	1.4	0.253	0.0017	1,041.2	151.0	23,814	175	23,538	226
BG6LR	1019	80	377	298.5	2.1	0.252	0.2525	887.3	107.4	23,753	221	23,430	276
BG6LR	1001	68	1,464	288.7	1.5	0.256	0.2558	196.1	4.3	24,291	156	22,789	765
BG6LR	944	76	1,896	329.3	2.1	0.233	0.2330	154.8	3.9	21,131	196	19,450	861
BG6LR	899	79	4,209	294.0	3.4	0.227	0.2266	70.6	1.3	21,074	283	17,360	1,863
BG6LR	883	91	233	330.3	2.0	0.168	0.1684	1,082.0	213.7	14,806	165	14,633	189
BG6LK	843	100	1,409	201.1	4.0	0.162	0.1623	190.4	6.7 11C 0	14,718	164	13,738	516
BGOLR	027 810	75	332 101	295.0	2.9	0.152	0.1521	703.5	22.8	13,645	124	13,424	255
BG6LR	783	95	525	283.8	2.2	0.130	0.1301	418.7	35.3	12 661	150	12 275	246
BG6LR	774	107	1.351	271.4	1.4	0.130	0.1304	169.8	5.7	11.795	119	10.901	463
BG6LR	759	135	4,177	251.5	1.5	0.121	0.1210	64.7	1.0	11,071	117	8,846	1,113
BG6LR	657	86	2,566	212.9	1.4	0.112	0.1120	62.1	0.9	10,540	96	8,326	1,106
BG6LR	139	172	323	204.2	1.7	0.031	0.0010	272.6	41.0	2,790	96	2,651	121
BG6LR	86	155	80	207.9	1.7	0.022	0.0007	720.9	312.3	1,987	62	1,949	67
BG6LR	10	122	43	196.7	18.9	0.014	0.0019	677.5	519.3	1,271	173	1,245	174
GCL6	439	91	2,815	76.3	2.3	0.862	0.0029	461.2	9.3	185,093	1,779	184,255	1,815
GCL6	394	86	3,009	125.7	2.0	0.881	0.0032	415.9	6.9	179,002	1,692	178,095	1,739
GCL6	335 202	70	4,5/9	02./ 78.2	3.U 2.0	0.826	0.0029	214.9 481 0	2.3	174,406	1,794	173 051	1,977
GCL0	256	116	1 019	86.2	2.9	0.045	0.0033	1 574 3	71 R	167 617	1 102	167 382	1 1 0 5
GCL 6	165	94	2.507	122.4	4.2	0.847	0.0049	526.3	13.4	162,712	2.368	162.022	2.550

Table 1. U/Th Isotopic Ratios and ²³⁰Th Ages

 $^{a} \ \delta^{234} U_{meas'd} = \left[\left(^{234} U / ^{238} U \right)_{meas'd} / \left(^{234} U / ^{238} U \right)_{eq} - 1 \right] \times 10^{3}, \text{ where } \left(^{234} U / ^{238} U \right)_{eq} \text{ is secular equilibrium activity ratio: } \lambda_{238} / \lambda_{234} = 1.0. \text{ Values reported as permil.} \right)$ ^b Errors are at the 2*o* level. ^c Present is defined as the vear AD 1950. ^d Initial ²³⁰Th/²³²Th atomic ratio of 13.5x10⁻⁶ \pm 6.75x10⁻⁶ used to correct for unsupported ²³⁰Th in BG stalagmites. GCL stalagmites use 4.4x10⁻⁶ \pm 2.2x10⁻⁶.